

South Dakota State University  
**Open PRAIRIE: Open Public Research Access Institutional  
Repository and Information Exchange**

---

Extension Extra

SDSU Extension

---

2-1-1991

## Safflower Production Tips

Robert G. Hall  
*South Dakota State University*

Follow this and additional works at: [http://openprairie.sdstate.edu/extension\\_extra](http://openprairie.sdstate.edu/extension_extra)

---

### Recommended Citation

Hall, Robert G., "Safflower Production Tips" (1991). *Extension Extra*. Paper 276.  
[http://openprairie.sdstate.edu/extension\\_extra/276](http://openprairie.sdstate.edu/extension_extra/276)

This Other is brought to you for free and open access by the SDSU Extension at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in Extension Extra by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact [michael.biondo@sdstate.edu](mailto:michael.biondo@sdstate.edu).



# Extension Extra

37ExEx 8069  
February 1991  
Plant Science

COLLEGE OF AGRICULTURE & BIOLOGICAL SCIENCES / SOUTH DAKOTA STATE UNIVERSITY / USDA

## Safflower Production Tips

*by JRobert G. Hall, Extension Agronomist-Crops*

The 1990 Farm Bill has designated safflower as a minor oilseed crop which is permitted on "flex" and 0/92 acres. This publication describes the agronomic characteristics of safflower.

### GENERAL

Safflower, an annual broadleaf plant is "thistle-like" in appearance. Soon after emergence it forms a rosette which lasts for about 4 weeks. The main stem then elongates rapidly to a 24-36" height. Thereafter, the plant branches extensively.

Safflower seed looks similar to sunflower but is white. Safflower meal is used for feed and the oil is processed and utilized as a cooking oil. Nearly all safflower is grown and sold under contract.

### SEEDING

Plant safflower from about mid-April to mid-May, after the soil temperatures have reached 40 degrees F. Soil crusting can greatly reduce stands; in some cases, therefore, no-till seeding into small grain stubble may enhance stand establishment. No-till seeding also may enhance stands by conserving moisture needed for seedling development (Table 3.).

Row spacings of 6-14 inches are common. Data for Highmore indicates 14-inch rows yielded higher than 7 or 21-inch spacings (Table 1.). In South Dakota a 20 lb/A seeding rate is suggested.

### ADAPTATION

Safflower tolerates spring temperatures down to 20 degrees F. Most varieties take around 120 days from planting to maturity. In some cases safflower may not mature if planted later than mid-May. Rotation with small grains is best. If volunteer plants are a concern, plant into wheat or oats as first choice and barley as second choice. Safflower is similar to sunflower in drought tolerance. Do not plant fields to safflower in consecutive years.

### HARVESTING

Safflower is generally straight combined. The combine cylinder speed should not exceed 3,000 FPM. This would be about 500 RPM for a 22-inch cylinder. Suggested concave clearances are 5/8-inch (front) and 1/2-inch (back).

Maintain high shaker speeds to prevent plant residues from clogging the combine. A clogged combine is extremely hard to unclog. During harvest a white fuzz from the seed may clog combine radiators and air intakes. Small meshed screen enclosures should minimize the problem. Monitor any buildup of white fuzz on the combine because it may create a fire hazard.

Swath safflower if green weeds or uneven ripening are a problem. Harvest when the moisture is 8% or less and the leaves are dry, seeds are white, and the seed threshes easily by hand. Wet weather at harvest may cause seed sprouting in the head.

## FERTILITY

Nitrogen (N) is the most limiting element on nonfallow land. Phosphorous can be limiting on fallow ground. On most recrop ground only a limited amount of N is needed. Unless a deep-rooted crop like sunflower, winter wheat, or safflower has been grown in the past 3-5 years, only about 1 lb of N is needed for each 20 lbs/acre of safflower grain.

## WEED CONTROL

Safflower, during its early stages, is a poor competitor with weeds.

Few herbicides are registered by EPA for this crop. These products are effective for annual grasses, but broadleaved plants are a problem. Approved herbicides include: Trifluralin (Treflan) at .5 to 1 lb/A active preplant incorporated; EPTC (Eptam) at 3 lb/A active preplant incorporated and metolachlor (Dual) at 2 to 3 lb/A active preemergence or shallow preplant incorporated. Gramoxone Extra at .6 to .9 lb/A active can be used for burndown of emerged weeds for no-till planting. Read and follow label directions.

Avoid planting safflower where carryover from residual herbicides such as atrazine or picloram may cause injury. Drift from applications of broadleaved herbicides to adjacent fields also can cause considerable damage.

## DISEASES

**Bacterial blight** occurs during cool wet weather but safflower often recovers in dry weather.

**Alternaria** may occur when heavy dew or rains persist during flowering and seed development. This disease can cause large safflower losses.

**Rust** is both seed and soil borne. Treating for rust is suggested. Dithane M-45 and Manzate 200 give excellent rust control on safflower.

**Sclerotinia and Botrytis** occur when wet weather persists during flowering.

## AGRONOMIC PERFORMANCE

Seed yields and agronomic traits of safflower in South Dakota and North Dakota are in Tables 1 and 2.

Table 1. Yields of four safflower varieties at three row spacings at Highmore in 1982.

Variety	Rowspacing		
	7"	14"	21"
	----- lbs./acre -----		
Hartman	1432	1604	1346
Rehbein	1379	1773	924
S-208	1377	1418	1146
S-541	1428	1876	1659
Average:	1404	1667	1268

Table 2. Yields of four safflower varieties at three locations in North Dakota, 1988-90.

Variety	Location		
	Williston	Langdon	Carrington
	----- lbs./acre -----		
Girard	621	1358	1071
S-208	669	1591	1061
S-541	679	1483	1148
Finch	657	1429	986
Average:	657	1465	1067

## POSSIBLE SEED SOURCES

- SVO Enterprises, Culbertson, MT  
(1-800-321-8489)
- Mindak Growers Ltd., Box 1307, Dickinson, ND 58601  
(701-227-4528)

## INFORMATION SOURCES

Kingsley, Q and M. Volek. 1982. Safflower trials. Pl. Sci. Pam. #71: Ann. Progr. Rep., Cent. Crops & Soils Res. Stn., Highmore, South Dakota, p. 10.

Crop Production Guide 1991. NDSU Extension Service. p. 289-293.

This publication and others can be accessed electronically from the SDSU College of Agriculture & Biological Sciences publications page, which is at <http://agbiopubs.sdstate.edu/articles/ExEx8069.pdf>



Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the USDA. Larry Tidemann, Director of Extension, Associate Dean, College of Agriculture & Biological Sciences, South Dakota State University, Brookings. SDSU is an Affirmative Action/Equal Opportunity Employer (Male/Female) and offers all benefits, services, and educational and employment opportunities without regard for ancestry, age, race, citizenship, color, creed, religion, gender, disability, national origin, sexual preference, or Vietnam Era veteran status.