

South Dakota State University
**Open PRAIRIE: Open Public Research Access Institutional
Repository and Information Exchange**

Fact Sheets

SDSU Extension

1-1-2011

Managing Crop Diseases with Seed Treatments

Kay R. Ruden
South Dakota State University

Lawrence E. Osborne

Follow this and additional works at: http://openprairie.sdstate.edu/extension_fact

Recommended Citation

Ruden, Kay R. and Osborne, Lawrence E., "Managing Crop Diseases with Seed Treatments" (2011). *Fact Sheets*. Paper 149.
http://openprairie.sdstate.edu/extension_fact/149

This Other is brought to you for free and open access by the SDSU Extension at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in Fact Sheets by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact michael.biondo@sdstate.edu.

Managing Crop Diseases with **Seed Treatments**

Kay R. Ruden
Research/Extension Associate

Lawrence E. Osborne
Extension Plant Pathologist

Extension Plant Pathology

South Dakota State University

Seed treatment fungicides are used in various agricultural crops and are useful tools in promoting stand establishment and seedling vigor. Seed treatments may also help preserve yield potential and prevent quality losses in grain by preventing development of seed and soil-borne diseases. Seed treatments addressed in this guide are those consisting of fungicides or fungicides in combination with insecticides for use in managing disease-causing organisms (pathogens). The development of effective seed treatments can be noted as one of the most significant advancements in plant disease management.

In general, fungicidal seed treatments are used for three primary reasons: (1) To control soil-borne fungal disease organisms such as those causing seed rots, damping-off, or seedling blights in many crops, as well as the agents of root rot complex, smuts, bunts, or downy mildews. (2) To control diseases caused by seed surface-borne fungal pathogens (e.g., covered smuts of barley and oats, bunt of wheat, safflower rust, and *Ascochyta* of legumes). (3) To manage diseases caused by seed-borne fungi, such as loose smut of cereals. Fungicide seed treatments are not effective against bacterial pathogens or in managing viral diseases. Most seed treatment products do not control all types of fungal pathogens.

Disease Management Using Seed Treatments

Disease management in agricultural crops requires a multifaceted approach as part of an integrated pest management (IPM) program. Effective components of an integrated plant disease management program include the following:

- Crop rotation, including rotation to non-host crops to reduce pathogen load.
- Residue and volunteer management for reduction of residue-borne and overwintering diseases.
- Use of high quality, disease-free seed to prevent the spread of seed-borne diseases and promote healthy stand establishment.
- Proper variety selection for host resistance and adaptation to the growing region.

- Proper plant health management. Healthy plants are more able to resist or tolerate the development of plant diseases.
- Judicious use of plant protectant products such as herbicides, insecticides, and fungicides to reduce losses, promote healthy plants, and prevent quality losses in seed.

Field history is a key component of the decision-making process for managing diseases with seed treatments. Keeping in mind the cropping sequence and the history of major disease or insect pests within the field can be important factors in seed treatment decisions. Proper identification of disease agents is also important. Local county Extension agronomy educators or the Plant Disease Diagnostic Clinic at SDSU can assist producers in identifying plant health problems throughout the growing season.

Effectiveness of control will vary with seed treatment product, rate, environmental conditions, and pests present. Seed treatments may provide some level of control for early season diseases as well as control seedling blights and seed- or soil-borne diseases.

Newly Opened Land— A Special Consideration

Newly opened land, such as CRP being returned to crop production, may present a special consideration and most certainly will be a situation where seed treatments should be considered. For example, small grains planted into these areas can be at high risk for the development of diseases and insect pests. Diseases such as root and crown rots, as well as seedling blights, can often be more severe when certain crops are planted into these high-residue situations. Also, insect pressure on newly cultivated lands may differ from a typical cropping situation.

Classification of Fungicidal Seed Treatments

Fungicidal seed treatments can be classified based on movement of the seed treatment product in relation to the seed. Fungicides used as protectants (contacts) are effective

SOUTH DAKOTA STATE UNIVERSITY

College of Agriculture & Biological Sciences • Cooperative Extension Service • U.S. Department of Agriculture

only on the seed surface, providing protection against seed surface-borne pathogens and providing some level of control of soil-borne pathogens. These products generally have a relatively short residual. Protectant fungicides such as captan, maneb, PCNB, thiram, or fludioxonil help control most types of soil-borne pathogens, with the exception of the root rotting organisms. Systemic seed treatment fungicides are absorbed into the emerging seedling and inhibit or kill the fungus inside host plant tissues. Systemic fungicides used for seed treatment include the following: azoxystrobin, carboxin, mefenoxam, metalaxyl, thiabendazole, trifloxystrobin, and various triazole fungicides, including difenoconazole, tebuconazole, and triticonazole. Mefenoxam and metalaxyl are primarily used to target the water mold fungi *Pythium* and *Phytophthora*. Biological agents as seed treatments are also available and may provide some level of protectant activity. Not all fungicides are available as seed treatments for every crop, and not all fungicides have activity against the same range of organisms. Refer to the specific crop-pest combinations listed in the text for product-use recommendations. Always read and follow label directions.

Proper Application and Use Precautions

Fungicide seed treatment products vary in formulation type, packaging, and use requirements. Products may be dry or liquid and in concentrate or ready-to-use formulations. While many seed treatments may be applied on-farm, several products are limited to use only by commercial applicators using closed application systems. Caution should be used when handling or working with seed treatment products. Fungicide seed treatments can be highly poisonous and many are irritants, so proper handling precautions must be taken when handling seed treatment chemicals, and producers or applicators must strictly adhere to all label directions regarding safe handling, mixing, storage, and disposal. Using personal protection, including an approved chemical respirator, goggles, and pesticide-resistant gloves, is recommended even if not specifically required by the fungicide label. Follow label rates, as over-application may result in unintentional damage to the seed, and under-application may reduce the effectiveness of products.

Properly calibrate all application equipment to assure uniform coverage. Uniform coverage of the seed is critical to optimize effectiveness of the seed treatment. Several seed treatment methods are available, though not all are appropriate for every situation. Commercial application or application through dedicated seed treatment equipment will likely provide the most uniform coverage. Grain auger-mounted treatment equipment is available, and may provide adequate coverage in an on-farm situation; however, an auger that has been used to treat seed may be unusable for moving grain intended for food or feed. Likewise, treated seed should not be allowed to contaminate equipment used to transport or store food or feed grains. Use caution when considering planter-applied (planter-box) seed treatments. Good disease control depends on uniform fungicide coverage of the seed, and this more difficult to accomplish in planter-applied situations.

Always read and follow label directions. Understand the product-specific guidelines for proper application: how and when to apply, pre-harvest intervals, feeding or grazing restrictions, as well as important safety precautions. Always dispose of pesticide containers properly.

Seed Treatments and Legume Inoculants

Seed treatments containing fungicides or fungicide/insecticide combinations may adversely affect microbial inoculants applied to legume seed, such as soybeans. Producers should carefully read and follow any label instructions and limitations for both the pesticide seed treatment and the inoculant. Liquid fungicides or fungicide/insecticide combinations should not be directly mixed with liquid inoculants prior to application, and care should be taken to limit the time that inoculants and pesticide seed treatments are in direct contact.

Do Not Use Treated Seed for Food or Feed!!

Following are the seed treatments fungicides or fungicide/insecticide combinations currently labeled for use in South Dakota. The list is dynamic and prone to frequent modifications. Always check the list of products currently registered with the South Dakota Department of Agriculture for legality of use in the state.



South Dakota
Cooperative Extension Service

South Dakota State University, South Dakota counties, and U.S. Department of Agriculture cooperating. South Dakota State University is an Affirmative Action/Equal Opportunity Employer and offers all benefits, services, education, and employment opportunities without regard for race, color, creed, religion, national origin, ancestry, citizenship, age, gender, sexual orientation, disability, or Vietnam Era veteran status.

FS949: 1000 revised at \$x.xx each, January 2011. Printed on Recycled Paper.

FS949 may be accessed on the web at <http://agbiopubs.sdstate.edu/articles/FS949.pdf>

SEED TREATMENTS FOR WHEAT

Diseases listed on label									
Common Bunt/ Covered Smut	Loose Smut	Seed & Seedling Rots	Common Root Rot	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products	Application Rate	Special Notes	
X		X			X	azoxystrobin Dynasty	0.153–0.382 fl oz/cwt		
		X				captan Captan 400 Captan 400–C	1.5–4 fl oz/cwt		
X	X	X				captan + carboxin Enhance	4 oz/cwt	Do not graze or feed livestock on treated areas for 42 days after planting.	
X	X	X	X	X	X	captan + carboxin + imidacloprid Enhance AW	4 oz/cwt	Do not graze or feed livestock on treated areas for 45 days after planting.	
X	X					carboxin Vitavax–34	2–3 oz/cwt	Do not graze or feed livestock on treated areas for 6 weeks after planting.	
X	X	X		X	X	carboxin + thiram RTU–Vitavax–Thiram	5–6.8 fl oz/cwt (Use the higher rate where smuts and bunts are severe.)	Do not graze or feed livestock on treated areas for 6 weeks after planting.	
						Vitaflo–280	3.5–5.0 fl oz/cwt (Use the high rate for control of loose smut.)		
X	X	X	X	X	X	difenoconazole + mefenoxam Dividend XL RTA	2.5 fl oz/cwt plus control of Fusarium seed scab.	Green wheat forage may not be grazed until 55 days after planting.	
						Incentive RTA	5 fl oz/cwt plus control of Fusarium seed scab and partial control of common root rot.	Do not plant any crop other than wheat or barley within 30 days to fields in which treated seeds were planted.	
							10 fl oz/cwt plus control of Fusarium seed scab and partial control of take-all and common root rot.		
X	X	X	X	X	X	difenoconazole + mefenoxam Dividend Extreme	1 fl oz/cwt plus control of Fusarium seed scab	Green wheat forage may not be grazed until 55 days after planting.	
							2 fl oz/cwt plus control of Fusarium seed scab and early season control of common root rot	Do not plant any crop other than wheat or barley within 30 days to fields in which treated seeds were planted.	
							4 fl oz/cwt plus control of Fusarium seed scab and early season control of take all and common root rot		

SEED TREATMENTS FOR WHEAT

Diseases listed on label								Application Rate	Special Notes
Common Bunt/ Covered Smut	Loose Smut	Seed & Seedling Rots	Common Root Rot	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products			
		X		X	X	fludioxonil Maxim 4 FS	0.08–0.16 fl oz/cwt	Do not graze forage until 30 days after planting.	
X	X	X		X	X	fludioxonil + mefenoxam Maxim XL	0.167–0.334 fl oz/cwt		
X	X	X		X	X	ipconazole Rancona 3.8 FS Rancona Apex	0.051–0.085 fl oz/cwt 5–8.33 fl oz/cwt plus suppression of common root rot		
X	X	X		X	X	ipconazole + metalaxyl Rancona Pinnacle	5–8.33 fl oz/cwt plus suppression of common root rot		
X	X	X		X		ipconazole + metalaxyl + imidacloprid Rancona Crest Rancona Crest WR	5–8.33 fl oz/cwt plus suppression of common root rot 5–8.33 fl oz/cwt plus suppression of common root rot Use higher rate when disease pressure will be high or if there is a history of high disease levels in the field.	Do not graze or feed livestock on treated areas for 45 days after planting.	
X	X					mancozeb Dithane M45 Penncozeb 75DF Penncozeb 80WP	2.2–3.3 oz/cwt 2.3–3.5 oz/cwt 2.2–3.3 oz/cwt		
X	X					mancozeb + surfactant Grain Guard Manzate Pro–Stick	2 oz/bu 2.2–3.3 oz/cwt		
X	X					mancozeb + surfactant + copper ManKocide	4 oz/cwt		
X		X				maneb Manex	3.5–5.2 fl oz/cwt		
		X				mefenoxam Apron XL Apron XL LS	0.0425–0.085 fl oz/cwt for Pythium damping off	Use the higher rate when the disease pressure is expected to be high.	

SEED TREATMENTS FOR WHEAT

Diseases listed on label							Application Rate	Special Notes
Common Bunt/ Covered Smut	Loose Smut	Seed & Seedling Rots	Common Root Rot	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products		
X	X	X		X		mefenoxam + difenoconazole + thiamethoxam Cruiser Maxx Cereals	5 fl oz/cwt plus suppression of common root rot and take all	Do not graze or feed livestock on treated areas for 45 days after planting.
						metalaxyl Acquire Allegiance Dry Allegiance FL Dyna—Shield Metalaxyl MetaStar ST Sebring 2.65 ST Sebring 480 FS	0.75 fl oz/cwt 1.5–2.0 oz/cwt 0.75 fl oz/cwt 0.75 fl oz/cwt 0.75 fl oz/cwt 0.75 fl oz/cwt 0.50 fl oz/cwt	For control of Pythium damping—Off only.
X	X	X	X	X	X	metalaxyl + tebuconazole + imidacloprid Gaucho XT Flowable	3.4 fl oz/cwt	Do not graze or feed livestock on treated areas for 45 days after planting.
X	X	X	X	X	X	prothioconazole + tebuconazole + metalaxyl Proceed Concentrate Proceed MD	1–1.5 fl oz/cwt 5–7.5 fl oz/cwt	Wheat forage may be grazed or harvested for hay 31 days after seeding.
X	X	X	X	X	X	prothioconazole + tebuconazole + metalaxyl + clothianidin Proceed Plus	5–7.5 fl oz/cwt	Wheat forage may be grazed or harvested for hay 31 days after seeding.
		X			X	pyraclostrobin Stamina	0.4–0.8 fl oz/cwt, plus suppression of common root rot	Use higher rates when disease pressure is expected to be high.
X	X	X		X	X	pyraclostrobin + triticonazole + metalaxyl Stamina F [®] HL	1.0 fl oz/cwt	

SEED TREATMENTS FOR WHEAT

Diseases listed on label

Common Bunt/ Covered Smut		Loose Smut	Seed & Seedling Rots	Common Root Rot	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products	Application Rate	Special Notes
X	X	X	X	X	X	X	tebuconazole Sativa 318 FS TebuStar 250 ST	0.08–0.10 fl oz/cwt 0.1 fl oz/cwt	Wheat forage may be grazed or harvested for hay 31 days after seeding.
X	X	X	X	X	X	X	tebuconazole + metalaxyl Dyna–Shield Foothold Dyna–Shield Small Grains Fungicide Raxil MD Raxil XT Sativa M Sativa M RTU	5–6.5 fl oz/cwt plus early season control of common root rot 5–6.5 fl oz/cwt plus early season control of common root rot 5–6.5 fl oz/cwt plus early season control of common root rot 0.16–0.20 oz/cwt plus early season control of common root rot 5 fl oz/cwt plus early season control of common root rot 5–6.5 fl oz/cwt plus early season control of common root rot	Wheat forage may be grazed or harvested for hay 31 days after seeding.
X	X	X	X	X	X	X	tebuconazole + metalaxyl + imazalil Raxil MD Extra	5 fl oz/cwt plus early season control of common root rot	Do not graze or feed livestock on treated areas for 6 weeks after planting.
X	X	X	X	X	X	X	tebuconazole + metalaxyl + imazalil + imidacloprid Raxil MD Extra W	5.14 fl oz/cwt plus early season control of common root rot	Do not graze or feed livestock on treated areas for 45 days after planting.
X	X	X	X	X	X	X	tebuconazole + metalaxyl + imidacloprid Dyna–Shield Foothold Extra Raxil MD W Sativa IM Max Sativa IM RTU	3.4–5 fl oz/cwt plus early season control of common root rot 5 fl oz/cwt plus early season control of common root rot 3.4–5 fl oz/cwt 5 fl oz/cwt	Do not graze or feed livestock on treated areas for 45 days after planting.

SEED TREATMENTS FOR WHEAT

Diseases listed on label								Special Notes
Common Bunt/ Covered Smut	Loose Smut	Seed & Seedling Rots	Common Root Rot	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products	Application Rate	
X	X	X	X	X	X	tebuconazole + thiram Raxil-Thiram	3.5-4.6 fl oz/cwt plus early season control of common root rot and control of Fusarium seed scab	Use low rate under dryland or low rainfall conditions. Wheat green forage may be grazed or harvested for hay 31 days after seeding.
X		X				thiram Signet 480 FS	2.0 fl oz/bu	
X	X			X		triticonazole Charter	3.1 fl oz/cwt plus suppression of common root rot and Rhizoctonia root rot.	Do not plant any crop not listed on the label in soil treated with Charter within 30 days after planting treated seed.
X	X	X		X	X	triticonazole + metalaxyl Charter F ^z	5.4 fl oz/cwt	Do not plant any crop not listed on the label in soil treated with Charter F ^z within 30 days after planting treated seed.
X	X	X	X	X	X	triticonazole + thiram Charter PB	5.5 fl oz/cwt plus suppression of Rhizoctonia root rot	Do not plant any crop not listed on the label within 30 days after planting.

SEED TREATMENTS FOR BARLEY

Diseases listed on label					Seed Treatment Products	Application Rate	Special Notes
Covered Smut	Loose Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases			
		X		X	azoxystrobin Dynasty	0.10–3.75 fl oz/cwt	
		X			captan Captain 400 Captain 400–C	2–3 fl oz/cwt 2–3 fl oz/cwt	
X	X	X			captan + carboxin Enhance	4 oz/cwt	Do not graze or feed livestock on treated areas for 42 days after planting.
X	X	X	X	X	captan + carboxin + imidacloprid Enhance AW	4 oz/cwt	Do not graze or feed livestock on treated areas for 45 days after planting.
X	X				carboxin Vitavax–34	2–3 oz/cwt	Do not graze or feed livestock on treated areas for 6 weeks after planting.
X	X	X	X	X	carboxin + thiram RTU–Vitavax–Thiram	5–6.8 fl oz/cwt (Use the higher rate where smuts and bunts are severe.) 3.5–5.0 fl oz/cwt (Use the high rate for control of loose smut.)	Do not graze or feed livestock on treated areas for 6 weeks after planting.
					Vitaflo–280		Green wheat forage may not be grazed until 55 days after planting.
					difenoconazole + mefenoxam Dividend Extreme	2 fl oz/cwt plus partial control of Fusarium seed scab and common root rot. 4 fl oz/cwt plus partial control of Fusarium seed scab and common root rot. 5 fl oz/cwt plus partial control of Fusarium seed scab and common root rot.	Do not plant any crop other than wheat or barley within 30 days to fields in which treated seeds were planted.
X		X	X	X	Dividend XL RTA		
					Incentive RTA	10 fl oz/cwt plus partial control of Fusarium seed scab, take–all and common root rot. 5 fl oz/cwt plus partial control of Fusarium seed scab and common root rot.	
		X	X	X	fludioxonil Maxim 4FS	0.08–0.16 fl oz/cwt	Cereal forage may not be grazed until 30 days after planting.
		X	X	X	fludioxonil + mefenoxam Maxim XL	0.167–0.334 fl oz/cwt	

SEED TREATMENTS FOR BARLEY

Diseases listed on label						Application Rate	Special Notes
Covered Smut	Loose Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products		
X	X	X	X	X	ipconazole Rancona 3.8 FS Rancona Apex	0.051–0.085 fl oz/cwt 5–8.33 fl oz/cwt plus suppression of common root rot	
X	X	X	X	X	ipconazole + metalaxyl Rancona Pinnacle	5–8.33 fl oz/cwt plus suppression of common root rot	
X	X	X	X	X	ipconazole + metalaxyl + imidacloprid Rancona Crest Rancona Crest WR	5–8.33 fl oz/cwt plus suppression of common root rot 5–8.33 fl oz/cwt plus suppression of common root rot Use higher rate when disease pressure will be high or if there is a history of high disease levels in the field.	Do not graze or feed livestock on treated areas for 45 days after planting.
X	X	X			mancozeb Dithane M45 Penncozeb 75DF Penncozeb 80WP	2.7–4.2 oz/cwt 2.9–4.5 oz/cwt 2.7–4.2 oz/cwt	
X	X	X			mancozeb + surfactant Grain Guard Manzate Pro–Stick	2 oz/bu 2.7–4.2 oz/cwt	
X	X	X			mancozeb + surfactant + copper ManKocide	4 oz/cwt	
X	X	X			maneb Manex	4.3–6.7 fl oz/cwt	
		X			mefenoxam Apron XL Apron XL LS	0.0425–0.085 fl oz/cwt for Pythium damping off	Use the higher rates when the disease pressure is expected to be high.

SEED TREATMENTS FOR BARLEY

Diseases listed on label						Application Rate	Special Notes
Covered Smut	Loose Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products		
X	X	X	X		mefenoxam + difenoconazole + thiamethoxam Cruiser Maxx Cereals	5 fl oz/cwt plus suppression of common root rot and take all	Do not graze or feed livestock on treated areas for 45 days after planting.
					metalaxyl Acquire	0.75 fl oz/cwt	For control of Pythium damping-off only.
					Allegiance Dry	1.5–2.0 oz/cwt	
					Allegiance FL	0.75 fl oz/cwt	
		X			Dyna–Shield Metalaxyl	0.75 fl oz/cwt	
					MetaStar ST	0.75 fl oz/cwt	
					Sebring 2.65 ST	0.75 fl oz/cwt	
					Sebring 480 FS	0.50 fl oz/cwt	
X	X	X	X	X	metalaxyl + tebuconazole + imidacloprid Gaucho XT Flowable	3.4 fl oz/cwt	Do not graze or feed livestock on treated areas for 45 days after planting.
X	X	X	X	X	prothioconazole + tebuconazole + metalaxyl Proceed Concentrate	1–1.5 fl oz/cwt	Barley forage may be grazed or harvested for hay 31 days after seeding.
					Proceed MD	5–7.5 fl oz/cwt	
X	X	X	X	X	prothioconazole + tebuconazole + metalaxyl + clothianindin Proceed Plus	5–7.5 fl oz/cwt	Barley forage may be grazed or harvested for hay 31 days after seeding.
					pyraclostrobin Stamina	0.4–0.8 fl oz/cwt plus suppression of common root rot (Use the higher rates when disease pressure is expected to be high.)	
X	X	X	X	X	pyraclostrobin + triticonazole+ metalaxyl Stamina F HL	1 fl oz/cwt	

SEED TREATMENTS FOR BARLEY

Diseases listed on label						Seed Treatment Products	Application Rate	Special Notes
Covered Smut	Loose Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases				
X	X	X	X	X	tebuconazole Sativa 318 FS TebuStar 250 ST	0.08–0.10 fl oz/cwt 0.1 fl oz/cwt	Barley forage may be grazed or harvested for hay 31 days after seeding.	
X	X	X	X	X	tebuconazole + metalaxyl Dyna–Shield Foothold Dyna–Shield Small Grains Fungicide Raxil MD Raxil XT Sativa M Sativa M RTU	5–6.5 fl oz/cwt plus early season control of common root rot. 5–6.5 fl oz/cwt plus early season control of common root rot. 5–6.5 fl oz/cwt plus early season control of common root rot. 0.16–0.20 oz/cwt plus early season control of common root rot. 5 fl oz/cwt plus early season control of common root rot. 5–6.5 fl oz/cwt plus early season control of common root rot.	Barley forage may be grazed or harvested for hay 31 days after seeding.	
X	X	X	X	X	tebuconazole + metalaxyl + imazalil Raxil MD Extra	5 fl oz/cwt plus early season control of common root rot.	Do not graze or feed livestock on treated areas for 6 weeks after planting.	
X	X	X	X	X	tebuconazole + metalaxyl + imazalil + imidacloprid Raxil MD Extra W	5.14 fl oz/cwt plus early season control of common root rot.	Do not graze or feed livestock on treated areas for 45 days after planting.	
X	X	X	X	X	tebuconazole + metalaxyl + imidacloprid Dyna–Shield Foothold Extra Raxil MD W Sativa IM Max Sativa IM RTU	3.4–5 fl oz/cwt plus early season control of common root rot. 5 fl oz/cwt plus early season control of common root rot. 3.4–5 fl oz/cwt 5 fl oz/cwt	Do not graze or feed livestock on treated areas for 45 days after planting.	

SEED TREATMENTS FOR BARLEY

Diseases listed on label							Application Rate	Special Notes
Covered Smut	Loose Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products	Application Rate		
X	X	X	X	X	tebuconazole + thiram Dyna-Shield Tebuconazole-Thiram	3.5–4.6 fl oz/cwt 3.5–4.6 fl oz/cwt	Do not graze or feed livestock on treated areas for 45 days after planting.	
X		X			thiram Signet 480 FS	2.0 fl oz/bu		
X	X		X		triticonazole Charter	3.1 fl oz/cwt plus suppression of common root rot and Rhizoctonia root rot.	Do not plant any crop not listed on the label in soil treated with Charter F ² within 30 days after planting treated seed.	
X	X	X	X	X	triticonazole + metalaxyl Charter F ²	5.4 fl oz/cwt	Do not plant any crop not listed on the label in soil treated with Charter F ² within 30 days after planting treated seed.	
X	X	X	X	X	triticonazole + thiram Charter PB	5.5 fl oz/cwt plus suppression of Rhizoctonia root rot	Do not plant any crop not listed on the label within 30 days after planting.	

SEED TREATMENTS FOR OATS

Diseases listed on label							Application Rate	Special Notes
Covered Smut	Loose Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products	Application Rate		
		X			captan Captan 400	2–4 fl oz/cwt		
					Captan 400–C	2–4 fl oz/cwt		
X	X	X			captan + carboxin Enhance	4 oz/cwt	Do not graze or feed livestock on treated areas for 42 days after planting.	
X	X	X	X	X	captan + carboxin + imidacloprid Enhance AW	4 oz/cwt	Do not graze or feed livestock on treated areas for 45 days after planting.	
X	X				carboxin Vitavax-34	2–3 oz/cwt	Do not graze or feed livestock on treated areas for 6 weeks after planting.	
X	X	X			carboxin + thiram RTU-Vitavax-Thiram	5–6.8 fl oz/cwt (Use the higher rate where smuts are severe)	Do not graze or feed livestock on treated areas for 6 weeks after planting.	
					Vitaflo-280	5.0 fl oz/cwt		
		X	X	X	fluidoxonil Maxim 4FS	0.08–0.16 fl oz/cwt	Cereal forage may not be grazed until 30 days after planting.	

SEED TREATMENTS FOR OATS

Diseases listed on label						Application Rate	Special Notes
Covered Smut	Loose Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products		
		X	X	X	fludioxonil + mefenoxam Maxim XL	0.167–0.334 fl oz/cwt	
X	X	X	X	X	ipconazole Rancona 3.8 FS Rancona Apex	0.051–0.085 fl oz/cwt 5–8.33 fl oz/cwt	
X	X	X	X	X	ipconazole + metalaxyl Rancona Pinnacle	5–8.33 fl oz/cwt	
X	X	X	X	X	ipconazole + metalaxyl + imidacloprid Rancona Crest Rancona Crest WR	5–8.33 fl oz/cwt 5–8.33 fl oz/cwt Use higher rate when disease pressure will be high or if there is a history of high disease levels in the field.	Do not graze or feed livestock on treated areas for 45 days after planting.
X	X	X			mancozeb Dithane M45	4.0–6.3 oz/cwt	
		X			Penncozeb 75DF	4.3–6.7 oz/cwt	
					Penncozeb 80WP	4.0–6.3 oz/cwt	
X		X			mancozeb + surfactant Grain Guard	2 oz/bu	
					Manzate Pro–Stick	4.0–6.3 oz/cwt	
X	X	X			maneb Manex	6.4–10 fl oz/cwt	
		X			mefenoxam Apron XL Apron XL LS	0.0425–0.085 fl oz/cwt for Pythium damping off	Use the higher rates when the disease pressure is expected to be high.
					metalaxyl Acquire	0.75 fl oz/cwt	For control of Pythium damping–off only.
					Allegiance Dry	1.5–2.0 oz/cwt	
					Allegiance FL	0.75 fl oz/cwt	
		X			Dyna–Shield Metalaxyl	0.75 fl oz/cwt	
					MetaStar ST	1.5 fl oz/cwt	
					Sebring 2.65 ST	0.75 fl oz/cwt	
					Sebring 480 FS	0.50 fl oz/cwt	

SEED TREATMENTS FOR OATS

Diseases listed on label						Application Rate	Special Notes
Covered Smut	Loose Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products		
X	X	X	X	X	metalaxy+ tebuconazole + imidacloprid Gaucho XT Flowable Sativa IM RTU	3.4 fl oz/cwt 5 fl oz/cwt	Do not graze or feed livestock on treated areas for 45 days after seeding.
		X	X	X	pyraclostrobin Stamina	0.4–0.8 fl oz/cwt	Use higher rates when disease pressure is expected to be high.
X	X	X	X		pyraclostrobin + triticonazole + metalaxy Stamina F HL	1 fl oz/cwt	
	X	X	X	X	tebuconazole Sativa 318 FS TebuStar 250 ST	0.08–0.10 fl oz/cwt 0.1 fl oz/cwt	Oat forage may be grazed or harvested for hay 31 days after seeding.
	X	X			tebuconazole + metalaxy Dyna–Shield Foothold Dyna–Shield Small Grains Raxil MD Raxil XT Sativa M Sativa M RTU	5–6.5 fl oz/cwt 5–6.5 fl oz/cwt 5–6.5 fl oz/cwt 0.16–0.20 oz /cwt 5 fl oz/cwt 5–6.5 fl oz/cwt	Oat forage may be grazed or harvested for hay 31 days after seeding.
	X	X	X	X	tebuconazole + thiram Dyna–Shield Tebuconazole–Thiram Raxil–Thiram	3.5–4.6 fl oz/cwt 3.5–4.6 fl oz/cwt	Use low rate under dryland or low rainfall conditions. Oat forage may be grazed or harvested for hay 31 days after seeding.
X		X			thiram Signet 480 FS	2.0 fl oz/bu	

SEED TREATMENTS FOR RYE

Diseases listed on label							Application Rate	Special Notes
Common Bunt/ Covered Smut	Loose Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products			
		X			captan Captan 400 Captan 400-C	2-3 fl oz/cwt 2-3 fl oz/cwt		
		X	X	X	fludioxonil Maxim 4 FS	0.08-0.16 fl oz/cwt	Cereal forage may not be grazed until 30 days after planting.	
		X	X	X	fludioxonil + mefenoxam Maxim XL	0.167-0.334 fl oz/cwt		
X	X	X	X	X	ipconazole Rancona 3.8 FS Rancona Apex	0.051-0.085 fl oz/cwt 5-8.33 fl oz/cwt plus suppression of common root rot		
		X	X		ipconazole + metalaxyl Rancona Pinnacle	5-8.33 fl oz/cwt		
		X			ipconazole + metalaxyl + imidacloprid Rancona Crest Rancona Crest WR	5-8.33 fl oz/cwt 5-8.33 fl oz/cwt	Do not graze or feed livestock on treated areas for 45 days after planting.	
		X			mancozeb Dirthane M45	2.3-3.6 oz/cwt		
X		X			mancozeb + surfactant Grain Guard Manzate Pro-Stick	2 oz/bu 2.3-3.6 oz/cwt		
		X			maneb Manex	3.6-5.7 fl oz/cwt		
		X			mefenoxam Apron XL Apron XL LS	0.0425-0.085 fl oz/cwt for Pythium damping off	Use the higher rates when the disease pressure is expected to be high.	

SEED TREATMENTS FOR RYE

Diseases listed on label							
Common Bunt/ Covered Smut	Loose Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products	Application Rate	Special Notes
		X			metalaxyl Acquire Allegiance Dry Allegiance FL Dyna—Shield Metalaxyl MetaStar ST Sebring 2.65 ST Sebring 480FS	0.75 fl oz/cwt 1.5–2.0 oz/cwt 0.75 fl oz/cwt 0.75 fl oz/cwt 0.75 fl oz/cwt 0.75 fl oz/cwt 0.50 fl oz/cwt	For control of Pythium damping-off only.
X	X	X	X	X	metalaxyl + tebuconazole + imidacloprid Gaucho XT Flowable	3.4 fl oz/cwt	Do not graze or feed livestock on treated areas for 45 days after seeding.
		X		X	pyraclostrobin Stamina	0.4–0.8 fl oz/cwt plus suppression of common root rot	Use the higher rates when disease pressure is expected to be high.
		X	X		pyraclostrobin + triticonazole + metalaxyl Stamina F HL	1 fl oz/cwt	

SEED TREATMENTS FOR CORN

Diseases listed on label						
Head Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases	Seed Treatment Products	Application Rate	Special Notes
				abamectin + thiamethoxam Avicta Duo Corn	See product label	For control of Corn nematodes.
				abamectin + one of the following: azoxystrobin, fludioxonil, mephenoxam and thiamethoxam. Avicta Complete Corn	See product label	For control of Corn nematodes.
	X		X	azoxystrobin Dynasty	0.153 fl oz/cwt (0.0025 mg ai/kernel)	Use Dynasty only in combination with labeled rates of Maxim 4FS, Maxim XL and Apron XL products.
				Bacillus firmus I-1582 + clothianidin Poncho/Vo Tivo	See product label	For control of Corn nematodes. Do not graze or feed forage and hay to livestock.

SEED TREATMENTS FOR CORN

Diseases listed on label				Seed Treatment Products	Application Rate	Special Notes
Head Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases			
	X	X		Bacillus subtilis GB03 Kodiak HB	4 oz/cwt	For suppression of Fusarium and Pythium root diseases.
	X			captan Captan 400	1.25–2.375 fl oz/cwt	
				Captan 400–C	1.25–2.375 fl oz/cwt	
				Captan Moly	2.4 oz/cwt	
X	X		X	carboxin Vitavax–34	2–4 oz/cwt	Do not graze or feed livestock on treated areas for 6 weeks after planting.
	X		X	carboxin + metalaxyl + imidacloprid Latitude	1.5 oz/42 lbs	Do not graze or feed livestock on treated areas for 45 days.
	X		X	carboxin + PCNB + metalaxyl Prevail	1.5–3 oz/bu	Do not graze or feed livestock on treated areas for 6 weeks after planting.
	X			carboxin + permethrin Kernel Guard Supreme	1.5 oz/42lbs	Do not graze or feed livestock on treated areas for 6 weeks after planting.
X	X	X	X	carboxin + thiram Vitaflo 280	4.5 fl oz/cwt for control of seed & seedling rots, Fusarium and Rhizoctonia root diseases. 8.5–11 fl oz.cwt for control of seed–borne head smut.	Do not graze or feed livestock on treated areas for 6 weeks after planting.
	X	X	X	fludioxonil Maxim 4FS	0.08–0.16 fl oz/cwt (0.0064–0.128 mg ai/kernel)	Green forage may not be grazed until 30 days after planting.
	X	X	X	fludioxonil + mefenoxam Maxim XL	0.167–0.334 fl oz/cwt (0.009–0.018 mg ai/kernel)	Corn forage may not be grazed until 30 days after planting.
X	X	X	X	fludioxonil + mefenoxam + azoxystrobin + thiabendazole Maxim Quattro	0.46 fl oz/80,000 kernel count	Forage may not be grazed until 30 days after planting.
	X	X	X	fludioxonil + mefenoxam + azoxystrobin + thiamethoxam Cruiser Extreme	See product label	Forage may not be grazed until 30 days after planting.
	X	X	X	ipconazole Vortex	0.044 fl oz/cwt	
	X			mancozeb Dithane M45	2.7–5.4 oz/cwt	

SEED TREATMENTS FOR CORN

		Diseases listed on label				Seed Treatment Products	Application Rate	Special Notes
		Head Smut	Seed & Seedling Rots	Fusarium root diseases	Rhizoctonia root diseases			
	X				mancozeb + surfactant Dithane F-45	4.3–8.6 fl oz/cwt		
	X				Manzate Pro–Stick	2.7–5.4 oz/cwt		
	X				maneb Manex	4.3–8.6 fl oz/cwt		
					mefenoxam Apron XL	0.0425–0.085 fl oz/cwt (0.0025–0.005 mg ai/kernel)	For control of Pythium damping–off only.	
					Apron XL LS			
					metalaxyl Aquire	0.75 fl oz/cwt	For control of Pythium damping–off only.	
					Allegiance Dry	1.5–2.0 oz/cwt		
	X				Allegiance FL	0.75 fl oz/cwt		
					Dyna–Shield Metalaxyl	0.75 fl oz/cwt		
					MetaStar ST	0.75 fl oz/cwt		
					Sebring 2.65 ST	0.75 fl oz/cwt		
					Sebring 480 FS	0.50 fl oz/cwt		
	X				metalaxyl + imidacloprid Concur	1.5 oz/42 lbs	For control of Pythium seedling diseases.	
	X			X	pyraclostrobin Stamina	0.4–0.8 fl oz/cwt Use the higher rates when disease pressure is expected to be high.		
	X			X	pyraclostrobin + triconazole + metalaxyl Stamina F HL	1 fl oz/cwt		
	X		X	X	trifloxystrobin Triflex	0.32–0.64 fl oz/cwt	Do not plant any crop without trifloxystrobin tolerances until 30 days after previous planting.	
	X		X	X	trifloxystrobin + metalaxyl Triflex 2000	0.5 fl oz/cwt	Do not plant any crop without trifloxystrobin tolerances until 30 days after previous planting.	

SEED TREATMENTS FOR SOYBEANS

Diseases listed on label				Seed Treatment Products	Application Rate	Special Notes
Seed & Seedling Rots	Fusarium root diseases	Pythium root diseases	Rhizoctonia root diseases			
X		X	X	abamectin + one of the following: metenoxam, fludioxonil, and thiamethoxam. Avicta Complete Beans azoxystrobin Dynasty Bacillus firmus I-1582 + clothianidin Poncho/NoTivo	See product label 0.153-0.459/cwt plus suppression of white mold. See product label	For control of soybean nematodes.
X				captan Captan 400 Captan 400-C Captan Moly captan + carboxin Enhance Vitavax M DC captan + carboxin + imidacloprid Enhance AW	1.5-2.5 fl oz/cwt 1.5-2.5 fl oz/cwt 3.5 oz/cwt 5 oz/bu 2 oz/bu 5 oz/cwt	For control of soybean nematodes. Do not graze or feed forage and hay to livestock.
X	X (Enhance)	X (Enhance)	X (Enhance)	captan + carboxin + metalaxyl Bean Guard Allegiance captan + molybdenum Hi Moly/Captan D carboxin Vitavax-34 carboxin + metalaxyl + imidacloprid Latitude carboxin + PCNB + metalaxyl Prevail carboxin + permethrin Kernel Guard Supreme	2 oz/bu 3-4 fl oz/cwt 3-4 fl oz/cwt 4 oz/cwt 2-4 oz/bu	Do not graze or feed livestock on soybean forage or hay. Do not graze or feed livestock on forage or hay grown from treated seed. Do not graze or feed livestock on forage and hay on treated areas for 6 weeks after planting. Do not graze or feed livestock on forage or hay grown from treated seed.
X	X	X	X	carboxin + thiram RTU-Vitavax-Thiram Vitaflo 280 Vitavax CT	1.5 oz/50 lbs 6.8 fl oz/cwt 4 fl oz/cwt 12 fl oz/cwt	Do not graze or feed livestock on treated areas for 6 weeks after planting. Do not graze or feed livestock on forage and hay grown on treated areas.

SEED TREATMENTS FOR SOYBEANS

Diseases listed on label				Seed Treatment Products	Application Rate	Special Notes
Seed & Seedling Rots	Fusarium root diseases	Pythium root diseases	Rhizoctonia root diseases			
			X	carboxin + thiram + molybdenum Vitavax M	12 fl oz/cwt	Do not graze or feed livestock on forage and hay grown on treated areas.
X	X		X	fludioxonil Maxim 4FS	0.08–0.16 fl oz/cwt	
X	X		X	ipconazole Rancona 3.8 FS	0.085 fl oz/cwt	
X	X	X	X	ipconazole + metalaxyl Rancona Summit	4 fl oz/cwt	Do not graze or feed livestock on soybean forage or hay.
				Rancona Xxtra	3.5 fl oz/cwt	
		X		mefenoxam Apron XL	0.16–0.64 fl oz/cwt (Use the higher rate for best early season Phytophthora protection.)	
				Apron XL LS		
				mefenoxam + fludioxonil ApronMaxx RFC	1.5 fl oz/cwt plus suppression of seedborne Sclerotinia.	
X	X	X	X	ApronMaxx RTA	5 fl oz/cwt plus control of early season Phytophthora and suppression of seedborne Sclerotinia.	
				Maxim XL	0.167–0.334 fl oz/cwt plus early season Phytophthora control.	
				Warden RTA	5 fl oz/cwt plus control of early season Phytophthora and suppression of seedborne Sclerotinia.	
X	X	X	X	mefenoxam + fludioxonil + thiamethoxam Cruiser Maxx	3 fl oz/cwt	
				Cruiser Maxx Plus	3.2 fl oz/cwt	
				Warden CZ	3.2 fl oz/cwt	

SEED TREATMENTS FOR SOYBEANS

Diseases listed on label				Seed Treatment Products	Application Rate	Special Notes
Seed & Seedling Rots	Fusarium root diseases	Pythium root diseases	Rhizoctonia root diseases			
				metalaxyl Aquire	0.75–1.5 fl oz/cwt plus early season control of Phytophthora.	
				Allegiance Dry	1.5–2.0 oz/cwt plus early season control of Phytophthora.	
				Allegiance FL	0.2–1.5 fl oz/cwt plus early season control of Phytophthora.	
		X		Dyna–Shield Metalaxyl	0.75–1.5 fl oz/cwt plus early season control of Phytophthora.	
				MetaStar ST	0.75–1.5 fl oz/cwt plus early season control of Phytophthora.	
				Sebring 2.65 ST	0.75–1.5 fl oz/cwt plus early season control of Phytophthora.	
				Sebring 480 FS	0.50–1.00 fl oz/cwt plus early season control of Phytophthora.	
X				thiabendazole Mertect 340–F	0.08–0.16 fl oz/cwt for control of pod and stem blight.	LSP flowable is for use only by Commercial Seed treaters.
				LSP Flowable	0.125–0.25 fl oz/cwt for control of pod and stem blight.	
		X	X	thiram + metalaxyl + molybdenum Protector–L–Allegiance	6.7 fl oz/cwt	
X				thiram + molybdenum Protector–D	3.3 oz/cwt	
X			X	trifloxystrobin Trilex	0.32 fl oz/cwt	Do not plant any other crop without trifloxystrobin tolerances until 30 days after planting.
X		X	X	trifloxystrobin + metalaxyl Trilex 2000	1 fl oz/cwt	Do not plant any other crop without trifloxystrobin tolerances until 30 days after planting.
				Trilex AL Flowable	5.7 fl oz/cwt	

SEED TREATMENTS FOR SUNFLOWERS

Diseases listed on label		Systemic Downy mildew	Seed Treatment Products	Application Rate	Special Notes
Seed & Seedling Rots					
X	X		azoxystrobin Dynasty	3.75–37.5 fl oz/cwt (0.025–0.1 mg ai/ kernel)	For suppression of downy mildew.
X			captan Captan 400 Captan 400–C	2–4 fl oz/cwt 2–4 fl oz/cwt	
X			fludioxonil Maxim 4FS	0.08–0.16 fl oz/cwt	
X	X		fludioxonil + mefenoxam Maxim XL	0.167–0.334 fl oz/cwt	
X	X		fludioxonil + mefenoxam + azoxystrobin + thiamethoxam CruiserMaxx Sunflower	See product label	
X			ipconazole Rancona 3.8FS	0.051–0.085 fl oz/cwt	
	X		mefenoxam Apron XL Apron XL LS	1.28 fl oz/cwt (0.029 mg ai/seed)	
		X	metalaxyl Aquire Allegiance Dry Allegiance FL Dyna–Shield Metalaxyl MetaStar ST Sebring 2.65 ST Sebring 480FS	1.5–3.0 fl oz/cwt 4.0 oz/cwt 1.5–3.0 fl oz/cwt 1.5–3.0 fl oz/cwt 1.5–3.0 fl oz/cwt 1.5–3.0 fl oz/cwt 1–2 fl oz/cwt	
X			pyraclostrobin Stamina	0.8–2.3 fl oz/cwt	Use higher rates when disease pressure is expected to be high.

SEED TREATMENTS FOR SAFFLOWER

Diseases listed on label		Seed Treatment Products	Application Rate	Special Notes
Seed & Seedling Rots	Seedborne Rust			
X		carboxin Vitavax-34	2 fl oz/cwt	
X		fludioxonil Maxim 4FS	0.08-0.16 fl oz/cwt	
	X	mancozeb Dithane M45	2.0 oz/cwt	
X	X	mancozeb + surfactant Dithane F-45 Grain Guard Manzate Pro-Stick	3.2 fl oz/cwt for control of seedborne rust only. 3 oz/cwt 2 oz/cwt	
	X	maneb Manex	3.2 fl oz/cwt	

SEED TREATMENTS FOR CHICKPEA

Diseases listed on label					
Seed & Seedling Rots	Seed borne Ascochyta blight	Seed Treatment Products	Application Rate	Special Notes	
X		azoxystrobin Dynasty	0.153–0.765 fl oz/cwt		
X		captan + carboxin + imidacloprid Enhance AW	5 oz/cwt	Do not allow livestock to graze or feed on forage until 60 days after planting.	
X		fludioxonil Maxim 4FS	0.08–0.16 fl oz/cwt		
X		fludioxonil + mefenoxam ApronMaxx RFC	1.5 fl oz/cwt plus early season Phytophthora control.		
		ApronMaxx RTA	5 fl oz/cwt plus early season Phytophthora control.		
		Maxim XL	0.167–0.334 fl oz/cwt plus early season Phytophthora control.		
X		ipconazole Rancona 3.8FS	0.085 fl oz/cwt		
X		ipconazole + metalaxyl Rancona Xxtra	3.5 fl oz/cwt		
X		mefenoxam Apron XL Apron XL LS	0.16–0.64 fl oz/cwt	For Pythium damping-off protection.	
X		mefenoxam + fludioxonil + thiamethoxam Cruiser Maxx	3.0 fl oz/cwt		

SEED TREATMENTS FOR CHICKPEA

Diseases listed on label		Seed Treatment Products	Application Rate	Special Notes
Seed & Seedling Rots	Seed borne Ascochyta blight			
X		metalaxyl Acquire Allegiance Dry Allegiance FL Dyna—Shield Metalaxyl MetaStar ST Sebring 2.65 ST Sebring 480 FS	0.75 fl oz/cwt 2 oz/cwt 0.75 fl oz/cwt for early season control of Phytophthora and 1.5 fl oz/cwt for Pythium damping-off. 0.75 fl oz/cwt for early season control of Phytophthora and 1.5 fl oz/cwt for Pythium damping-off. 0.75 fl oz/cwt for early season control of Phytophthora and 1.5 fl oz/cwt for Pythium damping-off. 0.75 fl oz/cwt for early season control of Phytophthora and Pythium damping-off. 0.50 fl oz/cwt for early season control of Phytophthora and Pythium damping-off.	For Pythium damping-off and early season Phytophthora control.
X		pyraclostrobin Stamina	0.4–1.5 fl oz/cwt (Use the higher rates when disease pressure is expected to be high.)	
X	X	thiabendazole Mertect® 430-F	2.04 fl oz/cwt	Mix with an equal amount of water to provide adequate coverage.
X		trifloxystrobin Trilex	0.32 fl oz/cwt	Do not plant any other crop without trifloxystrobin tolerances until 30 days after planting.
X		trifloxystrobin + metalaxyl Trilex 2000 Trilex AL Flowable	1 fl oz/cwt 5.7 fl oz/cwt	Do not plant any other crop without trifloxystrobin tolerances until 30 days after planting.

SEED TREATMENTS FOR FIELD PEA

Diseases listed on label		Seed Treatment Products	Application Rate	Special Notes
Seed & Seedling Rots	Seed borne Ascochyta blight			
X		captan Captan 400 Captan 400C	2 1/2 fl oz/cwt 2 1/2 fl oz/cwt	
X		captan + carboxin + imidacloprid Enhance AW	5 oz/cwt	Do not allow livestock to graze or feed on forage until 60 days after planting.
X		fludioxonil Maxim 4FS	0.08–0.16 fl oz/cwt	
X		fludioxonil + mefenoxam ApronMaxx RFC Apron Maxx RTA Maxim XL	1.5 fl oz/cwt plus early season Phytophthora control. 5 fl oz/cwt plus early season Phytophthora control. 0.167–0.334 plus early season Phytophthora control.	
X		ipconazole Rancona 3.8FS	0.085 fl oz/cwt	
X		ipconazole + metalaxyl Rancona Xxtra	3.5 fl oz/cwt	
X		mefenoxam Apron XL Apron XL LS	0.16–0.64 fl oz/cwt	For Pythium damping-off protection.
X		mefenoxam + fludioxonil + thiamethoxam Cruiser Maxx	1.5 fl oz/cwt	
		metalaxyl Acquire Allegiance Dry Allegiance FL Dyna-Shield Metalaxyl MetaStar ST Sebring 2.65 ST Sebring 480FS	0.75 fl oz/cwt 2 oz/cwt 0.75 fl oz/cwt for early season control of Phytophthora and 1.5 fl oz/cwt for Pythium damping off. 0.75 fl oz/cwt for early season control of Phytophthora and 1.5 fl oz/cwt for Pythium damping off. 0.75 fl oz/cwt for early season control of Phytophthora and 1.5 fl oz/cwt for Pythium damping off. 0.75 fl oz/cwt for early season control of Phytophthora and 1.5 fl oz/cwt for Pythium damping off and systemic downy mildew. 0.50 fl oz/cwt for early season control of Phytophthora and 1.00 fl oz/cwt for Pythium damping off and systemic downy mildew.	For Pythium damping-off and early season Phytophthora control.

SEED TREATMENTS FOR FIELD PEA

Diseases listed on label				
Seed & Seedling Rots	Seed borne Ascochyta blight	Seed Treatment Products	Application Rate	Special Notes
X		pyraclostrobin Stamina	0.4–1.5 fl oz/cwt (Use the higher rates when disease pressure is expected to be high.)	
	X	thiabendazole Mertect® 430-F	1.02 fl oz/cwt	
X		thiram + molybdenum Protector-D	4.5 oz/cwt	
X		trifloxystrobin Trilex	0.32 fl oz/cwt	Do not plant any other crop without trifloxystrobin tolerances until 30 days after planting.
X		trifloxystrobin + metalaxyl Trilex 2000	1 fl oz/cwt	Do not plant any other crop without trifloxystrobin tolerances until 30 days after planting.
		Trilex AL Flowable	5.7 fl oz/cwt	

SEED TREATMENTS FOR ALFALFA

Diseases listed on label		Downy mildew	Seed Treatment Products	Application Rate	Special Notes
Seed & Seedling Rots	Pythium/ Phytophthora root diseases				
X			captan Captan 400 Captan 400C	5-8 fl oz/cwt 5-8 fl oz/cwt	
X			fludioxonil Maxim 4FS	0.08-0.16 fl oz/cwt	Forage may not be grazed until 30 days after planting.
X	X	X	fludioxonil + metenoxam Maxim XL	0.167-0.334 fl oz/cwt	To be used with additional Apron XL (0.553-0.598 fl oz/cwt- see label for instructions).
X			mefenoxam Apron XL Apron XL LS	0.64 fl oz/cwt 0.64 fl oz/cwt	
			metalaxy Acquire Allegiance Dry Allegiance FL Dyna-Shield Metalaxy MetaStar ST Sebring 2.65 ST Sebring 480FS	0.75-1.5 fl oz/cwt 2 oz/cwt (Also for suppression of early season downy mildew.) 0.75-1.5 fl oz/cwt 0.75-1.5 fl oz/cwt 0.75-1.5 fl oz/cwt 0.75-1.5 fl oz/cwt 0.50-1.0 fl oz/cwt	
X			thiram + molybdenum Protector-D	8.0 oz/cwt	