

South Dakota State University
Open PRAIRIE: Open Public Research Access Institutional
Repository and Information Exchange

South Dakota Poultry Field Day Proceedings and
Research Reports, 1975

Animal Science Reports

1975

Egg Yolk Pigmentation with Dehydrated Alfalfa Meal, Pro-Xan and Xanthophyll-free Freeze Dried Alfalfa Juice

E. Guenther
South Dakota State University

O. E. Olson

C. W. Carlson

Follow this and additional works at: http://openprairie.sdstate.edu/sd_poultry_1975

Recommended Citation

Guenther, E.; Olson, O. E.; and Carlson, C. W., "Egg Yolk Pigmentation with Dehydrated Alfalfa Meal, Pro-Xan and Xanthophyll-free Freeze Dried Alfalfa Juice" (1975). *South Dakota Poultry Field Day Proceedings and Research Reports, 1975*. Paper 11.
http://openprairie.sdstate.edu/sd_poultry_1975/11

This Report is brought to you for free and open access by the Animal Science Reports at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in South Dakota Poultry Field Day Proceedings and Research Reports, 1975 by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact michael.biondo@sdstate.edu.

Egg Yolk Pigmentation with Dehydrated Alfalfa Meal,
Pro-Xan and Xanthophyll-free Freeze Dried Alfalfa Juice

E. Guenther, O. E. Olson and C. W. Carlson¹

This study concludes a series of tests in which natural and synthetic materials were used to produce deeply pigmented egg yolk. The most efficient pigmenter used was β -apo 8' carotenoic acid ethyl ester followed by β -apo 8' carotenal, alfalfa concentrates and meals of alfalfa, corn gluten and marigold petals. In this test pure lutein, dehydrated alfalfa meal and Pro-Xan, an alfalfa concentrate, were used as sources of pigments². The freeze dried alfalfa juice was tested for its possible effect in enhancing the utilization of pure lutein. Caged hens were depleted of body stored pigments by feeding an essentially pigment free, milo-soy diet for 12 months. Pigmenting materials were then added to the milo-soy basal to provide dietary pigment at levels of 10 and 20 ppm as shown in table 1.

The 17% protein dehydrated alfalfa meal analyzed 322 ppm xanthophyll. The Pro-Xan, a protein-xanthophyll concentrate of alfalfa, contained 40% crude protein and 1067 ppm xanthophyll. The lutein (a common name for xanthophyll) was extracted from the freeze dried alfalfa juice. The residue of the alfalfa juice was fed in diets 5, 6, 7 and 8 at levels equivalent to juice levels. The extracted lutein fed in diets 7, 8, 9 and 10 had a concentration of 6900 ppm.

The concentration of pigment in the egg yolk was chemically determined and expressed as micrograms of β -carotene equivalents (BCE) per gram of yolk. Utilization of the pigment was then calculated from data on feed consumption and weight of egg produced. The test covered a 49-day feeding period, being limited by the amount of lutein that was available.

The BCE values in table 1 show that doubling the amount of pigment in the diet does not double the amount of pigment found in the yolk. The efficiency of utilization decreases as the amount of pigment increases in the diet. This effect has been observed in previous tests. Other factors such as the source of the pigments and the rate of egg production also affect deposition of pigment in the egg yolk.

The concentration of pigment in the yolk from the dehydrated alfalfa more than doubled when the concentration of dietary pigment increased from 10 to 20 ppm. The hens in group 2 were producing at the rate of 34%, while production of the other groups was within the range of 50%. Similar effects have been observed in other tests. Apparently, more pigment accumulates in the yolk when the rate of ovulation declines.

¹Assistant Professor; Professor, Station Biochemistry; and Professor and Leader, Poultry Research and Extension, respectively.

²All products were obtained from the Western Regional Laboratories, U.S.D.A., Berkeley, California, through the courtesy of Dr. G. O. Kohler.

There were no significant differences in egg production, efficiency of pigment utilization and yolk BCE values for Pro-Xan, the freeze dried alfalfa juice, the lutein extract or the latter combination. Based on the conditions of this test, xanthophyll utilization from Pro-Xan and lutein would be considered to be superior to that from dehydrated alfalfa.

Table 1. Utilization and Effects of Dehydrated Alfalfa, Pro-Xan and Freeze Dried Alfalfa Juice on Yolk Pigmentation and Egg Production

| Treatment ¹ | Dietary xanthophyll ppm | Yolk BCE ppm | Utilization efficiency % | Hen-day production % |
|----------------------------|----------------------------|--------------------|-----------------------------|-------------------------|
| 1. Dehydrated alfalfa | 10 | 20.0c ⁵ | 19.0 | 51.3 |
| 2. Dehydrated alfalfa | 20 | 48.0b | 14.5 | 34.4 |
| 3. Pro-Xan ² | 10 | 44.3b | 35.9 | 45.8 |
| 4. Pro-Xan | 20 | 70.4a | 33.3 | 51.6 |
| 5. F.D. juice ³ | 10 | 0.9c | -- | 45.6 |
| 6. F.D. juice | 20 | 2.7c | -- | 44.3 |
| 7. F.D. + lutein | 10 | 43.8b | 43.8 | 46.3 |
| 8. F.D. + lutein | 20 | 63.3a | 32.5 | 52.5 |
| 9. Lutein ⁴ | 10 | 45.8b | 43.6 | 52.9 |
| 10. Lutein | 20 | 64.4a | 32.9 | 55.5 |
| 11. Basal | -- | 1.9c | -- | 43.8 |
| 12. Basal | -- | 0.4c | -- | 43.6 |

¹Basal plus indicated source of pigment.

²Alfalfa protein-xanthophyll concentrate.

³Freeze dried alfalfa juice.

⁴Crystalline lutein in corn oil.

⁵Values with same subscript not significantly different, P<.05.