

Syracuse University

SURFACE

Physics

College of Arts and Sciences

11-9-2001

Anomaly Induced QCD Potential and Quark Decoupling

Joseph Schechter

Department of Physics, Syracuse University, Syracuse, NY

Stephen D.H. Hsu

Yale University

Francesco Sannino

Yale University

Follow this and additional works at: <https://surface.syr.edu/phy>

 Part of the [Physics Commons](#)

Recommended Citation

Schechter, Joseph; Hsu, Stephen D.H.; and Sannino, Francesco, "Anomaly Induced QCD Potential and Quark Decoupling" (2001). *Physics*. 299.

<https://surface.syr.edu/phy/299>

This Article is brought to you for free and open access by the College of Arts and Sciences at SURFACE. It has been accepted for inclusion in Physics by an authorized administrator of SURFACE. For more information, please contact surface@syr.edu.

Anomaly induced QCD potential and Quark Decoupling

Stephen D.H. HSU * Francesco SANNINO †

*Department of Physics, Yale University, New Haven, CT 06520-8120,
USA.*

Joseph SCHECHTER ‡

*Department of Physics, Syracuse University, Syracuse, NY
13244-1130, USA.*

Abstract

We explore the anomaly induced effective QCD meson potential in the framework of the effective Lagrangian approach. We suggest a decoupling procedure, when a flavored quark becomes massive, which mimics the one employed by Seiberg for supersymmetric gauge theories. It is seen that, after decoupling, the QCD potential naturally converts to the one with one less flavor. We study the N_c and N_f dependence of the η' mass.

PACS numbers:11.30.Rd, 12.39.Fe,11.30.Pb.

*Electronic address: hsu@hsunext.physics.yale.edu

† Electronic address : sannino@genesis1.physics.yale.edu

‡ Electronic address : schechte@suhep.phy.syr.edu

I. INTRODUCTION

In the last few years there has been a great flurry of interest in the effective Lagrangian approach to supersymmetric gauge theories. This was stimulated by some papers of Seiberg [1] and Seiberg and Witten [2] in which a number of fascinating “exact results” were obtained. In particular Seiberg has provided a very interesting picture for different phases of supersymmetric gauge theories. There are already several interesting review articles [3–5].

It is natural to hope that information obtained from the more highly constrained supersymmetric gauge theories can be used to learn more about ordinary gauge theories, notably QCD. In a recent paper [6] it was shown how the effective Lagrangian for super QCD might go over to the one for ordinary QCD by a mechanism whereby the gluinos and squarks in the fundamental theory decouple below a given supersymmetric breaking scale m . To accomplish this goal a suitable choice of possible supersymmetry breaking terms was used. An amusing feature of the model was the emergence of the ordinary QCD degrees of freedom which had been hidden in the auxiliary fields of the supersymmetric effective Lagrangian. Constraints on the supersymmetry breaking terms were obtained by requiring the trace of the energy momentum tensor to agree (at one loop level), once supersymmetry was broken, with that of ordinary QCD. It was also noticed that a reasonable initial picture resulted by neglecting the Kähler terms in the original supersymmetric effective Lagrangian. This feature is analogous to Seiberg’s treatment [1] of supersymmetric effective Lagrangians with different numbers of flavors. It led to a dominant piece of the QCD effective Lagrangian possessing a kind of tree level “holomorphicity”. Physically this corresponds to the explicit realization of the axial and trace anomalies by the model.

In this letter we will further explore the “holomorphic” part of the potential for QCD with $N_f (< N_c)$ massless quarks as obtained by breaking super QCD according to the scheme above (see section III of [6]). In particular, we shall study the analog of the decoupling procedure used in [1] when one flavor becomes heavy. In the present case we no longer have the protection of supersymmetry for deriving exact results so the procedure will be carried out at the one loop level. The analog of the Affleck-Dine-Seiberg (ADS) superpotential [7] turns out to be [6] the holomorphic part of the potential:

$$V(M) = -C(N_c, N_f) \left[\frac{\Lambda^{\frac{11}{3}N_c - \frac{2}{3}N_f}}{\det M} \right]^{\frac{12}{11(N_c - N_f)}} + \text{h.c.} , \quad (1.1)$$

where Λ is the invariant QCD scale and the meson matrix M contains N_f^2 scalars and N_f^2 pseudoscalars. A particular choice of the dimensionless positive quantity $C(N_c, N_f)$ was

made in [6]. Here we find the constraints on $C(N_c, N_f)$ which follow from decoupling. The result can be used to suggest that the well known [8,9] large N_c behavior of the η' (pseudoscalar singlet) meson mass should also include an N_f dependence of the form:

$$M_{\eta'}^2 \propto \frac{N_f}{N_c - N_f} \Lambda^2 \quad (N_f < N_c) . \quad (1.2)$$

This could be considered an indication that, while $M_{\eta'}$ is suppressed in the large N_c limit, there is an enhancement mechanism for N_f near $N_c - 1$ to explain its relatively large numerical value.

II. MESON POTENTIAL AND QUARK DECOUPLING

The various super QCD (SQCD) effective Lagrangians [7] are essentially derived using the anomaly structure of the underlying gauge theory as well as supersymmetry. It is thus interesting to first see how much of the effective QCD Lagrangians we get after decoupling the super-partner fields, follows just from anomaly considerations. In ordinary QCD the two relevant anomalies are the trace and Adler-Bell-Jackiw $U_A(1)$ anomalies:

$$\theta_m^m = -b \frac{g^2}{32\pi^2} F_a^{mn} F_{mn;a} \equiv 2bH , \quad (2.1)$$

$$\delta_{U_A(1)} V_{QCD} = N_f \alpha \frac{g^2}{32\pi^2} \epsilon_{mnr s} F_a^{mn} F_a^{rs} \equiv 4N_f \alpha G . \quad (2.2)$$

N_f is the number of flavors, $b = \frac{11}{3}N_c - \frac{2}{3}N_f$ is the coefficient of the one loop beta function and left handed quarks have unit axial charge. H and G can be interpreted as composite operators describing, in the confining regime, the scalar and pseudoscalar glueball fields [10]. The general effective (i.e. in terms of composite operators) potential which, at tree level, saturates the one loop anomalies is [10]:

$$V = -bF \sum_n \frac{c_n}{n} \ln \left(\frac{\mathcal{O}_n}{\Lambda^n} \right) + h.c. + V_I , \quad (2.3)$$

where \mathcal{O}_n is an operator built out of the relevant degrees of freedom at a given scale μ (glueballs, mesons, baryons, ..) with mass dimension n and axial charge q_n , $F = H + i\delta G$ and V_I is a scale and $U_A(1)$ invariant potential. The anomaly constraints are:

$$\sum_n c_n = 1 , \quad \sum_n \frac{c_n}{n} q_n = \frac{2N_f}{b\delta} . \quad (2.4)$$

Λ is connected with the invariant scale of the theory and the θ parameter (at one loop) via $\Lambda^b = \mu^b e^{-\tau}$ with $\tau = \frac{8\pi^2}{g^2(\mu)} - i\theta$. At high scales ($\mu \gg \Lambda$) we expect to approach the classical

regime where the glue potential displays the holomorphic structure $-\tau(H + iG) + h.c.$. This suggests choosing $\delta = 1$ and hence $F = H + iG$. Here we assume that the lightest meson $(M)^*$ and glueball (H and G) fields are the relevant degrees of freedom. It is amusing to notice that the field associated with the $U_A(1)$ transformation (η') can only appear in the first term of the potential in Eq. (2.3).

We can arrange the anomaly potential (first term in Eq. (2.3)) to be "holomorphic" by setting $\mathcal{O}_n = \mathcal{O}_n(F, \det M)$. By holomorphic we mean a potential of the form $\chi(\Phi) + h.c.$ where χ is a function of the generic complex field Φ . Now, in Ref. [6] it was shown that a suitable decoupling of the holomorphic superpotential in the Taylor Veneziano Yankielowicz (TVY) model [7] yielded an important "holomorphic" contribution to the ordinary potential for QCD. For $N_f (< N_c)$ flavors the result was of the form

$$V(F, M) = -\frac{11}{12} (N_c - N_f) F \left[\ln \left(\frac{-N_c F}{\gamma \Lambda^4} \right) - 1 \right] - F \ln \left(\frac{\det M}{\Lambda^{3N_f}} \right) + A(N_f, N_c) F + h.c. , \quad (2.5)$$

where $\gamma = \frac{12N_c - 4N_f}{11N_c - 2N_f}$ and $A(N_c, N_f)$ is a dimensionless constant which cannot be fixed by saturating the QCD anomalies and assuming holomorphicity. This is because the term AF does not contribute to θ_m^m and is also a chiral singlet. The potential in Eq. (2.5) is easily seen to be consistent with the general form in Eq. (2.3), and the associated constraints in Eq. (2.4) when the scale dimension of M is fixed to be 3 and $\delta = 1$. (The constants are $c_4 = 1 - 3\frac{N_f}{b}$ and $c_{3N_f} = 3\frac{N_f}{b}$).

By integrating out the "heavy" glueball field F one obtains the "light" meson field potential for M . This is given in Eq. (1.1) wherein the unknown coefficient $C(N_c, N_f)$ is related to $A(N_c, N_f)$ by the following expression:

$$A(N_c, N_f) = \frac{11}{12} (N_c - N_f) \ln \left(\frac{12}{11} \frac{N_c}{\gamma (N_c - N_f)} C(N_c, N_f) \right) . \quad (2.6)$$

Similar effective potential models and some related phenomenological questions have already been discussed in the literature [11–15].

The potential for the meson variables in Eq. (1.1) is similar to the effective ADS superpotential for massless super QCD theory with $N_f < N_c$ [7]

$$W_{ADS}(T) = - (N_c - N_f) \left[\frac{\Lambda_S^{3N_c - N_f}}{\det T} \right]^{\frac{1}{N_c - N_f}} , \quad (2.7)$$

*In Ref. [6] $M \simeq -\bar{q}RqL$ was denoted F_T .

where $T \simeq Q\tilde{Q}$ is the composite meson superfield, Q and \tilde{Q} are the quark super fields and Λ_S is the invariant scale of SQCD. In the instanton generated super potential the exponent of Λ_S is the coefficient of the super symmetric beta function. In the potential for the ordinary meson variables Eq. (1.1) the exponent inside the square brackets of the QCD invariant scale Λ is similarly the coefficient of the one loop QCD beta function $b = \frac{11}{3}N_c - \frac{2}{3}N_f$. As in the SQCD case the ordinary QCD potential Eq. (1.1) can also be constructed if we assign to Λ^b charge $2N_f$ under the matter field axial transformation, as prescribed by the θ -angle variation, and if holomorphicity in the coupling constant is assumed.

An important difference with respect to the ADS superpotential, as already explained in Ref. [6], is the fact that the QCD potential displays a fall to the origin while the ADS potential does not have a minimum for finite values of the squark condensate. In [6] it was shown that the fall to the origin can be cured by introducing non holomorphic terms which also trigger spontaneous chiral symmetry breaking.

The mesonic holomorphic potential, due to the presence of the appropriate beta function, is expected to hold for any $N_f < N_c$. In particular a non trivial check would be to decouple a single heavy flavor and show that the potential reduces to the potential for one less flavor fermion. A similar procedure, at the supersymmetric level, was used by Seiberg to check the validity of the ADS superpotential. This is in the same spirit as the well known [16] criterion for decoupling a heavy flavor (at the one loop level). For a small mass m of the N_f -th quark the perturbative contribution, prescribed in the fundamental lagrangian, of the mass operator to the potential is

$$V(M) = \dots - mM_{N_f}^{N_f} + \text{h.c.} . \quad (2.8)$$

To achieve a complete decoupling we generalize the perturbative mass operator to

$$V_m = -m^\Delta M_{N_f}^{N_f \Gamma} + \text{h.c.} , \quad (2.9)$$

where dimensional analysis requires $\Delta = 4 - 3\Gamma$. We interpret the departure from unity of Δ as an effective dynamical evolution of the chiral symmetry breaking operator for large values of m . The total potential for large m ,

$$V_T(M) = -C(N_c, N_f) \left[\frac{\Lambda^{\frac{11}{3}N_c - \frac{2}{3}N_f}}{\det M} \right]^{\frac{12}{11(N_c - N_f)}} - m^\Delta M_{N_f}^{N_f \Gamma} + \text{h.c.} , \quad (2.10)$$

where Λ is the appropriate invariant scale for N_f flavors and N_c colors, must convert to the potential for $N_f - 1$ light quarks. Notice that the generalized mass term preserves the holomorphic structure. Here it is being assumed that the non-holomorphic terms in the potential

can be treated as small perturbations. A similar mass operator was already introduced in Ref. [6] to appropriately decouple the gluino in the underlying theory. Eliminating the heavy degrees of freedom by their equations of motion and substituting back in the potential gives

$$V(\hat{M}) = - \left[\frac{\Gamma + w}{w} \right] \left[\frac{C(N_c, N_f) w}{\Gamma} \right]^{\frac{\Gamma}{\Gamma+w}} \left[\frac{\Lambda^{b(N_c, N_f)} m^{\frac{\Delta}{\Gamma}}}{\det \hat{M}} \right]^{\frac{\Gamma w}{\Gamma+w}} + \text{h.c.} , \quad (2.11)$$

where $w = \frac{12}{11(N_c - N_f)}$, $b(N_c, N_f)$ is the coefficient of the beta function for the theory with N_c colors and N_f flavors and \hat{M} is the meson matrix for $N_f - 1$ flavors. In the standard physical picture the gauge coupling constant evolves according to the QCD beta-function for N_f flavors above scale m and according to the QCD beta-function for $N_f - 1$ flavors below scale m . Since the coupling constant at scale μ is given by $\left(\frac{\Lambda}{\mu}\right)^b = \exp\left(-\frac{8\pi^2}{g^2(\mu)}\right)$, the matching at $\mu = m$ requires $\left(\frac{\Lambda_{N_c, N_f}}{m}\right)^{b(N_c, N_f)} = \left(\frac{\Lambda_{N_c, N_f-1}}{m}\right)^{b(N_c, N_f-1)}$, which yields

$$\Lambda_{N_c, N_f-1}^{b(N_c, N_f-1)} = \Lambda_{N_c, N_f}^{b(N_c, N_f)} m^{\frac{2}{3}} . \quad (2.12)$$

If we require the presence of the chiral symmetry breaking term in Eq. (2.10) to convert the N_f theory into the $N_f - 1$ theory, the ratio Δ/Γ is fixed to $\frac{2}{3}$ and by using the relation $\Delta = 4 - 3\Gamma$ one derives

$$\Delta = \frac{8}{11}, \quad \Gamma = \frac{12}{11} . \quad (2.13)$$

This value[†] of Γ is exactly what is needed to get the correct exponent

$$\frac{\Gamma w}{\Gamma + w} = \frac{12}{11(N_c - N_f + 1)} , \quad (2.14)$$

in Eq. (2.11). A consistent decoupling is obtained provided that the coefficient $C(N_c, N_f)$ obeys the following recursive relation

$$\left[\frac{C(N_c, N_f)}{N_c - N_f} \right]^{N_c - N_f} = \left[\frac{C(N_c, N_f - 1)}{N_c - N_f + 1} \right]^{N_c - N_f + 1} . \quad (2.15)$$

The general solution of Eq. (2.15) is finally,

[†]These are the same exponents found in section II of [6] for decoupling a massive gluino. Here we can also use, as in [6], the one loop matching of the anomaly of the traced energy momentum tensor to derive the same exponents.

$$C(N_c, N_f) = (N_c - N_f) D(N_c)^{\frac{1}{N_c - N_f}} . \quad (2.16)$$

Using Eq. (2.6), $A(N_c, N_f)$ can be expressed as the function of D :

$$A(N_c, N_f) = \frac{11}{12} (N_c - N_f) \ln \left(\frac{12 N_c}{11 \gamma} \right) + \frac{11}{12} \ln D(N_c) . \quad (2.17)$$

It is interesting to contrast the result in Eq. (2.16) for the coefficient of the "holomorphic" part of the QCD potential Eq. (1.1) with Seiberg's result [1] $C(N_c, N_f) = (N_c - N_f)$ for the coefficient of the ADS superpotential in Eq. (2.7). Clearly, in the SUSY case the analog of $D(N_c)$ is just a constant (which turns out, in fact, to be unity by an instanton analysis [1]). This feature arises in the SUSY case because of the existence of squark fields which can break the gauge and flavor symmetries by the Higgs mechanism; it leads to $C(N_c, N_f) = C(N_c - N_f)$. Since there are no appropriate scalar fields in ordinary QCD we do not expect such a feature. The possibility of a non-constant $D(N_c)$ factor can thus be taken as an indication that there is no Higgs mechanism present.

III. PHYSICAL EFFECTS

As mentioned above, we must add to the "holomorphic" piece of the potential (Eq. (1.1) with Eq. (2.6)) a "non-holomorphic" piece in order to prevent a "fall to the origin". This would also yield a non-zero value for $\langle M_a^b \rangle$. The non-holomorphic piece will be required to neither contribute to the trace anomaly nor to the axial anomaly. At the level of Eq. (2.3), where the "heavy" glueball fields have not yet been integrated out, such a potential could be of the form

$$V_I = \sum_n A_n (F^* F)^{\frac{1}{2} - \frac{3}{4}n} \text{Tr} \left[(M^\dagger M)^n \right] , \quad (3.1)$$

where the A 's are real constants. (For definiteness the terms have been restricted to those which are leading in the large N_c limit). For our initial analysis we assume that V_I is small compared to the holomorphic piece and can be treated as a perturbation. An example of a single term with $n > \frac{2}{3}$ was presented in Ref. [6]. An analogous term without the heavy F fields is

$$V'_I = \sum_n A'_n \text{Tr} \left[(M^\dagger M)^n \right] , \quad (3.2)$$

which should be added to Eq. (1.1). The new feature of the present model is that the piece which mocks up the anomalies can be thought of as inheriting a specific holomorphic structure from the SQCD theory.

Since the model of Eq. (1.1) plus Eq. (3.2) contains a great deal of arbitrariness it is natural to wonder what can be learned from it. The key point is that Eq. (3.2) possesses an axial $U(N_f)_L \times U(N_f)_R$ symmetry and cannot contribute to the η' (pseudoscalar singlet) mass. Thus we may learn something about this quantity from Eq. (1.1) directly. Now the η' field may be isolated from the meson matrix M as

$$M = K\tilde{U} \exp \left[i \frac{\eta'}{\alpha\Lambda\sqrt{N_f}} \right], \quad (3.3)$$

where $K = K^\dagger$, $\tilde{U}^\dagger = \tilde{U}^{-1}$, $\det\tilde{U} = 1$ and α is a dimensionless real constant. K contains the scalar fields and \tilde{U} the non flavor-singlet pseudoscalars. For our present purpose we may set $K = \beta\Lambda^3\mathbf{1}$, where β is another dimensionless real constant. For simplicity we shall make the approximation $\alpha = \beta = 1$, which will not affect our results in the large N_c limit. Then substituting Eq. (3.3) into Eq. (1.1), using Eq. (2.16) and expanding the result to quadratic order in η' yields the mass

$$M_{\eta'}^2 = \frac{2N_f}{N_c - N_f} \left(\frac{12}{11} \right)^2 \Lambda^2 D(N_c)^{\frac{1}{N_c - N_f}}. \quad (3.4)$$

Here it has been assumed that the η' has the canonical Lagrangian mass term $-\frac{1}{2}\partial_m\eta'\partial^m\eta'$. If the factors α and β were to be included the right hand side of Eq. (3.4) would get multiplied by $\alpha^{-2}\beta^{-\frac{12N_f}{11(N_c - N_f)}}$. α does not have any large N_c dependence and that of β gets washed out.

$D(N_c)$ is unspecified by our model so we cannot predict $M_{\eta'}$. We can however check the dependence on N_c of the quantities in Eq. (3.4) for large N_c . It is well known [8,9] that $M_{\eta'}^2 \sim 1/N_c$ for large N_c . Furthermore, $\langle F \rangle$ (defined after Eq. (2.3)) should behave like N_c for large N_c by standard counting arguments. Hence (see the discussion after Eq. (2.18) of Ref. [6]) Λ is expected to behave as $(N_c\langle F \rangle)^{\frac{1}{4}} \sim N_c^{\frac{1}{2}}$ at large N_c . For consistency we thus expect the large N_c behavior

$$D(N_c) \sim \left(\frac{1}{N_c} \right)^{N_c}. \quad (3.5)$$

It is amusing to observe that when N_f is close to N_c the resulting pole in Eq. (3.4) suggests a possible enhancement mechanism for the η' mass. This could explain the unusually large value of this quantity in the realistic three flavor case.

Of course the $N_f = N_c$ case is rather non-trivial. Equation (1.1) shows that the overall coefficient of our model, $C(N_c, N_f)$ should vanish for $N_f = N_c$. This is what happens in the SQCD situation. There, Seiberg [1] shows that the effective superpotential should

be replaced by a quantum constraint which also involves the baryon superfields. Before considering the analog of these results in our context, let us try to understand the situation a little better by going back to the holomorphic part of the potential in Eq. (2.5), in which the heavy glueball field F has not yet been integrated out. Focusing on the pieces of Eq. (2.5) which are related to generating an η' mass we get in a *schematic* form

$$V(F, M) \sim -(N_c - N_f) G^2 + \eta' G , \quad (3.6)$$

where G is the pseudoscalar glueball field defined in Eq. (2.2) and inessential factors have been dropped. The first term is a wrong sign (for $N_f < N_c$) mass term for G and is obtained by expanding the first term of Eq. (2.5). Integrating out G according to the equation of motion results in a correct sign mass term for the η' as long as $N_f < N_c$. It is clear that this mechanism will not work for $N_f \geq N_c$. From our present treatment we can see what is likely to be happening in the case of ordinary QCD. Here the non-holomorphic piece in Eq. (3.1) may contain a $-G^2$ piece when it is expanded and may thus generate an η' mass. We have been treating V_I in Eq. (3.1) as a small perturbation but, for consistency, it must be non-negligible especially for $N_f \geq N_c$. The usual phenomenology is consistent with the large N_c approximation to QCD so perhaps it is easiest to use the present model in that case too. In that event we would always have $N_f \ll N_c$. However it is certainly of interest to examine the small N_c case. The present model suggests that something different is occurring as we let N_f vary near N_c . It is noteworthy that recent preliminary computer simulations [17] indicate that there is a rather noticeable change in the quark condensate for N_f near N_c .

Now let us return to the discussion of the holomorphic part of the potential and point out that formally the $N_f = N_c$ case can be handled as in SQCD. When we put $N_f = N_c$ in Eq. (2.5)

$$V(F, M) \rightarrow -F \ln \left(\frac{\det M}{e^{A(N_c, N_c)} \Lambda_{N_c, N_c}^{3N_c}} \right) + h.c. . \quad (3.7)$$

Integrating out F then gives the *constraint* on the field M ,

$$\det M = e^{A(N_c, N_c)} \Lambda_{N_c, N_c}^{3N_c} , \quad (3.8)$$

rather than a potential $V(M)$. In fact the holomorphic potential vanishes as we argued. Nevertheless this vanishing potential is consistent (as in the SQCD case) with decoupling to the $N_f = N_c - 1$ theory. Following Eq. (2.9) we consider the $N_c = N_f$ potential with a heavy N_f -th quark of mass m :

$$V = 0 - m^\Delta \left(M_{N_f}^{N_f} \right)^\Gamma + h.c. , \quad (3.9)$$

where $\Delta = 8/11$ and $\Gamma = 12/11$ as before. Now the constraint in Eq. (3.8) enables us to write

$$M_{N_f}^{N_f} \det \hat{M} \rightarrow e^{A(N_c, N_c)} \Lambda_{N_c, N_c}^{3N_c}, \quad (3.10)$$

where \hat{M} is the meson matrix for $N_c - 1$ flavors. Substituting this back into Eq. (3.9) yields

$$V = -e^{A(N_c, N_c)\Gamma} \left(\frac{m_{\tilde{F}}^{\Delta} \Lambda_{N_c, N_c}^{3N_c}}{\det \hat{M}} \right)^{\Gamma} + h.c. . \quad (3.11)$$

When one uses Eq. (2.12) and Eq. (2.17) this is recognized as precisely Eq. (1.1) together with Eq. (2.16) for the $N_f = N_c - 1$ case. This argument may also serve to clarify what is "happening" in the SQCD situation.

A final question concerns the role of baryons when $N_f = N_c$. Baryon superfields were shown to play an important role [1] for $N_f = N_c$ in SQCD. One might hope to decouple the superpartners of the baryon superfields B and \tilde{B} [1] to obtain information on the QCD three quark baryons. However it is easy to see that none of the components of B and \tilde{B} contain any three quark composite fields. Thus the quick answer would be that the baryon fields are irrelevant for the ordinary QCD effective Lagrangian. This point of view is perhaps reinforced by the usual treatment (for large N_c) of baryons as solitons of the effective meson Lagrangian. Nevertheless, the arguments in the first part of section II show that ordinary baryon fields may readily be added to the holomorphic potential in a manner consistent with the anomaly structure of QCD. They may very well be relevant for a small N_c treatment of the theory. We hope to return to this question elsewhere.

ACKNOWLEDGMENTS

One of us (F.S.) is happy to thank Thomas Appelquist and Noriaki Kitazawa for interesting discussions. We would also like to thank Amir Fariborz for helpful comments on the manuscript. The work of S.H. and F.S. has been partially supported by the US DOE under contract DE-FG-02-92ER-40704. The work of J.S. has been supported in part by the US DOE under contract DE-FG-02-85ER 40231.

[1] N. Seiberg, Phys. Rev. D **49**, 6857 (1994). N. Seiberg, Nucl. Phys. **B435**, 129 (1995).

[2] N. Seiberg and E. Witten, Nucl. Phys. **B426**, 19 (1994), E **430**, 495(1994); **B431**, 484 (1994).

- [3] K. Intriligator and N. Seiberg, Nucl. Phys. Proc. Suppl. **45BC**, 1, 1996.
- [4] M.E. Peskin, *Duality in Supersymmetric Yang-Mills Theory*, (TASI 96): Fields, Strings, and Duality, Boulder, CO, 2-28 Jun 1996, hep-th/9702094.
- [5] P. Di Vecchia, *Duality in Supersymmetric Gauge Theories*, Surveys High Energy Phys. **10**, 119 (1997), hep-th/9608090.
- [6] F. Sannino and J. Schechter, *Toy Model for Breaking Supergauge Theories at the Effective Lagrangian Level*, Report Numbers: SU-4240-663, hep-th/9708113. To appear in Phys. Rev. D.
- [7] I. Affleck, M. Dine and N. Seiberg, Nucl. Phys. **B241**, 493 (1984); **B256**, 557 (1985). T.R. Taylor, G. Veneziano and S. Yankielowicz, Nucl. Phys. **B218**, 493 (1983). See also G. Veneziano and S. Yankielowicz, Phys. Lett. **113B**, 321 (1982).
- [8] E. Witten, Nucl. Phys. **B156**, 269 (1979).
- [9] G. Veneziano, Nucl. Phys. **B159**, 213 (1979).
- [10] The two anomaly case was discussed in J. Schechter, Phys. Rev. **D21**, 3393 (1980). The axial anomaly effective Lagrangian was discussed in the following papers: C. Rosenzweig, J. Schechter and G. Trahern, Phys. Rev. **D21**, 3388 (1980). P. Di Vecchia and G. Veneziano, Nucl. Phys. **B171**, 253 (1980). E. Witten, Ann. of Phys. **128**, 363 (1980). P. Nath and A. Arnowitt, Phys. Rev. **D23**, 473 (1981). A. Aurilia, Y. Takahashi and D. Townsend, Phys. Lett. **95B**, 265 (1980). K. Kawarabayashi and N. Ohta, Nucl. Phys. **B175**, 477 (1980).
- [11] A.A. Migdal and M.A. Shifman, Phys. Lett. **114B**, 445 (1982). J.M. Cornwall and A. Soni, Phys. Rev. **D29**, 1424 (1984); **32**, 764 (1985).
- [12] A. Salomone, J. Schechter and T. Tudron, Phys. Rev. **D23**, 1143 (1981). J. Ellis and J. Lanik, Phys. Lett. **150B**, 289 (1985). H. Gomm and J. Schechter, Phys. Lett. **158B**, 449 (1985).
- [13] H. Gomm, P. Jain, R. Johnson and J. Schechter, Phys. Rev. **D33**, 801 (1986).
- [14] H. Gomm, P. Jain, R. Johnson and J. Schechter, Phys. Rev. **D33**, 3476 (1986).
- [15] A QCD Lagrangian with a similar “holomorphic” piece plus a needed non holomorphic piece was discussed in sect. VII of Ref. [13]. This was partly motivated by a complex form for the two QCD coupling parameters which had been speculated in J. Schechter, *Proceedings of the third annual MRST meeting*, University of Rochester, April 30 - May 1 (1981).

- [16] E. Witten, Nucl. Phys. **B104**, 445 (1976); **B122**, 109 (1977). M. Shifman, A. Vainshtein and V. Zakharov, Nucl. Phys. **B147**, 385 (1979); **B147**, 448 (1979).
- [17] R.D. Mawhinney, *Evidence for pronounced quark loop effects in QCD* Report Numbers: CU-TP-839, hep-lat/9705031; *Dependence of QCD Hadron Masses on the Number of Dynamical Quarks* CU-TP-802, hep-lat/9705030; D. Chen, R.D. Mawhinney, Nucl. Phys. Proc. Suppl. **53**, 216, 1997.