

Chapman University Chapman University Digital Commons

Student Research Day Abstracts and Posters

Office of Undergraduate Research and Creative
Activity

12-10-2014

Toxicity and Cosmesis Outcomes for Single Fraction Intra-Operative Electron Radiotherapy (IOERT) for Breast Cancer

Monica Hanna

Chapman University, hanna130@mail.chapman.edu

Robert Ash

ST. JOSEPH HOSPITAL

Wesley Babaran

ST. JOSEPH HOSPITAL, wbabaran@yahoo.com

Michele M. Carpenter


ST. JOSEPH HOSPITAL

Afshin Forouzannia

ST. JOSEPH HOSPITAL

See next page for additional authors

Follow this and additional works at: http://digitalcommons.chapman.edu/cusrd_abstracts

 Part of the [Cancer Biology Commons](#), [Oncology Commons](#), [Other Analytical, Diagnostic and Therapeutic Techniques and Equipment Commons](#), and the [Radiology Commons](#)

Recommended Citation


Hanna, Monica; Ash, Robert; Babaran, Wesley; Carpenter, Michele M.; Forouzannia, Afshin; Harness, Jay K.; Kaltenecker, Brian; Maddula, Snehith; Williams, Venita; and Wagman, Lawrence, "Toxicity and Cosmesis Outcomes for Single Fraction Intra-Operative Electron Radiotherapy (IOERT) for Breast Cancer" (2014). *Student Research Day Abstracts and Posters*. Paper 75.
http://digitalcommons.chapman.edu/cusrd_abstracts/75

This Poster is brought to you for free and open access by the Office of Undergraduate Research and Creative Activity at Chapman University Digital Commons. It has been accepted for inclusion in Student Research Day Abstracts and Posters by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Toxicity and Cosmesis Outcomes for Single Fraction Intra-Operative Electron Radiotherapy (IOERT) for Breast Cancer

Authors

Monica Hanna, Robert Ash, Wesley Babaran, Michele M. Carpenter, Afshin Forouzannia, Jay K. Harness, Brian Kaltenecker, Snehith Maddula, Venita Williams, and Lawrence Wagman



TOXICITY AND COSMESIS OUTCOMES FOR SINGLE FRACTION INTRA-OPERATIVE ELECTRON RADIOTHERAPY (IOERT) FOR BREAST CANCER



Kaltenecker, B.; **Hanna, M.**; Babaran, W. MD; Maddula, S.; Forouzannia, A. MD; Carpenter, M. MD; Harness, J. MD; Ash, R. MD; Williams, V. MD; Wagman, L. MD
The Center for Cancer Prevention and Treatment, St. Joseph Hospital, Orange, CA

BACKGROUND: Adjuvant radiation therapy is proven to reduce local recurrence in patients with early stage breast cancer. The major toxicity of standard external beam radiation therapy is sunburn like acute effects, long term tanning of the skin, and changes in breast size. After the completion of surgery and closure of the surgical incision, the precise location of the tumor is difficult to access. Currently the standard of external beam radiation therapy is 3-6 weeks often accompanied by an additional one-week boost. To reduce toxicity by delivering the dose only to the area with breast cancer, to improve geographic accuracy by delivering radiation to the tumor bed prior to incision closure, and to reduce treatment time to approximately 30 minutes, IOERT was utilized as an alternative to external beam radiation therapy.


PRIMARY OBJECTIVE: The study's objective was to determine the short term toxicity and cosmesis profile of single fraction IOERT given as definitive treatment in a community setting.

STUDY POPULATION: From Mar 2012 to Jul 2014, 84 patients (3 bilateral), ages 45-91 y.o. with stage 0-II were treated with IOERT.

TREATMENT DETAILS: At the time of breast conservation surgery, a single 21 Gy fraction was administered to the tumor bed after resection. IOERT was delivered using 4.5-6 cm applicators (cylindrical shaped tube) with electron energies from 6-12 MeV. The median pathologic tumor size was 13 mm (4 tumors >25mm). Median follow up was 57.1 weeks.



Sentinel Lymph Node Biopsies



IOERT TOXICITY AND COSMESIS MEASUREMENT: At 2w, 6mo and 12mo, toxicity was assessed according to CTCAE Version 4.0 (range 0-4) and cosmesis based on a scale derived for national trials.

RESULTS:

	Scale	2 Week (%)	6 Month (%)	12 Month (%)
Toxicity Grade	0	49	69	62
	1	44	29	35
	2	7	2	3
Cosmesis Appearance	Excellent	71	86	79
	Good	28	14	21
	Fair	1	0	0

CONCLUSION: Single fraction IOERT was well tolerated by all patients with no grade 3+ toxicity up to 12 months. At one year, 97% of patients had 0-1 grade toxicity and 100% of patients had excellent or good cosmesis. This treatment, consistent with current reports, meets critical criteria for incorporation into practice. It reduces treatment by 3-6 weeks by including the surgical procedure and radiation treatment at a single episode. This single fraction IOERT combined with surgery is completed in approximately two hours. This is a vast improvement upon separate surgical and radiation procedures done at different times which lasts almost 3 months to accomplish both. SF IOERT, in addition to yielding low toxicity and high cosmesis, improves compliance and cost to the patient and the healthcare system.

REFERENCES:

Fisher B, Anderson S, Bryant J, et al. Twenty-year follow-up of a randomized trial comparing total mastectomy, lumpectomy and lumpectomy plus irradiation for the treatment of invasive breast cancer. *N Engl J Med* 2002; 347: 1233-41

Ash, R, Williams, R, Wagman, L, et al. Intraoperative Radiotherapy for Breast Cancer: Its Perceived Simplicity. *Journals oncology* Volume 27 Number 2 Feb 15, 2013

Veronesi U, Orecchia R, Luini A, et al. Intraoperative radiotherapy during breast conserving surgery: a study of 1,822 cases treated with electrons. *Breast Cancer Res Treat* 2010; 124: 141-151