

2015

Choice of Human–Computer Interaction Mode in Stroke Rehabilitation

Hossein Mousavi Hondori
University of California - Irvine

Maryam Khademi
University of California - Irvine


Lucy Dodakian
University of California - Irvine

Alison Mackenzie
Chapman University, amckenzi@chapman.edu

Cristina V. Lopes
University of California - Irvine

See next page for additional authors

Follow this and additional works at: http://digitalcommons.chapman.edu/pt_articles

 Part of the [Graphics and Human Computer Interfaces Commons](#), [Movement and Mind-Body Therapies Commons](#), [Other Computer Sciences Commons](#), [Other Rehabilitation and Therapy Commons](#), [Physical Therapy Commons](#), [Physiotherapy Commons](#), and the [Recreational Therapy Commons](#)

Recommended Citation

Hondori, H.M., Khademi, M., Dodakian, L., McKenzie, A., Lopes, C.V., Cramer, S.C., (2015). Choice of Human–Computer Interaction Mode in Stroke Rehabilitation. *Neurorehabil Neural Repair*, 1545968315593805.
DOI:10.1177/1545968315593805

This Article is brought to you for free and open access by the Physical Therapy at Chapman University Digital Commons. It has been accepted for inclusion in Physical Therapy Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Choice of Human–Computer Interaction Mode in Stroke Rehabilitation

Comments

This is a pre-copy-editing, author-produced PDF of an article accepted for publication in *Neurorehabilitation and Neural Repair* in 2015 following peer review. The definitive publisher-authenticated version is available online DOI: [10.1177/1545968315593805](https://doi.org/10.1177/1545968315593805).

Copyright

The authors

Authors

Hossein Mousavi Hondori, Maryam Khademi, Lucy Dodakian, Alison Mackenzie, Cristina V. Lopes, and Steven C. Cramer

Choice of Human-Computer Interaction Mode in Stroke Rehabilitation

Hossein Mousavi Hondori, PhD^{1*}, Maryam Khademi, MEng^{2*}, Lucy Dodakian, MA, OTR¹,
Alison McKenzie, DPT, PhD³, Cristina V. Lopes, PhD² and Steven C. Cramer, MD¹

Depts. ¹Neurology and ²Information and Computer Sciences, University of California, Irvine;
³Dept Physical Therapy, Chapman University

*These authors made equal contributions

FOR CORRESPONDENCE:

Steven C. Cramer, MD
UC Irvine Medical Center
200 S. Manchester Ave.
Suite 206
Orange, CA 92868-4280

Phone: (714) 456-6876
Fax: (714) 456-8805
Email: scramer@uci.edu

Number of words: 2560

Number of figures: 3

Number of color figures: 0

Number of tables: 2

Running title: Human computer interaction after stroke

Abstract

Background and objective: Advances in technology are providing new forms of human-computer interaction. The current study examined one form of human-computer interaction, augmented reality (AR), whereby subjects train in the real world workspace with virtual objects projected by the computer. Motor performances were compared with those obtained while subjects used a traditional human-computer interaction, i.e., a personal computer (PC) with a mouse.

Methods: Patients used goal-directed arm movements to play AR and PC versions of the Fruit Ninja video game. The two versions required the same arm movements to control the game but had different cognitive demands. With AR, the game was projected onto the desktop, where subjects viewed the game plus their arm movements simultaneously, in the same visual coordinate space. In the PC version, subjects used the same arm movements but viewed the game by looking up at a computer monitor.

Results: Among 18 patients with chronic hemiparesis after stroke, the AR game was associated with 21% higher game scores ($p=0.0001$), 19% faster reaching times ($p=0.0001$), and 15% less movement variability ($p=0.0068$), as compared to the PC game. Correlations between game score and arm motor status were stronger with the AR version.

Conclusions: Motor performances during the AR game were superior to those during the PC game. This result is due in part to the greater cognitive demands imposed by the PC game, a feature problematic for some patients but preferred for others. Mode of human-computer interface influences rehabilitation therapy demands and can be individualized for patients.

Keywords: stroke / recovery / direct interaction / indirect interaction / augmented reality

Introduction

Computer-assisted technology is being increasingly used to enhance physical and occupational therapy after stroke^{1,2}. Many current applications rely on a traditional human-computer interaction, in which a person uses a game controller such as a mouse with a desktop personal computer (PC) while looking up at a computer monitor (Figure 1A)³⁻⁵. There are potential limitations with this type of human-computer interaction, for example, a subject must perform a visuospatial transform from the coordinates of arm movement space to the coordinates of computer monitor space, and interaction with the computer's output is indirect.

Other human-computer interfaces⁶⁻⁹ have been studied, such as spatial augmented reality (AR, Figure 1B)¹⁰. With an AR-based approach, subjects interact in the real world workspace with virtual objects projected by a computer¹¹. No visuospatial transform by the subject is required due to the fact that the subject interacts directly with the computer's output. The extent to which a subject's interaction with computer output is direct vs. indirect may be important at many levels¹²⁻¹⁵. With an indirect interaction, subjects do not handle the computer's output medium; with a direct interaction, they do. For instance, using a mouse to play a game is an indirect interaction between the user and the output medium (computer monitor), whereas using a finger on an iPad surface or on a virtual object projected by a computer in order to control a game is a direct interaction between user and computer output.

These differences between a PC and an AR approach provide an opportunity to study how choice of human-computer interface affects motor performance, and ultimately stroke rehabilitation. To examine this issue, motor performance of patients with chronic stroke was measured as each played two different versions of a popular video game, one PC-based and one AR-based. The two versions required the same arm movement kinematics to control the game but differed in level of cognitive demand, which was higher with the PC approach because it

1
2
3 uncouples a subject's eye and hand movements and requires the subject to perform a visuospatial
4
5 transform^{16, 17}. Increased cognitive demand is an important consideration when designing
6
7
8 computerized therapies to promote stroke rehabilitation¹⁸⁻²⁰. The current study hypothesized that
9
10 the higher level of cognitive demand during the PC-based game would be associated with lower
11
12 game scores and less efficient movements as compared to the AR-based game.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Methods

Subjects

Patients with chronic stroke were recruited from the community. This study was approved by the Institutional Review Board, and all subjects signed informed consent. Entry criteria were age >18 years, history of stroke with onset >6 months prior, and arm weakness operationally defined as Fugl-Meyer (FM) total arm motor score²¹ of 15-55 (out of 66) plus at least 5 degrees active range of motion in either wrist or index finger metacarpophalangeal joint on the stroke-affected side. Exclusion criteria were severe cognitive, attentional, or language deficits; a non-stroke diagnosis that affected arm or hand function; and co-existent major neurological or psychiatric disease.

After signing consent, medical history was reviewed, followed by a brief exam that included a measure of loss of body/function, the FM arm motor scale²¹ as well as a measure of activity limitations, the Box & Blocks test (BBT)^{22, 23}. Subjects then played the PC-based and the AR-based versions of the game with the paretic arm, as below.

Game design

Design of the computerized PC and AR games was modeled after the movements required by the BBT^{22, 23} which asks subjects to coordinate proximal and distal arm activity during a rapidly repeated task with high movement precision. We applied to this framework a simple version of the popular Fruit Ninja game, which by 2012 had been downloaded to 1 in 3 U.S. iPhones²⁴. Thus, in both our PC and AR versions of this game, subjects were seated at a table and asked to perform precise, rapid, repeated goal-directed reaching tasks requiring coordinated hand/shoulder movements (Figure 1). A small plastic cup was held in the stroke-affected hand. This cup served as a color-marker that was tracked by the camera and guided the fruit-slicing

1
2
3 cursor during game play; during the PC game this cup guided the on-screen cursor, while during
4
5 the AR game, this cup was the cursor.
6
7

8 Subjects were seated and introduced to the two versions of the game. A fruit target would
9
10 then appear in one of the four corners of the tabletop and start moving around the desktop
11
12 workspace. The subject was to use the stroke-affected hand to move the cursor onto the fruit
13
14 (which would slice the fruit in half, make a noise, and score 1 point) before the fruit disappeared
15
16 (which would score 0 points). Once the fruit was sliced, or disappeared unsliced, a new fruit
17
18 target would appear in one of the four corners, the sequence of which was pseudo-randomized.
19
20 Subjects were instructed to slice as many fruits as possible within the allotted time. The score
21
22 was presented throughout each game. One round of each game was 90 seconds long.
23
24
25
26

27 Subjects played one practice round each of the PC and the AR game, which were not
28
29 recorded or analyzed. Next, study data were acquired as the subject played three rounds of either
30
31 the PC or the AR game, the order of which was randomly assigned across subjects, followed by
32
33 three rounds of the other version of the game. Subjects were provided with a brief break between
34
35 each round.
36
37
38
39

40 *PC vs. AR game setup*

41
42 All game features and arm movement demands were identical across the two versions of the
43
44 game and required that the subject move the stroke-affected hand around the tabletop to slice
45
46 fruit. Subjects sat in the same chair at the same table. Both game versions used the same
47
48 overhead camera (Microsoft LifeCam VX-2000 Webcam, 30 fps, 720p HD capture) to capture
49
50 the subject's tabletop hand movements, which moved the game's fruit-slicing cursor. These
51
52 camera images were fed to a computer running a program that tracked the color-marker and that
53
54 provided real time audiovisual feedback.
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
PC set-up (Figure 1A): The subject was instructed to gaze straight ahead at a 15-inch computer monitor whose position was fixed throughout the experiments. This monitor displayed the fruit targets as well as the cursor, a circle that represented the real time position of the subject's hand and moving in a 15-inch tabletop workspace. With the PC setup, the subject's hand was 45 degrees away from his/her direction of gaze.

15
16
17
18
19
20
21
22
23
24
25
26
27
AR set-up (Figure 1B): The subject was instructed to gaze directly at the tabletop, onto which images appeared from a projector (P4-X Pico Projector, AAXA Technologies, Tustin, CA); the monitor was not used and remained blank. The projector displayed (720p HD) fruit targets, and the 15-inch workspace with four corners, directly onto the tabletop. With the AR setup, the subject gazed directly at his/her hand as it moved to slice the virtual fruit targets.

28 29 *Performance metrics*

30
31
32
33
34
The primary metric was game score. Four secondary metrics were extracted from game performance to aid interpretation of game scores:

35
36
37
38
39
1. *Game score*: The number of fruit targets successfully sliced during a 90-second round of game play. The 3 PC rounds were averaged to produce a single score, as were the 3 AR rounds.

40
41
42
43
44
2. *Reaching time*: The amount of time between reaching one fruit target and reaching the next fruit target.

45
46
3. *Movement time consistency*: The coefficient of variation of reaching times.

47
48
49
50
51
4. *Response latency*: The amount of time between display and target reaching, recorded for the first fruit target only in each round of the game.

52
53
54
55
56
57
58
59
60
5. *Movement efficiency*: The ratio of the ideal travel distance (the distance that would be travelled if the same number of targets were reached using perfect straight lines) over the actual

1
2
3 travel distance (the actual distance travelled by each subject reaching the respective number of
4
5 targets).
6
7
8
9
10

11 12 *Statistical Data Analysis* 13

14
15 Data were analyzed using *JMP 10.0* (SAS, Cary, NC) and used two-tailed testing at alpha of
16
17 0.05. Parametric methods were used to analyze data that were normally distributed or could be
18
19 transformed to a normal distribution, otherwise non-parametric methods were used. The primary,
20
21 plus three secondary, game performance metrics were compared between the PC and AR
22
23 versions of the game. Next, the relationship between game scores and motor status (BBT score
24
25 and FM score) were calculated, with secondary analyses considering FM sub-scores. Finally, in
26
27 an exploratory analysis, reaching times were compared according to the quadrant of visual space
28
29 in which the fruit target first appeared, separately for the PC and for the AR game versions, and
30
31 according to side of stroke.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

A total of 18 patients were recruited. All completed the study protocol, with no adverse events reported including no fatigue. Enrollees were on average 57 years old, 70 months post-stroke, and had mild-moderate motor deficits (Table 1).

Motor performance with PC vs. AR game versions

By all measures, motor performance while playing the AR-based version of the Fruit Ninja game was superior to the PC-based version. Game score, the primary metric, averaged 58 ± 10 targets/game (mean \pm SD; range 42-78) with the AR-based version, 21% higher ($p=0.0001$) than when the same patients played the same game but using the PC-based version (48 ± 10 targets, range 34-71). All three secondary metrics were also significantly different between AR and PC versions and provided insight into these game scores. Reaching times with AR were 19% faster than with PC (870 ± 350 vs. 1070 ± 510 msec, $p=0.0001$); consistent with faster movements in the AR-based version, subjects traveled longer distances during 90 seconds of playing the AR-based version as compared to the PC-based version (22.113 ± 0.433 vs. 19.998 ± 0.549 meters, $p=0.0002$). Movement time consistency was also better with AR, showing 15% less variability compared to PC (0.41 vs. 0.48, $p=0.0068$). Response latencies were 15% shorter with AR as compared to PC (1.71 ± 0.66 vs. 2.01 ± 0.79 sec, $p=0.0249$). Movements were more efficient with AR as compared to PC ($60.33 \pm 14.8\%$ vs. $55.28 \pm 14.4\%$, $p=0.0019$). Visual inspection was consistent, indicating that movements during the AR-based game were more consistent and less erratic vs. the PC-based game (Figure 2).

The rate of task learning was similar between AR and PC based on improvement in scores over time. Analysis of the scores in consecutive rounds within each version shows that in both the AR and PC versions, performances improved significantly from round 1 to 2 (from

Hondori et al

54.88±9.24 to 60.28±9.18 in AR, $p= 0.0010$; and from 45.11±7.78 to 48.72±9.98 in PC, $p= 0.0257$), but not from round 2 to 3 (from 60.28±9.18 to 61.94±9.66 in AR, $p= 0.2795$; and from 48.72±9.98 to 50.22±9.47 in PC, $p= 0.2991$). This suggests similar learning curves for the AR-based and the PC-based versions of the game.

Behavioral correlates of PC and AR scores

AR scores correlated significantly with all four motor assessments of the stroke-affected arm (Table 2 and Figure 3), particularly the BBT score. By contrast PC scores were related to motor status in only one instance, and to a weaker degree as compared to AR scores. If a formal Bonferroni correction was applied to correct for multiple comparisons, only AR scores would remain significantly related to motor status.

Motor performances in relation to quadrant of visual space

The quadrant into which each fruit target appeared during game play was classified in egocentric space, i.e., either near or far from the body, plus either ipsilateral or contralateral to the stroke-affected arm. During game play with the AR setting, patients with right arm motor deficits ($n=12$) showed the fastest reaching times when targets appeared in the near-contralateral quadrant, significantly faster vs. targets appearing in any of the three other quadrants ($p = 0.0001-0.0002$). The same was true when patients with left arm motor deficits ($n=6$) played the game with the AR setting, with reaching times being fastest for targets appearing in the near-contralateral quadrant as compared to each of the other three quadrants ($p = 0.0001-0.0007$). For both right arm and left arm groups, reaching times did not differ in relation to quadrant of space of target origin when the game was played using the PC setting.

Discussion

In the current study, patients with chronic post-stroke hemiparesis used the affected arm to play the Fruit Ninja game using two different human-computer interfaces, one PC and one AR. During the AR-based version of the game, as compared to PC, scores were higher, movements were faster and more consistent, and performances were more tightly linked with arm motor function. These differences are likely due to the additional cognitive demands imposed when playing the PC-based game. The current findings underscore how choice of human-computer interface can influence task demands and thus behavioral performances and is thus likely to be important when designing a stroke rehabilitation protocol.

Although movement demands were identical across the two versions of the game, cognitive demands were not. Both versions of the Fruit Ninja game required subjects to make arm movements that rapidly integrated visual and somatosensory inputs^{25, 26}. During AR version play, subjects looked directly at their hand as it moved to control gameplay, but during PC version play, subjects instead looked at a computer screen, which increased cognitive demand. This modulation of cognitive demand can be understood in terms of a difference in the spatial congruence between movement goal and motor output²⁷, an uncoupling between eye and hand movements¹⁶, or the imposition of an added sensorimotor transformation²⁸. Regardless, the higher cognitive demand associated with playing the PC-based version of the game reduced game scores by 21% compared with the AR-based version ($p=0.0001$), in association with slower and less consistent arm movements. Differences in cognitive requirements are particularly important after stroke, as demand increases having a negligible effect in healthy subjects can exceed cognitive reserve and degrade behavioral performance in subjects with stroke^{16, 18-20}.

1
2
3 Although the increased cognitive demand in the PC-based game reduced movement quality
4 (lower scores, slower reaching, less consistent movement), this might nonetheless be the
5 preferred rehabilitation approach for some subjects such as those needing to strengthen
6 visuospatial skills. The brain constructs a stable spatial representation of the workspace
7 environment²⁹, transforming hand and target positions into a common body-centered space³⁰⁻³².
8 The AR game exists in this same space--spatial transforms for game play are the same as those
9 used during real world activities--but the PC game plays out on the computer monitor, which has
10 its own coordinate space. As such, with an PC-based approach, the patient must perform at least
11 one extra spatial transform to convert the virtual world's coordinates into body-centered
12 coordinates. A PC approach may thus be useful to promote cognitive recovery after stroke.
13 Furthermore, increased cognitive demand modulates activity in brain motor networks and can
14 enhance motor learning in healthy subjects and in subjects with stroke³³⁻³⁶, and so might have a
15 role in some types of motor rehabilitation. Therefore, future studies might test the hypothesis that
16 sustained practice of the PC game, as compared to the AR game, may be associated with a larger
17 degree of motor improvement after stroke.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 Advances in technology are providing many new types of computer interface to promote
40 brain repair after stroke. The current study examined two forms, PC and AR, incorporating the
41 popular Fruit Ninja game into each. A PC approach to gaming substitutes a non-immersive
42 virtual environment in place of the real world¹¹, shifting the subject's attention towards the
43 virtual world (computer monitor) rather than to the real world (the tabletop upon which the
44 paretic hand is moving)³⁷. An AR approach to gaming overlays computer-generated virtual
45 objects into the real world, thereby offering a direct interaction between user and computer
46 output. Such real world activities may be more intuitive than many PC-based approaches, which
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 might promote massed practice. An AR approach might also allow patients to practice real-life
4
5 functional tasks safely, for example, a patient might work on being independent in the kitchen by
6
7 working with computer-generated objects that have zero risk of spilling, burning, or electric
8
9 shock. It also provides standardization and consistency across repeated trials, features that are
10
11 difficult to achieve in traditional therapy involving real objects; furthermore, adopting an AR
12
13 approach allows tasks to be practiced in a structured way, using highly controlled environments,
14
15 and can be gamified to encourage and intensify practice. Also, feedback (multimedia, error
16
17 augmentation etc.) and task difficulty are readily adjustable to the level of disability and/or
18
19 personal preferences of individual patients. A PC-based approach might nonetheless have
20
21 advantages, for example, to promote recovery of visuospatial skills, or by using modulation of
22
23 cognitive demands to promote other forms of recovery. The optimal choice of human-computer
24
25 interface likely varies across individuals, over time, and according to specific goals. The current
26
27 study provides insights into how choice of human-computer interface affects motor performance
28
29 and thus might influence stroke rehabilitation and motor recovery.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Hondori et al

Page 14

Acknowledgments

This work was funded by grants from the American Heart Association (13GRNT16990060) and from the National Institutes of Health (R01 NS059909, NIH/NCRR U1 TR000153, and K24 HD074722).

Conflict of interest disclosure

H.M. Hondori: none, M. Khademi: none, A. McKenzie: none, L. Dodakian: none, C.V. Lopes: none, S.C. Cramer: consultant/advisory board; modest; GlaxoSmithKline, Microtransponder, Dart Neuroscience, RAND Corporation, and Roche. Dr. Cramer is a cofounder of personalRN.

References

1. Ong S-K, Shen Y, Zhang J, Nee AY. Augmented reality in assistive technology and rehabilitation engineering. *Handbook of augmented reality*. Springer; 2011:603-630.
2. Fluet GG, Deutsch JE. Virtual reality for sensorimotor rehabilitation post-stroke: The promise and current state of the field. *Current physical medicine and rehabilitation reports*. 2013;1:9-20
3. Krebs HI, Ferraro M, Buerger SP, Newbery MJ, Makiyama A, Sandmann M, et al. Rehabilitation robotics: Pilot trial of a spatial extension for mit-manus. *J Neuroeng Rehabil*. 2004;1:5
4. Takahashi C, Der-Yeghiaian L, Le V, Motiwala R, Cramer S. Robot-based hand motor therapy after stroke. *Brain: a journal of neurology*. 2008;131:425
5. Housman S, Scott K, Reinkensmeyer D. A randomized controlled trial of gravity-supported, computer-enhanced arm exercise for individuals with severe hemiparesis. *Neurorehabil Neural Repair*. 2009;23:505
6. Saposnik G, Levin M. Virtual reality in stroke rehabilitation a meta-analysis and implications for clinicians. *Stroke*. 2011;42:1380-1386
7. Viaud-Delmon I, Gaggioli A, Ferscha A, Dunne S. Human computer confluence applied in healthcare and rehabilitation. *Stud Health Technol Inform*. 2012;181:42-45
8. Herz DM, Haagensen BN, Christensen MS, Madsen KH, Rowe JB, Lokkegaard A, et al. The acute brain response to levodopa heralds dyskinesias in parkinson disease. *Annals of neurology*. 2014;75:829-836
9. Paolucci S, Antonucci G, Grasso MG, Bragoni M, Coiro P, De Angelis D, et al. Functional outcome of ischemic and hemorrhagic stroke patients after inpatient rehabilitation: A matched comparison. *Stroke*. 2003;34:2861-2865
10. Mousavi HH, Khademi M, Dodakian L, Cramer S, Lopes C. A spatial augmented reality rehab system for post-stroke hand rehabilitation. *Studies in health technology and informatics*. 2013;184:279
11. Baus O, Bouchard S. Moving from virtual reality exposure-based therapy to augmented reality exposure-based therapy: A review. *Front Hum Neurosci*. 2014;8:112

12. Knoedel S, Hachet M. Multi-touch rst in 2d and 3d spaces: Studying the impact of directness on user performance. *3D User Interfaces (3DUI), 2011 IEEE Symposium on*. 2011:75-78
13. Schmidt D, Block F, Gellersen H. A comparison of direct and indirect multi-touch input for large surfaces. *Human-computer interaction–interact 2009*. Springer; 2009:582-594.
14. Cheng K, Pulo K. Direct interaction with large-scale display systems using infrared laser tracking devices. *Proceedings of the Asia-Pacific symposium on Information visualisation-Volume 24*. 2003:67-74
15. Forlines C, Balakrishnan R. Evaluating tactile feedback and direct vs. Indirect stylus input in pointing and crossing selection tasks. *Proceedings of the SIGCHI Conference on Human Factors in Computing Systems*. 2008:1563-1572
16. Lajoie Y, Teasdale N, Bard C, Fleury M. Attentional demands for static and dynamic equilibrium. *Experimental brain research*. 1993;97:139-144
17. Faust-Socher A, Kenett YN, Cohen OS, Hassin-Baer S, Inzelberg R. Enhanced creative thinking under dopaminergic therapy in parkinson disease. *Annals of neurology*. 2014;75:935-942
18. Hondori HM, Khademi M, McKenzie A, Dodakian L, Lopes CV, Cramer SC. Abstract t mp43: Utility of augmented reality in relation to virtual reality in stroke rehabilitation. *Stroke*. 2014;45:ATMP43-ATMP43
19. Knaut L, Subramanian S, McFadyen B, Bourbonnais D, Levin M. Kinematics of pointing movements made in a virtual versus a physical 3-dimensional environment in healthy and stroke subjects. *Archives of physical medicine and rehabilitation*. 2009;90:793
20. Subramanian S, Knaut LA, Beaudoin C, McFadyen BJ, Feldman AG, Levin MF. Virtual reality environments for post-stroke arm rehabilitation. *J Neuroeng Rehabil*. 2007;4:20
21. See J, Dodakian L, Chou C, Chan V, McKenzie A, Reinkensmeyer DJ, et al. A standardized approach to the fugl-meyer assessment and its implications for clinical trials. *Neurorehabilitation and neural repair*. 2013;27:732-741
22. Mathiowetz V, Volland G, Kashman N, Weber K. Adult norms for the box and block test of manual dexterity. *The American Journal of Occupational Therapy*. 1985;39:386-391

23. Desrosiers J, Bravo G, Hebert R, Dutil E, Mercier L. Validation of the box and block test as a measure of dexterity of elderly people: Reliability, validity, and norms studies. *Archives of physical medicine and rehabilitation*. 1994;75:751-755
24. Wikipedia Contributors. Fruit ninja. Last accessed september 26 2014. [Http://en.Wikipedia.Org/wiki/fruit_ninja](http://en.Wikipedia.Org/wiki/fruit_ninja). 2014
25. Sober SJ, Sabes PN. Multisensory integration during motor planning. *J Neurosci*. 2003;23:6982-6992
26. Shumway-Cook A, Woollacott M. Attentional demands and postural control: The effect of sensory context. *The journals of gerontology. Series A, Biological sciences and medical sciences*. 2000;55:M10-16
27. Fitts PM, Seeger CM. S-r compatibility: Spatial characteristics of stimulus and response codes. *J Exp Psychol*. 1953;46:199-210
28. Messier J, Kalaska JF. Differential effect of task conditions on errors of direction and extent of reaching movements. *Experimental brain research*. 1997;115:469-478
29. Fiehler K, Rösler F, Henriques DY. Interaction between gaze and visual and proprioceptive position judgements. *Exp Brain Res*. 2010;203:485-498
30. Crawford JD, Henriques DY, Medendorp WP. Three-dimensional transformations for goal-directed action. *Annual review of neuroscience*. 2011;34:309-331
31. Flanders M, Tillery SIH, Soechting JF. Early stages in a sensorimotor transformation. *Behavioral and Brain Sciences*. 1992;15:309-320
32. Soechting J, Tillery SH, Flanders M. Transformation from head-to shoulder-centered representation of target direction in arm movements. *Journal of Cognitive Neuroscience*. 1990;2:32-43
33. Turolla A, Dam M, Ventura L, Tonin P, Agostini M, Zucconi C, et al. Virtual reality for the rehabilitation of the upper limb motor function after stroke: A prospective controlled trial. *Journal of neuroengineering and rehabilitation*. 2013;10:85
34. Todorov E, Shadmehr R, Bizzi E. Augmented feedback presented in a virtual environment accelerates learning of a difficult motor task. *Journal of motor behavior*. 1997;29:147-158

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
35. Dennis A, Bosnell R, Dawes H, Howells K, Cockburn J, Kischka U, et al. Cognitive context determines dorsal premotor cortical activity during hand movement in patients after stroke. *Stroke*. 2011;42:1056-1061
36. Dodakian L, Sharp KG, See J, Abidi NS, Mai K, Fling BW, et al. Targeted engagement of a dorsal premotor circuit in the treatment of post-stroke paresis. *NeuroRehabilitation*. 2013;33:13-24
37. Imam B, Jarus T. Virtual reality rehabilitation from social cognitive and motor learning theoretical perspectives in stroke population. *Rehabilitation research and practice*. 2014;2014:594540

Figure legends

Figure 1. A patient playing the [A] PC and [B] AR versions of the Fruit Ninja game. Game features and movement demands were identical across the two versions of the game. Both required that the subject move the stroke-affected hand across the tabletop in order to control the movement of a cursor that earned points by slicing fruit targets. Both versions of the game leveraged the same camera; however in the PC version, game activity was displayed using a computer monitor, while in the AR version, game activity was displayed on the tabletop using a projector. This difference imposed two key differences. First, during PC play but not AR play, the subject needed to perform an extra spatial transform in order to convert the coordinates of tabletop arm movements to the coordinates of the cursor movements seen on the monitor. Second, during PC play, eye and hand movements were uncoupled, with subjects receiving proprioceptive feedback but no visual feedback (gaze was directed at the monitor not their hand), while during AR play, visual and proprioceptive feedback were coupled. These differences likely account for the significantly poorer motor performances seen with PC play.

Figure 2. Distribution of hand position for [A] all patients playing the AR version of the game and [B] all patients playing the PC versions of the Fruit Ninja game. The target locations and straight trajectories between targets are more prominent in the AR version, while movements are more inconsistent and erratic during the PC version, consistent with quantitative values for game metrics. Also pictured are the distribution of hand position for a single exemplary subject during the [C] AR version and [D] the PC version of the game.

Figure 3. The top row shows correlation of AR game scores with the proximal Fugl-Meyer subscores, distal Fugl-Meyer subscores, and the Box & Blocks Test scores, from left to right; the bottom row shows correlation of PC game scores with the same. FM=Fugl-Meyer.

Table 1. Subject characteristics

n	18
Age	57 ± 14 years
Gender (F/M)	8/10
Dominant hand (R/L)	12/6
Stroke-affected Side (R/L)	12/6
Time post-stroke	70 ± 73 months
Hypertension	61%
Diabetes mellitus	22%
Atrial fibrillation	5%
FM total score	56 ± 11
FM hand/wrist subscore	21 ± 5
FM proximal subscore	30 ± 7
BBT score	
Affected arm	38 ± 14
Affected/unaffected arm	0.66 ± 0.25


Values are mean ± SD or percentages.

Table 2. Behavioral correlates of AR and PC game scores


Behavioral Measure	AR game		PC game	
	r	p	r	p
FM hand/wrist subscore	0.56	0.0149	0.36	0.148
FM proximal subscore	0.52	0.0266	0.32	0.1957
BBT score, affected arm	0.64	0.0042	0.48	0.0433

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A. Personal computer version of the game




B. Augmented reality version of the game



C. camera
P. projector

53x56mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



101x83mm (300 x 300 DPI)

view

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




fig 3
101x75mm (300 x 300 DPI)

Review