

## Chapman University Chapman University Digital Commons

Physical Therapy Faculty Articles and Research

Physical Therapy

2014

# The Influence of Hip Strength on Lower-Limb, Pelvis, and Trunk Kinematics and Coordination Patterns During Walking and Hopping in Healthy Women

Jo Armour Smith

*Chapman University*, [josmith@chapman.edu](mailto:josmith@chapman.edu)


John M. Popovich

*Michigan State University*

Kornelia Kulig

*University of Southern California*

Follow this and additional works at: [http://digitalcommons.chapman.edu/pt\\_articles](http://digitalcommons.chapman.edu/pt_articles)

 Part of the [Musculoskeletal System Commons](#), [Physical Therapy Commons](#), and the [Women's Health Commons](#)

### Recommended Citation

Armour Smith J, Popovich JM Jr, Kulig K. The influence of hip strength on lower limb, pelvis and trunk kinematics during walking and hopping in healthy women. *Journal of Orthopaedic and Sports Physical Therapy*. 2014; 44 (7): 525-531.  
DOI: 10.2519/jospt.2014.5028

This Article is brought to you for free and open access by the Physical Therapy at Chapman University Digital Commons. It has been accepted for inclusion in Physical Therapy Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact [laughtin@chapman.edu](mailto:laughtin@chapman.edu).

---

# The Influence of Hip Strength on Lower-Limb, Pelvis, and Trunk Kinematics and Coordination Patterns During Walking and Hopping in Healthy Women

## **Comments**

This is a pre-copy-editing, author-produced PDF of an article accepted for publication in *Journal of Orthopaedic and Sports Physical Therapy*, volume 44, issue 7, in 2014 following peer review. The definitive publisher-authenticated version is available online at [DOI: 10.2519/jospt.2014.5028](https://doi.org/10.2519/jospt.2014.5028)

## **Copyright**

Journal of Orthopaedic & Sports Physical Therapy 2014. For personal use only. No other uses without permission. All rights reserved.

24 Key Words: coordination, muscle performance, **hopping**, gait.

Review Copy

## 25 INTRODUCTION

26 Musculoskeletal disorders of the lower limbs are often associated with both poor hip  
27 muscle performance and altered kinematics during dynamic tasks. However, it is still unclear  
28 whether altered lower limb or pelvis/trunk motion as a result of hip weakness contributes to the  
29 development of musculoskeletal pathology and pain.<sup>13,25</sup> During the stance phases of activities  
30 such as walking, running or hopping, the hip extensors and abductors play a complex role in  
31 control of the lower extremities, pelvis and trunk. This includes deceleration of hip internal  
32 rotation and adduction<sup>16</sup> and maintenance of the equilibrium of the pelvis and trunk over the  
33 stance limb.<sup>8</sup> Additionally, motion at the hip, pelvis and trunk influences kinematics and kinetics  
34 at the knee.<sup>13,25</sup> Therefore, weakness of the hip musculature may be associated with altered  
35 kinematics at the knee, hip, pelvis or trunk.

36  
37 A number of studies have examined the relationship between diminished hip muscle  
38 performance and kinematics in patients with musculoskeletal dysfunction. For example, females  
39 with patellofemoral pain syndrome (PFP) have decreased maximum hip abductor and extensor  
40 torque and increased peak knee external rotation and increased hip adduction during the stance  
41 phase of running compared with healthy controls.<sup>30,34</sup> Similarly, hip osteoarthritis is associated  
42 with decreased hip abductor strength as well as increased pelvic drop and hip internal rotation  
43 during the stance phase of walking.<sup>2,33</sup> However, cross sectional studies of patient populations do  
44 not discriminate between weakness resulting from musculoskeletal pain or pathology and  
45 weakness that may have contributed to the original development of the disorder.<sup>4,25</sup>

46

47 Existing studies that have investigated the relationship between hip strength and single  
48 joint/segment kinematics in healthy subjects have failed to account for the confounding influence  
49 of trunk motion in persons with weak hip musculature.<sup>4,14,19,25</sup> In the frontal plane, subjects with  
50 weak hip abductors often demonstrate increased trunk motion towards the stance limb,<sup>23,25</sup>  
51 resulting in altered moments at the hip and knee.<sup>4,21</sup> In addition, existing studies utilizing mixed  
52 samples of male and female subjects may also have been confounded by sex-specific differences  
53 in kinematics during dynamic tasks.<sup>4,6,16,25,26</sup> Therefore, the effect of hip muscle performance on  
54 peak **kinematics** of the lower limbs, pelvis and trunk in the absence of musculoskeletal  
55 pathology remains unclear.

56  
57 **Analysis of the relative timing, or coordination, of motion occurring between joints**  
58 **or segments may facilitate identification of subtle adaptations in lower limb, pelvis or trunk**  
59 **motion associated with diminished hip muscle performance during sub-maximal tasks.<sup>11,12</sup>**  
60 **Adaptations in patterns of joint or segmental coordination have the potential to alter joint**  
61 **loading during the stance phase of dynamic activities and therefore may also be associated**  
62 **with the development of lower limb pathologies.<sup>5,11,32</sup> Continuous methods of analyzing**  
63 **coordination, such as the vector coding method, quantify patterns of coordination between**  
64 **segments (inter-segmental coordination) or joints (inter-joint coordination) across the time-**  
65 **series of a task.<sup>1,27</sup> These types of coordination analyses may have greater sensitivity to**  
66 **detect subtle kinematic differences between groups of subjects, or between modes of gait**  
67 **with varying mechanical demands, than single joint/segment kinematics.**

68

69           The purpose of this study was to investigate kinematics in healthy women with strong  
70 and weak hip muscle performance during the stance phase of walking at self-selected speed and  
71 rate controlled single-legged hopping. We hypothesized that during both walking and hopping,  
72 women with weak hip musculature would demonstrate greater peak lower limb, trunk and pelvis  
73 **angular motion** in the frontal and transverse planes in addition to **different** patterns of  
74 coordination compared to women with strong musculature.

75

## 76 METHODS

77           All participants provided written informed consent and the **University of Southern**  
78 **California** Institutional Review Board approved the study procedures. Eligible participants were  
79 free from any history of injury or surgery to the lower extremities and spine or other medical  
80 conditions affecting physical activity.

81

82           Isometric hip abductor and extensor strength were tested bilaterally in healthy women  
83 using a dynamometer (Primus RS, BTE Technologies, Hanover, MD). Hip abduction strength  
84 was tested in a side lying position with the test limb in neutral hip alignment and full knee  
85 extension. Hip extension strength was tested in a prone position with 30° and 90° of hip and knee  
86 flexion respectively.<sup>23,30</sup> Participants performed three trials with each leg. Peak torque was  
87 averaged across the three trials and was normalized to participant body mass. **Participants were**  
88 **given three practice trials prior to testing, and consistent verbal encouragement was**  
89 **provided during each trial. This protocol has high test-retest reliability.**<sup>17</sup>

90

91 Participants were stratified to a weak or strong group if the normalized peak torque of  
92 both hip abduction and extension on their dominant limb fell outside of a 95% confidence  
93 interval. This confidence interval was calculated from the distribution of abduction and extension  
94 torque values from a database of the first 30 female participants tested in this study (age  $25.8 \pm$   
95  $1.8$  years, height  $1.68 \pm 0.01$  m, weight  $64.3 \pm 8.2$  kg). Threshold values for the strong group  
96 (SG) were  $2.74$  and  $1.63 \text{ N} \cdot \text{m} \cdot \text{kg}^{-1}$  for extension and abduction respectively. Threshold values for  
97 the weak group (WG) were  $1.35$  and  $0.77 \text{ N} \cdot \text{m} \cdot \text{kg}^{-1}$ . The dominant limb was defined as the  
98 preferred leg for kicking a ball.<sup>23,28</sup> **The hip performance of 150 women was tested in order to**  
99 **find 22 that met the criteria for either the SG or the WG. These women were retained for**  
100 **the second phase of the study, consisting of the complete biomechanical assessment. These**  
101 **data were collected as part of a broader study investigating kinematics and EMG during a**  
102 **number of dynamic activities that included drop jumps and running in addition to walking**  
103 **and hopping. The EMG and kinematic data from the drop jump task have been presented**  
104 **elsewhere.<sup>23</sup> A-priori power analysis was completed for the drop jump task utilizing pilot**  
105 **data for lumbopelvic excursion and indicated that a total sample of 16 participants was**  
106 **required to achieve a power of 80% at an alpha level of 0.05. A conservative recruitment**  
107 **goal of 22 participants was selected to account for attrition.**

108

109 Instrumentation

110 Lower extremity, pelvis and trunk kinematic data were collected using a ten-camera  
111 three-dimensional motion capture system sampling at 250 Hz (Qualisys AB, Gothenburg,  
112 Sweden). Retro-reflective markers were placed on bony landmarks to define the local coordinate  
113 frames of the lower extremities, pelvis and trunk. Motion of the pelvis segment was tracked by

114 markers on the bilateral anterior superior iliac spines, iliac crests and at the L5/S1 interspinous  
115 space. A rigid cluster of markers placed over the spinous process of T3 was used to track the  
116 motion of the trunk, and clusters of markers on the heel counter of the shoe, shanks and lateral  
117 thighs were used to track segmental motion of the lower extremities.

118

119 Experimental tasks

120 For walking gait, participants walked along a walkway at self-selected speed. Average  
121 speed during the walking trials was calculated from the time taken to pass between two  
122 photoelectric triggers. **For the hopping task, participants performed consecutive hops on a**  
123 **46cm by 51cm force plate (AMTI OR-6, Watertown, MA, sampling rate 1500Hz) in time**  
124 **with a metronome.** Hops were performed at a rate of 100 hops per minute. This hopping rate is  
125 slower than typical self-selected hopping rate, and induces greater demand on the knee than self-  
126 selected hopping.<sup>29</sup> Participants were required to land with the support foot fully within the force  
127 plate for at least 20 consecutive hops. All hops were performed on the participant's dominant leg  
128 and the arms were crossed over the chest for the duration of the trial.

129

130

131 Data processing

132 Marker coordinates and force plate data were processed using Visual 3D™ (C-Motion  
133 Inc., MD). For walking, stance phase initiation and termination on the dominant leg were  
134 identified using the heel marker trajectories. For hopping, support phase initiation and  
135 termination were identified as the moment the vertical ground reaction force exceeded or  
136 dropped below 20 N respectively. A model consisting of the feet, shanks, femurs, pelvis and



137 trunk was constructed. Motion of the lower extremity segments was referenced to the proximal  
138 segment. Motion of the trunk and pelvis segments was referenced to the global coordinate frame  
139 and was normalized to a static calibration trial to account for individual postural alignment.<sup>23</sup>  
140 Peak **angles** of the knee and hip joints and the pelvis and trunk segments in the frontal and  
141 transverse planes were calculated for ten stance phases on the dominant leg for walking and for  
142 the first ten hops for hopping and were averaged across the repeated trials for each subject. **The**  
143 **first ten hops were selected in order to maximize the consistency of the task performance.**  
144 Coordination between lower extremity joints and between the trunk and pelvis segments was  
145 quantified using the vector coding technique.<sup>5,10,20</sup> **Vector coding is based on methods**  
146 **originally described by Sparrow et al.<sup>31</sup> to quantify coordination behavior using angle-**  
147 **angle plots. Coordination between two segments or joints is calculated as the angle of the**  
148 **vector between successive points on the angle-angle plot relative to the right horizontal.**  
149 **This provides an angle, called the coupling angle, between 0 and 360 degrees for each**  
150 **successive interval on the time series. The pattern of coordination for each time interval**  
151 **across the time series can then be defined as in-phase (both segments/joints moving in the**  
152 **same direction at the same time,); anti-phase (both segments/joints moving in the opposite**  
153 **direction at the same time); proximal phase (motion occurring primarily in the proximal**  
154 **joint/segment); distal phase (motion occurring primarily in the distal joint/segment) using**  
155 **45° bin widths (Figure 1a).<sup>1,5</sup> In-phase coordination is represented by coupling angles**  
156 **between 22.5 – 67.5° and 202.5 – 247.5°. Anti-phase coordination is represented by**  
157 **coupling angles between 112.5 – 157.5° and 292.5 – 337.5°. Proximal phase coordination is**  
158 **represented by coupling angles between 157.5 – 202.5° and 337.5 – 360°. Distal phase**  
159 **coordination is represented by coupling angles between 67.5 – 112.5° and 247.5 to 292.5°.<sup>5</sup>**

160 The vector coding technique was utilized in this study as, unlike other continuous methods  
161 of coordination analysis such as continuous relative phase, it does not require amplitude  
162 normalization of kinematic data and therefore can be more easily interpreted relative to  
163 the original kinematics, and is appropriate for both discrete and oscillatory motor tasks.<sup>22</sup>  
164 For both walking and hopping, coordination was quantified between the following  
165 joint/segment pairs: Coupling 1: Hip/knee motion in the frontal plane (positive values =  
166 abduction); Coupling 2: Hip/knee motion in the transverse plane (positive values = rotation  
167 ipsilateral to the stance limb); Coupling 3: Pelvis/trunk motion in the frontal plane  
168 (positive values = tilt towards the side of the stance limb); Coupling 4: Pelvis/trunk motion  
169 in the transverse plane (positive values = rotation towards the side of the stance limb)  
170 (Figure 1b). The amount of each coordination pattern utilized during walking and hopping  
171 for each coupling segment/joint pair was quantified as a percentage of the total  
172 coordination. This indicates the amount of each movement cycle that was spent in each of  
173 the four coordination patterns.

174

175 Statistics

176 Individual two-way repeated measures ANOVA were used to examine the main effects of  
177 group (between subjects factor; SG, WG) and the interaction effects of group by task (within  
178 subjects factor; walk, hop) on the dependent variables. Post-hoc comparisons on significant  
179 group main effects were made using t-tests for independent samples with a Bonferroni correction  
180 for multiple comparisons, with statistical significance set at  $p \leq 0.05$ . Effect sizes for pairwise  
181 comparisons were calculated using Cohen's d (PASW Statistics 18, IBM Corp., Armonk, NY).

182

## 183 RESULTS

184           There was no significant difference in age, height or weight between the groups (Table  
185 1). Hip abductor and extensor strength was significantly greater in the SG than in the WG on  
186 both the dominant and the non-dominant limb (Table 1). Kinematic data from three participants  
187 were excluded due to technical issues leaving a total of 19 subjects (SG n = 10, WG n = 9).  
188 Mean (SD) self-selected walking speed for the entire sample was 1.32 (0.18) m·s<sup>-1</sup> and was not  
189 significantly different between groups (p = 0.49).

190

## 191 Single joint/segment kinematics

192           The only significant main effect of group for peak single-joint/segment kinematics was in  
193 frontal plane trunk motion (F = 13.19, p = 0.002). Post-hoc analyses indicated that there was no  
194 significant difference between groups during walking (WG = 2.5 (1.6)°, SG = 1.3 (1.5)°,  
195 adjusted p = 0.234). However, the WG had significantly greater trunk lateral bend towards the  
196 stance limb during the hopping task than the SG (WG = 7.9 (2.1)°, SG = 4.1 (2.0)°; adjusted p =  
197 0.002, effect size d = 1.88). In addition, the WG demonstrated a significantly greater change in  
198 peak trunk motion during hopping compared with walking than the strong group (ordinal  
199 interaction, F = 8.657, p = 0.009). A **disordinal** group by task interaction was also evident for  
200 ipsilateral pelvic tilt. The WG demonstrated less ipsilateral pelvic tilt than the SG during walking  
201 and a greater amount of ipsilateral tilt during hopping (Walking, WG = 2.0 (1.3)°, SG = 2.5  
202 (1.1)°; Hopping, WG = 11.0 (2.1)°, SG = 9.0 (2.0)°, F = 8.079, p = 0.011).

203

## 204 Coordination

205           There was a significant effect of group for hip/knee transverse plane coordination  
206 (coupling 2; anti-phase ( $F = 7.376$ ,  $p = 0.015$ ), in-phase ( $F = 8.22$ ,  $p = 0.011$ ), hip phase ( $F =$   
207  $10.311$   $p = 0.005$ )). During walking, the WG utilized less in-phase coordination between the hip  
208 and knee in the transverse plane (WG = 22.4 (6.4)%, SG = 29.4 (2.7)%, adjusted  $p = 0.036$ ,  $d =$   
209  $1.45$ ) and greater primarily hip motion than the SG (WG = 23.2 (6.1)%, SG = 15.7 (2.0)%,  
210 adjusted  $p = 0.036$ ,  $d = 1.70$ ) (Figure 2). The WG had significantly greater anti-phase  
211 coordination between the hip and knee in the transverse plane during hopping than the SG (WG  
212 = 30.2 (7.1)%; SG = 17.0 (10.4)%, adjusted  $p = 0.03$ ,  $d = 1.47$ ) (Figures 2 and 3). There was also  
213 a significant effect of group for coordination between the pelvis and the trunk in the frontal plane  
214 (coupling 3; in-phase coordination,  $F = 5.44$ ,  $p = 0.032$ ). The WG tended to utilize more in-phase  
215 coordination between the trunk and the pelvis in the frontal plane than the SG during hopping  
216 (WG = 10.0 (5.3)%, SG = 5.4 (1.8)%, adjusted  $p = 0.066$ ,  $d = 1.19$ ). In addition, the WG  
217 demonstrated a smaller change in the amount of in-phase coordination utilized in the transverse  
218 plane between the pelvis and the thorax between walking and hopping than the SG (ordinal  
219 interaction,  $p = 0.026$ ).

220

## 221 DISCUSSION

222           **This study indicates that in healthy young women, hip muscle performance does not**  
223 **affect peak kinematics of the hip or knee during walking or rate-controlled hopping.**  
224 **However, women with strong or weak hip musculature do demonstrate significantly**  
225 **different patterns of coordination between the hip and knee and the trunk and pelvis.**

226

227 By demonstrating little relationship between isometric strength and peak hip and knee  
228 joint **kinematics**, this present research **supports** the findings of other studies investigating  
229 subjects at the extremes of typical hip muscle performance.<sup>4,15,19</sup> Some previous studies using  
230 healthy subjects have demonstrated changes in lower extremity kinematics after the hip  
231 musculature is fatigued.<sup>3,9</sup> **However, the kinematics observed after fatigue in these studies**  
232 **may in part represent a short-term response to a novel, localized loss of muscle**  
233 **performance rather than the purely habitual movement strategy for that subject.**

234  
235 In this study the weaker participants did demonstrate increased frontal plane trunk  
236 **motion** in the direction of the stance limb during hopping. It is possible that if this trunk lateral  
237 bend had been constrained during hopping a greater group difference in peak lower limb  
238 **kinematics** would have emerged. The fact that this strategy was not evident during walking gait  
239 is reflective of the higher mechanical demands of the rate-controlled slow hopping task.

240  
241 The quantification of coordination patterns in this study permitted greater insight into  
242 differences between groups than the single joint/segment peak kinematics. During weight-  
243 bearing tasks, the coordination between joints or segments is in part constrained by the  
244 morphology of the joints and associated soft tissues.<sup>7,32</sup> However, the kinematics of multiple  
245 segments or joints are also coordinated as part of a motor control strategy or synergy.<sup>18</sup> Despite  
246 the lack of group differences in peak hip or knee **kinematics**, the coordination analyses indicated  
247 differences in patterns of lower extremity coordination between the SG and the WG. The weak  
248 subjects demonstrated significantly greater anti-phase coordination between the hip and knee in  
249 the transverse plane compared with the SG during hopping. The anti-phase coordination pattern,

250 consisting of simultaneous hip internal rotation and knee external rotation, occurred during both  
251 the deceleration and acceleration components of the hop stance phase in the WG. This pattern of  
252 coordination may result in increased patellofemoral joint stress<sup>25,34</sup> and therefore suggests a  
253 mechanism for the development of PFP in subjects with weak hip musculature.

254

255 Interestingly, in this present study there were also differences between the groups in transverse  
256 plane lower extremity coordination during the less mechanically demanding walking task. The  
257 WG used less in-phase hip and knee rotation than the SG, and also spent a greater amount of  
258 time utilizing primarily hip motion (hip phase) than the SG. These differences were driven  
259 primarily by a pattern of relative external rotation of the hip during mid-stance in the WG that  
260 did not occur in the SG. Powers et al.,<sup>24</sup> also demonstrated decreased hip internal rotation during  
261 walking in subjects with PFP compared with controls. They suggested that this may be a  
262 compensatory mechanism to minimize the lateral forces on the patella. **This present study**  
263 **indicates that this finding may also be related to hip muscle performance.**

264

265

#### 266 **Limitations**

267 **This study utilized a relatively small sample size. However, the large effect sizes for group**  
268 **differences in a number of variables suggest that the study was adequately powered. As our**  
269 **study aimed to investigate women with contrasting hip muscle performance, the**  
270 **generalizability of these results to individuals with less extreme muscle performance may**  
271 **be limited. The strength thresholds for inclusion in the study were calculated a priori after**  
272 **testing only an initial 30 participants. However, utilizing strength data calculated from all**

273 **150 study participants would have resulted in a smaller sample due to larger standard**  
274 **deviations in the entire cohort data. Further, due to the time required to screen all 150**  
275 **subjects, retaining subjects for biomechanical testing might have been difficult. It should**  
276 **also be noted that as the criterion for stratification to the SG and WG in this study was the**  
277 **performance of the hip extensors and abductors, it is possible that differing performance in**  
278 **other lower extremity or trunk musculature may have contributed to the group differences.**  
279 **In particular, the adaptations in transverse plane coordination patterns may also be**  
280 **associated with poor hip rotator performance. In addition, this study did not control for**  
281 **habitual physical or sporting activity in the participants and did not investigate the non-**  
282 **dominant limb.**

283

284 This study helps to clarify the relationship between hip muscle performance and lower  
285 limb, pelvis, and trunk kinematics in young women. In the absence of the confounding  
286 influences of pain or pathology, hip weakness is not associated with significant differences in  
287 peak **kinematics** in the lower limbs, pelvis, or trunk during walking. Compensatory frontal plane  
288 trunk motion in weak subjects may reduce the effect of weak hip musculature on lower limb  
289 kinematics during hopping. The **significantly different** lower limb and pelvis/trunk coordination  
290 patterns during both walking and hopping in the weak participants suggest subtle adaptations to  
291 diminished hip performance even in young, healthy women during sub-maximal motor tasks.  
292 Further research is needed to establish the relationships between these coordination adaptations  
293 and joint loading or the development of musculoskeletal pathology.

294

295

296

297 Key Points:

298 Findings: Healthy women with poor hip muscle performance have **different** coordination, but  
299 not **different** peak lower limb **kinematics** during walking and hopping compared with women  
300 with strong hip muscle performance.

301 Implications: The differences in **kinematics** previously observed in patients with  
302 musculoskeletal disorders may be more related to pain or pathology than hip muscle weakness.  
303 However, the adaptations in trunk motion and in patterns of lower limb and trunk coordination  
304 evident in this study may contribute to the development of musculoskeletal disorders.

305 Caution: This study only investigated young, healthy women performing sub-maximal tasks. In  
306 addition, the interpretation of the data relies on a premise that functional tasks require a common  
307 pattern of coordination.

308



## REFERENCES

1. Armour Smith J, Siemienski A, Popovich JM Jr, Kulig K. Intra-task variability of trunk coordination during a rate-controlled bipedal dance jump. *J Sports Sci.* 2012;30(2):139–147.
2. Arokoski MH, Arokoski JP, Haara M, et al. Hip muscle strength and muscle cross sectional area in men with and without hip osteoarthritis. *J Rheumatol.* 2002;29(10):2185–2195.
3. Carcia C, Eggen J, Shultz S. Hip-abductor fatigue, frontal-plane landing angle, and excursion during a drop jump. *J Sport Rehabil.* 2005;14(4):321–331.
4. Cashman GE. The effect of weak hip abductors or external rotators on knee valgus kinematics in healthy subjects: a systematic review. *J Sport Rehabil.* 2012;12:273–284.
5. Chang R, van Emmerik REA, Hamill J. Quantifying rearfoot–forefoot coordination in human walking. *J Biomech.* 2008;41(14):3101–3105.
6. Chumanov ES, Wall-Scheffler C, Heiderscheit BC. Gender differences in walking and running on level and inclined surfaces. *Clin Biomech.* 2008;23(10):1260–1268.
7. DeLeo AT, Dierks TA, Ferber R, Davis IS. Lower extremity joint coupling during running: a current update. *Clin Biomech.* 2004;19(10):983–991.
8. Eng JJ, Winter DA. Kinetic analysis of the lower limbs during walking: what information can be gained from a three-dimensional model? *J Biomech.* 1995;28(6):753–758.
9. Geiser CF, O’Connor KM, Earl JE. Effects of Isolated Hip Abductor Fatigue on Frontal Plane Knee Mechanics. *Med Sci Sports Exerc.* 2010;42(3):535–545.
10. Hamill J, Haddad J, McDermott WJ. Issues in quantifying variability from a dynamical systems perspective. *J Appl Biomech.* 2000;16:407–418.
11. Hamill J, van Emmerik REA, Heiderscheit BC, Li L. A dynamical systems approach to lower extremity running injuries. *Clin Biomech.* 1999;14(5):297–308.
12. Heiderscheit B, Hamill J, van Emmerik REA. Variability of stride characteristics and joint coordination among individuals with unilateral patellofemoral pain. *J Appl Biomech.* 2002;18:110–121.
13. Heiderscheit BC. Lower Extremity Injuries: Is It Just About Hip Strength? *J Orthop Sports Phys Ther.* 2010;40(2): 39–41.
14. Hollman JH, Hohl JM, Kraft JL, Strauss JD, Traver KJ. Effects of hip extensor fatigue on lower extremity kinematics during a jump-landing task in women: A controlled laboratory study. *Clin Biomech.* 2012;27(9):903–909.

15. Homan KJ, Norcross MF, Goerger BM, Prentice WE, Blackburn JT. The influence of hip strength on gluteal activity and lower extremity kinematics. *J Electromyogr Kinesiol.* 2012.
16. Jacobs CA, Uhl TL, Mattacola CG, Shapiro R, Rayens WS. Hip abductor function and lower extremity landing kinematics: sex differences. *J Athl Train.* 2007;42(1):76–83.
17. Kulig K, Popovich JM, Noceti-Dewit LM, Reischl SF, Kim D. Women with posterior tibial tendon dysfunction have diminished ankle and hip muscle performance. *J Orthop Sports Phys Ther.* 2011;41(9):687–694.
18. Latash ML, Anson JG. Synergies in health and disease: relations to adaptive changes in motor coordination. *Phys Ther.* 2006;86(8):1151–1160.
19. Lawrence RK III, Kernozek TW, Miller EJ, Torry MR, Reuteman P. Influences of hip external rotation strength on knee mechanics during single-leg drop landings in females. *Clin Biomech.* 2008;23(6):806–813.
20. Miller RH, Chang R, Baird JL, van Emmerik REA, Hamill J. Variability in kinematic coupling assessed by vector coding and continuous relative phase. *J Biomech.* 2010;43(13):2554–2560.
21. Mündermann A, Asay JL, Mündermann L, Andriacchi TP. Implications of increased medio-lateral trunk sway for ambulatory mechanics. *J Biomech.* 2008;41(1):165–170.
22. Peters BT, Haddad JM, Heiderscheit BC, van Emmerik REA, Hamill J. Limitations in the use and interpretation of continuous relative phase. *J Biomech.* 2003;36(2):271–274.
23. Popovich JM, Kulig K. Lumbopelvic landing kinematics and EMG in women with contrasting hip strength. *Med Sci Sports Exerc.* 2012;44(1):146–153.
24. Powers CM, Chen P-Y, Reischl SF, Perry J. Comparison of foot pronation and lower extremity rotation in persons with and without patellofemoral pain. *Foot Ankle Int.* 2002;23(7):634–640.
25. Powers CM. The influence of abnormal hip mechanics on knee injury: a biomechanical perspective. *J Orthop Sports Phys Ther.* 2010;2:42–51.
26. Schache A, Blanch P, Rath D, Wrigley T, Bennell K. Differences between the sexes in the three-dimensional angular rotations of the lumbo-pelvic-hip complex during treadmill running. *J Sports Sci.* 2003;21(2):105–118.
27. Seay JF, van Emmerik REA, Hamill J. Influence of Low Back Pain Status on Pelvis-Trunk Coordination During Walking and Running. *Spine.* 2011;36(16):E1070–E1079.
28. Snyder KR, Earl JE, O'Connor KM, Ebersole KT. Resistance training is accompanied by increases in hip strength and changes in lower extremity biomechanics during running. *Clin Biomech.* 2009;24(1):26–34.

29. Souza RB, Arya S, Pollard CD, Salem G, Kulig K. Patellar tendinopathy alters the distribution of lower extremity net joint moments during hopping. *J Appl Biomech*. 2010;26(3):249–255.
30. Souza RB, Powers CM. Differences in hip kinematics, muscle strength, and muscle activation between subjects with and without patellofemoral pain. *J Orthop Sports Phys Ther*. 2009;39(1):12–19.
31. Sparrow W, Donovan E, van Emmerik REA, Barry EB. Using relative motion plots to measure changes in intra-limb and inter-limb coordination. *J Mot Behav*. 1987; 19(1):115-129.
32. Tiberio D. The effect of excessive subtalar joint pronation on patellofemoral mechanics: a theoretical model. *J Orthop Sports Phys Ther*. 1987;9(4):160–165.
33. Watelain E, Dujardin F, Babier F, Dubois D, Allard P. Pelvic and lower limb compensatory actions of subjects in an early stage of hip osteoarthritis. *Arch Phys Med Rehabil*. 2001;82(12):1705–1711.
34. Willson JD, Davis IS. Lower extremity mechanics of females with and without patellofemoral pain across activities with progressively greater task demands. *Clin Biomech*. 2008;23(2):203–211.


**TABLE 1.** Subject demographics and hip strength. (WG = weak group, SG = strong group).

	<b>WG (n=11)</b>	<b>SG (n=11)</b>	<b>p</b>
Age (years)	23.91 (3.96)	23.91 (3.21)	1.00
Weight (kg)	60.43 (6.18)	59.44 (6.82)	0.70
Height (m)	1.66 (0.04)	1.66 (0.06)	0.77
Dominant hip abduction (N·m·kg <sup>-1</sup> )	0.67 (0.08)	1.81 (0.13)	<0.001
Dominant hip extension (N·m·kg <sup>-1</sup> )	1.14 (0.19)	2.96 (0.13)	<0.001
Non-dominant hip abduction (N·m·kg <sup>-1</sup> )	0.75 (0.13)	1.84 (0.20)	<0.001
Non-dominant hip extension (N·m·kg <sup>-1</sup> )	1.18 (0.22)	2.93 (0.34)	<0.001


All values mean (± SD)




**FIGURE 1(a).** Exemplar angle-angle plot and detail from plot demonstrating calculation of coupling angle and categorization of coupling angles for a single coupling pair into coordination patterns using 45° bin widths. In-phase coordination, coupling angles between 22.5 – 67.5° and 202.5 – 247.5°; anti-phase coordination, coupling angles 112.5 – 157.5° and 292.5 – 337.5°; proximal phase coordination, coupling angles 157.5 – 202.5° and 337.5 – 360°; distal phase coordination, coupling angles 67.5 – 112.5° and 247.5 to 292.5°. **FIGURE 1(b).** Coupling joint/segment pairs in the frontal (1 & 3) and transverse (2 & 4) planes. Direction of arrows indicates direction of motion with positive values.



**FIGURE 2.** Coordination pattern between the hip and knee in the transverse plane during stance phase of hopping and walking; weak group (WG, n = 9) and strong group (SG, n = 10), each coordination pattern expressed as a % of total stance phase. \* = significant difference between groups.



**FIGURE 3:** Angle-angle plots between the hip and knee in the transverse plane during hopping; weak group (WG, n = 9) and strong group (SG, n = 10). IC = initial contact, TO = toe-off, arrows indicate direction of motion.



	WG (n=11)	SG (n=11)	p
Age (years)	23.91 (3.96)	23.91 (3.21)	1.00
Weight (kg)	60.43 (6.18)	59.44 (6.82)	0.70
Height (m)	1.66 (0.04)	1.66 (0.06)	0.77
Dominant hip abduction (N·m·kg <sup>-1</sup> )	0.67 (0.08)	1.81 (0.13)	<0.001
Dominant hip extension (N·m·kg <sup>-1</sup> )	1.14 (0.19)	2.96 (0.13)	<0.001
Non-dominant hip abduction (N·m·kg <sup>-1</sup> )	0.75 (0.13)	1.84 (0.20)	<0.001
Non-dominant hip extension (N·m·kg <sup>-1</sup> )	1.18 (0.22)	2.93 (0.34)	<0.001

All values mean (± SD)

TABLE 1. Subject demographics and hip strength. (WG = weak group, SG = strong group).  
1083x804mm (72 x 72 DPI)






FIGURE 1(a). Exemplar angle-angle plot and detail from plot demonstrating calculation of coupling angle and categorization of coupling angles for a single coupling pair into coordination patterns using 45° bin widths. In-phase coordination, coupling angles between 22.5 – 67.5° and 202.5 – 247.5°; anti-phase coordination, coupling angles 112.5 – 157.5° and 292.5 – 337.5°; proximal phase coordination, coupling angles 157.5 – 202.5° and 337.5 – 360°; distal phase coordination, coupling angles 67.5 – 112.5° and 247.5 to 292.5°. FIGURE 1(b). Coupling joint/segment pairs in the frontal (1 & 3) and transverse (2 & 4) planes. Direction of arrows indicates direction of motion with positive values.

1083x804mm (72 x 72 DPI)






FIGURE 2. Coordination pattern between the hip and knee in the transverse plane during stance phase of hopping and walking; weak group (WG, n = 9) and strong group (SG, n = 10), each coordination pattern expressed as a % of total stance phase. \* = significant difference between groups.  
1083x804mm (72 x 72 DPI)




FIGURE 3: Angle-angle plots between the hip and knee in the transverse plane during hopping; weak group (WG, n = 9) and strong group (SG, n = 10). IC = initial contact, TO = toe-off, arrows indicate direction of motion.

1083x804mm (72 x 72 DPI)