

Chapman University Chapman University Digital Commons

Pharmacy Faculty Articles and Research

School of Pharmacy


2008

Drug Selection and Adverse Drug Reactions: Balancing Outcomes and Costs

Enrique Seoane-Vazquez

Chapman University, seoanevazquez@chapman.edu

Follow this and additional works at: http://digitalcommons.chapman.edu/pharmacy_articles

 Part of the [Health and Medical Administration Commons](#), and the [Pharmacy Administration, Policy and Regulation Commons](#)

Recommended Citation

Seoane-Vazquez E. Drug selection and adverse drug reactions: Balancing outcomes and costs. *Electronic Journal of Biomedicine*. 2008;2:3-5.

This Editorial is brought to you for free and open access by the School of Pharmacy at Chapman University Digital Commons. It has been accepted for inclusion in Pharmacy Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Drug Selection and Adverse Drug Reactions: Balancing Outcomes and Costs

Comments

This editorial was originally published in *Electronic Journal of Biomedicine*, volume 2, in 2008.

Copyright

The author

Editorial:

DRUG SELECTION AND ADVERSE DRUG REACTIONS: BALANCING OUTCOMES AND COSTS

Enrique Seoane-Vazquez, Ph.D.

Assistant Professor, College of Pharmacy and College of Public Health, The Ohio State University, Columbus, Ohio, USA

[pharmacoeconomics @ osu.edu](mailto:pharmacoeconomics@osu.edu)

[Version española](#)

Regulatory agencies approve drugs for marketing when the ratio of risk to benefit of the drugs is positive, i.e. the potential benefit of using the drug justifies the risk of drug related adverse reactions (ADR). At the point of approval of a drug there is not adequate information about all possible negative outcomes arising from its use in clinical practice. The information is even more limited for patient subgroups, diseases and treatment combinations that were not evaluated in the clinical trials required for approval. Therefore, performing post-marketing pharmacoepidemiological studies is necessary for a better understanding of drug use in usual clinical practice conditions and the effect of the drugs on the general patient population.

The incidence of ADRs has been considered high. A study pooling the incidence of ADR related hospital admissions of multiple studies derived a median of 4.1% ADR related hospital admissions in the USA¹. Higher ADR related hospital admissions were reported among specific subpopulations at risk, e.g. cardiovascular patients¹. Another study found that 6.5% of all hospitalizations in the UK were related to ADR². Severe ADRs may lead to hospital intensive care unit admission³. The intensive care units also have a higher rate of ADRs than other hospital units⁴. A high incidence of ADRs was also identified in the ambulatory setting with an estimated 180.000 life threatening or fatal adverse drug effects per year in the USA⁵.

[The article by Alfonso Orta et al⁶](#) is a pharmacoepidemiologic study that presents a series of case reports of ADRs identified in the intensive care unit of a Cuban hospital. The study identifies streptokinase, a thrombolytic drug used for dissolving blood clots, as the drug involved in 61% of the ADRs reported. Problems associated with the safety of streptokinase have been known to be caused by its potential antigenicity, short half-life,

and lack of fibrin specificity^{7,8}. These problems motivated a drastic reduction of streptokinase use in the USA.

Streptokinase is available as a generic drug and its therapeutic alternatives have higher cost⁹. Drug cost is the main reason why streptokinase continues to be used around the world in spite of its potential risks, especially in health care systems with fewer resources available for health care.

Rational drug utilization requires consideration of drug cost as a factor in the therapy selection decision-making process. But other costs should also be considered in the process. Health care costs such as physician visits or hospital care, and not health care costs such as informal caregiver time and patient time for treatment should also be considered. Consideration of health care costs is especially important when a drug of lower cost may generate a higher rate of ADRs that could require expensive hospital care or intensive care.

While ADRs generate negative health outcomes, costs are also associated with the prevention, identification and treatment of these reactions. Hospital care represents an important part of the costs of ADRs due to an increase in admissions and in the length of stay^{1, 10, 11}.

Rational drug utilization requires an evaluation of the risk/benefit ratio and the cost of pharmaceuticals. Pharmacoepidemiologic studies of ADRs allow for the identification of potential drugs and therapeutic classes where prevention efforts could best improve clinical outcomes and/or reduce the costs of those events. The fact that the majority of the ADR identified by [Alfonso Orta et al⁶](#) are related to the use of streptokinase suggests that prevention efforts in Cuba should first focus on this drug. Additionally, the cost-benefit of streptokinase from the perspective of the Cuban health care system should be assessed.

Several strategies have been proposed to reduce the incidence of ADRs^{1,10}. These strategies include educational activities, risk group identification programs, implementation of clinical guidelines, clinical and laboratory monitoring of ADR, and drug safety monitoring. Special mention should be given to the need of more pharmacoepidemiologic research, such as the one conducted by Alfonso Orta et al⁶, and more organized pharmacoepidemiologic studies (e.g. case-control studies).

REFERENCES

1. Muehlberger N, Schneeweiss S, Hasford J. Adverse drug reaction monitoring-cost and benefit considerations. Part I: frequency of adverse drug reactions causing hospital admissions. *Pharmacoepid Drug Safety* 1997; 6(S3):S71-S77.

2. Pirmohamed M, James S, Meakin S, Green C, et al. Adverse drug reactions as cause of admission to hospital: prospective analysis of 18 820 patients. *BMJ* 2004;329:15-19.
3. Grenouillet-Delacre M, Verdoux H, Moore N, Haramburu F, Miremont-Salamé G, et al. Life-threatening adverse drug reactions at admission to medical intensive care: a prospective study in a teaching hospital. *Intensive Care Med* 2007;33(12):2150-7.
4. Kane-Gill SL, Devlin JW. Adverse Drug Event Reporting in Intensive Care Units: A Survey of Current Practices. *Ann Pharmacother* 2006;40(7):1267-1273.
5. Gurwitz JH, Field TS, Harrold LR, Rothschild J, et al. Incidence and preventability of adverse drug events among older persons in the ambulatory setting. *JAMA* 2003;289:1107-16.
6. Alfonso Orta I, Garcia Arnao O, Triolet Galvez A, Gómez C, Ruiz K. Reacciones adversas a medicamentos en una unidad de terapia intensiva. *Electron J Biomed* 2008;2.
7. Sura AC, Kelemen MD. Early management of ST-segment elevation myocardial infarction. *Cardiol Clin* 2006;1(24):37-51.
8. Khan IA, Gowda RM Clinical perspectives and therapeutics of thrombolysis. *Int J Cardiol* 2003;91,115-127
9. Hernández L, Marrero MA. Streptokinase: about a thrombolytic patented in Cuba. *Biotechnologia Aplicada*, 2005;22:191-9. Available at: <http://elfosscientiae.cigb.edu.cu/PDFs/BA/2005/22/3/BA002203RV191-198.pdf>. Accessed on May 15, 2008.
10. Gautier S, Bachelet H, Bordet R, Caron J. The cost of adverse drug reactions. *Expert Opin Pharmacother* 2003;4 (3):319-26.
11. Bordet R, Gautier S, Le Louet H, Dupuis B, Caron J. Analysis of the direct cost of adverse drug reactions in hospitalised patients. *Eur J Clin Pharmacol* 2001;56(12):935-41.