

Chapman University Chapman University Digital Commons

Pharmacy Faculty Articles and Research

School of Pharmacy

2010

Localization of Angiotensin Converting Enzyme in Rabbit Cornea and Its Role in Controlling Corneal Angiogenesis *in vivo*

Ajay Sharma

Chapman University, sharma@chapman.edu

Daniel I. Bettis

Harry S. Truman Memorial Veterans' Hospital

John W. Cowden

University of Missouri-Columbia

Rajiv R. Mohan

Harry S. Truman Memorial Veterans' Hospital

Follow this and additional works at: https://digitalcommons.chapman.edu/pharmacy_articles

 Part of the [Animal Experimentation and Research Commons](#), [Animals Commons](#), [Musculoskeletal, Neural, and Ocular Physiology Commons](#), [Ophthalmology Commons](#), and the [Other Pharmacy and Pharmaceutical Sciences Commons](#)

Recommended Citation

Sharma A, Bettis DI, Cowden JW, Mohan RR. Localization of angiotensin converting enzyme in rabbit cornea and its role in controlling corneal angiogenesis *in vivo*. *Mol Vis*. 2010;16:720-728.

This Article is brought to you for free and open access by the School of Pharmacy at Chapman University Digital Commons. It has been accepted for inclusion in Pharmacy Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Localization of Angiotensin Converting Enzyme in Rabbit Cornea and Its Role in Controlling Corneal Angiogenesis *in vivo*

Comments

This article was originally published in *Molecular Vision*, volume 16, in 2010.

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 3.0 License](https://creativecommons.org/licenses/by-nc-nd/3.0/).

Copyright

Molecular Vision

Localization of angiotensin converting enzyme in rabbit cornea and its role in controlling corneal angiogenesis in vivo

Ajay Sharma,^{1,2} Daniel I. Bettis,^{1,2} John W. Cowden,² Rajiv R. Mohan^{1,2,3}

¹Harry S. Truman Memorial Veterans' Hospital, Columbia, MO; ²Mason Eye Institute, School of Medicine, University of Missouri-Columbia, MO; ³Department of Ophthalmology, College of Veterinary Medicine, University of Missouri-Columbia, MO

Purpose: The renin angiotensin system (RAS) has been shown to modulate vascular endothelial growth factor and angiogenesis. In this study we investigated (i) the existence of the RAS components angiotensin converting enzyme (ACE) and angiotensin II receptors (AT₁ and AT₂) in the rabbit cornea using in vitro and ex vivo models and (ii) the effect of enalapril, an ACE inhibitor, to inhibit angiogenesis in rabbit cornea in vivo.

Methods: New Zealand White rabbits were used. Cultured corneal fibroblasts and corneal epithelial cells were used for RNA isolation and cDNA preparation using standard molecular biology techniques. PCR was performed to detect the presence of *ACE*, *AT₁*, and *AT₂* gene expression. A corneal micropocket assay to implant a vascular endothelial growth factor (VEGF) pellet in the rabbit cornea was used to induce corneal angiogenesis. Rabbits of the control group received sterile water, and the treated group received 3 mg/kg enalapril intramuscularly once daily for 14 days starting from day 1 of pellet implantation. The clinical eye examination was performed by slit-lamp biomicroscopy. We monitored the level of corneal angiogenesis in live animals by stereomicroscopy at days 4, 9, and 14 after VEGF pellet implantation. Collagen type IV and lectin immunohistochemistry and fluorescent microscopy were used to measure corneal angiogenesis in tissue sections of control and enalapril-treated corneas of the rabbits. Image J software was used to quantify corneal angiogenesis in the rabbit eye in situ.

Results: Our data demonstrated the presence of *ACE*, *AT₁*, and *AT₂* expression in corneal fibroblasts. Cells of corneal epithelium expressed *AT₁* and *AT₂* but did not show *ACE* expression. Slit-lamp examination did not show any significant difference between the degree of edema or cellular infiltration between the corneas of control and enalapril-treated rabbits. VEGF pellet implantation caused corneal angiogenesis in the eyes of vehicle-treated control rabbits, and the mean area of corneal neovascularization was 1.8, 2.8, and 3.2 mm² on days 4, 9, and 14, respectively. Enalapril treatment caused a notable decrease in corneal neovascularization of 44% (1 mm²), 28% (2.1 mm²), and 31% (2.2 mm²) on the three tested time points, respectively. The immunostaining of corneal tissue sections with collagen type IV and lectin confirmed the presence of blood vessels, with enalapril-treated rabbit corneas showing a lesser degree of blood vessel staining.

Conclusions: Corneal cells show expression of tissue RAS components, such as *ACE*, *AT₁*, and *AT₂*. Treatment with ACE inhibitor enalapril markedly decreased corneal angiogenesis in a rabbit model of VEGF-induced corneal neovascularization, suggesting that ACE inhibitors may represent a novel therapeutic strategy to treat corneal angiogenesis.

Corneal clarity is necessary for normal vision. Corneal insult, such as trauma, chemical injury, infections, or immune disorders, can lead to corneal neovascularization and loss of corneal transparency [1,2]. Clinical management of corneal neovascularization is challenging and preexisting neovascularization in acceptor eye also significantly increases the risk of corneal transplant rejection [3]. A variety of agents, such as corticosteroids [4], cyclosporine [5], indomethacin [6], methotrexate [7], rapamycin [8], low-molecular weight heparin sulfate [9], and thalidomide [10], have been evaluated in animal models of angiogenesis to treat corneal neovascularization but have shown limited success. Bevacizumab, an anti-vascular endothelial growth factor

(VEGF) antibody, has recently been successfully tested in patients of corneal neovascularization [11]. Although bevacizumab is well tolerated, multiple applications may be required for effective treatment, and this can be expensive. Therefore, there is an unmet medical need for clinically effective agents that can inhibit corneal neovascularization.

The renin angiotensin system (RAS) is among the most powerful regulators of blood pressure and plasma volume [12]. The RAS consists of kidney renin (converts circulating plasma protein angiotensinogen into angiotensin I) and angiotensin-converting enzyme (ACE) that converts angiotensin I to angiotensin II (Ang II). Besides the circulating RAS, the presence of a tissue-specific local RAS is also well documented for brain, heart, pancreas, kidney, blood vessels, and gonads [12]. In ocular tissue, multiple studies have tested the presence of the tissue RAS, demonstrating mRNA and protein expression of renin, angiotensinogen, and Ang II receptors in ciliary bodies [13,14], choroid [15], ganglion cells

Correspondence to: Rajiv R. Mohan, Ph.D., Mason Eye Institute, School of Medicine, EC-210, University of Missouri-Columbia, 1 Hospital Drive, Columbia, MO, 65212; Phone: (573) 884-1449; FAX: (573) 814 6551; email: mohanr@health.missouri.edu

[14], pigmented epithelial cells [16,17], and Müller cells of the retina [18]. The presence of angiotensin and ACE enzyme activity have also been detected in the vitreous fluid [19]. Among retina cells, the presence of renin has been shown in pigmented epithelial cells but not in neural retina, whereas angiotensinogen and ACE could be detected in both pigmented epithelial cells and neural retina [16]. These findings suggest that RAS expression is not constant, even in the various cell types of a single tissue such as retina. A few researchers have tested the presence of the RAS in cornea, but these, performed in rodent, dog, and primate eyes, have shown contradictory findings [14,15,17]. Savaskan et al. [14] and Usui et al. [20] reported expression of ACE and Ang II receptors (AT₁) in the primate and mouse cornea, whereas Sheota et al. [15] and Murata et al. [17] did not detect ACE or AT₁ in dog and rat cornea. The detailed investigation of RAS components in different corneal cell types is still missing.

Typically, Ang II is implicated in controlling plasma volume, electrolyte balance, and blood pressure [12]. Recently, Ang II has been ascribed to modulate many more pro-angiogenic activities, including the proliferation and chemotaxis of vascular smooth muscle and endothelial cells and increased transcription of VEGF [21-24]. These reports suggest that Ang II may have a role in promoting angiogenesis. Thus, it is reasonable to speculate that suppression of Ang II can inhibit angiogenesis. Numerous studies have demonstrated that ACE inhibitors (known to block Ang II formation) decrease cellular proliferation, angiogenesis, and VEGF expression in different tumor cell lines [25-27]. Furthermore, the anti-angiogenic effect of ACE inhibitors has been shown in various experimental animal models of cancer. In these studies, ACE inhibitors showed a decrease in tumor growth and VEGF levels [25-30]. In the eye, ACE inhibitors have shown a significant decrease in choroidal [31] and retinal vascularization [32] in rodent models *in vivo*. A recent clinical trial performed with an ACE inhibitor, enalapril, showed 65% reduction in progression of diabetic retinopathy in enalapril-treated patients [33]. We hypothesized that inhibition of RAS activity with ACE inhibitor can prevent angiogenesis in rabbit cornea *in vivo*. Therefore, we evaluated the effect of enalapril on corneal neovascularization using a well-defined VEGF-induced rabbit model of corneal neovascularization. The expression of ACE enzyme and the angiotensin receptors AT₁ and AT₂ in rabbit corneal epithelium and fibroblasts was also investigated.

METHODS

Rabbits and corneal cultures: Female New Zealand White rabbits (Myrtle laboratories Inc., Thompson's Station, TN) weighing 2.5–3.0 kg were used in this study. The study was approved by the Institutional Animal Care and Use Committee (Harry S. Truman Memorial Veterans' Hospital, Columbia, MO). Animals were treated in accordance with the

tenets of the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research. The cells of corneal epithelium were collected by scraping the epithelium with a surgical blade. Primary corneal fibroblasts were generated from the rabbit corneas as previously described [34]. Briefly, cornea was washed with cell culture medium, and epithelium and endothelium were removed by gentle scraping with a scalpel blade. Corneal stroma was cut into small pieces and incubated in a humidified (90%) CO₂ incubator at 37 °C in Dulbecco's modified Eagle's medium, containing 10% fetal bovine serum, to obtain rabbit corneal fibroblasts. Seventy percent confluent cultures of rabbit corneal fibroblasts were used for experiments.

RNA extraction, cDNA synthesis, and PCR: Total RNA from the cells was extracted using the RNeasy kit (Qiagen Inc., Valencia, CA) and was reverse-transcribed to cDNA, following the vendor's instructions (Promega, Madison, WI). Briefly, cells were lysed in 350 µl RLT buffer (Qiagen Inc., Valencia, CA) containing β-mercaptoethanol, followed by equal volume of ethanol addition. The solution was then applied onto RNeasy column and RNA was eluted from the column using 30 µl of RNase free water. The eluted RNA was reverse transcribed to cDNA using oligo(dT) primers, AMV reverse transcriptase (Promega), reverse transcriptase buffer, dNTPs, RNasin. The reverse transcription reaction was carried out at 42 °C for 30 min followed by enzyme inactivation by heating at 90 °C for 2 min. PCR was performed for the detection of ACE, AT₁, and AT₂. A 50-µl reaction mixture containing 3 µl cDNA, 2 µl forward (200 nM) and 2 µl reverse (200 nM) primer, 3.125 mM of deoxynucleotide triphosphates (dNTPs) and Taq polymerase was run one cycle at 95 °C for 3 min, then 40 cycles of 95 °C 30 s, followed by 55 °C 30 s and 55 °C for 60 s, using a thermocycler (Bio-Rad Laboratories, Hercules, CA). For ACE, the forward primer sequence 5'-ACG AGC ACG ACA TCA ACT TCC TCA-3' and reverse primer sequence 5'-AGT AGT TCA TCA TGG CCG AGG CT-3' were used. For AT₁, the forward primer sequence was 5'-AGG ATG ACT GTC CCA AAG CTG GAA-3', and the reverse primer sequence was 5'-ACG TTT CGG ATG ATA GCT GGT-3'. For AT₂, the forward primer sequence was 5'-TGA GAA ATA TGC CCA GTG GTC GGT-3', and the reverse primer sequence was 5'-ATA ATC CAG ATG GGC CTC AAG CCA-3'. β-actin (ACTB) was used as a positive control gene: forward primer (5'-AGG CCA ACC GCG AGA AGA TGA CC-3'), reverse primer (5'-GAA GTC CAG GGC GAC GTA GCA C-3'). cDNA samples were prepared from two different rabbits. Each PCR was repeated at least three times.

Corneal angiogenesis in rabbits: Eight female New Zealand White rabbits (2.5–3.0 kg) were divided in two groups. Animals were anesthetized by intramuscular injection of ketamine hydrochloride (50 mg/kg) and xylazine hydrochloride (10 mg/kg). In addition, topical proparacaine



Figure 1. Representative image showing detection of *ACE*, *AT₁*, and *AT₂* mRNA expression in rabbit corneal fibroblasts with PCR. Appropriate size amplification products for *ACE* (469 bp), *AT₁* (463 bp), and *AT₂* (551 bp) were detected in two independent cDNA samples of corneal fibroblast prepared from different rabbits. -ve con denotes negative controls that contained *ACE*, *AT₁*, or *AT₂* primers but water instead of cDNA. + ve con represents the positive control of β-actin (*ACTB*; 350 bp). L denotes 1 kb plus the DNA ladder.




Figure 2. Representative image showing *AT₁* and *AT₂* mRNA expression in rabbit corneal epithelium detected with PCR. Appropriate size amplification products for *AT₁* (463 bp) and *AT₂* (551 bp) were detected in two independent rabbit corneal epithelium cDNA samples prepared from different rabbits. No signal for *ACE* was detected. -ve con denotes negative controls that contained *ACE*, *AT₁*, or *AT₂* primers but water instead of cDNA. + ve con represents the positive control of β-actin (*ACTB*; 350 bp). L denotes 1 kb plus the DNA ladder.

hydrochloride 0.5% (Alcon, Ft. Worth, TX) was applied to each eye just before surgery. A corneal micropocket assay was performed in the rabbit eyes under general and local anesthesia. Only one eye of each animal was used for the surgical procedure, and the contralateral eye served as a naive control. A wire speculum was positioned in the eye, and a sucralfate-hydrone pellet containing 650 ng of VEGF (PeproTech, Rocky Hill, NJ) was implanted in the cornea micropocket. Rabbits of the control group received sterile water, and the treated group received 3 mg/kg enalapril (Sigma Aldrich Inc., St. Louis, MO) via intramuscular injection once daily for 14 days starting from day 1 of pellet implantation.

Stereomicroscopy and slit-lamp examination: The level of neovascularization in the cornea was monitored with a micrometer-calibrated stereomicroscope (Leica, Wetzlar, Germany) equipped with a digital camera (SpotCam RT KE; Diagnostic Instruments Inc., Sterling Heights, MI). Rabbit eyes were imaged at 4, 9, and 14 days after pellet implantation for quantitative analysis of corneal neovascularization. Slit-lamp biomicroscopy was also performed on naive and VEGF-implanted untreated and enalapril-treated rabbit eyes. Qualitative clinical evaluation data about the redness, edema, and inflammation were recorded.

Tissue collection: Rabbits were humanely euthanized with pentobarbitone (150 mg/kg) under general anesthesia. Corneas were removed with forceps and sharp Westcott scissors, embedded in liquid optimal temperature cutting compound (OCT) compound (Sakura FineTek, Torrance, CA) within a 24 mm×24 mm×5 mm mold (Fisher, Pittsburgh, PA), and snap frozen. The frozen tissue blocks were maintained at -80 °C. Tissue sections (7 μm) were cut with a cryostat (HM 525M; Microm GmbH, Walldorf, Germany). Sections were placed on 25 mm×75 mm×1 mm microscope Superfrost Plus slides (Fisher), and maintained frozen at -80 °C until staining.

Immunostaining for collagen type IV and lectin: Collagen type IV is a major component of the basal lamina of blood vessels. We therefore performed collagen type IV immunohistochemistry to detect the presence of blood vessels. Blood vessel staining was also confirmed with lectin (obtained from *Lycopersicon esculatum*). Briefly, the tissue sections were equilibrated at room temperature to thaw OCT. Tissue sections were washed with 1× HEPES buffer for 15 min, blocked with 5% bovine serum albumin for 30 min, and incubated with 1:50 dilution (in 1× HEPES buffer) of goat polyclonal antibody for collagen type IV (catalog no. sc9302; Santa Cruz Biotechnology Inc., Santa Cruz, CA) for 90 min. Donkey anti-goat Alexa 594 secondary antibodies (catalog no.




Figure 3. Representative broad-beam (A) and narrow-beam (B) slit-lamp biomicroscopy images of VEGF-implanted rabbit corneas. Corneas did not show any cellular infiltrate or edema in the area inferior to VEGF pellet (as shown by arrow). A mild level of edema is noticeable around the area of VEGF pellet. The scale bar denotes 2 mm.

A-11058; Invitrogen, Carlsbad, CA) were used at a dilution (in $1\times$ HEPES buffer) of 1:500 for 1 h. For lectin staining, sections were incubated with a solution of 20 $\mu\text{g/ml}$ Texas red-conjugated tomato lectin (Vector laboratories, Burlingame, CA) for 30 min. The sections were washed three times in HEPES buffer, mounted in vectashield containing 4'-6-diamidino-2-phenylindole (DAPI; Vector laboratories), and viewed and photographed with a Leica fluorescent microscope (Leica DM 4000B; Leica) equipped with a digital camera (SpotCam RT KE).

Quantification of corneal neovascularization: Adobe Photoshop CS2 (Adobe systems, San Jose, CA) and National Institutes of Health Image J 1.38X (NIH, Bethesda, MD) software were used for quantifying corneal neovascularization. The quantification of corneal vasculature in the eye of the live rabbit at different selected days was performed with stereo and slit-lamp microscopy. Images of the eye covering corneal neovascularization were taken with a digital camera fitted in a microscope for quantification analysis.

Statistical analysis: The results were expressed as mean \pm standard error of the mean (SEM). Statistical analysis was performed using two-way analysis of variance (ANOVA) followed by Bonferroni multiple comparisons test. A value of $p < 0.05$ was considered statistically significant.

RESULTS

Detection of ACE, AT₁, and AT₂ in rabbit corneal epithelium and fibroblasts: Figure 1 shows the presence of ACE, AT₁, and AT₂ in rabbit corneal fibroblasts. An appropriate size amplification product of 469 bp was detected for ACE in two independent rabbit corneal fibroblast samples. Amplification products for AT₁ and AT₂, corresponding to the product size of 463 bp and 551 bp, respectively, were also present in the rabbit corneal fibroblast cDNA samples. No PCR product was detected in the negative controls that contained primers for ACE, AT₁, or AT₂ but no cDNA. ACTB was used as a positive

control. The appropriate size amplification product of ACTB at 350 bp was detected in all PCRs (Figure 1).

Figure 2 shows the presence of AT₁ and AT₂ in the epithelial cells of rabbit cornea. Anticipated size PCR products of 463 bp and 551 bp for AT₁ and AT₂, respectively, were detected in the two independent cDNA samples of rabbit epithelium. On the other hand, no amplification product for ACE was detected, suggesting that rabbit epithelium does not express ACE enzyme. Negative controls containing appropriate primers and water instead of cDNA samples were also analyzed and did not show any amplification products. ACTB was used as a positive control and showed a PCR product band of 350 bp (Figure 2).

Slit-lamp examination of rabbit eyes: Figure 3 shows broad beam and narrow beam slit-lamp pictures of the eyes of control and enalapril-treated rabbits. A qualitative comparison between the control and enalapril-treated corneas did not reveal any significant difference in the degree of redness, edema, or cellular infiltration.

Effect of enalapril on corneal angiogenesis: Figure 4 shows stereomicroscopic images depicting the localization and extent of corneal neovascularization in control and enalapril-treated rabbit eyes. VEGF pellet implantation produced corneal neovascularization in both control and enalapril-treated rabbits. The images captured on day 4, 9, and 14 show a time-dependent increase in corneal blood vessels in both the control and enalapril-treated groups. However, a notable decrease in corneal neovascularization can be seen in eyes of rabbits treated with enalapril (Figure 4).

For quantitative comparisons between control and treated animals, we calculated the area of corneal neovascularization using Image J software. Figure 5 shows the mean area of corneal neovascularization in control and enalapril-treated rabbits at the three tested time points. Control corneas showed a mean corneal neovascularization area of 1.8 mm² at day 4. The area of corneal neovascularization noted at day 9 and day 14 was 2.8 and 3.2 mm², respectively, and it was significantly higher compared to day 5 ($p < 0.01$; shown by ψ in Figure 5).




Figure 4. Representative stereomicroscopy images showing VEGF-induced neovascularization in the cornea of control and enalapril-treated rabbit eyes. The rabbit eyes were imaged on day 4, 9, and 14 days after VEGF implantation to monitor corneal angiogenesis in control and enalapril treated rabbits.

Corneas of enalapril-treated rabbits showed a mean corneal neovascularization area of 1 mm² on day 4, which increased significantly ($p < 0.01$; shown by ψ in Figure 5) to 2.1–2.2 mm² on day 9 or day 14. A relative comparison between control and enalapril-treated rabbits revealed a mean decrease in corneal neovascularization of 44%, 28%, and 31% on day 4, 9, and 14, respectively, and the decrease was statistically significant at day 4 and day 14 ($p < 0.01$; shown by the asterisk in Figure 5).

The effect of enalapril on corneal angiogenesis was further confirmed by collagen type IV and lectin staining of the blood vessels in the corneal tissue sections of control and enalapril-treated rabbits. Figure 6 demonstrates the localization and level of collagen type IV-stained blood vessels in corneal tissue sections of control and enalapril-treated rabbits. Collagen type IV and lectin staining (data not shown) revealed fewer blood vessels in corneal sections obtained from enalapril-treated rabbits compared to control corneal sections (Figure 6).

DISCUSSION

The presence of the RAS in a wide variety of nonocular tissues, including brain, heart, pancreas, and gonads, has been reported [12]. In the eyes, renin, angiotensin, and angiotensinogen protein and mRNA expression have been reported in retina [14,16,17], choroid [15], and ciliary bodies [13,14] but were not tested in various cells of the cornea




Figure 5. Digital quantification of VEGF-induced corneal neovascularization in enalapril-treated (white bar) and enalapril-untreated control (black bar) eyes of live rabbit performed on days 4, 9, and 14. ψ denotes a $p < 0.01$ and shows a significant value of corneal neovascularization on day 9 and 14 compared to day 4; * denotes a $p < 0.05$ and shows a significant value of corneal neovascularization in enalapril-treated rabbit compared to enalapril-untreated controls. Bars represent standard error.

[13-17]. Among various cell types of retina, renin expression has been detected in pigmented epithelial cells but not in neural retina, whereas angiotensinogen and ACE expression were demonstrated in both pigmented epithelial cells and neural retinal cells [13]. Similarly, expression of AT_1 has been mainly reported in retinal ganglion and photoreceptor cells but not in pigmented epithelial cells [14,15,17]. These studies suggest that expression of RAS components in different cells of a tissue may vary and may possess unique cellular distributions. Our study demonstrates that both corneal epithelium and fibroblasts express AT_1 and AT_2 , whereas ACE is expressed in corneal fibroblasts but not in corneal epithelium. Our data are in agreement with earlier studies that detected the presence of AT_1 and ACE in whole corneal homogenates of the mouse and human cornea [14,20]. It is also evident from the literature that corneal RAS expression varies by species, as the presence of AT_1 and ACE was not detected in rat [17] and dog [15] cornea, respectively. To the best of our knowledge, this is the first study to report cellular localization of ACE, AT_1 , and AT_2 expression in two different cells of the rabbit cornea.

Transdifferentiation of fibroblast to myofibroblast is thought to be a central event in the pathogenesis of corneal fibrosis. In nonocular tissues, myofibroblasts are shown to express high levels of ACE that can generate large amounts of Ang II [35]. The elevated level of Ang II is reported to modulate expression of collagen, fibronectin, and many other extracellular matrix proteins, suggesting that the RAS expressed locally in fibroblasts may participate in fibrosis and wound healing [36-38]. Elevated levels of angiotensin receptors were reported in skin fibroblasts during wound healing, which further supports this notion [39]. Based on these studies in nonocular tissues, we speculate that the presence of corneal RAS components demonstrated in our study may play a role in corneal wound healing and fibrosis. Future studies will investigate the role of the RAS in corneal wound healing and fibrosis. Besides fibrosis, the corneal RAS may participate in regulation of the inflammatory response, as suggested by a recent study demonstrating a marked increase in ACE, AT_1 , and angiotensinogen gene expression in a mouse corneal model of a nylon suture-induced corneal injury [20].

Ang II is the biologically active end product of the RAS. Accumulating evidence suggests that Ang II promotes endothelial and vascular smooth muscle cell proliferation and chemotaxis besides increasing VEGF expression [21-24]. Further support for the pro-angiogenic effect of Ang II comes from in vivo studies that showed Ang II implantation induces blood vessel formation in rabbit cornea and in the chick chorio-allantoic membrane. These investigations provide strong support about the role of Ang II in angiogenesis [40, 41]. Most importantly, these observations led to a novel interventional strategy to control angiogenesis by blocking




Figure 6. Representative immunohistochemistry image showing collagen type IV staining of blood vessels (red staining) in tissue sections of rabbit corneas collected 14 days after sterile water (A) or enalapril treatment (B). The images show angiogenesis in the central cornea region. Enalapril-treated corneas showed fewer blood vessels. Nuclei are stained blue with DAPI. The scale bar denotes 100 μ m.

Ang II formation. ACE inhibitors are known to block Ang II formation, and numerous studies have demonstrated that ACE inhibitors decrease cellular proliferation, VEGF expression, and inhibit angiogenesis in culture and in animal models of cancer [25-30]. These studies and our data showing expression of the RAS locally in the cornea led us to hypothesize that ACE inhibitors can inhibit corneal neovascularization in rabbit eye in vivo. We tested this hypothesis using a well-defined rabbit model of corneal neovascularization, and our results showed that enalapril treatment significantly prevents VEGF-induced corneal neovascularization in rabbit eye in vivo. A relatively low dose (3 mg/kg) of enalapril was used in our experiments to avoid the potential risk of hypotension [42]. However, it is likely that higher or multiple doses of enalapril may show a greater anti-angiogenic effect in the cornea. The intramuscular administration of enalapril over topical application was selected based on pharmacokinetic information available in the literature.

In conclusion, this study demonstrates the presence of AT₁ and AT₂ in rabbit corneal fibroblasts and epithelial cells and ACE expression in corneal fibroblasts of rabbit cornea. Furthermore, we found a significant decrease in VEGF-induced corneal neovascularization in rabbit in vivo with enalapril, an ACE inhibitor. ACE inhibitors represent a novel therapeutic strategy to treat corneal neovascularization; however, this requires further investigation.

ACKNOWLEDGMENTS

The work was supported by grants RO1EY17294 (R.R.M.) and RO1EY17294S2 (R.R.M.) from the National Eye Institute, National Institutes of Health, Bethesda, MD, and an unrestricted grant to the Mason Eye Institute, University of Missouri-Columbia from Research to Prevent Blindness, New York, NY. Thanks are also due to James Fox (M4), Eric Hansen (M3) and Rodger Wilhite (undergrad) students for their help in animal and immunohistochemistry experiments.

REFERENCES

1. Chang JH, Gabison EE, Kato T, Azar DT. Corneal neovascularization. *Curr Opin Ophthalmol* 2001; 12:242-9. [PMID: 11507336]
2. Azar DT. Corneal angiogenic privilege: angiogenic and antiangiogenic factors in corneal avascularity, vasculogenesis, and wound healing. *Trans Am Ophthalmol Soc* 2006; 104:264-302. [PMID: 17471348]
3. Williams KA, Esterman AJ, Barlett C, Holland H, Hornsby NB, Coster DJ. How effective is penetrating corneal transplantation? Factors influencing long-term outcome in multivariate analysis. *Transplantation* 2006; 81:896-901. [PMID: 16570014]
4. BenEzra D, Griffin BW, Maftzir G, Sharif NA, Clark AF. Topical formulations of novel angiostatic steroids inhibit rabbit corneal neovascularization. *Invest Ophthalmol Vis Sci* 1997; 38:1954-62. [PMID: 9331259]
5. Lipman RM, Epstein RJ, Hendricks RL. Suppression of corneal neovascularization with cyclosporine. *Arch Ophthalmol* 1992; 110:405-7. [PMID: 1543461]
6. Haynes WL, Proia AD, Klintworth GK. Effect of inhibitors of arachidonic acid metabolism on corneal neovascularization in the rat. *Invest Ophthalmol Vis Sci* 1989; 30:1588-93. [PMID: 2473047]
7. Jousseaume AM, Kruse FE, Völcker HE, Kirchhof B. Topical application of methotrexate for inhibition of corneal angiogenesis. *Graefes Arch Clin Exp Ophthalmol* 1999; 237:920-7. [PMID: 10541903]
8. Shi W, Gao H, Xie L, Wang S. Sustained intraocular rapamycin delivery effectively prevents high-risk corneal allograft rejection and neovascularization in rabbits. *Invest Ophthalmol Vis Sci* 2006; 47:3339-44. [PMID: 16877400]
9. Lepri A, Benelli U, Bernardini N, Bianchi F, Lupetti M, Danesi R, Del Tacca M, Nardi M. Effect of low molecular weight heparan sulphate on angiogenesis in the rat cornea after chemical cauterization. *J Ocul Pharmacol* 1994; 10:273-80. [PMID: 8207330]
10. Kruse FE, Jousseaume AM, Rohrschneider K, Becker MD, Völcker HE. Thalidomide inhibits corneal angiogenesis induced by

- vascular endothelial growth factor. Graefes Arch Clin Exp Ophthalmol 1998; 236:461-6. [PMID: 9646092]
11. Dastjerdi MH, Al-Arfaj KM, Nallasamy N, Hamrah P, Jurkunas UV, Pineda R 2nd, Pavan-Langston D, Dana R. Topical bevacizumab in the treatment of corneal neovascularization: results of a prospective, open-label, noncomparative study. Arch Ophthalmol 2009; 127:381-9. [PMID: 19365012]
 12. Vinson GP, Ho MM, Puddefoot JR. The distribution of angiotensin II type 1 receptors, and the tissue renin-angiotensin systems. Mol Med Today 1995; 1:35-9. [PMID: 9415136]
 13. Cullinane AB, Leung PS, Ortego J, Coca-Prados M, Harvey BJ. Renin-angiotensin system expression and secretory function in cultured human ciliary body non-pigmented epithelium. Br J Ophthalmol 2002; 86:676-83. [PMID: 12034692]
 14. Savaskan E, Löffler KU, Meier F, Müller-Spahn F, Flammer J, Meyer P. Immunohistochemical localization of angiotensin-converting enzyme, angiotensin II and AT1 receptor in human ocular tissues. Ophthalmic Res 2004; 36:312-20. [PMID: 15627831]
 15. Shiota N, Saegusa Y, Nishimura K, Miyazaki M. Angiotensin II-generating system in dog and monkey ocular tissues. Clin Exp Pharmacol Physiol 1997; 24:243-8. [PMID: 9131292]
 16. Wagner J, Danser AHJ, Derkx FH, de Jong TV, Paul M, Mullins JJ, Schalekamp MA, Ganten D. Demonstration of renin mRNA, angiotensinogen mRNA, and angiotensin converting enzyme mRNA expression in the human eye: evidence for an intraocular renin-angiotensin system. Br J Ophthalmol 1996; 80:159-63. [PMID: 8814748]
 17. Murata M, Nakagawa M, Takahashi S. Expression and localization of angiotensin II type 1 receptor mRNA in rat ocular tissues. Ophthalmologica 1997; 211:384-6. [PMID: 9380359]
 18. Berka JL, Stubbs AJ, Wang DZ, DiNicolantonio R, Alcorn D, Campbell DJ, Skinner SL. Renin-containing Muller cells of the retina display endocrine features. Invest Ophthalmol Vis Sci 1995; 36:1450-8. [PMID: 7775123]
 19. Danser AH, Derkx FH, Admiraal PJ, Deinum J, de Jong PT, Schalekamp MA. Angiotensin levels in the eye. Invest Ophthalmol Vis Sci 1994; 35:1008-18. [PMID: 8125711]
 20. Usui T, Sugisaki K, Iriyama A, Yokoo S, Yamagami S, Nagai N, Ishida S, Amano S. Inhibition of corneal neovascularization by blocking the angiotensin II type 1 receptor. Invest Ophthalmol Vis Sci 2008; 49:4370-6. [PMID: 18829859]
 21. Chua CC, Hamdy RC, Chua BH. Upregulation of vascular endothelial growth factor by angiotensin II in rat heart endothelial cells. Biochim Biophys Acta 1998; 1401:187-94. [PMID: 9531974]
 22. Lyall F, Morton JJ, Lever AF, Cragoe EJ. Angiotensin II activates Na⁺-H⁺ exchange and stimulates growth in cultured vascular smooth muscle cells. J Hypertens Suppl 1988; 6:S438-41. [PMID: 3241232]
 23. Bell L, Madri JA. Influence of the angiotensin system on endothelial and smooth muscle cell migration. Am J Pathol 1990; 137:7-12. [PMID: 2164777]
 24. Tamarat R, Silvestre JS, Durie M, Levy BI. Angiotensin II angiogenic effect *in vivo* involves vascular endothelial growth factor- and inflammation-related pathways. Lab Invest 2002; 82:747-56. [PMID: 12065685]
 25. Yasumatsu R, Nakashima T, Masuda M, Ito A, Kuratomi Y, Nakagawa T, Komune S. Effects of the angiotensin-I converting enzyme inhibitor perindopril on tumor growth and angiogenesis in head and neck squamous cell carcinoma cells. J Cancer Res Clin Oncol 2004; 130:567-73. [PMID: 15449186]
 26. Volpert OV, Ward WF, Lingen MW, Chesler L, Solt DB, Johnson MD, Molteni A, Polverini PJ, Bouck NP. Captopril inhibits angiogenesis and slows the growth of experimental tumors in rats. J Clin Invest 1996; 98:671-9. [PMID: 8698858]
 27. Yoshiji H, Kuriyama S, Kawata M, Yoshii J, Ikenaka Y, Noguchi R, Nakatani T, Tsujinoue H, Fukui H. The angiotensin-I-converting enzyme inhibitor perindopril suppresses tumor growth and angiogenesis: possible role of the vascular endothelial growth factor. Clin Cancer Res 2001; 7:1073-8. [PMID: 11309359]
 28. Hii SI, Nicol DL, Gotley DC, Thompson LC, Green MK, Jonsson JR. Captopril inhibits tumour growth in a xenograft model of human renal cell carcinoma. Br J Cancer 1998; 77:880-3. [PMID: 9528828]
 29. Arafat HA, Gong Q, Chipitsyna G, Rizvi A, Saa CT, Yeo CJ. Antihypertensives as novel antineoplastics: angiotensin-I-converting enzyme inhibitors and angiotensin II type 1 receptor blockers in pancreatic ductal adenocarcinoma. J Am Coll Surg 2007; 204:996-1005. [PMID: 17481528]
 30. Yoshiji H, Kuriyama S, Fukui H. Angiotensin-I-converting enzyme inhibitors may be an alternative anti-angiogenic strategy in the treatment of liver fibrosis and hepatocellular carcinoma. Possible role of vascular endothelial growth factor. Tumour Biol 2002; 23:348-56. [PMID: 12677092]
 31. Nagai N, Oike Y, Izumi-Nagai K, Koto T, Satofuka S, Shinoda H, Noda K, Ozawa Y, Inoue M, Tsubota K, Ishida S. Suppression of choroidal neovascularization by inhibiting angiotensin-converting enzyme: minimal role of bradykinin. Invest Ophthalmol Vis Sci 2007; 48:2321-6. [PMID: 17460297]
 32. Nagai N, Noda K, Urano T, Kubota Y, Shinoda H, Koto T, Shinoda K, Inoue M, Shiomi T, Ikeda E, Tsubota K, Suda T, Oike Y, Ishida S. Selective suppression of pathological, but not physiological, retinal neovascularization by blocking angiotensin II type 1 receptor. Invest Ophthalmol Vis Sci 2005; 46:1078-84. [PMID: 15728568]
 33. Mauer M, Zinman B, Gardiner R, Suissa S, Sinaiko A, Strand T, Drummond K, Donnelly S, Goodyer P, Gubler MC, Klein R. Renal and retinal effects of enalapril and losartan in type 1 diabetes. N Engl J Med 2009; 361:40-51. [PMID: 19571282]
 34. Vij N, Sharma A, Thakkar M, Sinha S, Mohan RR. PDGF-driven proliferation, migration, and IL8 chemokine secretion in human corneal fibroblasts involve JAK2-STAT3 signaling pathway. Mol Vis 2008; 14:1020-7. [PMID: 18523665]
 35. Sun Y, Weber KT. Angiotensin converting enzyme and myofibroblasts during tissue repair in the rat heart. J Mol Cell Cardiol 1996; 28:851-8. [PMID: 8762025]
 36. Min LJ, Cui TX, Yahata Y, Yamasaki K, Shiuchi T, Liu HW, Chen R, Li JM, Okumura M, Jinno T, Wu L, Iwai M, Nahmias C, Hashimoto K, Horiuchi M. Regulation of collagen synthesis in mouse skin fibroblasts by distinct angiotensin II

- receptor subtypes. *Endocrinology* 2004; 145:253-60. [PMID: 14551224]
37. Lijnen PJ, Petrov VV. Role of intracardiac renin-angiotensin-aldosterone system in extracellular matrix remodeling. *Methods Find Exp Clin Pharmacol* 2003; 25:541-64. [PMID: 14571285]
38. Santos CF, Akashi AE, Dionísio TJ, Sipert CR, Didier DN, Greene AS, Oliveira SH, Pereira HJ, Becari C, Oliveira EB, Salgado MC. Characterization of a local renin-angiotensin system in rat gingival tissue. *J Periodontol* 2009; 80:130-9. [PMID: 19228099]
39. Min LJ, Cui TX, Yahata Y, Yamasaki K, Shiuchi T, Liu HW, Chen R, Li JM, Okumura M, Jinno T, Wu L, Iwai M, Nahmias C, Hashimoto K, Horiuchi M. Regulation of collagen synthesis in mouse skin fibroblasts by distinct angiotensin II receptor subtypes. *Endocrinology* 2004; 145:253-60. [PMID: 14551224]
40. Le Noble FA, Hekking JW, Van Straaten HW, Slaaf DW, Stryker Boudier HA. Angiotensin II stimulates angiogenesis in the chorio-allantoic membrane of the chick embryo. *Eur J Pharmacol* 1991; 195:305-6. [PMID: 1874278]
41. Fernandez LA, Twickler J, Mead A. Neovascularization produced by angiotensin II. *J Lab Clin Med* 1985; 105:141-5. [PMID: 2579174]
42. Hoshida S, Nishida M, Yamashita N, Igarashi J, Aoki K, Hori M, Kuzuya T, Tada M. Vascular angiotensin-converting enzyme activity in cholesterol-fed rabbits: effects of enalapril. *Atherosclerosis* 1997; 130:53-9. [PMID: 9126648]