

Dalibor Volný

The central limit problem for strictly stationary sequences [Abstract of thesis]

Commentationes Mathematicae Universitatis Carolinae, Vol. 26 (1985), No. 4, 840--842

Persistent URL: <http://dml.cz/dmlcz/106422>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1985

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ON THE REGULARITY OF WEAK SOLUTIONS TO NONLINEAR ELLIPTIC SYSTEMS
OF PARTIAL DIFFERENTIAL EQUATIONS

J. DANĚČEK, Math. Dept., Faculty of Civil Engineering, Technical Univ., Barvičova 85, 66237 Brno, Czechoslovakia
(13.5. 1985, supervisor J. Nečas)

A regularity of weak solution for nonlinear elliptic system of partial differential equations is proved for the case of weak solution gradient of the system being unbounded and belonging to Campanato's space $\mathcal{L}^{2,n}$. Existing nonlinear system whose weak solution has its gradient in $\mathcal{L}^{2,n}$ space is also presented.

The considered nonlinear elliptic system whose gradient weak solution locally belongs to $\mathcal{L}^{2,n}$ space is in the form of

$$- D_i((a_{ij}^{rs}(x)D_j u_s) + g_i^r(x,u,Du)) + g^r(x,u,Du) = f^r(x),$$

where $r,s=1,\dots,N$, $i,j=1,\dots,n$, $N>1$, $n\geq 3$ are natural numbers, $x \in \Omega$, Ω is bounded, open subset in \mathbb{R}^n and $u(x) \in W_{loc}^{1,2}(\Omega, \mathbb{R}^N)$ is a weak solution of this system. The functions $a_{ij}^{rs}(x)$, $g_i^r(x,u,p)$, $g^r(x,u,p)$, $f^r(x)$ of the system fulfil certain hypotheses on the smooth and growth conditions in the variables $(u,p) \in \mathbb{R}^N \times \mathbb{R}^{nN}$.

The major part of the work is devoted to the proof of a new statement describing $C^{1,\alpha}$ -regularity for the nonlinear elliptic system in divergent form

$$- D_i a_i^r(x,u,Du) + a^r(x,u,Du) = - D_i f_i^r(x) + f^r(x),$$

where $r = 1, \dots, N$, $N > 1$, $x \in \Omega$, Ω bounded, open subset in \mathbb{R}^n and $u(x) \in W_{loc}^{1,2}(\Omega, \mathbb{R}^N)$ is a weak solution. The system is further assumed to fulfil Liouville's condition and the gradient of weak solution is assumed to belong to Campanato's space $\mathcal{L}^{2,n}$.

THE CENTRAL LIMIT PROBLEM FOR STRICTLY STATIONARY SEQUENCES

D. VOLNÝ, Math. Inst. of Charles Univ., Sokolovská 83, 186 00 Praha 8, Czechoslovakia
(4.4. 1985, supervisor K. Winkelbauer)

Let $(X_i; i \in \mathbb{Z})$ be a strictly stationary sequence of random variables. Then there exists a function f and a bijective, bimeasurable and measure preserving transformation T on some probability space $(\Omega, \mathcal{A}, \mu)$ such that $(X_i; i \in \mathbb{Z})$ have the same distributions. In this thesis the central limit problem is investigated for strictly stationary sequences $(f \circ T^i; i \in \mathbb{Z})$ where $f \in L^2(\mu)$; more exactly, there is investigated the weak convergence of probability measures $\mu_n^{-1}(f)$ where $s_n(f) = \frac{1}{\sqrt{n}} \sum_{j=1}^n f \circ T^j$. As a supporting but relatively independent result, a theory of decompositions of $(f \circ T^i; i \in \mathbb{Z})$ into sums of martingale difference sequences is developed.

A σ -algebra $\mathcal{M} \subset \mathcal{A}$ such that $\mathcal{M} \subset T^{-1}\mathcal{M}$, will be called invariant. If Λ is a linearly ordered set and for each $\alpha \in \Lambda$, \mathcal{M}_α is an invariant σ -algebra and for $\alpha < \beta$ it is $T^i \mathcal{M}_\alpha \supset T^j \mathcal{M}_\beta$ (for each $i, j \in \mathbb{Z}$), we say that $(\mathcal{M}_\alpha; \alpha \in \Lambda)$ is a family of invariant σ -algebras. For an invariant σ -algebra \mathcal{M} , the projection operator onto $L^2(T^{-i-1}\mathcal{M}, \mu) \otimes L^2(T^{-i}\mathcal{M}, \mu)$ will be called a difference projection (generated by \mathcal{M} in $L^2(\mu)$) and denoted by P_i . Given a family of invariant σ -algebras $(\mathcal{M}_\alpha; \alpha \in \Lambda)$ we thus obtain a family of difference projections $(P_{-\infty}, P_{\alpha, i}; \alpha \in \Lambda, i \in \mathbb{Z})$ where $P_{-\infty}$ is the projection onto $L^2(\bigcap_{\alpha \in \Lambda} \bigcap_{i \in \mathbb{Z}} T^i \mathcal{M}_\alpha, \mu)$ and $P_{\alpha, i}$ are generated by \mathcal{M}_α . If $f \in L^2(\mu)$ and there exists a family of difference projections such that $f = \sum_{\alpha \in \Lambda} \sum_{i \in \mathbb{Z}} P_{\alpha, i} f \text{ mod } \mu$, we shall say that f is difference decomposable. Note that for a difference projection operator P and $f \in L^2(\mu)$, $(Pf) \circ T^i, i \in \mathbb{Z}$ is a martingale difference sequence.

Let \mathcal{P} denote the Pinsker σ -algebra. If $f \in L^2(\mathcal{P}, \mu)$, we shall say that f is absolutely undecomposable.

Theorem 1. Let $\mathcal{Q} \subset \mathcal{P}$ be an invariant σ -algebra and $\mathbb{P} = (\mathcal{M}_\alpha; \alpha \in \Lambda)$ be a family of invariant σ -algebras. $\mathbb{P}' = (\mathcal{M}_\alpha \vee \mathcal{Q}; \alpha \in \Lambda)$ is then also a family of invariant σ -algebras. Let $(P_{-\infty}, P_{\alpha, i}; \alpha \in \Lambda, i \in \mathbb{Z})$ and $(P'_{-\infty}, P'_{\alpha, i}; \alpha \in \Lambda, i \in \mathbb{Z})$ be the families of difference projections generated by \mathbb{P}, \mathbb{P}' . If $f \in L^2(\mu)$ and $f = \sum_{\alpha \in \Lambda} \sum_{i \in \mathbb{Z}} P_{\alpha, i} f + P_{-\infty} f \text{ mod } \mu$, then $P_{-\infty} f = P'_{-\infty} f \text{ mod } \mu$ and for each $\alpha \in \Lambda, i \in \mathbb{Z}$, it is $P_{\alpha, i} f = P'_{\alpha, i} f \text{ mod } \mu$.

Theorem 2. The set of all difference decomposable functions from $L^2(\mu)$ coincides with the Hilbert space $L^2(\mu) \otimes L^2(\mathcal{P}, \mu)$. For each difference decomposable function $f \in L^2(\mu)$ there exists an invariant σ -algebra \mathcal{M} such that $f = \sum_{i \in \mathbb{Z}} P_i f \text{ mod } \mu$ (where P_i are generated by \mathcal{M}).

Let \mathcal{J} be the σ -algebra of invariant sets from \mathcal{A} (i.e. $A \in \mathcal{J}$ iff $A = TA$). Let us suppose that \mathcal{A} has an at most countable generator and that there exist probability measures $m_\omega, \omega \in \Omega$ such that for $A \in \mathcal{A}, \omega \mapsto m_\omega(A)$ is a version of the conditional probability $\mu(A | \mathcal{J})$. For almost all $\omega \in \Omega[\mu]$, m_ω are ergodic measures and the following theorem holds.

Theorem 3. Let $\mathcal{M} \subset \mathcal{A}$ be an invariant σ -algebra and $P_i, P_i^\omega, i \in \mathbb{Z}$, be the difference projections generated by \mathcal{M} in $L^2(\mu), L^2(m_\omega), \omega \in \Omega$. Given $f \in L^2(\mu)$, for almost all $\omega \in \Omega[\mu]$, we have $P_i^\omega f = P_i f \text{ mod } m_\omega$.

Let \mathcal{X} denote the set of all functions $f \in L^2(\mu)$, such that

there exists a family $(P_{-\infty, i}^{\alpha}, P_{\alpha, i}; \alpha \in \Lambda, i \in \mathbb{Z})$ and a natural number k such that $f = \sum_{\alpha \in \Lambda} \sum_{i=0}^k P_{\alpha, i} f \text{ mod } \mu$. If

$\inf_{g \in \mathcal{G}} \limsup_{n \rightarrow \infty} \|s_n(f-g)\|_2 = 0$, we shall say that f is finitely approximable.

Theorem 4. If $f \in L^2(\mu)$ is a finitely approximable function, there exists a limit $\eta^2 = \lim_{n \rightarrow \infty} E(s_n^2(f) | \mathcal{J})$ in $L^1(\mu)$ and the measures $\mu_{s_n^{-1}(f)}$ weakly converge to a distribution with the characteristic function $\varphi: t \mapsto E \exp(-\frac{1}{2} \eta^2 t^2)$. If $\eta^2 > 0$ a.s., then $\mu_{(\frac{1}{\eta} s_n(f))^{-1}}$ weakly converge to the standard normal distribution.

Theorem 5. In the Hilbert space of difference decomposable functions, there exists a dense set of functions f such that $\mu_{s_n^{-1}(f)}$ converge to some probability measure and provided there exists a nonzero difference decomposable function, the set of difference decomposable functions f such that the sequence $\mu_{s_n^{-1}(f)}$, $n=1, 2, \dots$ has at least two different limit points, is dense (in the space of difference decomposable functions).

In deriving these results, several facts from ergodic theory (namely from entropic ergodic theory) and from martingale limit theory were used. I consider this thesis to be a continuation of the research made by M.I. Gordin, C.C. Heyde, G.K. Eagleson (preceded by the limit theorem of P. Billingsley and I.A. Ibragimov) and others. Several counterexamples to the CLP for strictly stationary processes (different from that one in Theorem 5) are given by R.C. Bradley.

References

1. Billingsley P.: The Lindeberg-Lévy theorem for martingales, Proc. Amer. Math. Soc. 12(1961), 788-792.
2. Bradley R.C. Jr.: Information regularity and the central limit question, Rocky Mountain Journ. of Math. 13(1983), 77-97.
3. Eagleson G.K.: On Gordin's central limit theorem for stationary processes, J. Appl. Probab. 12(1975), 176-179.
4. Gordin M.I.: The central limit theorem for stationary processes, Soviet Math. Dokl. 10(1969), 1174-1176. Doklady AN SSSR 188 (1969), 739-741.
5. Hall P. and Heyde C.C.: Martingale Limit Theory and its Application, Academic Press New York (1980).
6. Heyde C.C.: On the central limit theorem for stationary processes, Z. Wahrsch. Verw. Gebiete 30(1974), 315-320.
7. Ibragimov I.A.: A central limit theorem for a class of dependent random variables, Theory Probab. Appl. 8(1963), 83-89.
8. Volný D.: A negative answer to the central limit problem for strictly stationary sequences, Proc. 3rd Prague Symp. on Asymptotic Statistics, 433-441, Elsevier Science Publ. B.V., Amsterdam (1984)