

Jan Trlifaj

Non-perfect rings and a theorem of Eklof and Shelah

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 32 (1991), No. 1, 27--32

Persistent URL: <http://dml.cz/dmlcz/116939>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1991

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Non-perfect rings and a theorem of Eklof and Shelah

JAN TRLIFAJ

*Abstract.* We prove a stronger form,  $A^+$ , of a consistency result,  $A$ , due to Eklof and Shelah.  $A^+$  concerns extension properties of modules over non-left perfect rings. We also show (in ZFC) that  $A$  does not hold for left perfect rings.

*Keywords:* perfect ring, Ext, uniformization

*Classification:* 16A50, 16A51

Recently, a significant extension of the theory of Whitehead modules from domains to arbitrary non-left perfect rings has been performed by Paul C. Eklof and Sharon Shelah ([3]). In [3, Theorem 2.1 and Corollary 2.2], they proved that the assertion

$A$ : “for any non-left perfect ring  $R$  and any uncountable cardinal  $\kappa$  such that  $\text{cf}(\kappa) = \aleph_0$  and  $\kappa \geq \text{card}(R)$  there is a non-projective  $\kappa^+$ -free module  $M$  such that  $\text{card}(M) = \kappa^+$  and  $\text{Ext}_R(M, N) = 0$  whenever  $N$  is a module with  $\text{card}(N) < \kappa$ ” is consistent with ZFC + GCH. Their proof consists of two parts: the set theoretic one showing consistency of the existence of certain  $\omega$ -trees and the algebraic one inferring  $A$  from the existence of the trees.

Independently, using consistency of a uniformization principle due to Shelah, we proved a weaker form of  $A$  is consistent in the particular case of von Neumann regular rings ([5, Lemma 2.4]). In the present paper, we show our approach can be modified to obtain a simple proof of the consistency of  $A$ . Moreover, we show that a stronger form of  $A$ , denoted by  $A^+$ , is consistent, namely the expression “ $\kappa^+$ -free” can be replaced by “strongly  $\kappa^+$ -free” (see Corollary 1.6 below). The point here is that we use the definition of Ext via Hom-groups rather than via exact sequences. We also work directly with the defining relations of modules rather than with the tree-module structures.

The result of Eklof and Shelah is the best possible: we show in ZFC that for any left perfect ring  $R$  there is a proper class  $C$  consisting of pairwise non-isomorphic modules such that  $\text{Ext}_R(M, N) \neq 0$  for all  $N \in C$  and all non-projective modules  $M$  (Theorem 1.10).

Let  $M$  be a module. Then  $\text{gen}(M)$  denotes the minimum of cardinalities of  $R$ -generating subsets of  $M$ . Further,  $M$  is said to be  $\kappa$ -free provided for each submodule  $N \subseteq M$  with  $\text{gen}(N) < \kappa$  there is a free module  $P \subseteq M$  such that  $N \subseteq P$  and  $\text{gen}(P) < \kappa$ . Moreover,  $M$  is strongly  $\kappa$ -free provided for each submodule  $N \subseteq M$  with  $\text{gen}(N) < \kappa$  there is a free module  $P \subseteq M$  such that  $N \subseteq P$ ,  $\text{gen}(P) < \kappa$  and  $M/P$  is  $\kappa$ -free. A sequence  $(M_\alpha \mid \alpha < \kappa)$  is said to be a  $\kappa$ -filtration of  $M$ , if for

all  $\alpha < \kappa$ ,  $M_\alpha$  is a submodule of  $M_{\alpha+1}$  such that  $\text{gen}(M_\alpha) < \kappa$ ,  $M_\alpha = \bigcup_{\beta < \alpha} M_\beta$  for all limit  $\alpha < \kappa$ , and  $M = \bigcup_{\alpha < \kappa} M_\alpha$ .

Let  $R$  be a ring. Then  $R$  is said to be completely reducible provided  $R$  is a ring direct sum of a finite number of full matrix rings over skew fields.

Homomorphisms of (left  $R$ -)modules are written as acting on the right. Further concepts and notation can be found e.g. in [1] and [2].

**Definition 1.1.** Let  $R$  be a non-left perfect ring. By [1, Theorem 28.4], there exist elements  $a_i \in R$ ,  $i < \aleph_0$ , such that  $(a_0 \dots a_i R \mid i < \aleph_0)$  is a strictly decreasing chain of principal right ideals of  $R$ . Let  $\kappa$  be an infinite cardinal and  $E$  be a subset of  $\kappa^+$  such that  $E \subseteq \{\alpha < \kappa^+ \mid \text{cf}(\alpha) = \aleph_0\}$ . Let  $(n_\nu \mid \nu \in E)$  be a ladder system, i.e. for each  $\nu \in E$ , let  $(n_\nu(i) \mid i < \aleph_0)$  be a strictly increasing sequence of non-limit ordinals less than  $\nu$  such that  $\sup_{i < \aleph_0} n_\nu(i) = \nu$ .

Let  $(R_\alpha \mid \alpha < \kappa)$  be a system of free modules defined as follows:  $R_\alpha = R$  provided  $\alpha \in \kappa^+ \setminus E$ , and  $R_\alpha = R^{(\aleph_0)}$  provided  $\alpha \in E$ . For  $\alpha \in \kappa^+ \setminus E$ , denote by  $1_\alpha$  the canonical generator of  $R_\alpha$ , and for  $\alpha \in E$  let  $\{1_{\alpha,i} \mid i < \aleph_0\}$  be the canonical basis of  $R_\alpha$ . Note that by [1, Lemmas 28.1 and 28.2], for every  $\nu \in E$ , the module

$$S_\nu = \sum_{i < \aleph_0} R(-1_{\nu,i} + a_i \cdot 1_{\nu,i+1})$$

is a free submodule of  $R_\nu$  such that  $R_\nu/S_\nu$  is not projective. Put  $P = \bigoplus_{\alpha < \kappa^+} R_\alpha$  and  $Q = \sum_{\alpha \in E} Q_\alpha$ , where  $Q_\alpha = \sum_{i < \aleph_0} R g_{\alpha i}$  and  $g_{\alpha i} = (1_{n_\alpha(i)} - 1_{\alpha,i} + a_i \cdot 1_{\alpha,i+1}) \in P$ , for all  $\alpha \in E$  and  $i < \aleph_0$ . Finally, put  $M = P/Q \in R\text{-mod}$ .

**Lemma 1.2.** (i)  $\text{gen}(M) = \kappa^+$ .

(ii) If  $E$  is a stationary subset of  $\kappa^+$ , then  $M$  is not projective.

(iii) If  $E$  is non-reflecting (i.e.  $E \cap \sigma$  is not stationary in  $\sigma$  for all limit ordinals  $\sigma < \kappa^+$ ), then  $M$  is strongly  $\kappa^+$ -free.

PROOF: (i) This follows easily from the fact that  $\{1_\alpha + Q \mid \alpha \in \kappa^+ \setminus E\} \cup \{1_{\alpha,i} + Q \mid \alpha \in E, i < \aleph_0\}$  is an  $R$ -generating subset of  $M$ .

(ii) Put  $M_0 = 0$  and, for each  $0 < \alpha < \kappa^+$ ,  $M_\alpha = (\bigoplus_{\beta < \alpha} R_\beta + Q)/Q$ . Then  $(M_\alpha \mid \alpha < \kappa^+)$  is a  $\kappa^+$ -filtration of  $M$ .

Assume  $M$  is projective. By [1, Corollary 26.2] there exist modules  $(P_\alpha \mid \alpha < \kappa^+)$  such that  $\text{gen}(P_\alpha) \leq \aleph_0$  for all  $\alpha < \kappa^+$  and  $M = \bigoplus_{\alpha < \kappa^+} P_\alpha$ . Put  $N_0 = 0$  and, for each  $0 < \alpha < \kappa^+$ ,  $N_\alpha = \bigoplus_{\beta < \alpha} P_\beta$ . Clearly,  $(N_\alpha \mid \alpha < \kappa^+)$  is a  $\kappa^+$ -filtration of  $M$ . Since the set  $C = \{\alpha < \kappa^+ \mid M_\alpha = N_\alpha\}$  is closed and cofinal in  $\kappa^+$ , there exists  $\nu \in E \cap C$ . Of course,  $D = C \cap \{\alpha < \kappa^+ \mid \nu < \alpha\}$  is also closed and cofinal in  $\kappa^+$ , whence there is some  $\mu \in E \cap D$ . Then  $X = N_\mu/N_\nu$  is a projective module. On the other hand, put  $Y = \bigoplus_{\nu < \alpha < \mu} R_\alpha$ . Then  $X = M_\mu/M_\nu = M_{\nu+1}/M_\nu + (Y + M_\nu)/M_\nu$ . By 1.1,  $(Y + M_\nu) \cap M_{\nu+1} \subseteq M_\nu$ , whence  $M_{\nu+1}/M_\nu \simeq R_\nu/S_\nu$  is a non-projective direct summand of  $X$ , a contradiction.

(iii) First, we prove by induction on  $\nu < \kappa^+$  that for any  $\emptyset \neq A \subseteq E$  such that  $\sup(A) = \nu$  there is a sequence  $(p_a \mid a \in A)$  such that  $p_a < \aleph_0$  for all  $a \in A$ , and  $\{\{n_a(i) \mid p_a < i < \aleph_0\} \mid a \in A\}$  is a set of disjoint subsets of  $\nu$

(cp. with [3, p. 15]). For  $\nu = \min(E)$ , put  $p_a = 0$ . If  $\nu > \min(E)$ , there is a closed and cofinal subset  $C \subseteq \nu$  such that  $C \cap E \cap \nu = \emptyset$  and  $0 \in C$ . Let  $f$  be a strictly increasing function  $f : \text{card}(\nu) \rightarrow C$ . For each  $\alpha < \text{card}(\nu)$ , put  $B_\alpha = \{\beta \mid f(\alpha) < \beta < f(\alpha + 1)\}$ . If  $A \cap B_\alpha \neq \emptyset$ , then by induction there are  $(q_a \mid a \in A \cap B_\alpha)$  such that  $\{\{n_a(i) \mid q_a < i < \aleph_0\} \mid a \in A \cap B_\alpha\}$  is a set of disjoint subsets of  $f(\alpha + 1)$ . For  $a \in A \cap B_\alpha$ , put  $s_a = \min\{i < \aleph_0 \mid f(\alpha) < n_a(i)\}$ . Since  $A$  is a disjoint union of the sets  $A \cap B_\alpha, \alpha < \text{card}(\nu)$ , it suffices to put  $p_a = \max(q_a, s_a)$ , for all  $a \in A \cap B_\alpha$  and  $\alpha < \text{card}(\nu)$ . To complete the proof, we show that for all  $\alpha < \kappa^+$ , the module  $M_\alpha = (\oplus \sum_{\beta < \alpha} R_\beta + Q)/Q$  is free, and for all  $\alpha < \beta < \kappa^+$ , the module  $M_\beta/M_{\alpha+1}$  is free. Put  $A = E \cap \alpha$ . By 1.1 and the construction of  $(p_a \mid a \in A)$ , we see that  $\{1_{a,i} + Q \mid a \in A \& p_a < i < \aleph_0\} \cup \{1_b + Q \mid b < \alpha \& b \notin A \& \text{non}(\exists a \in A \exists i < \aleph_0 : p_a < i \& b = n_a(i))\}$  is a free  $R$ -basis of the module  $M_\alpha$ . Finally, put  $A = E \cap \beta$ . For each  $a \in A$  such that  $a > \alpha$ , let  $r_a < \aleph_0$  be such that  $p_a \leq r_a$  and  $\alpha < n_a(i)$  for all  $r_a < i < \aleph_0$ . Then by 1.1,  $\{1_{a,i} + M_{\alpha+1} \mid a \in A \& a > \alpha \& r_a < i < \aleph_0\} \cup \{1_b + M_{\alpha+1} \mid \alpha < b < \beta \& b \notin A \& \text{non}(\exists a \in A \exists i < \aleph_0 : a > \alpha \& r_a < i \& b = n_a(i))\}$  is a free  $R$ -basis of the module  $M_\beta/M_{\alpha+1}$ .  $\square$

**Lemma 1.3.** *Let  $\kappa$  be a cardinal such that  $\text{cf}(\kappa) = \aleph_0$ . Consider the following assertion*

UP $_\kappa$ : “there exist a non-reflecting stationary subset  $E$  of  $\kappa^+$  satisfying  $E \subseteq \{\alpha < \kappa^+ \mid \text{cf}(\alpha) = \aleph_0\}$  and a ladder system  $(n_\nu \mid \nu \in E)$  such that for each cardinal  $\lambda < \kappa$  and each sequence  $(h_\nu \mid \nu \in E)$  of mappings from  $\aleph_0$  to  $\lambda$  there is a mapping  $f : \kappa^+ \rightarrow \lambda$  such that  $\forall \nu \in E \exists j < \aleph_0 \forall j < i < \aleph_0 : f(n_\nu(i)) = h_\nu(i)$ ”.

Then the assertion “UP $_\kappa$  holds for every uncountable cardinal  $\kappa$  such that  $\text{cf}(\kappa) = \aleph_0$ ” is consistent with ZFC + GCH.

PROOF: By [4, §2] or [3, §2].  $\square$

**Lemma 1.4.** *Let  $\kappa$  be a cardinal such that  $\text{cf}(\kappa) = \aleph_0$  and  $\text{card}(R) \leq \kappa$ . Assume UP $_\kappa$  holds. Let  $M = P/Q$  be the module corresponding to the  $E$  and  $(n_\nu(i) \mid \nu \in E)$  from UP $_\kappa$  by 1.1. Then  $\text{Ext}_R(M, N) = 0$  for all  $N \in R\text{-mod}$  such that  $\text{card}(N) < \kappa$ .*

PROOF: Since  $P$  is a free module, we have  $\text{Ext}_R(M, N) = \text{Hom}_R(Q, N)/\tau \circ \text{Hom}_R(P, N)$ ,  $\tau$  being the inclusion of  $Q$  into  $P$ . Hence, we are to prove that every  $x \in \text{Hom}_R(Q, N)$  is a restriction of some  $y \in \text{Hom}_R(P, N)$ , i.e.  $x = \tau y$ . Take  $x \in \text{Hom}_R(Q, N)$ . Let  $b : N \rightarrow \lambda$  be a bijection of  $N$  onto  $\lambda = \text{card}(N)$ . Using the notation of 1.1, for each  $\nu \in E$ , we define  $h_\nu : \aleph_0 \rightarrow \lambda$  by  $h_\nu(i) = b(g_{\nu i}x)$  for all  $i < \aleph_0$ . By UP $_\kappa$ , there exists  $f : \kappa^+ \rightarrow \lambda$  such that  $\forall \nu \in E \exists j_\nu < \aleph_0 \forall j_\nu < i < \aleph_0 : h_\nu(i) = f(n_\nu(i))$ . Define  $y \in \text{Hom}_R(P, N)$  as follows: Take  $\alpha < \kappa^+$ .

(I) If  $\alpha = n_\nu(i)$  for some  $\nu \in E$  and  $j_\nu < i < \aleph_0$ , put  $1_\alpha y = b^{-1}f(\alpha)$ ;

(II) If  $\alpha$  does not satisfy (I) and  $\alpha \notin E$ , put  $1_\alpha y = 0$ ;

(III) If  $\alpha \in E$ , put  $1_{\alpha,i}y = 0$  provided  $i > j_\alpha$ . For  $0 \leq i \leq j_\alpha$ , define  $1_{\alpha,i}y$  by induction on  $i$  (downwards): If there exist  $\nu \in E$  and  $k > j_\nu$  such that  $n_\alpha(i) = n_\nu(k)$ , put  $1_{\alpha,i}y = b^{-1}f(n_\alpha(i)) - g_{\alpha i}x + a_i \cdot 1_{\alpha,i+1}y$ . If there are no  $\nu \in E$  and  $k > j_\nu$  such that  $n_\alpha(i) = n_\nu(k)$ , put  $1_{\alpha,i}y = -g_{\alpha i}x + a_i \cdot 1_{\alpha,i+1}$ .

It remains to prove that  $g_{\alpha i}x = g_{\alpha i}y$  for all  $\alpha \in E$  and  $i < \aleph_0$ . Put  $\beta = n_\alpha(i)$ . Of course,  $g_{\alpha i}y = 1_\beta y - 1_{\alpha, i}y + a_i \cdot 1_{\alpha, i+1}y$ . We distinguish the following three cases:

- (1)  $i > j_\alpha$ . Then  $1_\beta y = b^{-1}f(\beta) = b^{-1}h_\alpha(i) = g_{\alpha i}x$  and  $1_{\alpha, i}y = 1_{\alpha, i+1}y = 0$ , whence  $g_{\alpha i}y = g_{\alpha i}x$ ;
- (2)  $i \leq j_\alpha$ , but there exist  $\nu \in E$  and  $k > j_\nu$  such that  $\beta = n_\nu(k)$ . Then  $1_\beta y = b^{-1}f(\beta)$  and  $1_{\alpha, i}y = b^{-1}f(\beta) - g_{\alpha i}x + a_i \cdot 1_{\alpha, i+1}y$ , whence  $g_{\alpha i}y = g_{\alpha i}x$ ;
- (3)  $i \leq j_\alpha$  and there are no  $\nu \in E$  and  $k > j_\nu$  such that  $\beta = n_\nu(k)$ . Then  $1_\beta y = 0$  and  $1_{\alpha, i}y = -g_{\alpha i}x + a_i \cdot 1_{\alpha, i+1}y$  whence  $g_{\alpha i}y = g_{\alpha i}x$ , q.e.d.  $\square$

**Theorem 1.5.** *Let  $\kappa$  be a cardinal such that  $\text{cf}(\kappa) = \aleph_0$  and  $\text{UP}_\kappa$  holds. Let  $R$  be a non-left perfect ring with  $\text{card}(R) \leq \kappa$ . Then there is a non-projective strongly  $\kappa^+$ -free module  $M$  such that  $\text{card}(M) = \kappa^+$  and  $\text{Ext}_R(M, N) = 0$  for all  $N \in R\text{-mod}$  with  $\text{card}(N) < \kappa$ .*

PROOF: By 1.2 and 1.4.  $\square$

**Corollary 1.6.** *Consider the following assertion*

$A^+$ : “for any non-left perfect ring  $R$  and any uncountable cardinal  $\kappa$  such that  $\text{cf}(\kappa) = \aleph_0$  and  $\kappa \geq \text{card}(R)$  there is a non-projective strongly  $\kappa^+$ -free module  $M$  such that  $\text{card}(M) = \kappa^+$  and  $\text{Ext}_R(M, N) = 0$  for all  $N \in R\text{-mod}$  with  $\text{card}(N) < \kappa$ ”.

Then  $A^+$  is consistent with ZFC + GCH.

PROOF: By 1.3 and 1.5.  $\square$

The following proposition shows (in ZFC) that the extension properties of “small” non-projective modules may depend strongly on the particular structure of the non-left perfect ring  $R$ .

**Proposition 1.7.** (i) *Let  $R = k[y, D]$  be the ring of all differential polynomials in one indeterminate  $y$  over a universal differential field  $k$  with the differentiation  $D$ . Then  $R$  is not left perfect, but  $\text{Ext}_R(M, N) \neq 0$  for all non-injective modules  $N$  and all finitely generated non-projective modules  $M$ .*

(ii) *Let  $R$  be a simple countable non-completely reducible von Neumann regular ring. Then  $R$  is not left perfect, but  $\text{Ext}_R(M, N) \neq 0$  for all non-projective modules  $M$  such that  $\text{gen}(M) \leq \aleph_0$  and all non-zero modules  $N$  such that  $\text{gen}(N) \leq \aleph_0$ . However, there exist a simple non-projective module  $S$  and a non-injective module  $N$  such that  $\text{Ext}_R(S, N) = 0$ .*

(iii) *Let  $R$  be a self-injective non-left perfect ring (e.g. let  $R$  be the maximal left quotient ring of a non-completely reducible von Neumann regular ring). Then there exists a non-projective module  $M$  such that  $\text{gen}(M) = \aleph_0$  and  $\text{Ext}_R(M, N) = 0$  for all finitely generated modules  $N$ .*

PROOF: (i) By [6, Theorem 9.3].

(ii) By [6, Theorem 10.4].

(iii) Let  $a_i, i < \aleph_0$  be as in 1.1. Let  $1_i, i < \aleph_0$  be the canonical basis of  $F = R^{(\aleph_0)}$  and let  $G = \sum_{i < \aleph_0} R(1_i - a_i \cdot 1_{i+1}) \subseteq F$ . Put  $M = F/G$ . By [1, Lemmas 28.1 and 28.2],  $F$  and  $G$  are free modules,  $M$  is not projective, and  $\text{gen}(M) = \aleph_0$ . If  $\text{gen}(N) < \aleph_0$ , we have  $N \simeq R^{(n)}/X$  for some  $n < \aleph_0$  and a submodule  $X \subseteq R^{(n)}$ .

As the sequence  $0 \rightarrow G \rightarrow F \rightarrow M \rightarrow 0$  is exact, we get  $0 = \text{Ext}_R(G, X) \rightarrow \text{Ext}_R^2(M, X) \rightarrow \text{Ext}_R^2(F, X) = 0$ , whence  $\text{Ext}_R^2(M, X) = 0$ . Since the sequence  $0 \rightarrow X \rightarrow R^{(n)} \rightarrow N \rightarrow 0$  is exact and  $R$  is left self-injective, we have  $0 = \text{Ext}_R(M, R^{(n)}) \rightarrow \text{Ext}_R(M, N) \rightarrow \text{Ext}_R^2(M, X) = 0$ , whence  $\text{Ext}_R(M, N) = 0$ .  $\square$

**Theorem 1.8.** *Let  $R$  be a left perfect ring.*

- (i) *For any non-projective module  $M$  there is a simple module  $S_M$  such that  $\text{Ext}_R(M, S_M) \neq 0$ .*
- (ii) *There exists a module  $N$  such that  $\text{Ext}_R(M, N) \neq 0$  for all non-projective modules  $M$ .*

PROOF: (i) Since  $R$  is left perfect, there exists a projective cover of  $M$ , i.e. a projective module  $P$  and a non-zero superfluous submodule  $K \subseteq P$  such that  $M \simeq P/K$ . By [1, Theorem 28.4], there exists a maximal submodule  $L$  of  $K$ . Put  $S_M = K/L$ . Let  $x \in \text{Hom}_R(K, S_M)$  be the projection of  $K$  onto  $K/L$ . Assume there exists  $y \in \text{Hom}_R(P, S_M)$  such that  $\tau y = x$ ,  $\tau$  being the inclusion of  $K$  into  $P$ . Then  $\text{Ker}(y)$  is a maximal submodule of  $P$  and by [1, Proposition 9.13],  $K \subseteq \text{Rad}(P) \subseteq \text{Ker}(y) \subset P$ . Thus  $\tau y = 0$ , a contradiction.

Hence  $\text{Hom}_R(K, S_M)/\tau \circ \text{Hom}_R(P, S_M) = \text{Ext}_R(M, S_M) \neq 0$ .

(ii) Denote by  $V$  a representative set of the class of all simple modules. Put  $N = \bigoplus_{S \in V} S$ . Then  $\text{Ext}_R(M, N) \simeq \text{Ext}_R(M, S_M) \dot{+} X$ , for an abelian group  $X$ . Thus, by (i),  $\text{Ext}_R(M, N) \neq 0$ .  $\square$

**Definition 1.9.** Let  $R$  be a ring. Define  $W = \{N \in R\text{-mod} \mid \text{Ext}_R(M, N) \neq 0 \text{ for all non-projective } M \in R\text{-mod}\}$ .

**Theorem 1.10.** *Let  $R$  be a ring. Consider the following assertions:*

- (i)  *$R$  is left perfect;*
- (ii)  *$W \neq \emptyset$ ;*
- (iii) *There exists a proper class  $C$  such that  $C \subseteq W$  and no two distinct elements of  $C$  are isomorphic.*

*Then (i) implies (ii), and (ii) is equivalent to (iii). The implication (iii)  $\Rightarrow$  (i) is independent of ZFC + GCH.*

PROOF: (i) implies (ii) by 1.8 (ii). If  $N \in W$ , then also  $\{N^{(\kappa)} \mid \kappa \geq \text{card}(N)\} \subseteq W$  and  $N^{(\kappa)} \not\cong N^{(\lambda)}$  for all cardinals  $\kappa \neq \lambda \geq \text{card}(N)$ . Hence (ii) is equivalent to (iii). By 1.6, the implication (iii)  $\Rightarrow$  (i) is consistent with ZFC + GCH. On the other hand, by [6, Theorem 10.8 (ii)], (non-(i) & (ii)) is consistent with ZFC + GCH.  $\square$

**Remark 1.11.** Let  $R$  be a left perfect ring. Denote by  $I$  the class of all injective modules. Clearly, always  $W \subseteq R\text{-mod} \setminus I$ . Despite 1.10 (iii), almost never  $W = R\text{-mod} \setminus I$ . Indeed, if  $R$  is left non-singular, then  $W = R\text{-mod} \setminus I$ , if and only if either  $R = S$  or  $R = T$  or  $R = S \boxplus T$ , where  $S$  is a completely reducible ring and there exists a skew field  $K$  such that  $T$  is Morita equivalent to the upper triangular matrix ring of degree two over  $K$  (see [6, Theorems 3.4 and 8.1]).

## REFERENCES

- [1] Anderson F.W., Fuller K.R., *Rings and Categories of Modules*, Springer, New York 1974.
- [2] Eklof P.C., *Set Theoretic Methods in Homological Algebra and Abelian Groups*, Montreal Univ. Press, Montreal 1980.
- [3] Eklof P.C., Shelah S., *On Whitehead modules*, preprint 1990.
- [4] Shelah S., *Diamonds, uniformization*, J. Symbolic Logic **49** (1984), 1022–1033.
- [5] Trlifaj J., *Von Neumann regular rings and the Whitehead property of modules*, Comment. Math. Univ. Carolinae **31** (1990), 621–625.
- [6] Trlifaj J., *Associative Rings and the Whitehead Property of Modules*, R. Fischer, Munich 1990.

DEPARTMENT OF ALGEBRA, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY,  
SOKOLOVSKÁ 83, 186 00 PRAHA 8, CZECHOSLOVAKIA

(Received November 5, 1990)