

Antonio de Padua Franco-Filho

Topological characterization of the small cardinal i

Commentationes Mathematicae Universitatis Carolinae, Vol. 44 (2003), No. 4, 745--750

Persistent URL: <http://dml.cz/dmlcz/119429>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2003

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Topological characterization of the small cardinal i

ANTONIO DE PADUA FRANCO-FILHO

Abstract. We show that the small cardinal number $i = \min\{|\mathcal{A}| : \mathcal{A} \text{ is a maximal independent family}\}$ has the following topological characterization: $i = \min\{\kappa \leq c : \{0, 1\}^\kappa \text{ has a dense irresolvable countable subspace}\}$, where $\{0, 1\}^\kappa$ denotes the Cantor cube of weight κ . As a consequence of this result, we have that the Cantor cube of weight c has a dense countable submaximal subspace, if we assume (ZFC plus $i = c$), or if we work in the Bell-Kunen model, where $i = \aleph_1$ and $c = \aleph_{\omega_1}$.

Keywords: independent family, irresolvable, submaximal

Classification: Primary 54A05, 54A35, 54C25; Secondary 54A25, 54B05, 54B10

1. Introduction

In this paper we will explore the relationship between the independent families of the power set of ω and the canonical subbasis of the Cantor cubes of weight $\leq c$. Let $\{0, 1\}^I$ be the Cantor cube of weight $\aleph_0 \leq |I| \leq c$. The elements of the canonical basis of this topological product space will be denoted by $W(p)$, which by definition, is $W(p) = \{s \in {}^I 2 : s \upharpoonright \text{dom}(p) = p\}$, for each $p \in \text{Fn}(I, 2)$, where $\text{Fn}(I, 2)$ is the set of all finite partial functions from the set I into 2.

Let us recall the definitions of independent families and irresolvable spaces.

Definition 1.1. $\mathcal{A} \subset \mathcal{P}(\omega)$ is an independent family if and only if for all $n, m \in \omega$ and all pairwise distinct elements $a_0, \dots, a_n, b_0, \dots, b_m$ of \mathcal{A} we have $|a_0 \cap \dots \cap a_n \cap (\omega \setminus b_0) \cap \dots \cap (\omega \setminus b_m)| = \omega$.

We will always assume $|\mathcal{A}| > \omega$ and we say that \mathcal{A} is a maximal independent family if for all $x \in \mathcal{P}(\omega) \setminus \mathcal{A}$, $\mathcal{A} \cup \{x\}$ is not an independent family.

Definition 1.2. Let (X, τ) be a Hausdorff dense-in-itself space. X is an irresolvable space if and only if for all dense subset $D \subset X$ we have $\text{int}(D) \neq \emptyset$.

Definition 1.3. Let us define the following small cardinal numbers: $i = \min\{|\mathcal{A}| : \mathcal{A} \subset \mathcal{P}(\omega) \text{ is a maximal independent family}\}$ and $\lambda = \min\{\kappa \leq c : \exists \mathcal{A} \subset \{0, 1\}^\kappa \text{ dense irresolvable countable subspace}\}$.

The author was supported by the University of São Paulo and CAPES.

2. Main theorem

Definition 2.1. (1) If $\mathcal{A} = \{A_i : i \in I\}$ is any family of subsets of ω , we define the mapping $\psi_{\mathcal{A}} : \omega \longrightarrow \{0, 1\}^I$ by: $(\forall x \in \omega) (\forall i \in I)$

$$\psi_{\mathcal{A}}(n)(i) = \begin{cases} 0 & \text{if } n \in A_i \\ 1 & \text{if } n \in \omega \setminus A_i. \end{cases}$$

(2) If $A = \{a_n : n \in \omega\} \subset \{0, 1\}^I$, let $\mathcal{A}^*(A) = \{A_i : i \in I\}$, where $A_i = \{n : a_n(i) = 0\}$.

To show that $i = \lambda$ we need to prove the following lemma.

Lemma 2.2. (a) If $\mathcal{A} \subset \mathcal{P}(\omega)$, then $\mathcal{A}^*(\psi_{\mathcal{A}}(\omega)) = \mathcal{A}$.

(b) If $A = \{a_n : n \in \omega\} \subset \{0, 1\}^I$, then $\psi_{\mathcal{A}^*(A)}(\omega) = A$.

(c) A is dense in $\{0, 1\}^I$ if and only if $\mathcal{A}^*(A)$ is an independent family. Further, if A is dense, then $|I| = |\mathcal{A}^*(A)|$.

(d) A is dense and irresolvable if and only if $\mathcal{A}^*(A)$ is maximal independent.

PROOF: (a) We define $A = \{\psi_{\mathcal{A}}(n) : n \in \omega\}$. Given any $A_i \in \mathcal{A}$, then we have $A_i = \{n : \psi_{\mathcal{A}}(n)(i) = 0\}$.

(b) As $\psi_{\mathcal{A}^*(A)}(n)(i) = a_n(i)$, $\forall i \in I$ we have that $\psi_{\mathcal{A}^*(A)}(n) = a_n$.

(c) Assume that A is dense in $\{0, 1\}^I$.

Given $n, m \in \omega$ and $\mathcal{B} = \{U_1, \dots, U_n, V_1, \dots, V_m\}$ a set of pairwise distinct elements of $\mathcal{A}^*(A)$, pick $p \in \text{Fn}(I, 2)$ such that: $\text{dom}(p) = \{i \in I : A_i \in \mathcal{B}\}$, and

$$p(i) = \begin{cases} 0 & \text{if } A_i \in \{U_1, \dots, U_n\} \\ 1 & \text{if } A_i \in \{V_1, \dots, V_m\}. \end{cases}$$

Since A is dense in $\{0, 1\}^I$ we have $|W(p) \cap A| = \omega$ and

$$\{n \in \omega : a_n \in W(p)\} = U_1 \cap \dots \cap U_n \cap (\omega \setminus V_1) \cap \dots \cap (\omega \setminus V_m)$$

which shows that $\mathcal{A}^*(A)$ is independent.

If A is not dense, there is $p \in \text{Fn}(I, 2)$ such that $W(p) \cap A = \emptyset$. Then $\bigcap \{A_i : p(i) = 0\} \cap \bigcap \{A_i : p(i) = 1\} = \emptyset$, which shows that if $\mathcal{A}^*(A)$ is independent, then A is dense in $\{0, 1\}^I$. If A is dense, then $A_i \neq A_j$ whenever $i \neq j$. Indeed, if $i \neq j$ and $A_i = A_j$, then $a_n(i) = a_n(j)$ for each $n \in \omega$, hence A is not dense.

(d) Assume that there exist two dense subsets D_0, D_1 in the space A such that $D_0 \cup D_1 = A$ and $D_0 \cap D_1 = \emptyset$. Then, for instance, $\{n : a_n \in D_0\} \notin \mathcal{A}^*(A)$ and $\mathcal{A}^*(A) \cup \{\{n : a_n \in D_0\}\} = \mathcal{B}$ would be an independent family.

Now suppose that $B \subset \omega$, $B \notin \mathcal{A}^*(A)$ and $\mathcal{A}^*(A) \cup \{B\}$ is an independent family. Then for all $p \in \text{Fn}(I, 2) \setminus \{\emptyset\}$ we have $W(p) \cap \{a_n : n \in B\} \neq \emptyset \neq W(p) \cap \{a_n : n \in A \setminus B\}$.

This shows that the space A is resolvable if and only if the independent family \mathcal{A} is not maximal. □

Theorem 2.3. $i = \lambda$.

PROOF: We first prove $i \leq \lambda$. Pick $A \subset \{0, 1\}^\lambda$ a countable dense irresolvable subspace of the Cantor cube of weight λ . By Lemma 2.2(d), we have that $\mathcal{A}^*(A)$ is a maximal independent family of cardinality $|\mathcal{A}^*(A)| = \lambda$, hence $i \leq \lambda$.

Now we prove $\lambda \leq i$. Let $\mathcal{A} \subset \mathcal{P}(\omega)$ be a maximal independent family of cardinality $|\mathcal{A}| = i$. By Lemma 2.2(d), $A = \psi_{\mathcal{A}}(\omega)$ is a dense, countable, irresolvable subspace of the Cantor cube $\{0, 1\}^i$. Hence $\lambda \leq i$. \square

3. Submaximal spaces

We say that a topological space (X, τ) is a submaximal space if and only if every dense subset of X is open in X .

In [ASTTW] the authors show that the Tychonoff cube $[0, 1]^c$ in a model of ZFC plus BL (Booth's Lemma or equivalently $\mathbf{p} = c$), has a dense countable submaximal subspace. It is well known that $\mathbf{p} \leq i$ and that it is consistent that $\mathbf{p} < i$. In [Ma], Malykhin shows that in the Bell-Kunen's model, the Cantor cube of weight ω_1 has a dense countable irresolvable subspace. Thus, in this model, by Theorem 2.3, it is true that $i = \omega_1$. Also we will show that the Cantor cube of weight $c = \aleph_{\omega_1}$ has a dense countable submaximal subspace. Hence the existence of these dense countable subspaces of these cubes is independent of $i = c$ or $\mathbf{p} = c$.

Let \mathcal{A} be an independent family of $\mathcal{P}(\omega)$. Follows from Lemma 2.2(c) that $\psi_{\mathcal{A}}(\omega)$ is the dense subspace of the Cantor cube $\{0, 1\}^{\mathcal{A}}$.

Definition 3.1. We say that $M \subset \omega$ is dense (open) in \mathcal{A} if $\psi_{\mathcal{A}}(M)$ is dense (open) in $\psi_{\mathcal{A}}(\omega)$.

Lemma 3.2. (a) D is dense in \mathcal{A} if and only if $\forall p \in \text{Fn}(\mathcal{A}, 2) \setminus \{\emptyset\}$ holds $D \cap V(p) \neq \emptyset$.

(b) G is open in \mathcal{A} if and only if for each $x \in G$ there exists $p \in \text{Fn}(\mathcal{A}, 2)$ such that $x \in V(p) \subset G$.

PROOF: Let $X = (\omega, \tau)$ be the topological space whose basis of open set is: $\{V(p) : p \in \text{Fn}(\mathcal{A}, 2) \setminus \{\emptyset\}\}$, and each $V(p)$ is defined by:

$$V(p) = \bigcap \{V : p(V) = 0\} \cap \bigcap \{\omega \setminus V : p(V) = 1\}.$$

By Lemma 2.2(c), $\psi_{\mathcal{A}}$ is a continuous open mapping from X onto the dense subspace $\psi_{\mathcal{A}}(\omega)$ of $\{0, 1\}^{\mathcal{A}}$. \square

The following example is important for construction subsets of ω which are dense and open in a given independent family, like in Lemma 3.5 and Lemma 3.7(ii).

Example 3.3. Let $Z = \{z \in 2^\omega : \{n : z(n) = 1\} \text{ is finite}\}$; Z is a countable dense subspace of the Cantor set.

Choose $s \in 2^\omega$ such that $|\{n : s(n) = 0\}| = |\{n : s(n) = 1\}| = \omega$, and define the family $\mathcal{A} = \{A_\alpha : \alpha \in \omega\}$, where $A_\alpha = \{z \in Z : z(\alpha) = s(\alpha)\}$, $\forall \alpha \in \omega$. Then follows easily from the definition, that \mathcal{A} satisfies: $\bigcap \mathcal{A} = \emptyset$ and $\bigcup \mathcal{A} = Z$. Now we define the independent family $\mathcal{T} = \{T_\alpha : \alpha \in \omega\}$, where for each $\alpha \in \omega$, $T_\alpha = \{n \in \omega : z_n \in A_\alpha\}$ and $Z = \{z_n : n \in \omega\}$ is an enumeration of the set Z .

Lemma 3.4. *Let $\mathcal{A} \subset \mathcal{B}$ be two independent families. Then*

- (i) $D \subset \omega$, dense in \mathcal{B} , implies D is also dense in \mathcal{A} ,
- (ii) $G \subset \omega$, open in \mathcal{A} , implies G is also open in \mathcal{B} .

PROOF: If we pick $p \in \text{Fn}(\mathcal{B}, 2)$ such that $\text{dom}(p) \subset \mathcal{A}$, then both (i) and (ii) follow easily from the Lemma 3.2. □

Lemma 3.5. *Let D be a subset of ω such that:*

- (i) $D = \bigcup_{n \in \omega} D_n$ and for all $n \in \omega$, $|D_n| \geq 1$,
- (ii) $n \neq m \implies D_n \cap D_m = \emptyset$.

Let \mathcal{T} be the independent family of Example 3.3. Define $\mathcal{D} = \{V_\alpha : \alpha < \omega\}$, where $V_\alpha = \bigcup \{D_n : n \in T_\alpha\}$. Then the family \mathcal{D} is independent as the family \mathcal{T} is independent; D is open in \mathcal{D} as $\bigcup \mathcal{T} = \omega$.

Lemma 3.6. *Let \mathcal{A} be an independent family and suppose that $|\mathcal{A}| < i$. If $D \subset \omega$ is dense in \mathcal{A} , then there exists a sequence $\{D_n : n \in \omega\}$ of subsets of D such that:*

- (i) for each $n \in \omega$, D_n is dense in \mathcal{A} ,
- (ii) $n \neq m \implies D_n \cap D_m = \emptyset$,
- (iii) $\bigcup_{n \in \omega} D_n = D$.

PROOF: Follows from the Main Theorem, $|\mathcal{A}| < i \implies |\mathcal{A}| < \lambda$ and hence any dense countable subset of $\{0, 1\}^\mathcal{A}$ is resolvable. □

Lemma 3.7. *Let $D \subset \omega$ be dense in \mathcal{A} and suppose that $|\mathcal{A}| < i$. Then there exists an independent family $\mathcal{B} \subset \mathcal{P}(\omega)$ such that:*

- (i) $\mathcal{A} \subset \mathcal{B}$ and $|\mathcal{B} \setminus \mathcal{A}| = \omega$,
- (ii) D is dense and open in \mathcal{B} .

PROOF: Let $\{D_n : n \in \omega\}$ be a sequence of subsets of D , as in Lemma 3.6. By Lemma 3.5, we may define an independent family \mathcal{D} , such that D is open in \mathcal{D} . The family $\mathcal{B} = \mathcal{A} \cup \mathcal{D}$ is independent by definition of \mathcal{D} and density of each D_n , which also imply that D is dense in \mathcal{B} . □

Theorem 3.8 ($i = c$). *There exists an independent family $\mathcal{A} \subset \mathcal{P}(\omega)$ such that:*

- (i) $|\mathcal{A}| = c$ and
- (ii) if $D \subset \omega$ is dense in \mathcal{A} , then D is open in \mathcal{A} .

PROOF: Let $\{E_\alpha : \alpha < c\}$ be an enumeration of all infinite subsets of ω with $E_0 = \omega$, and choose an independent family \mathcal{A}_0 such that $|\mathcal{A}_0| < c$ and $\bigcup \mathcal{A}_0 = \omega$. By transfinite induction on $c \setminus \{0\}$, we can choose a sequence of independent families $\{\mathcal{A}_\alpha : \alpha \in c \setminus \{0\}\}$, such that, by Lemma 3.7:

- (i) $\mathcal{A}_\alpha \subset \mathcal{A}_{\alpha+1}$,
- (ii) $|\mathcal{A}_{\alpha+1} \setminus \mathcal{A}_\alpha| = \omega$,
- (iii) if E_α is dense in \mathcal{A}_α , then E_α is dense and open in $\mathcal{A}_{\alpha+1}$, and
- (iv) if β is a limit ordinal, then $\mathcal{A}_\beta = \bigcup_{\alpha \in \beta} \mathcal{A}_\alpha$.

The independent family $\mathcal{A} = \bigcup_{\alpha \in c} \mathcal{A}_\alpha$ has $|\mathcal{A}| = c$. Also, if $D \subset \omega$ is dense in \mathcal{A} then $\exists \alpha \in c$ such that $D = E_\alpha$ and this set is dense in \mathcal{A}_α as it is dense in \mathcal{A} , by Lemma 3.4(i). Therefore by construction, it is open in $\mathcal{A}_{\alpha+1}$, so it is also open in \mathcal{A} , by Lemma 3.4(ii). □

Corollary 3.9 ($i = c$). *The Cantor cube $\{0, 1\}^c$ has a dense countable submaximal subspace.*

PROOF: Let \mathcal{A} be an independent family which satisfies Theorem 3.8 and let $\psi_{\mathcal{A}}(\omega)$ be a dense countable subspace of the Cantor cube of weight c . If D is dense in \mathcal{A} , then $\psi_{\mathcal{A}}^{-1}(D)$ is dense in \mathcal{A} , so it is also open in \mathcal{A} , which implies that D is open in $\psi_{\mathcal{A}}(\omega)$. □

4. Bell-Kunen’s model

Let M be a countable transitive model of ZFC plus GCH. In [BK], Bell and Kunen construct in M an increasing family of partial ordered sets $\{P_\alpha : \alpha \leq \omega_1\}$ such that:

- (i) each P_α has c.c.c.,
- (ii) if β is limit, $P_\beta = \bigcup \{P_\alpha : \alpha < \beta\}$,
- (iii) if α is not a limit ordinal, then P_α is such that both MA (Martin’s axiom) and $2^\omega = \aleph_\alpha$ hold in M^{P_α} .

Let $G = G_{\omega_1}$ be a P_{ω_1} – generic over M and $G_\alpha = G \cap P_\alpha$ for each $\alpha \leq \omega_1$. In M_{ω_1} there is a transfinite increasing sequence of models $\{M_\alpha = M[G_\alpha] : \alpha \leq \omega_1\}$, and if $\alpha > 0$ is a non-limit ordinal, then the assertion “MA plus $c = \aleph_\alpha$ ” is true in M_α . Let us also note that, in M_{ω_1} , the power set of all subset of ω is the union of the increasing sequence $\{\mathcal{P}(\omega) \cap M_{\alpha+1} : \alpha < \omega_1\}$.

Theorem 4.1. *In the Bell-Kunen’s model there is an independent family $\mathcal{A} \subset \mathcal{P}(\omega)$ such that every $D \subset \omega$ dense in \mathcal{A} is also open in \mathcal{A} . Further $|\mathcal{A}| = c$.*

PROOF: By transfinite induction in M_{ω_1} , we construct an increasing sequence of independent families $\{\mathcal{A}_\alpha : \alpha < \omega_1\}$ such that, in $M_{\alpha+1}$, the independent family \mathcal{A}_α satisfies Theorem 3.8. This is possible because MA holds in $M_{\alpha+1}$, thus $i = c = \aleph_{\alpha+1}$. Now, we look at this family in $M_{\alpha+2}$, and in this model we have that $|\mathcal{A}_{\alpha+1}| \leq \aleph_{\alpha+1} < \aleph_{\alpha+2} = c$. Then, by the prove of the Theorem 3.8, in $M_{\alpha+2}$,

we can choose an independent family $\mathcal{A}_{\alpha+2} \supset \mathcal{A}_{\alpha+1}$, such that (in $M_{\alpha+2}$) if D is dense in $\mathcal{A}_{\alpha+2}$, then it is open too. (For a limit ordinal we take the union.) The family $\mathcal{A} = \bigcup \{\mathcal{A}_\alpha : \alpha < \omega_1\}$ is an independent family in M_{ω_1} . Also, if $D \subset \omega$ is dense in \mathcal{A} , then $\exists \alpha < \omega_1$ such that $D \in \mathcal{P}(\omega) \cap M_{\alpha+1}$ and D is open in $\mathcal{A}_{\alpha+2}$ (in $M_{\alpha+2}$), so it is open in \mathcal{A} in M_{ω_1} . Further, for each $\alpha \in \omega_1$ we have that $|\mathcal{A}_{\alpha+2}| = c$ (in $M_{\alpha+2}$) hence, follows $|\mathcal{A}| = c$ in Bell-Kunen's model. \square

Corollary 4.2. *In the Bell-Kunen's model there is a countable dense submaximal subspace X of the Cantor cube $\{0, 1\}^c$.*

PROOF: Let \mathcal{A} be an independent family as in Theorem 4.1 and take $X = \psi_{\mathcal{A}}(\omega)$ be the countable dense submaximal subspace of $\{0, 1\}^c$. \square

Acknowledgment. The author wishes to acknowledge the helpful comments of the referee.

REFERENCES

- [ASTTW] Alas O.T., Sanchis M., Tkačenko M.G., Tkachuk V.V., Wilson R.G., *Irresolvable and submaximal spaces: homogeneity vs σ -discreteness and new ZFC examples*, Topology Appl. **107** (2000), 259–278.
- [BK] Bell M., Kunen K., *On the Pi-character of ultrafilters*, C.R. Math. Rep. Acad. Sci. Canada **3** (1981), 351–356.
- [Ma] Malykhin V.I., *Irresolvable countable spaces of weight less than c* , Comment. Math. Univ. Carolinae **40.1** (1999), 181–185.

INSTITUTO DE MATEMATICA E ESTATISTICA, UNIVERSITY OF SÃO PAULO,
CAIXA POSTAL 66281, 5311-970 SÃO PAULO, BRAZIL

E-mail: apaduaff@ig.com.br

(Received November 20, 2002)