

J. Schröder

Delannoy and tetrahedral numbers

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 48 (2007), No. 3, 389--394

Persistent URL: <http://dml.cz/dmlcz/119667>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2007

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Delannoy and tetrahedral numbers

JOACHIM SCHRÖDER

*Abstract.* We establish an identity between Delannoy numbers and tetrahedral numbers of arbitrary dimension.

*Keywords:* Delannoy numbers, tetrahedral numbers, king’s walk, coordination number, crystal ball

*Classification:* Primary 05A10; Secondary 05A15, 05A19

The numbers  $D(n, m)$  of Henri Auguste Delannoy (1833–1915), [4], count the lattice paths in  $\mathbb{Z} \times \mathbb{Z}$  from  $(0, 0)$  to  $(m, n)$  with set of permitted steps  $\{(0, 1), (1, 0), (1, 1)\}$ , i.e. north, east and north-east steps. They can conveniently be described as minimal king walks from the bottom left corner to the upper right corner on a  $m \times n$  chess board (see Figure 3 for  $m = n = 2$ ). It is known since the times of Delannoy that

$$(1) \quad D(n, m) = \sum_{\nu=0}^m \binom{m}{\nu} \binom{n+\nu}{m} = \sum_{\nu=0}^{\min\{m,n\}} 2^\nu \binom{m}{\nu} \binom{n}{\nu}.$$

Tetrahedral numbers have, by definition, a simple geometric meaning, too. In two dimensions they are the number of lattice points in an equilateral triangle, correspondingly in three dimension they count the number of points in a regular tetrahedron, see Figure 1. Generalization to higher dimensions is apparent and involves the  $d$ -dimensional simplex (hypertetrahedron). Their determination is even easier than the determination of Delannoy numbers, because they are built up inductively. If  $T_d(n)$  denotes the number of points with integer coordinates in the  $d$ -dimensional hypertetrahedron of edge length<sup>1</sup>  $n - 1$ , then  $T_1(n) = n$  and  $T_{d+1}(n) = \sum_{v=1}^n T_d(v)$ . It happens that  $T_1(n) = \binom{n}{1}$  and the triangular number

---

I wish to thank Prof Preuß and the Mathematics Department of the Free University for their hospitality during my stay in Berlin.

web: [www16.brinkster.com/jodis](http://www16.brinkster.com/jodis)

<sup>1</sup>The edge length is the number of intervals created by  $n$  equidistant points, where start and end point of the edge carries a point. It is naturally 1 less than the number of points.




FIGURE 1: An example of the simplest 3-dimensional polytop, a discrete tetrahedron or 3-simplex, of edge length 3.

$T_2(n)$  equals  $\frac{n(n+1)}{2} = \binom{n+1}{2}$ . The discrete antiderivative of  $\binom{x}{v}$  is  $\binom{x}{v+1}$ , i.e.  $\Delta \binom{x}{v+1} = \binom{x}{v}$ . Hence discrete integration can be used to show

$$(2) \quad T_d(n) = \binom{n+d-1}{d}.$$

In order to establish a link between Delannoy and tetrahedral numbers, we will need a result about crystal balls, which was first discovered by Vassilev & Atanassov [10].

- Definition 1.** (1) Let  $\mathbf{x} = (x_1, x_2, \dots, x_d) \in \mathbb{Z}^d$ ,  $d \in \mathbb{N}$ . The  $L^1$ -norm  $|\mathbf{x}|_1$  of  $\mathbf{x}$  is defined by  $|\mathbf{x}|_1 := \sum |x_i|$ .
- (2)  $\mathcal{S}_d(n) := \{\mathbf{x} \mid \mathbf{x} \in \mathbb{Z}^d \text{ and } |\mathbf{x}|_1 = n\}$  is called  $d - 1$ -dimensional crystal sphere of radius  $n$ . We set  $S_d(n) := |\mathcal{S}_d(n)|$ . The sequence  $(S_d(n))_{n \in \mathbb{N}}$  is called coordination-sequence (or -numbers). The union  $\bigcup_{\nu=0}^n \mathcal{S}_d(\nu) =: \mathcal{G}_d(n)$  is called  $d$ -dimensional crystal ball of radius  $n$ , see Figure 2. We put  $|\mathcal{G}_d(n)| =: G_d(n) = \sum_{\nu=0}^n S_d(\nu)$ .
- (3) If  $f : D \rightarrow \mathbb{Z}$ ,  $D \subseteq \mathbb{Z}$ , is a function then the (forward) difference operator  $\Delta$  is defined by  $\Delta f(n) = f(n+1) - f(n)$ .  $\Delta^m$  is defined by  $\Delta^1 = \Delta$  and  $\Delta^m = \Delta \circ \Delta^{m-1}$ .

**Theorem 2** ([10], [8]).

$$D(n, m) = \sum_{\mu=0}^m S_n(\mu) = G_n(m),$$

see Figure 3.

PROOF: We will follow the proof in Schröder [8], because it is considerably shorter. The GF of the Delannoy numbers  $D(n, m)$  is known to be (cf. [9])

$$\sum_{n,m \geq 0} D(n, m) x^n y^m = \frac{1}{1 - x - y - xy}.$$




FIGURE 2: The shape of a crystal ball in 3 dimensions, also called regular octahedron (the discrete points are not drawn).

We have

$$\begin{aligned} \frac{1}{1-x-y-xy} &= \frac{1}{1-y} \frac{1}{1-x\frac{1+y}{1-y}} = \frac{1}{1-y} \sum_{n \geq 0} \left(\frac{1+y}{1-y}\right)^n x^n \\ &= \frac{1}{1-y} \sum_{n,m \geq 0} S_n(m) y^m x^n = \sum_{n,m \geq 0} \sum_{\mu=0}^m S_n(\mu) y^\mu x^n = \sum_{n,m \geq 0} G_n(m) y^m x^n. \end{aligned}$$

Indeed, Conway & Sloane show in [3, p. 9, Equation (16)], that

$$S_d(n) = \sum_{k=0}^d \binom{d}{k} \binom{n+d-k-1}{d-1} \quad (= \sum_{k=0}^d \binom{d}{k} \binom{n+k-1}{d-1})$$

is the coordination number of distance  $n$  in  $\mathbb{Z}^d$  and their generating function is

$$\sum_{n \geq 0} S_d(n) y^n = \left(\frac{1+y}{1-y}\right)^d.$$




FIGURE 3:  $D(2, 2) = 13$  and  $G_2(2) = 13$ .

See also [6, p. 4, Aufg. 29]. □

We are ready to state and prove the main theorem. The idea is to decompose a crystal ball into a number of hypertetrahedra. For instance, if we look at Figure 2, we can see that the top pyramid consists of 4 tetrahedra of type as depicted in Figure 1. Overall  $\mathcal{G}_3(n)$  is composed out of 8 tetrahedra. Unfortunately these tetrahedra overlap. They have some faces and edges in common. Therefore, to get the correct number of points, we have to apply the principle of inclusion-exclusion. For instance, if we stay with Figure 1 and Figure 2, the first approximation to  $G_3(3)$  is obtained by taking 8 times the number of lattice points in Figure 1. Then we have to subtract 12 times the number of points of a tetrahedral face, because 12 faces are common to 2 tetrahedra, add 6 times the number of points of a tetrahedral edge, because 6 edges in the coordinate axes' are contained in 4 tetrahedra and finally subtract 1 for the point in the center, which is common to all tetrahedra. We have shown  $D(3, 4-1) = G_3(4-1) = 8\binom{4+2}{3} - 12\binom{4+1}{2} + 6\binom{4}{1} - 1 = 8 \times 20 - 12 \times 10 + 6 \times 4 - 1 = 63$  and more generally

$$D(3, n - 1) = G_3(n - 1) = 8\binom{n + 2}{3} - 12\binom{n + 1}{2} + 6\binom{n}{1} - 1,$$

see Equation 2.

**Theorem 3.**

$$(3) \quad D(n, m) = \sum_{v=0}^m (-1)^{m-v} 2^v \binom{m}{v} T_v(n + 1) = \sum_{v=0}^m (-1)^{m-v} 2^v \binom{m}{v} \binom{n + v}{v}$$

$$(4) \quad D(n, m) = \Delta^m 2^x \binom{n + x}{x} \Big|_{x=0} = \Delta^m 2^x T_x(n + 1) \Big|_{x=0}$$

$$(5) \quad 2^m T_m(n + 1) = 2^m \binom{n + m}{m} = \sum_{v=0}^m \binom{m}{v} D(n, v)$$

$$(6) \quad 2^m \binom{n}{m} = \sum_{v=0}^m (-1)^{m-v} \binom{m}{v} D(n, v) = \Delta^m D(n, x) \Big|_{x=0}$$

PROOF: Equations 3, 4 and 5 are equivalent via binomial inversion and difference formula. Binomial inversion again shows the (known) equivalence of Equation 1 and Equation 6, which was added for completeness. In order to prove Equation 3, we have to determine the number of simplices with a given, common sub-simplex. The easiest method might be to use a linear scheme in which we record the quadrant of the point and the varying coordinates, as in Schröder [7]. Given a  $d$ -dimensional crystal ball  $\mathcal{G}_d$ , let  $e = e_1e_2e_3 \dots e_d$  be a finite sequence, where  $e_i \in \{+, 0, -\}$ . To every point  $p = (x_1, x_2, \dots, x_d) \in \mathcal{G}_d$  we assign a sequence  $e$  by

$$e_i := \begin{cases} + & \text{if } x_i > 0 \\ - & \text{if } x_i < 0 \\ 0 & \text{if } x_i = 0 \end{cases}$$

Vice versa, every  $e$  defines a sub-simplex of  $\mathcal{G}_d$ , modulo size. An entry 0 in  $e$  has a special meaning, because it indicates a set of points which are common to more than 1 sub-simplex. For instance, in 3 dimensions,  $e = + - +$  stands for the 3-simplex which lies in the octant with positive x- and z- coordinates and negative y- coordinate.  $e = +0+$  describes the sub-simplex (triangle) with positive x- and z- coordinates and vanishing y- coordinate. It is the common face of  $+++$  and  $+ - +$ , see Figure 2. In  $d$  dimensions, the first approximation to  $G_d(n)$  is  $2^d T_d(n + 1)$ , because there are  $2^d$  different  $+, -$  sequences of length  $d$ , i.e.  $\mathcal{G}_d$  is composed out of  $2^d$   $d$ -simplices. We have to subtract points in the common faces to get the second approximation. There are  $\binom{d}{1}$  possibilities to insert 0 in a sequence of length  $d$  and  $2^{d-1}$  possibilities to fill the remaining places with  $+$  and  $-$ . Each case accounts for  $T_{d-1}(n + 1)$  points. In the next step we have to add  $2^{d-2} \binom{d}{2} T_{d-2}(n + 1)$  points, etc. Eventually we arrive at

$$D(n, d) = \sum_{v=0}^d (-1)^v 2^{d-v} \binom{d}{v} T_{d-v}(n + 1) = \sum_{v=0}^d (-1)^v 2^{d-v} \binom{d}{v} \binom{n + d - v}{d - v}.$$

Substitution  $d - v \rightarrow v$  produces Equation 3. □

### REFERENCES

- [1] Aigner M., *Diskrete Mathematik*, Vieweg, Braunschweig, 1993.
- [2] Banderier C., Schwer S.R., *Why Delannoy numbers?*, J. Statist. Plann. Inference **135** (2005), no. 1, 40–54; CoRR math.CO/0411128: (2004).
- [3] Conway J.H., Sloane N.J.A., *Low dimensional lattices VII: Coordination sequences*, Proc. Royal Soc. London Ser. A 453 (1997), no. 1966, 2369–2389; [citeseer.ist.psu.edu/article/conway96lowdimensional.html](http://citeseer.ist.psu.edu/article/conway96lowdimensional.html).
- [4] Delannoy H., *Emploi de l'échiquier pour la résolution de certains problèmes de probabilités*, Association Francaise pour l'Avancement des Sciences, Bordeaux XXIV (1895), 70–90.
- [5] *Handbook of Discrete and Combinatorial Mathematics*, Eds. K.H. Rosen et al., CRC Press, Boca Raton etc., 2000.

- [6] Pólya G., Szegő G., *Aufgaben und Lehrsätze aus der Analysis, I*, Springer, Berlin, 1970.
- [7] Schröder J., *Filling boxes densely and disjointly*, Comment. Math. Univ. Carolin. **44** (2003), no. 1, 187–196.
- [8] Schröder J., *Generalized Schröder numbers and the rotation principle*, preprint, 2007.
- [9] Stanley R.P., *Enumerative Combinatorics*, Vol. 2, Cambridge University Press, Cambridge, 1999.
- [10] Vassilev M., Atanassov K., *On Delan[n]oy numbers*, Annuaire Univ. Sofia Fac. Math. Inform. **81** (1987), 153–162.

DEPARTEMENT VAN WISKUNDE, UNIVERSITEIT VAN DIE VRYSTAAT, POSBUS 339, BLOEM-FONTEIN 9300, SUID AFRIKA

*E-mail:* schroderjd@qwa.uovs.ac.za

(Received February 8, 2007, revised May 21, 2007)