

Takashi Noiri

Weak-continuity and closed graphs

Časopis pro pěstování matematiky, Vol. 101 (1976), No. 4, 379--382

Persistent URL: <http://dml.cz/dmlcz/117936>

Terms of use:

© Institute of Mathematics AS CR, 1976

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

WEAK-CONTINUITY AND CLOSED GRAPHS

TAKASHI NOIRI, Yatsushiro

(Received October 9, 1975)

I. INTRODUCTION

The concept of weak-continuity was first introduced by N. LEVINE [4]. In 1968, M. K. SINGAL and A. R. SINGAL [7] defined almost-continuous functions and showed that every continuous function is almost-continuous and every almost-continuous function is weakly-continuous, but the converses are not necessarily true in general. Recently, P. E. LONG and L. L. HERRINGTON [6] have obtained several properties concerning almost-continuous functions and have given two sufficient conditions for almost-continuous functions to be continuous. The purpose of the present note is to give some sufficient conditions for weakly-continuous functions to be continuous.

II. DEFINITIONS

Let S be a subset of a topological space X . The closure of S and the interior of S are denoted by $\text{Cl}_X(S)$ and $\text{Int}_X(S)$, respectively. Throughout this note, X and Y denote topological spaces, and by $f : X \rightarrow Y$ we represent a function f of a space X into a space Y .

Definition 1. A function $f : X \rightarrow Y$ is said to be *almost-continuous* [7] (resp. *weakly-continuous* [4]) if for each point $x \in X$ and each open set $V \subset Y$ containing $f(x)$, there exists an open set $U \subset X$ containing x such that $f(U) \subset \text{Int}_Y(\text{Cl}_Y(V))$ (resp. $f(U) \subset \text{Cl}_Y(V)$).

Definition 2. A subset S of a space X is said to be *N -closed relative to X* (briefly *N -closed*) [1] if for each cover $\{U_\alpha \mid \alpha \in \mathcal{A}\}$ of S by open sets of X , there exists a finite subfamily $\mathcal{A}_0 \subset \mathcal{A}$ such that

$$S \subset \bigcup \{ \text{Cl}_X(U_\alpha) \mid \alpha \in \mathcal{A}_0 \}.$$

Definition 3. A space X is said to be *rim-compact* [8, p. 276] if each point of X has a base of neighborhoods with compact frontiers.

III. WEAK-CONTINUITY AND CLOSED GRAPHS

It is well known that if $f : X \rightarrow Y$ is continuous and Y is Hausdorff, then the graph $G(f)$ is closed in the product space $X \times Y$. P. E. Long and L. L. Herrington showed that "continuous" in this result can be replaced by "almost-continuous" [6, Theorem 9]. Moreover, we shall show that "almost-continuous" can be replaced by "weakly-continuous".

Theorem 1. *If $f : X \rightarrow Y$ is weakly-continuous and Y is Hausdorff, then f has the following property:*

(P) *For each $(x, y) \notin G(f)$, there exist open sets $U \subset X$ and $V \subset Y$ containing x and y , respectively, such that $f(U) \cap \text{Int}_Y(\text{Cl}_Y(V)) = \emptyset$.*

Proof. Let $(x, y) \notin G(f)$, then $y \neq f(x)$. Since Y is Hausdorff, there exist disjoint open sets V and W containing y and $f(x)$, respectively. Thus, we have $\text{Int}_Y(\text{Cl}_Y(V)) \cap \text{Cl}_Y(W) = \emptyset$. Since f is weakly-continuous, there exists an open set $U \subset X$ containing x such that $f(U) \subset \text{Cl}_Y(W)$. Therefore, we obtain $f(U) \cap \text{Int}_Y(\text{Cl}_Y(V)) = \emptyset$.

Remark 1. It is obvious that if a function has the property (P), then the graph is closed. The converse is not necessarily true, however, as the following example due to P. KOSTYRKO [3] shows.

Example 1. Let X and Y be the sets of positive integers. Let X have the discrete topology, Y have the cofinite topology and $f : X \rightarrow Y$ be the identity mapping. Then, although $G(f)$ is closed, f does not hold the property (P).

Corollary 1. *If $f : X \rightarrow Y$ is weakly-continuous and Y is Hausdorff, then $G(f)$ is closed.*

R. V. FULLER showed that if $f : X \rightarrow Y$ has the closed graph, then the inverse image $f^{-1}(K)$ of each compact set K of Y is closed in X [2, Theorem 3.6]. We shall obtain an analogous result to this theorem.

Theorem 2. *If $f : X \rightarrow Y$ has the property (P), then the inverse image $f^{-1}(K)$ of each N -closed set K of Y is closed in X .*

Proof. Assume that there exists a N -closed set $K \subset Y$ such that $f^{-1}(K)$ is not closed in X . Then, there exists a point $x \in \text{Cl}_X(f^{-1}(K)) - f^{-1}(K)$. Since $f(x) \notin K$, for each $y \in K$ we have $(x, y) \notin G(f)$. Therefore, there exist open sets $U_y(x) \subset X$ and $V(y) \subset Y$ containing x and y , respectively, such that $f(U_y(x)) \cap \text{Int}_Y(\text{Cl}_Y(V(y))) = \emptyset$. The family $\{V(y) \mid y \in K\}$ is a cover of K by open sets of Y . Since K is N -closed, there exist a finite number of points y_1, y_2, \dots, y_n in K such that $K \subset \bigcup_{j=1}^n \text{Int}_Y(\text{Cl}_Y(V(y_j)))$. Now, put $U = \bigcap_{j=1}^n U_{y_j}(x)$. Then we obtain $f(U) \cap K = \emptyset$. On the other hand, since $x \in \text{Cl}_X(f^{-1}(K))$, we have $f(U) \cap K \neq \emptyset$ because U is an open set containing x . This is a contradiction.

Remark 2. The converse to Theorem 2 is not always true, as Example 1 shows.

Corollary 2. *Let Y be a Hausdorff space such that every closed set is N -closed. If $f : X \rightarrow Y$ is weakly-continuous, then it is continuous.*

Proof. This is an immediate consequence of Theorem 1 and Theorem 2.

In [6, Theorem 7], it is shown that if Y is a rim-compact space and $f : X \rightarrow Y$ is an almost-continuous function with the closed graph, then f is continuous. We shall show that “almost-continuous” in this theorem can be replaced by “weakly-continuous”.

Theorem 3. *If Y is a rim-compact space and $f : X \rightarrow Y$ is a weakly-continuous function with the closed graph, then f is continuous.*

Proof. Let $x \in X$ and V be any open set of Y containing $f(x)$. Since Y is rim-compact, there exists an open set $W \subset Y$ such that $f(x) \in W \subset V$ and the frontier $\text{Fr}(W)$ is compact. It is obvious that $f(x) \notin \text{Fr}(W)$. Thus, for each $y \in \text{Fr}(W)$, we have $(x, y) \notin G(f)$. Since $G(f)$ is closed, there exist open sets $U_y(x) \subset X$ and $V(y) \subset Y$ containing x and y , respectively, such that $f(U_y(x)) \cap V(y) = \emptyset$. The family $\{V(y) \mid y \in \text{Fr}(W)\}$ is a cover of $\text{Fr}(W)$ by open sets of Y . Since $\text{Fr}(W)$ is compact, there exist a finite number of points y_1, y_2, \dots, y_n in $\text{Fr}(W)$ such that $\bigcup_{j=1}^n V(y_j) \supset \text{Fr}(W)$. Now, since f is weakly-continuous, there exists an open set $U_0 \subset X$ containing x such that $f(U_0) \subset \text{Cl}_Y(W)$. Put $U = U_0 \cap [\bigcap_{j=1}^n U_{y_j}(x)]$, then U is an open set containing x such that

$$f(U) \cap (Y - W) = f(U) \cap \text{Fr}(W) \subset \bigcup_{j=1}^n f(U) \cap V(y_j) \subset \bigcup_{j=1}^n f(U_{y_j}(x)) \cap V(y_j) = \emptyset.$$

This shows that $f(U) \subset V$ and hence f is continuous.

Theorem 4. *Every rim-compact Hausdorff space is regular.*

Proof. This proof is similar to that of Theorem 3.

Corollary 3. *If Y is rim-compact Hausdorff and $f : X \rightarrow Y$ is weakly-continuous, then f is continuous.*

Proof. This follows immediately from [4, Theorem 2].

In [6, Theorem 8], it is shown that if f is an almost-continuous function of a first countable space into a countably compact Hausdorff space, then f is continuous. The following theorem shows that “almost-continuous” in this result can be replaced by “weakly-continuous”.

Theorem 5. *Let X be a first countable space and Y a countably compact Hausdorff space. If $f : X \rightarrow Y$ is weakly-continuous, then f is continuous.*

Proof. This is an immediate consequence of Corollary 1 and [5, Theorem 2].

References

- [1] *D. Carnahan*: Locally nearly-compact spaces, *Boll. Un. Mat. Ital.* (4), 6 (1972), 146–153.
- [2] *R. V. Fuller*: Relations among continuous and various non-continuous functions, *Pacific J. Math.*, 25 (1968), 495–509.
- [3] *P. Kostyrko*: A note on the functions with closed graphs, *Časopis Pěst. Mat.*, 94 (1969), 202–205.
- [4] *N. Levine*: A decomposition of continuity in topological spaces, *Amer. Math. Monthly*, 68 (1961), 44–46.
- [5] *P. E. Long*: Functions with closed graphs, *Amer. Math. Monthly*, 76 (1969), 930–932.
- [6] *P. E. Long* and *L. L. Herrington*: Properties of almost-continuous functions, *Boll. Un. Mat. Ital.* (4), 10 (1974), 336–342.
- [7] *M. K. Singal* and *A. R. Singal*: Almost-continuous mappings, *Yokohama Math. J.*, 16 (1968), 63–73.
- [8] *S. Willard*: *General Topology*, Addison-Wesley, Reading, 1970.

Author's address: Department of Mathematics, Yatsushiro College of Technology, Yatsushiro, Kumamoto, 866 Japan.