

## WEIGHTS AND $L \log L$

A. CARBERY, S.-Y. A. CHANG AND J. GARNETT

*In memory of Irving L. Glicksberg*  
 April 1, 1925–November 30, 1983

**Let  $\omega(x)$  be a positive locally integrable weight on  $[0, 1]$ . Discussed are conditions on  $\omega$  necessary and sufficient for the (dyadic) Hardy-Littlewood maximal function to map  $L \log L(\omega dx)$  into  $L^1(\omega dx)$  or into weak  $L^1$ .**

**1. Introduction.** Let  $Mf$  denote the (dyadic) Hardy-Littlewood maximal function of  $f$ , for  $f$  locally integrable on  $\mathbf{R}^n$ . That is,

$$Mf(x) = \sup_{x \in Q} \frac{1}{|Q|} \int_Q |f|,$$

the sup being taken over all dyadic cubes in  $\mathbf{R}^n$  containing  $x$ . It is well-known that  $M$  is a bounded operator from  $L^p(\mathbf{R}^n)$  to  $L^p(\mathbf{R}^n)$  when  $p > 1$ , takes  $L^1$  to weak  $L^1$ , and for functions  $f$  supported in a dyadic cube  $Q_0$  satisfies

$$\int_{Q_0} Mf \leq C \int_{Q_0} |f| \log^+ |f| + C|Q_0|.$$

More recently, Muckenhoupt and others have studied the behaviour of  $M$  when the  $L^p$  spaces with respect to Lebesgue measure are replaced by those with respect to the measure  $\omega(x) dx$ ,  $\omega \in L^1_{\text{loc}}$ . A nonnegative locally integrable function  $\omega$  is said to be in Muckenhoupt's (dyadic)  $A_p$  class for  $1 \leq p < \infty$  if

$$(1.1) \quad \sup_{Q \text{ dyadic}} \left\| \frac{\omega_Q}{\omega(x)} \right\|_{L^q(\omega dx/\omega(Q))} = A_p(\omega) = A_p < \infty.$$

Here  $1/p + 1/q = 1$ ,  $\omega_Q$  is the average  $(1/|Q|) \int_Q \omega$  of  $\omega$  over  $Q$  and  $\omega(Q) = \int_Q \omega$ . (More generally, if  $E$  is a measurable set in  $\mathbf{R}^n$  we denote  $\int_E \omega$  by  $\omega(E)$ .) In [3], Muckenhoupt proved that given  $\omega \in L^1_{\text{loc}}$  there exists a constant  $C = C_{p,\omega,n}$  such that

$$(1.2) \quad \int |Mf|^p \omega \leq C \int |f|^p \omega$$

if and only if  $\omega$  is in the  $A_p$  class, and that  $\omega\{Mf > \lambda\} \leq (C/\lambda) \int |f| \omega$  if and only if  $\omega$  is in the  $A_1$  class. It would therefore seem reasonable to

expect that a necessary and sufficient condition on  $\omega \in L^1_{\text{loc}}$  for the existence of a  $C$  such that

$$(1.3) \quad \int_{Q_0} Mf\omega \leq C \int_{Q_0} |f| \log^+ |f| \omega + C\omega(Q_0)$$

should hold whenever  $\text{supp } f \subseteq Q_0$  would be obtained by taking the “exponential limit” as  $q \rightarrow \infty$  in (1.1). That is,  $\omega$  should satisfy what we shall call the  $A^*$  condition:

( $A^*$ ) There exists an  $\varepsilon > 0$  and a  $C > 1$  such that

$$\sup_{Q \text{ dyadic}} \int_Q \exp\left\{\frac{\varepsilon\omega_Q}{\omega(x)}\right\} \frac{\omega(x) dx}{\omega(Q)} \leq C.$$

Unfortunately, while  $A^*$  is necessary for (1.3) to hold, it is not sufficient (as we shall see in §4). In §3 we give a necessary and sufficient condition  $A^{**}$  that (1.3) hold. This condition may be realized as the “exponential limit” as  $q \rightarrow \infty$  of certain other expressions, which, for each  $p > 1$  are equivalent with the  $A_p$  condition. The equivalence breaks down in the limit, however, and  $A^*$  turns out to be necessary and sufficient only for  $M$  to take  $L \log L(\omega dx)$  to weak  $L^1(\omega dx)$ . In §4 we prove this fact, and we compare our rather complicated condition  $A^{**}$  to growth conditions on  $A_p(\omega)$  as  $p \downarrow 1$ , and see that none of these conditions is adequate to describe  $A^{**}$ . It would be of interest to find a more concise, easy-to-verify form of  $A^{**}$ .

Central to our proof of the equivalence of  $A^{**}$  with (1.3) is a new proof of the weighted  $L^p$  theorem which does not rely upon interpolation or upon the step “ $\omega \in A_p \Rightarrow \omega \in A_{p-\varepsilon}$  for some  $\varepsilon > 0$ ”, used both by Muckenhoupt and Coifman and C. Fefferman, [2]. E. Sawyer [4] and M. Christ and R. Fefferman [1] have recently given other such proofs, but neither proof has a counterpart in the  $L \log L$  setting. Section 2 is devoted to this new proof.

For simplicity, we choose to work with the dyadic Hardy-Littlewood maximal operator in this note. However, our results can easily be extended to those for the full maximal operator by standard techniques. Finally,  $C$  denotes a constant depending only possibly on the dimension (but not necessarily the same at each occurrence), and dependence of constants upon other quantities is indicated by subscripts.

**2. The weighted  $L^p$  theorem.** In this section we give a new proof of the weighted  $L^p$  theorem of Muckenhoupt. We shall need the insights it provides when we treat the case of  $L \log L$  in §3.

We first prove the elementary fact that if  $\omega \in A_p$ , then  $(1/\omega)^{q-1}$  satisfies the doubling condition.

LEMMA 2.1. *Suppose  $\omega \in A_p$ . Then there exists an  $a < 1$  such that whenever  $E$  is a measurable subset of a cube  $Q$  and  $|E| \leq \frac{1}{2}|Q|$ , we have*

$$\int_E \left( \frac{\omega_Q}{\omega} \right)^q \frac{\omega dx}{\omega(Q)} \leq a \int_Q \left( \frac{\omega_Q}{\omega} \right)^q \frac{\omega dx}{\omega(Q)}.$$

*Proof.* By Hölder's inequality we have

$$\omega(Q) \frac{|Q-E|}{|Q|} = \int_{Q-E} \omega_Q \leq \left( \int_{Q-E} \left( \frac{\omega_Q}{\omega} \right)^q \omega dx \right)^{1/q} \left( \int_Q \omega dx \right)^{1/p},$$

that is,

$$\omega(Q)^{1/q} \frac{|Q-E|}{|Q|} \leq \left( \int_{Q-E} \left( \frac{\omega_Q}{\omega} \right)^q \omega dx \right)^{1/q}.$$

Thus if  $\omega \in A_p$  and  $|E| \leq \frac{1}{2}|Q|$  we obtain

$$\frac{1}{2^q A_p^q} \int_Q \left( \frac{\omega_Q}{\omega} \right)^q \omega dx \leq \left( \frac{1}{2} \right)^q \omega(Q) \leq \int_{Q-E} \left( \frac{\omega_Q}{\omega} \right)^q \omega dx.$$

Hence

$$\int_E \left( \frac{\omega_Q}{\omega} \right)^q \omega dx \leq a \int_Q \left( \frac{\omega_Q}{\omega} \right)^q \omega dx,$$

with  $a = 1 - (2A_p)^{-q}$ . □

THEOREM 2.2. *Let  $p > 1$ . Then  $\omega \in A_p$  if and only if there exists a constant  $C_{p,\omega}$  such that*

$$\int |Mf|^p \omega \leq C_{p,\omega} \int |f|^p \omega \quad \text{for all } f \in L^p(\omega).$$

*Proof.* As in [3], setting  $f = \chi_Q \omega^{-1/p-1}$ , we see that (1.2) implies that  $\omega \in A_p$ . To see that  $A_p$  is sufficient for (1.2), we apply a Calderón-Zygmund decomposition to  $Mf$ , and choosing  $R_k = 2^{k(n+1)}$ , we write  $D_k = \{Mf > R_k\} = \cup_j Q_j^k$ , where the  $Q_j^k$  are the maximal dyadic cubes satisfying

$$2^n R_k \geq \frac{1}{|Q_j^k|} \int_{Q_j^k} |f| > R_k.$$

Notice that  $|Q_j^k \cap D_{k+1}| \leq \frac{1}{2}|Q_j^k|$  by our choice of  $R_k$ . Let  $E_k = D_k - D_{k+1}$ . Then

$$\begin{aligned}
[1 - 2^{-(n+1)p}] \sum_k R_k^p \omega(D_k) &\leq \int |Mf|^p \omega \leq 2^{(n+1)p} \sum_k R_k^p \omega(D_k) \\
&\leq 2^{(n+1)p} \sum_k R_k^{p-1} \sum_j \frac{1}{|Q_j^k|} \int_{Q_j^k} |f| \omega(Q_j^k) \\
&= 2^{(n+1)p} \sum_k R_k^{p-1} \sum_j \sum_{l=0}^{\infty} \int_{Q_j^k \cap E_{k+l}} |f| \omega_{Q_j^k} \\
&\leq 2^{(n+1)p} \sum_k R_k^{p-1} \sum_j \sum_{l=0}^{\infty} \left( \int_{Q_j^k \cap E_{k+l}} |f|^p \omega \, dx \right)^{1/p} \\
&\quad \cdot \left( \int_{Q_j^k \cap E_{k+l}} \left( \frac{\omega_{Q_j^k}}{\omega} \right)^q \omega \, dx \right)^{1/q}.
\end{aligned}$$

By repeated application of Lemma 2.1, we see that

$$\int_{Q_j^k \cap D_{k+l}} \frac{1}{\omega^q} \omega \, dx \leq a^l \int_{Q_j^k} \frac{1}{\omega^q} \omega \, dx.$$

Consequently,

$$\begin{aligned}
\int |Mf|^p \omega &\leq 2^{(n+1)p} \sum_k R_k^{p-1} \sum_j \sum_{l=0}^{\infty} \left( \int_{Q_j^k \cap E_{k+l}} |f|^p \omega \, dx \right)^{1/p} (a^l A_p^q \omega(q_j^k))^{1/q} \\
&\leq 2^{(n+1)p} \sum_k R_k^{p-1} \sum_{l=0}^{\infty} (a^l A_p^q)^{1/q} \left( \int_{E_{k+l}} |f|^p \omega \right)^{1/p} (\omega(D_k))^{1/q} \\
&= 2^{(n+1)p} A_p \sum_{l=0}^{\infty} a^{l/q} \sum_k \left( \int_{E_{k+l}} |f|^p \omega \right)^{1/p} (R_k^p \omega(D_k))^{1/q} \\
&\leq 2^{(n+1)p} A_p \sum_{l=0}^{\infty} a^{l/q} \left( \int_{\mathbb{R}^n} |f|^p \omega \right)^{1/p} \left( \sum_k R_k^p \omega(D_k) \right)^{1/q}.
\end{aligned}$$

We therefore obtain

$$\int |Mf|^p \omega \leq C_{p,\omega} \int |f|^p \omega$$

with

$$C_{p,\omega} = C_p A_p^p (1 - a^{1/q})^{-p} \leq C_p A_p^{p(q+1)}. \quad \square$$

**3. The  $L$  log  $L$  theorem.** In this section we give a necessary and sufficient condition on  $\omega$  for the maximal function to be bounded from  $L$  log  $L(\omega)$  to  $L^1(\omega)$  locally. By a Calderón-Zygmund decomposition of a dyadic cube  $Q$  we shall mean a collection of dyadic subcubes  $\{Q_j^k\}_{k \geq 0}$  of

$Q$  such that (a) in  $Q_j^k \cap \text{int } Q_l^k = \emptyset$  if  $j \neq l$ , (b) each  $Q_j^{k+1}$  is contained in some  $Q_l^k$ ,  $k \geq 0$ , and (c) the 0-th generation of cubes consists of the single cube  $Q$ . In such a situation, we write  $D_k = \bigcup_j Q_j^k$  and  $E_k = D_k - D_{k+1}$ .

**THEOREM 3.1.** *The following conditions on a locally integrable  $\omega$  are equivalent:*

(i) *There exists a constant  $C_\omega$  such that whenever  $f$  is supported in a dyadic cube  $Q$ , then*

$$\int_Q Mf\omega \leq C_\omega \int_Q |f| \log^+ |f| \omega + C_\omega \omega(Q).$$

(ii) *(A\*\*) There exists an  $\varepsilon > 0$  and a  $C_\omega$  such that whenever  $Q$  is a dyadic cube and  $\{Q_j^k\}$  is a Calderón-Zygmund decomposition of  $Q$ , then*

$$\int_Q \exp \left\{ \varepsilon \sum_{k,j} \frac{\omega(Q_j^k \cap E_k)}{\omega(x)|Q_j^k|} \chi_{Q_j^k}(x) \right\} \frac{\omega dx}{\omega(Q)} \leq C_\omega.$$

(iii) *There exists an  $\varepsilon > 0$  and a  $C_\omega$  such that whenever  $T$  is a positive linear operator satisfying  $|Tf(x)| \leq Mf(x)$ , and  $Q$  is a dyadic cube, then*

$$\int_Q \exp \left\{ \varepsilon \frac{T^*(\chi_Q \omega)(x)}{\omega(x)} \right\} \frac{\omega dx}{\omega(Q)} \leq C_\omega.$$

(Here,  $T^*$  is the adjoint of  $T$  with respect to  $L^2(dx)$ , i.e.  $\int (Tf)g dx = \int f(T^*g) dx$ .)

*Proof.* We shall show (i)  $\Rightarrow$  (iii)  $\Rightarrow$  (ii)  $\Rightarrow$  (i).

(i)  $\Rightarrow$  (iii). Suppose  $Q$  is a dyadic cube and  $T$  is a positive linear operator with  $|Tf(x)| \leq Mf(x)$ . Let  $E = \{x \in Q | T^*(\chi_Q \omega)(x) > \omega(x)\}$ . Let

$$g = e^{\varepsilon(T^*(\chi_Q \omega)/\omega)} \cdot \frac{\omega}{T^*(\chi_Q \omega)} \chi_E,$$

for a small  $\varepsilon$  to be determined later. Then

$$\begin{aligned} & \int_E \exp \left\{ \varepsilon \frac{T^*(\chi_Q \omega)}{\omega} \right\} \omega dx \\ &= \int g \frac{T^*(\chi_Q \omega)}{\omega} \omega dx = \int_Q Tg\omega dx \leq \int_Q Mgw dx \\ &\leq C_\omega \int_Q g \log^+ g \omega dx + C_\omega \omega(Q) \quad (\text{by (i)}) \\ &\leq C_\omega \int_E e^{\varepsilon(T^*(\chi_Q \omega)/\omega)} \frac{\omega}{T^*(\chi_Q \omega)} \varepsilon \frac{T^*(\chi_Q \omega)}{\omega} \omega dx + C_\omega \omega(Q). \end{aligned}$$

If we now choose  $\varepsilon$  such that  $C_\omega \varepsilon < 1$  we obtain (iii).

(iii)  $\Rightarrow$  (ii). If  $Q$  is a dyadic cube and  $\{Q_j^k\}$  is a Calderón-Zygmund decomposition of  $Q$ , let

$$Tf(x) = \sum_k \sum_j \frac{1}{|Q_j^k|} \int_{Q_j^k} f(t) dt \chi_{Q_j^k \cap E_k}(x).$$

Then clearly  $|Tf(x)| \leq Mf(x)$  and

$$T^*(\chi_Q \omega)(x) = \sum_{k,j} \frac{\omega(Q_j^k \cap E_k)}{|Q_j^k|} \chi_{Q_j^k}(x).$$

Thus (ii) is a special case of (iii).

(ii)  $\Rightarrow$  (i). We assume that  $f$  is supported inside a dyadic cube  $Q$  and proceed as in the positive part of Theorem 2.2 with the same Calderón-Zygmund decomposition used there. Then the collection  $\{Q_j^k \cap Q\}_{k \geq 1}$  together with  $Q$  forms a Calderón-Zygmund decomposition of  $Q$  in the sense above. So we have

$$\begin{aligned} \int_Q Mf \omega dx &\leq C \sum_{k=1}^{\infty} R_k \omega(E_k \cap Q) + C\omega(Q) \\ &\leq C \sum_{k=1}^{\infty} \sum_j \frac{1}{|Q_j^k|} \int_{Q_j^k} |f| \omega(Q_j^k \cap Q \cap E_k) + C\omega(Q) \\ &\leq C \int_Q |f(x)| \left\{ \sum_{k=1}^{\infty} \sum_j \frac{\omega(Q_j^k \cap Q \cap E_k)}{|Q_j^k \cap Q|} \chi_{Q_j^k \cap Q}(x) \right\} dx + C\omega(Q) \\ &\leq C \int_Q \frac{|f|}{\varepsilon} \log^+ \frac{|f|}{\varepsilon} \omega dx \\ &\quad + C \int_Q \exp \left\{ \varepsilon \sum_{k,j} \frac{\omega(Q_j^k \cap Q \cap E_k)}{\omega(x) |Q_j^k \cap Q|} \chi_{Q_j^k \cap Q}(x) \right\} \omega dx + C\omega(Q) \\ &\leq C_\omega \int_Q |f| \log^+ |f| \omega dx + C_\omega \omega(Q), \end{aligned}$$

by Young's inequality and (ii). We have finished the proof of the theorem.  $\square$

REMARK. We may re-work the above proof in the case of  $L^p$  to obtain a condition similar to  $A^{**}$  (involving Calderón-Zygmund decompositions

of cubes, but with the exponential replaced by an  $L^q$ -norm) which would necessarily be equivalent to  $A_p$ . As noted in the introduction, the equivalence fails in the  $L \log L$  case and  $A^{**}$  turns out to be strictly stronger than  $A^*$ . See the following section.

**4. A comparison of  $A^{**}$  with  $A^*$  and related conditions.** While it is clear that  $A_1 \Rightarrow A^{**} \Rightarrow A^*$ , we give in this section some examples to show that the reverse implications do not hold. We show the equivalence between  $A^*$  and the growth condition  $A_p = O(1/(p-1))$ , and show that no growth condition  $A_p = O((1/(p-1))^\beta)$  with  $\beta > 0$  is sufficient to imply  $A^{**}$ . Finally, although an inequality of the form  $\|Mf\|_{M^p(\omega)} \leq (C/(p-1))\|f\|_{L^p(\omega)}$  is sufficient for  $\omega$  to be in  $A^{**}$  by Yano's theorem [5], we give an example to show that such an inequality is not necessary.

If  $0 < \alpha < \infty$  and any of the equivalent conditions of Proposition 4.1 below is satisfied, we say that  $\omega$  belongs to  $A_\alpha^*$ .

**PROPOSITION 4.1.** *Let  $0 < \alpha < \infty$ . Then the following conditions on a locally integrable  $\omega$  are equivalent:*

(i) *There exists an  $\varepsilon > 0$ , and a  $C_\omega$  such that*

$$\sup_Q \int_Q \exp\left(\frac{\varepsilon\omega_Q}{\omega}\right)^\alpha \frac{\omega dx}{\omega(Q)} = C_\omega < \infty.$$

(ii) *There exists a  $C_\omega$  such that*

$$\omega \{Mf > \lambda\} \leq C_\omega \int \frac{|f|}{\lambda} \left[1 + \log^+\left(\frac{|f|}{\lambda}\right)\right]^{1/\alpha} \omega.$$

(iii)  $A_p(\omega) = O((1/(p-1))^{1/\alpha})$  as  $p \downarrow 1$ .

*Proof.* We show the implications (i)  $\Rightarrow$  (ii)  $\Rightarrow$  (iii)  $\Rightarrow$  (i), and, for simplicity, treat only the case  $\alpha = 1$ .

(i)  $\Rightarrow$  (ii). Suppose  $\varepsilon > 0$  is such that

$$\sup_Q \int_Q \exp\left(\frac{\varepsilon\omega_Q}{\omega}\right) \frac{\omega dx}{\omega(Q)} \leq C_\omega.$$

By Young's inequality, we see that

$$\begin{aligned} (4.1) \quad \frac{1}{|Q|} \int_Q |f| &= \int_Q \frac{\omega_Q}{\omega} |f| \frac{\omega dx}{\omega(Q)} \\ &\leq C \int_Q \exp\left(\frac{\varepsilon\omega_Q}{\omega}\right) \frac{\omega dx}{\omega(Q)} + C_\varepsilon \int_Q |f| (1 + \log^+ |f|) \frac{\omega dx}{\omega(Q)}. \end{aligned}$$

By homogeneity it suffices to show that

$$\omega \{Mf > \lambda_0\} \leq \int |f|(1 + \log^+ |f|) \omega \, dx$$

for some fixed  $\lambda_0$ , which we take to be  $C_\omega C + C_\varepsilon$ . Now, by (4.1),  $\{Mf > \lambda_0\} = \cup Q_j$ , where the  $Q_j$  are disjoint dyadic cubes satisfying  $\int_{Q_j} |f|(1 + \log^+ |f|) \omega \, dx / \omega(Q) > 1$ . Hence

$$\omega \{Mf > \lambda_0\} = \omega(\cup Q_j) \leq \int |f|(1 + \log^+ |f|) \omega \, dx.$$

(ii)  $\Rightarrow$  (iii). Using the inequality  $a(1 + \log^+ a) \leq (C/(p-1))a^p$  (valid when  $a \geq 1/2$  and  $p > 1$ ) and combining it with the fact that if  $\omega \{Mf > \lambda\} \leq (A/\lambda^p) \int |f|^p \omega$ , then  $A_p(\omega) \leq C^p A^{1/p}$ , we see the result immediately.

(iii)  $\Rightarrow$  (i). Expanding out the exponential,

$$\begin{aligned} \int_Q \exp\left(\frac{\varepsilon \omega_Q}{\omega}\right) \frac{\omega \, dx}{\omega(Q)} &= \int_Q \sum_{k=0}^{\infty} \frac{1}{k!} \left(\frac{\varepsilon \omega_Q}{\omega}\right)^k \frac{\omega \, dx}{\omega(Q)} \\ &\leq 1 + \varepsilon + \sum_{k=2}^{\infty} \frac{\varepsilon^k}{k!} A_{k'}(\omega)^k \\ &\leq 1 + \varepsilon + \sum_{k=2}^{\infty} \frac{\varepsilon^k}{k!} C_\omega^k k^k \leq C \quad \text{if } \varepsilon \leq (2eC_\omega)^{-1}. \quad \square \end{aligned}$$

We remark here that it can also be shown that the condition  $A_\alpha^*$  is equivalent to the seemingly stronger condition

$$\sup_Q \int_Q \exp\left\{\frac{\varepsilon M(\omega \chi_Q)}{\omega}\right\}^\alpha \frac{\omega \, dx}{\omega(Q)} < \infty.$$

For  $j \in \mathbf{Z}$ , let  $I_j = [2^{-j-1}, 2^{-j})$ , and from now on we consider weights of the form

$$\omega(x) = \sum_{j \in \mathbf{Z}} \omega_j \chi_{I_j}(x).$$

We shall restrict ourselves to sequences satisfying  $\omega_k = \omega_0$ ,  $k \leq 0$ ,  $\omega_j \downarrow 0$  as  $j \rightarrow \infty$ , and  $\omega_j \leq C\omega_{j+1}$  for all  $j$ .

LEMMA 4.2. *Let  $\omega(x)$  be a weight of the above form.*

(a)  $\omega \in A_\alpha^* \Leftrightarrow \exists \varepsilon > 0, C_\omega$  such that

$$\sum_{k=j}^{\infty} \exp\left\{\varepsilon \left(\frac{\omega_j}{\omega_k}\right)\right\}^\alpha \frac{\omega_k}{2^k} \leq C_\omega \frac{\omega_j}{2^j}$$



(b)  $\omega \in A^{**} \Rightarrow \exists \varepsilon > 0, C_\omega$  such that

$$\sum_{k=0}^{\infty} \exp\left\{\frac{\varepsilon}{\omega_k} \sum_{l=0}^k \omega_l\right\} \frac{\omega_k}{2^k} \leq C_\omega.$$

*Proof.* (a) We need only check the  $A_\alpha^*$  condition on dyadic intervals of the form  $[0, 2^{-m})$ ,  $m \in \mathbf{Z}$ , since  $\omega$  is constant on all others. Also, because of the form of  $\omega$ , only the positive integers are relevant. On such an interval  $I = [0, 2^{-m})$ ,  $m \geq 0$ ,  $\omega(I) \approx \omega_m/2^m$  and  $\omega_I \approx \omega_m$ , and the expression for  $A_\alpha^*$  follows.

(b) Since  $(1/x) \int_0^x |f(t)| dt \leq CMf(x)$ ,  $\omega \in A^{**}$  implies that

$$\begin{aligned} \int_0^1 \frac{1}{x} \int_0^x |f(t)| dt \omega(x) dx &= \int_0^1 \left( \int_t^1 \frac{\omega(x)}{x} dx \right) |f(t)| dt \\ &\leq C_\omega \int_0^1 |f| \log^+ |f| \omega + C_\omega \end{aligned}$$

whenever  $\text{supp } f \subseteq [0, 1]$ . By duality, we obtain

$$\int_0^1 \exp\left\{\frac{\varepsilon}{\omega(t)} \int_t^1 \frac{\omega(x)}{x} dx\right\} \omega(t) dt \leq C_\omega$$

which reduces to the above expression. □

**PROPOSITION 4.3.** Let  $\omega_j = (j + 1)^{-\beta}$  where  $0 \leq \beta < \infty$ .

(a)  $\omega \in A_\alpha^* \Leftrightarrow \alpha\beta \leq 1$

(b)  $\omega \in A^{**} \Leftrightarrow \beta < 1$

(c)  $\omega \in A_1 \Leftrightarrow \beta = 0$ .

Consequently no two of the conditions  $A_1, A^{**}, A_1^* = A^*$  coincide.

*Proof.* (c) is obvious; so is (a) and the forward implication of (b) once we have applied the previous lemma. To prove the reverse implication of (b), we reverse the steps of part (b) of the lemma. Let

$$M_0 f(x) = \sup_{\substack{x \in I \\ |I| < x}} \frac{1}{|I|} \int_I |f(t)| dt,$$

and for  $n \geq 1$ ,

$$M_n f(x) = \sup_{\substack{x \in I \\ 2^{n-1}x \leq |I| < 2^n x}} \frac{1}{|I|} \int_I |f(t)| dt.$$

(Here, we are assuming that  $\text{supp } f \subseteq [0, 2^{-m})$  for some integer  $m$ ,  $\omega$  being constant on all other dyadic cubes.) Then  $Mf(x) \leq \sum_{n=0}^{\infty} M_n f(x)$ . Now

$$\begin{aligned} \int_0^{2^{-m}} M_0 f(x) \omega(x) dx &= \sum_{k=m}^{\infty} \omega_k \int_{I_k} M_0 f(x) dx \\ &\leq C \sum_{k=m}^{\infty} \omega_k \left( \int_{I_k} |f| \log^+ |f| dx + 2^{-k} \right) \\ &\leq C \int_0^{2^{-m}} |f| \log^+ |f| \omega dx + C\omega([0, 2^{-m})), \end{aligned}$$

by the unweighted  $L \log L$  theorem and the property  $\omega_{j+1} \leq \omega_j \leq C\omega_{j+1}$  of  $\omega$ . Now let  $\tilde{M}f(x) = \sum_{n=1}^{\infty} M_n f(x)$ ; notice that

$$M_n f(x) \leq \frac{1}{2^{n-1}x} \int_0^{2^{n+1}x} |f(t)| dt.$$

Hence,

$$\begin{aligned} \int_0^{2^{-m}} \tilde{M}f(x) \omega(x) dx &\leq \int_0^{2^{-m}} \left( \sum_{n=1}^{\infty} \frac{1}{2^{n-1}} \int_{t/2^{n+1}}^{2^{-m}} \frac{\omega(x)}{x} dx \right) |f(t)| dt \\ &\leq C_\varepsilon \int_0^{2^{-m}} |f| \log^+ |f| \omega dx \\ &\quad + C_\varepsilon \int_0^{2^{-m}} \exp\left\{ \frac{\varepsilon}{\omega(t)} \sum_{n=1}^{\infty} \frac{1}{2^{n-1}} \int_{t/2^{n+1}}^{2^{-m}} \frac{\omega(x)}{x} dx \right\} \omega(t) dt. \end{aligned}$$

With a little calculation, the reader may verify that the last term is dominated by  $\omega(0, 2^{-m})$  in the case  $\omega_j = (j+1)^{-\beta}$ ,  $0 < \beta < 1$ .  $\square$

**EXAMPLE 4.4.** For each  $\alpha$ ,  $1 < \alpha < \infty$ , there exists  $\omega \in A_\alpha^* - A^{**}$ .

*Proof.* Let  $\lambda = 2^{-1/\alpha}$ , and let  $\omega_j = \lambda^r$  when  $n_{r-1} < j \leq n_r$ . Here,  $n_r$  is an increasing sequence of positive integers,  $n_0 = 0$ ; let  $\Delta_r = n_r - n_{r-1} = \#\{j | \omega_j = \lambda^r\}$ , so that  $n_r = \Delta_1 + \cdots + \Delta_r$ .

Let  $0 = r_0 < r_1 < r_1 \cdots$  be an increasing sequence of positive integers ( $r_j = j$  will do) and for  $\sum_{j=1}^{l-1} r_j < r \leq \sum_{j=1}^l r_j$ , let  $\Delta_r = 2^{2r_1+2r_2+\cdots+2r_{l-1}+r_l+2-r}$ . Thus  $\{\Delta_r\}$  is a rearrangement of the geometric progression  $\{2^r\}$ , and clearly any set of  $j$  consecutive  $\Delta_r$ 's must contain one of size greater than or equal to  $2^{j+1}$ ; thus

$$(4.2) \quad n_{q+j} - n_q = \Delta_{q+1} + \cdots + \Delta_{q+j} \geq 2^{j+1} \quad \forall j, \forall q.$$

On the other hand,

$$\sum_{j=1}^{r_1+\cdots+r_k} \Delta_j = 4(2^{r_1+\cdots+r_k} - 1)$$

and therefore

$$\begin{aligned} \sum_{\substack{j=1 \\ j \neq r_1 + \dots + r_{k+1}}}^{r_1 + \dots + r_{k+1}} \Delta_j &= 4(2^{4_1 + \dots + r_{k+1}} - 1) - 2^{r_1 + \dots + r_{k+1} + 1} \\ &= 2^{r_1 + \dots + r_{k+1} + 1} - 4 = \Delta_{r_1 + \dots + r_{k+1}} - 4. \end{aligned}$$

Therefore

$$(4.3) \quad \Delta_{r_1 + \dots + r_{k+1}} \geq \sum_{\substack{j=1 \\ j \neq r_1 + \dots + r_{k+1}}}^{r_1 + \dots + r_{k+1}} \Delta_j \quad \forall k.$$

Now our sequence  $\omega_j$  satisfies the conditions of Lemma 4.2; we shall use (4.2) to prove that  $\omega \in A_\alpha^*$ , and (4.3) to prove that  $\omega \notin A^{**}$ .

To see that  $\omega \in A_\alpha^*$ , we suppose that  $n_{q-1} < j \leq n_q$  and compute the expression in Lemma 4.2.(a):

$$\begin{aligned} \sum_{k=j}^{\infty} \exp \varepsilon \left( \frac{\omega_j}{\omega_k} \right)^\alpha \frac{\omega_k}{2^k} / \frac{\omega_j}{2^j} &= \sum_{k=j}^{n_q} 2^{j-k} e^\varepsilon + \frac{2^j}{\omega_j} \sum_{p=q+1}^{\infty} \sum_{k=n_{p-1}+1}^{n_p} e^{\varepsilon 2^{p-q}} \frac{\omega_k}{2^k} \\ &\leq 2e^\varepsilon + 2^{n_q} \sum_{p=q+1}^{\infty} e^{\varepsilon 2^{p-q}} 2^{-n_{p-1}} \\ &\leq 2e^\varepsilon + \sum_{j=0}^{\infty} 2^{n_q - n_{q+j}} e^{\varepsilon 2^{j+1}} \\ &\leq C \quad \text{if } \varepsilon \leq \frac{\log 2}{2}, \quad \text{by (4.2).} \end{aligned}$$

If  $\omega$  were in  $A^{**}$ , Lemma 4.2.(b) would give the existence of  $C_\omega$  and  $\varepsilon$  such that

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{\omega_k}{2^k} \exp \left( \varepsilon \sum_{j=1}^k \frac{\omega_j}{\omega_k} \right) &= \sum_{p=1}^{\infty} \sum_{k=n_{p-1}+1}^{n_p} \frac{\lambda^p}{2^k} e^{\varepsilon(k-n_{p-1})} \exp \left\{ \varepsilon \sum_{q=1}^{p-1} \Delta_q \lambda^{q-p} \right\} \\ &= \sum_{p=1}^{\infty} \frac{\lambda^p}{2^{n_{p-1}}} e^{\varepsilon(\Delta_1 \lambda^{1-p} + \dots + \Delta_{p-1} \lambda^{-1})} \sum_{k=n_{p-1}+1}^{n_p} \frac{e^{\varepsilon(k-n_{p-1})}}{2^{k-n_{p-1}}} \\ &\leq C_\omega. \end{aligned}$$

For  $\varepsilon \leq (\log 2)/2$ ,

$$C_1 \leq \sum_{k=n_{p-1}+1}^{n_p} \frac{e^{\varepsilon(k-n_{p-1})}}{2^{k-n_{p-1}}} \leq C_2,$$

and so if  $\omega$  were in  $A^{**}$ , we would have that for all  $\varepsilon$  sufficiently small

$$\alpha_p(\varepsilon) = \frac{\lambda^p}{2^{n_{p-1}}} e^{\varepsilon(\Delta_1 \lambda^{1-p} + \dots + \Delta_{p-1} \lambda^{-1})}$$

satisfies  $\alpha_p(\varepsilon) < 1$  for all but finitely many  $p$ . We shall now show that for arbitrarily small  $\varepsilon$ ,  $\log \alpha_p(\varepsilon)$  can be nonnegative for infinitely many  $p$ , and so  $\omega$  cannot be in  $A^{**}$ . Observe that

$$\frac{\log \alpha_p(\varepsilon)}{\Delta_1 + \dots + \Delta_{p-1}} = \frac{\varepsilon(\Delta_1 \lambda^{1-p} + \dots + \Delta_{p-1} \lambda^{-1})}{\Delta_1 + \dots + \Delta_{p-1}} - \log 2 - \frac{p \log(1/\lambda)}{n_{p-1}},$$

and so if we choose  $p = r_1 + r_2 + \dots + r_{k+1} + 1$ , (4.3) implies that

$$\begin{aligned} \frac{\log \alpha_p(\varepsilon)}{\Delta_1 + \dots + \Delta_{p-1}} &\geq \frac{\varepsilon(\Delta_1 \lambda^{1-p} + \dots + \Delta_{p-1} \lambda^{-1})}{2\Delta_{r_1 + \dots + r_{k+1}}} - \log 2 - \frac{p \log(1/\lambda)}{n_{p-1}} \\ &\geq \frac{\varepsilon}{2} \lambda^{-r_{k+1}} - \log 2 - \frac{p \log(1/\lambda)}{n_{p-1}} \geq 0 \end{aligned}$$

if  $k = k(\varepsilon)$  is chosen sufficiently large. This completes the example.  $\square$

**EXAMPLE 4.5.** Let  $\omega_j = (j+1)^{-\beta}$ , with  $0 < \beta < 1$ . While  $\omega \in A^{**}$  by Proposition 4.3,  $\omega$  does not satisfy  $\int |Mf|^p \omega \leq C^p / (p-1)^p \int |f|^p \omega$  as  $p \downarrow 1$ .

*Proof.* Let  $g(x) = (\log(1/x))^{\beta/q}$  ( $1/p + 1/q = 1$ ). Then  $\|g\|_{L^q(\omega)} \cong 1$ , and

$$\|Mf\|_{L^p(\omega)} \geq C \sup \left\{ \left| \int (Tf)g \omega dx \right| : \|g\|_{L^q(\omega)} \leq 1 \right\}$$

where

$$Tf(x) = \frac{1}{x} \int_0^{2x} f(t) dt.$$

But

$$\int (Tf)g \omega dx = \int f(\tilde{T}g) \omega dx$$

where

$$\tilde{T}g(t) = \frac{1}{\omega(t)} \int_{t/2}^1 g(x) \omega(x) \frac{dx}{x} \geq Cq \left( \log \frac{2}{t} \right)^{\beta/q+1},$$

with our choice of  $g$ . Hence

$$(p-1)M_p \geq \frac{C}{q} \|T^*g\|_{L^q(\omega)} \geq C \left( \int_0^1 \left( \log \frac{2}{t} \right)^q dt \right)^{1/q},$$

(where  $M_p = \sup\{\|Mf\|_{L^p(\omega)} \mid \|f\|_{L^p(\omega)} \leq 1\}$ ), and, since  $\log(2/t) \notin L^\infty$ ,  $M_p$  is not  $O(1/(p-1))$  as  $p \downarrow 1$ .

## REFERENCES

- [1] F. M. Christ and R. Fefferman, *A note on weighted norm inequalities for the Hardy-Littlewood maximal operator*, PAMS, 3, **87** (1983), 447–448.
- [2] R. R. Coifman and C. Fefferman, *Weighted norm inequalities for maximal functions and singular integrals*, Studia Math., **51** (1974), 241–250.
- [3] B. Muckenhoupt, *Weighted norm inequalities for the Hardy maximal function*, TAMS, **165** (1972), 207–226.
- [4] E. Sawyer, *Two Weight Norm Inequalities for Certain Maximal and Integral Operators*, in Springer Lecture Notes in Math., **908** (1982), 102–127.
- [5] S. Yano, *An extrapolation theorem*, J. Math. Soc. Japan, **3** (1951), 296–305.

Received February 18, 1984. Research supported in part by NSF grants.

CALIFORNIA INSTITUTE OF TECHNOLOGY  
PASADENA, CA 91125

AND

UNIVERSITY OF CALIFORNIA, LOS ANGELES  
LOS ANGELES, CA 90024

