



RESEARCH

Open Access

Developing a provisional, international Minimal Dataset for Juvenile Dermatomyositis: for use in clinical practice to inform research

Liza J McCann^{1*}, Katie Arnold², Clarissa A Pilkington^{2,4}, Adam M Huber⁵, Angelo Ravelli⁶, Laura Beard², Michael W Beresford^{1,7}, Lucy R Wedderburn^{2,3,4} and the UK Juvenile Dermatomyositis Research Group (JDRG)⁴

Abstract

Background: Juvenile dermatomyositis (JDM) is a rare but severe autoimmune inflammatory myositis of childhood. International collaboration is essential in order to undertake clinical trials, understand the disease and improve long-term outcome. The aim of this study was to propose from existing collaborative initiatives a preliminary minimal dataset for JDM. This will form the basis of the future development of an international consensus-approved minimum core dataset to be used both in clinical care and inform research, allowing integration of data between centres.

Methods: A working group of internationally-representative JDM experts was formed to develop a provisional minimal dataset. Clinical and laboratory variables contained within current national and international collaborative databases of patients with idiopathic inflammatory myopathies were scrutinised. Judgements were informed by published literature and a more detailed analysis of the Juvenile Dermatomyositis Cohort Biomarker Study and Repository, UK and Ireland.

Results: A provisional minimal JDM dataset has been produced, with an associated glossary of definitions. The provisional minimal dataset will request information at time of patient diagnosis and during on-going prospective follow up. At time of patient diagnosis, information will be requested on patient demographics, diagnostic criteria and treatments given prior to diagnosis. During on-going prospective follow-up, variables will include the presence of active muscle or skin disease, major organ involvement or constitutional symptoms, investigations, treatment, physician global assessments and patient reported outcome measures.

Conclusions: An internationally agreed minimal dataset has the potential to significantly enhance collaboration, allow effective communication between groups, provide a minimal standard of care and enable analysis of the largest possible number of JDM patients to provide a greater understanding of this disease. This preliminary dataset can now be developed into a consensus-approved minimum core dataset and tested in a wider setting with the aim of achieving international agreement.

Keywords: Juvenile dermatomyositis, Idiopathic Inflammatory myopathy, International, Collaboration, Dataset, Core set, Disease activity

* Correspondence: Liza.mccann@alderhey.nhs.uk

¹Alder Hey Children's NHS Foundation Trust, Eaton Road, Liverpool L12 2AP, UK
Full list of author information is available at the end of the article

Background

Juvenile dermatomyositis (JDM) is a severe, autoimmune, inflammatory myositis that can cause death and major long-term health problems [1-3]. There are major gaps in knowledge regarding epidemiology, pathogenesis, response to medication and long-term outcome of the disease [4]. Although most children with JDM now survive to adulthood (5 year survival > 95%) [5], a significant number (70-80%) have on-going disease activity [1-3]. JDM differs in many ways from myositis in adults [5]. As JDM is rare, affecting 2-3 per million children per year [6,7], international collaboration between paediatric and adult JDM research groups is essential to undertake clinical trials, understand the disease, and improve long-term outcome. The first prospective randomised trial in myositis (evaluating rituximab in refractory disease) enrolling both paediatric and adult patients through international collaboration was recently published [8]. A collaborative randomised trial of prednisolone, methotrexate and ciclosporin in JDM has recently been completed [9]. Aside from these, there remains a significant paucity of robust evidence regarding therapeutic interventions for JDM in the paediatric and adolescent age group.

To date, two parallel processes have led to the development of two 'core sets' for JDM in America (via the International Myositis and Clinical Studies [IMACS] group) [10-12] and in Europe (via Paediatric Rheumatology International Trials Organisation, PRINTO) [13-15]. These core sets overlap but differ in important details and lack international agreement. They were developed with the main purpose for clinical trials and therefore can be difficult to use in routine clinical practice due to time constraints.

Disease specific registries, particularly when linked to a biobank of biological specimens, can be powerful tools enabling epidemiological research, identification of risk factors to develop disease (such as gene-environment interaction) and identification of patients for clinical trials [16]. They can also drive a minimal standard of care by encouraging capture of important variables for disease monitoring within clinical use. In addition, registries have the potential of providing a platform for the development of robust pharmacovigilance and post-marketing drug safety monitoring, exemplified by the Childhood Arthritis and Rheumatology Research Alliance (CARRA) Consolidated Safety Registry (CoRe) [17]. The Juvenile Dermatomyositis Cohort Biomarker Study and Repository, UK and Ireland (JDCBS) provides a large prospective collection of juvenile idiopathic inflammatory myopathies (currently >430 patients) with a repository of linked biological and serological specimens [18]. In 2010, CARRA initiated a multi-centre observational registry in the United States to create a clinical database for the major rheumatic diseases, [19] which has reported data from 384 JDM patients enrolled within the first 2 years

[20]. In 2008, a European initiative was taken to develop a web-based registry for adult myositis patients, Euromyositis [16], which is currently being expanded to include JDM patients through collaboration with the UK Juvenile Dermatomyositis Research Group (JDRG) [18].

The concept underpinning disease-specific registries is a user-friendly, accessible database that can be useful in clinical practice and encourages data-entry for research purposes. In order to achieve this, databases need to collect important information that helps to define disease outcome and if possible inform treatment, but not be so extensive that it is too time-consuming. Ideally, it should be linked to the patient record to avoid duplication [16] and yet be protected so that patients are not identifiable within a secure web-based system when data are extracted for research purposes.

Currently, existing databases within JDCBS, CARRA and Euromyositis have dataset parameters that are partially overlapping but use different data items. Analysis of information held within these individual data collections has led to a greater understanding of the disease, resulting in numerous publications on JDM [18,20]. However, many important questions remain unanswered and depend on analysis of larger numbers of patients than are currently available in individual collections. To address this, this present study aimed to scrutinize the existing collaborative datasets to propose a common minimal core data set comprising clinical, laboratory and patient/parent reported measures for patients with JDM. We aimed for such a core data set to be clinically practical, incorporate key variables that allow accurate assessment of disease activity and measure change (such as response to treatment), and for the data set to be used in the development of future clinical trials in JDM. Such a common dataset, could form the basis for a future consensus-driven, internationally approved minimum core dataset that can be rapidly incorporated into national and international collaborative efforts, including existing prospective databases, and be available for use in randomised controlled trials and for treatment / protocol comparisons in cohort studies.

Methods

A group of five JDM experts from the UK, Italy and Canada (LM,CP,LW,AH,AR) representative of the major groups studying JDM and maintaining databases (CARRA, UK JDCBS, PRINTO) formed a working group which aimed to produce a provisional minimal dataset for JDM [21].

A literature search, carried out by one author (LM), aimed to identify any variables potentially missing from current database collections but used within validated research tools. The search (EMBASE, MEDLINE, September 2012; main search terms 'dermatomyositis' OR

'myositis' AND 'juvenile' OR 'childhood' AND 'database' or 'dataset' or 'consensus' or 'core set) identified 272 publications. Articles containing variables used within published 'core sets' for research purposes (PRINTO/IMACS) and variables used within articles describing disease classification, diagnostic criteria, clinical characteristics, disease monitoring, outcome assessment/response to treatment or definition of inactive disease were listed for initial consideration and presented to the group.

A detailed analysis of variables from three existing large registries of patients with Idiopathic Inflammatory Myopathies (IIM), namely JDCBS (www.juveniledermatomyositis.org.uk), CARRA (www.carragroup.org), and Euromyositis (www.euromyositis.eu), as well as those used in a Paediatric Rheumatology International Trials Organisation (PRINTO) coordinated retrospective study analyzing 490 JDM patients [1], was undertaken. Each variable common to at least two databases was listed and scrutinized for usefulness within its clinical context and for research studies by e-mail and teleconference discussion within the expert group [21]. Variables were only included in the provisional dataset when 100% of group members agreed that they were potentially important for clinical use and/or future research.

To inform discussion, a separate detailed analysis of the UK JDCBS database was carried out in July 2011 to determine the number of times that variables were completed accurately. The JDCBS had ethical approval from the North Yorkshire Multi-Centre Research Ethics Committee and this study was also approved by the Steering Committee of the UK JDM Cohort Study. At this time the JDCBS had recruited 285 children with sufficient data available for analysis on 275 cases [18] of which 175 (63.3%) had been recruited prospectively within 12 months of onset. Any variable completed <50% of the time was scrutinized and discussed within the expert group to determine importance of inclusion versus reliability of completion and to see if questions could be asked in a more steadfast way to encourage completion.

Definitions of variables used by the individual databases were reviewed. Those used within the Euromyositis database were modified for paediatric use. A combination of these definitions, with the aid of agreed and published descriptions of cutaneous manifestations of IIM [22], were used to form a glossary/definition of variables.

Results

Analysis of the four datasets (JDCBS, CARRA, Euromyositis, PRINTO) showed that 19 variables were common to all four datasets, 16 were common to three datasets, and 24 were common to two datasets (Table 1). Although only one database (JDCBS) included data on the use of physiotherapy, this variable was retained within the treatment listing as it was felt to be important by all JDM experts. Likewise,

malignancy was retained as a variable, despite being recorded by Euromyositis alone. Therefore a total of 61 variables were scrutinized for potential inclusion. An additional 29 variables were identified as only being in one of the four datasets and excluded from further consideration at this stage. Cross-referencing of variables listed from the literature search and included within research tools such as the Myositis Disease Activity Assessment (MDAA) and Disease Activity Score (DAS) [15] was carried out to ensure that members of the working groups were satisfied that all variables important for clinical use were included. This process informed discussion but did not lead to the inclusion of any additional variables.

All registries or data collections included patient demographics, date of onset of symptoms, date of diagnosis and growth indicators. They all included diagnostic indicators of JDM, including evidence of a characteristic skin rash, proximal muscle weakness, muscle enzyme elevation, electromyogram (EMG) and muscle biopsy evidence of myositis. There were some differences in how these data were documented, particularly when comparing adult and paediatric registries. All databases asked about the presence of skin ulceration, calcinosis, arthritis and cardiac/respiratory involvement although there were differences in how the questions were asked and data presented. All databases asked for medication history.

Two databases asked about neurological involvement. Capture of systemic symptoms (weight loss, fatigue, myalgia, irritability) varied between databases. There were differences in how disease activity over time was monitored, although the majority of databases included patient/parent Visual Analogue Scale (VAS), a Health Assessment Questionnaire (CHAQ/HAQ), physician global VAS, and a measure of muscle strength; either the Childhood Myositis Assessment Scale (CMAS) and/or Manual Muscle Testing (MMT) [23]. The PRINTO study used the Disease Activity Score (DAS) whereas Euromyositis included the Myositis Intention to Treat Activity Index (MITAX) and the Myositis Disease Activity Assessment Visual Analogue Scales (MYOACT) [23].

The presence of Raynaud's phenomenon was retained due to the potential significance of progression of childhood onset myositis into scleroderma or myositis/scleroderma overlap in adult life [24]. The presence of mechanic hands and malignancy were each retained despite being infrequent in paediatrics, in order to compare with adult datasets.

Scrutiny of the UK JDCBS database, which included routine collection of 54 out of a total of 61 possible variables in the provisional minimal database, showed that at baseline, 8/54 (15%) variables were completed in <50% of patients. Specific concerns were noted with completion of family history, Childhood Health Assessment Questionnaire,

Table 1 Variables common to more than two datasets, considered for inclusion in provisional minimal dataset

Variables common to four datasets (n = 19)	Variables common to three datasets (n = 16)	Variables common to two datasets (n = 24)
Date of birth and patient code	Patient name	Postcode
Gender	Patient centre	Death
Ethnicity	Diagnosis date	Fatigue due to JDM
Family history	JDM rash	Distribution of rash
Onset date	Skin erythema	Abdominal pain due to JDM
Height/Weight	MRI consistent with JDM at diagnosis	Neurological involvement
Symmetrical muscle weakness	CMAS	Weight loss due to JDM
Gottron's or heliotrope rash	Gottron's sign	Alopecia due to JDM
Skin ulcers	Shawl sign	Nail-fold changes
Dysphagia or dysphonia	Gastro-intestinal disease / ulceration due to JDM	Dysphagia
Calcinosis	Fever due to JDM	Dysphonia
Cardiac involvement	Abnormal pulmonary function tests	Respiratory symptoms
Arthritis	Contractures	Myalgia
Interstitial lung disease	Lipodystrophy	Eye disease (glaucoma/cataract)
MMT8	Autoantibody data (more than ANA)	Raynaud's
Muscle enzymes elevated	Physician global assessment	Parental Visual Analogue Scale (VAS)
EMG changes consistent with JDM at diagnosis		Pain VAS
Muscle biopsy consistent with JDM		CHAQ
Medication history		MITAX/MYOACT
		Hospitalisation since last visit
		School absence
		Other significant diagnosis
		Sporting activities
		Pubertal delay

Child Health Questionnaire, parental visual analogue scale of disease activity, pain visual analogue score, as well as nail-fold capillary changes, lipodystrophy and Raynaud's phenomenon. Analysis of completion rates for two of these variables (nail-fold changes and Raynaud's) could have been confounded by the fact that they were not included in the original data collection forms, but added at a later date. With prospective collection, parameters performing poorly (completed in <50% patients) were MRI, muscle biopsy, antibody and pulmonary function test results. Discussion within the expert group confirmed the importance of these parameters but led to rephrasing of questions to support future enhanced data completion. For example, the question 'pulmonary involvement due to myositis' was favoured as opposed to asking clinicians to document the results of pulmonary function tests.

A provisional minimal dataset was therefore established from the existing collaborative datasets for use in clinics and inform clinical trials and research studies. This provisional dataset was then configured into user-

friendly forms (Figures 1, 2, 3, 4 and 5) and organized into ten domains. Three of these domains (demographic data, diagnostic data and treatment prior to diagnosis) are questions that would be asked at first presentation only (Figures 1 and 2). The remaining seven domains (growth, clinical features, ongoing follow up, physician reported outcome measures, patient/parent reported outcome measures, investigations, treatment) are relevant for every patient visit (Figures 3, 4 and 5). A glossary for form A (Figures 1 and 2) and form B (Figures 3, 4 and 5) are available as Additional file 1 (Form A: provisional definitions, Form B: provisional definitions).

Our working group includes a member of the Euromyositis Steering Committee and in collaboration with the Euromyositis Research Group, we have used the provisional dataset as a platform to create a paediatric-specific webpage for the Euromyositis database, ensuring compatibility with the web-based system used to collect data on adult-onset idiopathic inflammatory (www.euromyositis.eu).

Further work will develop this preliminary dataset into a consensus-approved minimum core dataset, tested

Provisional JDM Minimal Dataset, Form A - first data entry only

Instructions on use / definitions available as supplementary material.

- For variables that are not available at time of data entry, but will be available at a later date, mark 'not available', (so that they can be chased and added at a later date).
- For variables that have not been examined / investigated, mark 'not recorded'.

Patient ID Number*		Patient Name*	
Patient Consultant / Lead Clinician? *		Patient Hospital* and country	

* Information to be completed or details omitted pending local ethics and legal considerations

Date of form completion	
--------------------------------	--

A. Demographic Data				
DOB*:		Gender:	Male / Female	
Ethnicity (<i>Details to be further defined by consensus</i>)				
Family History (1st degree relative)				
1. Neuromuscular disease	Yes	No	Not Known	Not Available
If yes, state disease _____				
2. Autoimmune disease	Yes	No	Not Known	Not Available
If yes, state disease _____				

B. Diagnostic Data				
<i>For definitions please refer to supplementary sheet</i>				
1. Disease type:	JDM JPM JDM overlap with: Arthritis / Lupus / Scleroderma / Sjogren's / MCTD (RNP +ve) <i>Mark all overlap features that apply</i> Amyopathic JDM Incomplete DM Inclusion body myositis (IBM)			
2. Date of first symptom (mm/yyyy)	____/____/____			
3. Date of diagnosis (mm/yyyy)	____/____/____			
4. Diagnostic Clinical Criteria				
Symmetrical proximal muscle weakness	Yes	No	Not Available <small>(at time of data entry but can be completed later)</small>	Not Done
Dermatological features of JDM:				
Heliotrope rash	Yes	No	Not Available	Not Done
Gottron's papules or Gottron's sign	Yes	No	Not Available	Not Done
Nailfold capillary changes	Yes	No	Not Available	Not Done
Other characteristic JDM rash	Yes	No	Not Available	Not Done

Figure 1 Provisional JDM minimal dataset Form A (first data entry only), page 1 –demographic and diagnostic data.

in a wider setting with the aim of achieving international agreement.

Discussion

International collaboration among myositis experts within IMACS and PRINTO has led to the development of two core set measures for JDM [10-15]. These are important tools predominantly designed for research and clinical trials. Time constraints limit the use of these in routine clinical practice. However, the tools capture important variables that measure disease activity and damage in JDM. In this study we have attempted to capture these individual variables

within a provisional minimal dataset with the aim of producing a platform for recording important information within routine clinical practice, and providing a minimum standard of care for JDM. This will have the added benefit of being able to be used to inform future research and will therefore compliment the existing IMACS/PRINTO core sets.

Our group has scrutinized variables collected within four existing databases, in addition to variables used within established research tools and published literature, to form a provisional minimal dataset for JDM. Further development is now needed through a formal consensus-driven process with the aim of ultimately achieving a unified

5. Diagnostic Investigations					
Muscle biopsy evidence of myositis:		Yes	No	Not Available	Not Done
If yes, date of biopsy (dd/mm/yyyy): _____					
Elevation of muscle enzyme:	CK	Yes	No	Not Available	Not Done
	LDH	Yes	No	Not Available	Not Done
	Adolase	Yes	No	Not Available	Not Done
	AST/SGOT	Yes	No	Not Available	Not Done
	ALT/SGPT	Yes	No	Not Available	Not Done
EMG changes of myopathy		Yes	No	Not Available	Not Done
MRI Changes of Myositis		Yes	No	Not Available	Not Done
Auto-antibodies positive:	ANA	Yes	No	Not Available	Not Done
	Anti-p155/140	Yes	No	Not Available	Not Done
	Anti-p140	Yes	No	Not Available	Not Done
	Anti-Jo1	Yes	No	Not Available	Not Done
	Anti-ARS	Yes	No	Not Available	Not Done
	Anti-PL12	Yes	No	Not Available	Not Done
	Anti-PL7	Yes	No	Not Available	Not Done
	Anti-EJ	Yes	No	Not Available	Not Done
	Anti-OJ	Yes	No	Not Available	Not Done
	Anti-Mi2	Yes	No	Not Available	Not Done
	Anti-SRP	Yes	No	Not Available	Not Done
	Anti-U1-RNP	Yes	No	Not Available	Not Done
	Anti-U3-RNP	Yes	No	Not Available	Not Done
	Anti-PM-Scl 75	Yes	No	Not Available	Not Done
	Anti-PM-Scl 100	Yes	No	Not Available	Not Done
	Anti-Ro52	Yes	No	Not Available	Not Done
	Anti-Ro60	Yes	No	Not Available	Not Done
	Anti-La	Yes	No	Not Available	Not Done
	Anti-Ku	Yes	No	Not Available	Not Done
	Anti-Topo	Yes	No	Not Available	Not Done
	Anti-SAE	Yes	No	Not Available	Not Done
	Anti-Zo	Yes	No	Not Available	Not Done
	Anti-CAM-140	Yes	No	Not Available	Not Done
	Anti-KS	Yes	No	Not Available	Not Done
	Anti-HA	Yes	No	Not Available	Not Done
	Anti-ZA	Yes	No	Not Available	Not Done
	Other MSA / MAA	Yes	No	Not Available	Not Done
C. Treatments					
Medication given prior to diagnosis of myositis and registration into database:					Yes No
If yes, medications given:					
Steroid	Yes	No	Not Available	Not Recorded	
Methotrexate / other DMARD (Disease modifying anti-rheumatic drug)	Yes	No	Not Available	Not Recorded	
Previous biologic	Yes	No	Not Available	Not Recorded	
If yes to steroid:	Oral	Intravenous	Topical	Intramuscular	

Figure 2 Provisional JDM minimal dataset Form A, (first data entry only), page 2 – diagnostic investigations and treatments.

internationally agreed minimal dataset for JDM for clinical use that can also inform research.

The development of an internationally agreed minimal dataset could have numerous potential benefits. Unification of data collection internationally will significantly enhance collaborative efforts and communication between groups (whilst respecting data protection/ownership) and allow analysis to be undertaken on the largest possible number of JDM patients. This has the potential

to improve understanding of disease phenotype and outcome in a way that will define research questions and underpin prioritisation of future clinical trials. Analysis of a large number of JDM patients with international collaboration has the potential to help answer fundamental questions including characteristics of patients who develop JDM, how patients develop the disease, time to diagnosis (and raise awareness leading to quicker diagnosis), which tests are most useful for diagnosis and

Provisional JDM Minimal Dataset, Form B: prospective data entry

Instructions on use / definitions available as supplementary material.

- For variables that are not available at time of data entry, but will be available at a later date, mark 'not available' (so that they can be chased and added at a later date).
- For variables that have not been examined / investigated, mark 'not recorded'.

*Details of patient ID to be completed / omitted as per local ethics / legal requirements.

Patient ID Number*		Patient Name*	
Patient Consultant / Lead Clinician		Patient Hospital and country	

Date of form completion	
--------------------------------	--

A. Growth	
Height (cm)	Weight (kg)

B. Disease activity / damage (clinical features)				
1. Active Muscle Disease				
Proximal muscle weakness	Yes	No	Not Available	Not Done
Other muscle weakness	Yes	No	Not Available	Not Done
CMAS	Score	_____	Not Available	Not Done
MMT8	Score	_____	Not Available	Not Done
2. Active Skin Disease				
Heliotrope rash	Yes	No	Not Available	Not Done
Gottron's Papules or Gottron's sign	Yes	No	Not Available	Not Done
Nailfold capillary changes	Yes	No	Not Available	Not Done
Lipodystrophy	Yes	No	Not Available	Not Done
Calcinosis	Yes	No	Not Available	Not Done
Cutaneous ulceration due to myositis	Yes	No	Not Available	Not Done
Subcutaneous oedema due to myositis				
Generalised	Yes	No	Not Available	Not Done
Peri-orbital	Yes	No	Not Available	Not Done
Other skin changes				
Malar / facial erythema due to myositis	Yes	No	Not Available	Not Done
Shawl sign / V sign due to myositis	Yes	No	Not Available	Not Done
Mechanic's hands	Yes	No	Not Available	Not Done
Alopecia due to myositis	Yes	No	Not Available	Not Done
Vasculopathic lesions due to myositis	Yes	No	Not Available	Not Done
Photosensitivity due to myositis	Yes	No	Not Available	Not Done
Livido reticularis due to myositis	Yes	No	Not Available	Not Done
Other erythema due to myositis	Yes	No	Not Available	Not Done
Panniculitis due to myositis	Yes	No	Not Available	Not Done

Figure 3 Provisional JDM minimal dataset, Form B, (prospective data collection), page 1 – disease activity and damage.

which features predict a more severe disease course. Increasingly, distinct clinical subtypes are recognised within IIM, associated with myositis specific antibodies, but differences are seen in adults and children [25]. Our understanding of the mechanisms that underlie these differences is limited by lack of studies directly comparing adults and children in sufficient numbers. A minimal dataset including disease characteristics and antibody profiling may help to answer these questions with the potential of directing subsequent clinical trials of treatment strategies on disease stratified by serology. It is increasingly recognized that patient reported perspectives reveal disease aspects not covered by traditional outcomes in myositis [26]. A minimal dataset will include patient/parent reported outcome measures and work is ongoing to develop new JDM-specific patient/parent

outcome measures (the Juvenile Dermatomyositis Multi-dimensional Assessment Report, JDMAR) [27]. With international collaboration, patient reported outcome measures, such as this, could be evaluated in large numbers of patients, helping to predict effectiveness for their use in future clinical trials.

A large international collection has the potential to allow detailed evaluation of 'real world' treatment interventions used in clinical practice – based on clinician's intention to treat, to see if one treatment is better than another. The CARRA database includes the option of clinicians choosing which of a predefined number of treatment strategies to use within the context of a consensus treatment plan [28]. If other studies contain the same minimal dataset, outcome of patients on different treatments can be compared and evaluated.

3. Major organ involvement due to myositis				
Musculoskeletal involvement due to myositis	Yes	No	Not Available	Not Done
If yes: Arthritis	Yes	No	Not Available	Not Done
If yes: Contractures due to myositis	Yes	No	Not Available	Not Done
Gastrointestinal involvement due to myositis	Yes	No	Not Available	Not Done
If yes: Dysphagia	Yes	No	Not Available	Not Done
If yes: Dysphonia	Yes	No	Not Available	Not Done
If yes: Abdominal pain or GI ulceration due to myositis (exclude peptic ulcer disease)	Yes	No	Not Available	Not Done
Pulmonary involvement due to myositis – interstitial lung disease	Yes	No	Not Available	Not Done
Cardiac involvement due to myositis	Yes	No	Not Available	Not Done
4. Constitutional				
Fever (>38°C) due to myositis	Yes	No	Not Available	Not Done
Weight loss (>5%) due to myositis	Yes	No	Not Available	Not Done
Fatigue due to myositis	Yes	No	Not Available	Not Done
Irritability due to myositis	Yes	No	Not Available	Not Done
Raynaud's phenomenon	Yes	No	Not Available	Not Done
5. Other				
Malignancy	Yes	No	Not Available	Not Recorded
C. Is patient having on-going follow up at this centre after this date?				
			Yes	No
If no, reason why:	Death	Yes	No	
	Transfer to another geographical area	Yes	No	
	Transfer to adult care	Yes	No	
	Patient in stable remission	Yes	No	
D. Global disease activity				
1. Physician VAS – global activity (overall)			Not Available	Not Done
Value: _____	(range 0.0-10.0cm)			
2. Physician VAS – extra-muscular global activity			Not Available	Not Done
Value: _____	(range 0.0-10.0cm)			
3. Parental / patient VAS			Not Available	Not Done
Value: _____	(range 0.0-10.0cm)			
E. Health Questionnaires				
CHAQ (range 0 - 3)	Not Available	Not Done	Value	_____
QoL Tool				
<i>(Exact tool to be determined by consensus)</i>				
QoL Tool used	Yes	No	Not Available	
Name of QoL Tool used			Score	

Figure 4 Provisional JDM minimal dataset, Form B, (prospective data collection), page 2 –disease activity/damage (continued).

International collaboration has the potential to enable comparative studies of differences between adult and juvenile dermatomyositis. Our group has collaborated with Euromyositis to produce a paediatric specific web page

for data entry for patients with JDM (www.euromyositis.eu). This web-page complements current data collection on adult myositis and will allow comparison between the two groups. New variables that are proposed as part of

F. Investigations				
1. Blood tests- muscle enzyme elevation				
CK	Yes	No	Not Available	Not Done
LDH	Yes	No	Not Available	Not Done
Adolase	Yes	No	Not Available	Not Done
AST/SGOT	Yes	No	Not Available	Not Done
ALT/SGPT	Yes	No	Not Available	Not Done
2. Abnormal investigation since last visit (consistent with flare of myositis)				
MRI	Yes	No	Not Available	Not Done
Muscle biopsy	Yes	No	Not Available	Not Done
EMG	Yes	No	Not Available	Not Done
Lung function tests	Yes	No	Not Available	Not Done
ECG / ECHO	Yes	No	Not Available	Not Done
G. Treatment				
1. Treatment start date (mm/yyyy)				
	/		Not Available	Not Recorded
2. Medication				
Steroids	Yes	No	Not Available	Not Recorded
<i>If yes to steroid:</i>	Oral	Intravenous	Topical	Intramuscular
Methotrexate	Yes	No	Not Available	Not Recorded
Ciclosporin A	Yes	No	Not Available	Not Recorded
Mycophenolate Mofetil	Yes	No	Not Available	Not Recorded
Azathioprine	Yes	No	Not Available	Not Recorded
Hydroxychloroquine	Yes	No	Not Available	Not Recorded
Leflunamide	Yes	No	Not Available	Not Recorded
Cyclophosphamide	Yes	No	Not Available	Not Recorded
Rituximab	Yes	No	Not Available	Not Recorded
Etanercept	Yes	No	Not Available	Not Recorded
Infliximab	Yes	No	Not Available	Not Recorded
Adalimumab	Yes	No	Not Available	Not Recorded
IVIg	Yes	No	Not Available	Not Recorded
Plasmaphoresis	Yes	No	Not Available	Not Recorded
Topical steroid	Yes	No	Not Available	Not Recorded
Topical tacrolimus	Yes	No	Not Available	Not Recorded
Other Treatment:				
3. Physiotherapy				
	Yes	No	Not Available	Not Recorded

Figure 5 Provisional JDM minimal dataset, Form B, (prospective data collection), page 3 – investigations and treatment.

this international minimal dataset could be incorporated into the Euromyositis data collection. An important part of this collaboration is to ensure that children entered into adult networks have paediatric-specific outcomes (such as growth) recorded. Collaborative efforts in JDM have led to data-driven approaches such as recently published PRINTO-driven criteria for defining clinically inactive disease in JDM [29]. A minimal dataset will allow validation of such approaches in large cohorts of patients across several prospective data collections.

A minimal dataset will provide a framework within which all children with JDM can be monitored in a similar way, including important variables, as a minimum standard of care. The dataset will prompt clinicians to

check important clinical features including muscle strength, skin activity and investigations each time a child with JDM is seen. This is important since JDM is a rare disease and some clinicians are not seeing large numbers of patients with the disease. The importance of defining minimal standards of care has been recognized internationally and members of our group are involved in a European effort to produce standards of care within a Single Hub and Access Point for Paediatric Rheumatology in Europe (SHARE) for JDM as well as other paediatric rheumatological conditions [30].

A provisional minimal dataset has been defined by consensus between a group of experts in JDM, representative of the major national and international JDM collaborative efforts. This now needs to be refined and tested for its

usefulness and acceptability in a wider international setting. In order to achieve this, a formal consensus-driven methodology is required of all stakeholders. This will include a Delphi process to gain widespread opinion via the established JDM networks that will help to inform a more detailed face-to-face nominal group consensus process by internationally representative JDM experts.

Conclusion

JDM is a rare but serious inflammatory condition with an appreciable risk of long-term morbidity. There are many unanswered questions around epidemiology, pathogenesis, treatment and outcome of the disease. The provisional dataset developed to date will be used as a template to aid a formal multi-stage consensus process, involving large groups of clinicians caring for children with JDM, with the aim of developing a feasible, internationally agreed minimal dataset for clinical use that can inform research.

Additional file

Additional file 1: Form A: provisional definitions, Form B: provisional definitions (to be refined further as part of on-going work) [31-37].

Abbreviations

JDM: Juvenile dermatomyositis; IIM: Idiopathic Inflammatory Myopathy; JDRG: Juvenile Dermatomyositis Research Group (UK and Ireland); JDCBS: Juvenile Dermatomyositis Cohort Biomarker Study and Repository, UK and Ireland; CARRA: Childhood Arthritis and Rheumatology Research Alliance; PRINTO: Paediatric Rheumatology International Trials Organisation; IMACS: International Myositis and Clinical Studies group; OMERACT: Outcome Measures in Rheumatology; EMG: Electromyogram; MRI: Magnetic Resonance Imaging; CHAQ: Childhood Health Assessment Questionnaire; VAS: Visual Analogue Scale; CMAS: Childhood Myositis Assessment Scale (CMAS); MMT: Manual Muscle Testing (MMT); DAS: Disease Activity Score (DAS); MITAX: Myositis Intention to Treat Activity Index (MITAX); MYOACT: Myositis Disease Activity Assessment Visual Analogue Scales.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

LM, CP, LW, AH, AR involved in all stages of the development of minimal dataset. LB was involved in scrutinising the UK JDCBS (carrying out data queries) and administrative tasks. KA was involved in scrutinising the UK JDCBS database, carrying out administrative tasks and developing the database format / clinical forms. LM, CP, MB, LW involved in intellectual content and obtaining funding to take this work forward. All authors read and approved the final manuscript.

Acknowledgements

We would like to acknowledge the Euromyositis Research Steering Committee for their collaboration and impetus for this work, in particular Professor Ingrid Lundberg, Dr Hector Chinoy, Professor Jiri Vencovsky and Niels Steen Krogh. We acknowledge the IMACS, CARRA, PRINTO and PReS working groups for their support and collaboration. The UK JDM Cohort and Biomarker study has been supported by generous grants from the Wellcome Trust UK (085860), Action Medical Research UK, (SP4252), The Myositis Support Group UK, Arthritis Research UK (14518), The Henry Smith Charity. LW is supported in part by Great Ormond Street Children's Charity. The JDM Cohort study is adopted onto the Comprehensive Research Network through the Medicines for Children Research Network (www.mcrn.org.uk) and is supported by the GOSH/ICH Biomedical Research Centre. Full list of

contributors to the JDCBS can be found on the JDRG website <http://www.juveniledermatomyositis.org.uk>

Author details

¹Alder Hey Children's NHS Foundation Trust, Eaton Road, Liverpool L12 2AP, UK
²Rheumatology Unit, UCL Institute of Child Health, University College London, London, UK
³Centre for Adolescent Rheumatology at University College London, University College London Hospital, London, UK
⁴Great Ormond Street Hospital, London, UK
⁵IWK Health Centre and Dalhousie University, 5850 University Avenue, Halifax, NS B3K 6R8, Canada.
⁶Università degli Studi di Genova and Istituto Giannina Gaslini, Largo G. Gaslini 5, 16147 Genoa, Italy.
⁷Department of Women's and Children's Health, Institute of Translational Medicine, University of Liverpool, Liverpool, UK

Received: 12 February 2014 Accepted: 8 July 2014

Published: 21 July 2014

References

1. Ravelli A, Trail L, Ferrari C, Ruperto N, Pistorio A, Pilkington C, Maillard S, Oliveira SK, Sztajnbock F, Cuttica R, Beltramelli M, Corona F, Katsicas MM, Russo R, Ferriani V, Burgos-Vargas R, Magni-Manzoni S, Solis-Vallejo E, Bandeira M, Zullian F, Baca V, Cortis E, Falcini F, Alessio M, Alpigiani MG, Gerloni V, Saad-Magalhaes C, Podda R, Silva CA, Lepore L, et al: **Long-term outcome and prognostic factors of juvenile dermatomyositis: a multinational, multicenter study of 490 patients.** *Arthritis Care Res (Hoboken)* 2010, **62**:63-72.
2. Mathiesen PR, Zak M, Herlin T, Nielsen SM: **Clinical features and outcome in a Danish cohort of juvenile dermatomyositis patients.** *Clin Exp Rheumatol* 2010, **28**:782-789.
3. Sanner H, Kirkhus E, Merckoll E, Tollisen A, Roisland M, Lie BA, Taraldsrud E, Gran JT, Flato B: **Long-term muscular outcome and predisposing and prognostic factors in juvenile dermatomyositis: A case-control study.** *Arthritis Care Res (Hoboken)* 2010, **62**:1103-1111.
4. Feldman BM, Rider LG, Reed AM, Pachman LM: **Juvenile dermatomyositis and other idiopathic inflammatory myopathies of childhood.** *Lancet* 2008, **371**:2201-2212.
5. Robinson AB, Reed AM: **Clinical features, pathogenesis and treatment of juvenile and adult dermatomyositis.** *Nat Rev Rheumatol* 2011, **7**:664-675.
6. Symmons DP, Sills JA, Davis SM: **The incidence of juvenile dermatomyositis: results from a nation-wide study.** *Br J Rheumatol* 1995, **34**:732-736.
7. Mendez EP, Lipton R, Ramsey-Goldman R, Roettcher P, Bowyer S, Dyer A, Pachman LM: **US incidence of juvenile dermatomyositis, 1995-1998: results from the National Institute of Arthritis and Musculoskeletal and Skin Diseases Registry.** *Arthritis Rheum* 2003, **49**:300-305.
8. Oddis CV, Reed AM, Aggarwal R, Rider LG, Ascherman DP, Levesque MC, Barohn RJ, Feldman BM, Harris-Love MO, Koontz DC, Fertig N, Kelley SS, Pryber SL, Miller FW, Rockette HE, RIM Study Group: **Rituximab in the treatment of refractory adult and juvenile dermatomyositis and adult polymyositis: a randomized, placebo-phase trial.** *Arthritis Rheum* 2013, **65**:314-324.
9. Ruperto N, Pistorio A, Oliveira S, Cuttica RJ, Ravelli A, Fischbach M, Hagelberg S, Avcin A, Cheuret E, Corona F, Couillaud G, Dressler F, Gerloni V, Sterba G, Zulian F, Apaz MT, Cespedes-Cruz A, Cimaz R, Bracaglia C, Joos R, Quartier P, Russo R, Tardieu M, Wulfraat N, Angioloni S, Martini A: **A randomised trial in new onset juvenile dermatomyositis: prednisolone versus prednisolone plus cyclosporine versus prednisolone plus methotrexate. [abstract].** *Arthritis Rheum* 2012, **64**:2473.
10. Rider LG, Giannini EH, Harris-Love M, Joe G, Isenberg D, Pilkington C, Lachenbruch PA, Miller FW: **Defining Clinical Improvement in Adult and Juvenile Myositis.** *J Rheumatol* 2003, **30**:603-617.
11. Rider LG, Giannini EH, Brunner HI, Ruperto N, James-Newton L, Reed AM, Lachenbruch PA, Miller FW: **International consensus on preliminary definitions of improvement in adult and juvenile myositis.** *Arthritis Rheum* 2004, **50**:2281-2290.
12. Oddis CV, Rider LG, Reed AM, Ruperto N, Brunner HI, Koneru B, Feldman BM, Giannini EH, Miller FW: **International consensus guidelines for trials of therapies in the idiopathic inflammatory myopathies.** *Arthritis Rheum* 2005, **52**:2607-2615.
13. Ruperto N, Ravelli A, Pistorio A, Ferriani V, Calvo I, Ganser G, Brunner J, Dannecker G, Silva CA, Stanevicha V, Cate RT, van Suijlekom-Smit LW,

- Voygjoyka O, Fischbach M, Foeldvari I, Hilario O, Modesto C, Saurenmann RK, Sauvain MJ, Scheibel I, Sommelet D, Tambic-Bukovac L, Barcellona R, Brik R, Ehl S, Jovanovic M, Rovensky J, Bagnasco F, Lovell DJ, Martini A, et al: **The provisional Paediatric Rheumatology International Trials Organisation/American College of Rheumatology/European League Against Rheumatism Disease activity core set for the evaluation of response to therapy in juvenile dermatomyositis: a prospective validation study.** *Arthritis Rheum* 2008, **59**:4–13.
14. Ruperto N, Pistorio A, Ravelli A, Rider LG, Pilkington C, Oliveira S, Wulffraat N, Espada G, Garay S, Cuttica R, Hofer M, Quartier P, Melo-Gomes J, Reed AM, Wierzbowska M, Feldman BM, Harjacek M, Huppertz HI, Nielsen S, Flato B, Lahdenne P, Michels H, Murray KJ, Punaro L, Rennebohm R, Russo R, Balogh Z, Rooney M, Pachman LM, Wallace C, et al: **The Paediatric Rheumatology International Trials Organisation provisional criteria for the evaluation of response to therapy in juvenile dermatomyositis.** *Arthritis Care Res (Hoboken)* 2010, **62**:1533–1541.
 15. Ruperto N, Ravelli A, Murray KJ, Lovell DJ, Andersson-Gare B, Feldman BM, Garay S, Kuis W, Machado C, Pachman L, Prieur AM, Rider LG, Silverman E, Tsitsami E, Woo P, Giannini EH, Martini A, Paediatric Rheumatology International Trials Organisation (PRINTO); Pediatric Rheumatology Collaborative Study Group (PRCSG): **Preliminary core sets of measures for disease activity and damage assessment in juvenile systemic lupus erythematosus and juvenile dermatomyositis.** *Rheumatology* 2003, **42**:1452–1459.
 16. Lundberg IE, Svensson J: **Registries in idiopathic inflammatory myopathies.** *Curr Opin Rheumatol* 2013, **25**:729–734.
 17. Lionetti G, Kimura Y, Schanberg LE, Beukelman T, Wallace CA, Ilowite NT, Winsor J, Fox K, Natter M, Sundry JS, Brodsky E, Curtis JR, Del Gaizo V, Iyasu S, Jahreis A, Meeker-O'Connell A, Mittleman BB, Murphy BM, Peterson ED, Raymond SC, Setoguchi S, Siegel JN, Sobel RE, Solomon D, Southwood TR, Vesely R, White PH, Wulffraat NM, Sandborg CI: **Using registries to identify adverse events in rheumatic diseases.** *Pediatrics* 2013, **132**:e1384–e1394.
 18. Martin N, Krol P, Smith S, Murray J, Pilkington CA, Davidson JE, Wedderburn LR: **A national registry for juvenile dermatomyositis and other paediatric idiopathic inflammatory myopathies: 10 years' experience; the Juvenile Dermatomyositis National (UK and Ireland) Cohort Biomarker Study and Repository for Idiopathic Inflammatory Myopathies.** *Rheumatology (Oxford)* 2011, **50**:137–145.
 19. Natter MD, Quan J, Ortiz DM, Bousvaros A, Ilowite NT, Inman CJ, Marsolo K, McMurry AJ, Sandborg CI, Schanberg LE, Wallace CA, Warren RW, Weber GM, Mandl KD: **An i2b2-based, generalizable, open source, self-scaling chronic disease registry.** *J Am Med Inform Assoc* 2013, **20**:172–179.
 20. Robinson AB, Hoeltzel MF, Wahezi DM, Becker ML, Kessler E, Schmeling H, Carrasco R, Huber AM, Feldman BM, Reed AM: **Clinical characteristics of children with juvenile dermatomyositis - the children's arthritis and rheumatology research alliance (CARRA) registry.** *Arthritis Care Res (Hoboken)* 2013. doi:10.1002/acr.22142[Epub ahead of print].
 21. McCann LJ, Beard L, Ravelli A, Huber A, Wedderburn LR: **Proposal for the development of an international minimal data collection for juvenile dermatomyositis [abstract].** *Arthritis Rheum* 2011, **61**:234.
 22. Dugan EM, Huber AM, Miller FW, Rider LG: **Photoessay of the cutaneous manifestations of the idiopathic inflammatory myopathies.** *Dermatol Online J* 2009, **15**:1.
 23. Rider LG, Werth VP, Huber AM, Alexanderson H, Rao AP, Ruperto N, Herbelin L, Barohn R, Isenberg D, Miller FW: **Measures of adult and juvenile dermatomyositis, polymyositis, and inclusion body myositis: Physician and Patient/Parent Global Activity, Manual Muscle Testing (MMT), Health Assessment Questionnaire (HAQ)/Childhood Health Assessment Questionnaire (C-HAQ), Childhood Myositis Assessment Scale (CMAS), Myositis Disease Activity Assessment Tool (MDAAT), Disease Activity Score (DAS), Short Form 36 (SF-36), Child Health Questionnaire (CHQ), physician global damage, Myositis Damage Index (MDI), Quantitative Muscle Testing (QMT), Myositis Functional Index-2 (FI-2), Myositis Activities Profile (MAP), Inclusion Body Myositis Functional Rating Scale (IBMFRS), Cutaneous Dermatomyositis Disease Area and Severity Index (CDASI), Cutaneous Assessment Tool (CAT), Dermatomyositis Skin Severity Index (DSSI), Skindex, and Dermatology Life Quality Index (DLQI).** *Arthritis Care Res (Hoboken)* 2011, **63**(Suppl 11):S118–S157.
 24. Wedderburn LR, McHugh NJ, Chinoy H, Cooper RG, Salway F, Ollier WE, McCann LJ, Varsani H, Dunphy J, North J, Davidson JE: **HLA class II haplotype and autoantibody associations in children with juvenile dermatomyositis and juvenile dermatomyositis-scleroderma overlap.** *Rheumatology (Oxford)* 2007, **46**:1786–1791.
 25. Tansley SL, McHugh NJ, Wedderburn LR: **Adult and juvenile dermatomyositis: are the distinct clinical features explained by our current understanding of serological subgroups and pathogenic mechanisms?** *Arthritis Res Ther* 2013, **15**:211.
 26. Alemo Munters L, van Vollenhoven RF, Alexanderson H: **Patient preference assessment reveals disease aspects not covered by recommended outcomes in polymyositis and dermatomyositis.** *ISRN Rheumatol* 2011, **2011**:463124.
 27. Filocamo G, Consolaro A, Ferrari C, Ravelli A: **Introducing new tools for assessment of parent- and child-reported outcomes in paediatric rheumatology practice: a work in progress.** *Clin Exp Rheumatol* 2013, **31**:964–968.
 28. Huber AM, Giannini EH, Bowyer SL, Kim S, Lang B, Lindsley CB, Pachman LM, Pilkington C, Reed AM, Rennebohm LG, Rider RM, Wallace CA, Feldman BA: **Protocols for the initial treatment of moderately severe juvenile dermatomyositis: results of a Children's Arthritis and Rheumatology Research Alliance Consensus Conference.** *Arthritis Care Res (Hoboken)* 2010, **62**:219–225.
 29. Lazarevic D, Pistorio A, Palmisani E, Miettunen P, Ravelli A, Pilkington C, Wulffraat NM, Malattia C, Garay SM, Hofer M, Quartier P, Dolezalova P, Penades IC, Ferriani VP, Ganser G, Kasapcopur O, Melo-Gomes JA, Reed AM, Wierzbowska M, Rider LG, Martini A, Ruperto N, Paediatric Rheumatology International Trials Organisation (PRINTO): **The PRINTO criteria for clinically inactive disease in juvenile dermatomyositis.** *Ann Rheum Dis* 2013, **72**:686–693.
 30. Wulffraat NM, Vastert B: **Time to share.** *Pediatr Rheumatol Online J* 2013, **11**:5.
 31. Bohan A, Peter JB: **Polymyositis and dermatomyositis. Parts 1 and 2.** *New Engl J Med* 1975, **292**:344–347. 403–407.
 32. **Washington Neuromuscular Disease Centre.** [http://neuromuscular.wustl.edu]
 33. Santmyre-Rosenberger B, Dugan EM: **Skin Involvement in Dermatomyositis.** *Curr Opin Rheumatol* 2003, **15**:714–722.
 34. Lovell DJ, Lindsley CB, Rennebohm RM, Ballinger SH, Bowyer SL, Giannini EH, Hicks JE, Levinson JE, Mier R, Pachman LM, Passo MH, Perez MD, Reed AM, Schikler KN, Smith M, Zemel LS, Rider LG: **Development of validated disease activity and damage indices for the juvenile idiopathic inflammatory myopathies: II. The childhood myositis assessment scale (CMAS): a quantitative tool for the evaluation of muscle function.** *Arthritis Rheum* 1999, **42**:2213–2219.
 35. Huber AM, Lovell DJ, Pilkington CA, Rennebohm RM, Rider LG: **Confusion concerning multiple versions of the Childhood Myositis Assessment Scale.** *Arthritis Care Res* 2013. doi:10.1002/acr.22239[Epub ahead of print].
 36. Nugent J, Ruperto N, Grainger J, Machado C, Sawhney S, Baildam E, Davidson J, Foster H, Hall A, Hollingworth P, Sills J, Venning H, Walsh JE, Landgraf JM, Roland M, Woo P, Murray KJ, Paediatric Rheumatology International Trials Organisation: **The British version of the Childhood Health assessment questionnaire (CHAQ) and the child health Questionnaire (CHQ).** *Clin Exp Rheumatol* 2001, **2001**(19):S163–S167.
 37. Huber AM, Hicks JE, Lachenbruch PA, Perez MD, Zemel LS, Rennebohm RM, Wallace CA, Lindsley CB, Passo MH, Ballinger SH, Bowyer SL, Reed AM, White PH, Katona IM, Miller FW, Rider LG, Feldman BM, Juvenile Dermatomyositis Disease Activity Collaborative Group: **Validation of the Childhood Health Assessment Questionnaire in the juvenile idiopathic myopathies.** *J Rheumatol* 2001, **28**:1106–1111.

doi:10.1186/1546-0096-12-31

Cite this article as: McCann et al.: Developing a provisional, international Minimal Dataset for Juvenile Dermatomyositis: for use in clinical practice to inform research. *Pediatric Rheumatology* 2014 **12**:31.