



# *C9orf72* amyotrophic lateral sclerosis and frontotemporal dementia: gain or loss of function?

Sarah Mizielinska and Adrian M. Isaacs

## Purpose of review

The molecular mechanisms that underlie chromosome 9 open reading frame 72 (*C9orf72*)-associated amyotrophic lateral sclerosis and frontotemporal dementia are rapidly emerging. Two potential disease mechanisms have been postulated – gain or loss of function. We provide an overview of recent advances that support or oppose gain-of-function and loss-of-function mechanisms.

## Recent findings

Since the discovery that a noncoding repeat expansion in *C9orf72* was responsible for chromosome 9-linked amyotrophic lateral sclerosis and frontotemporal dementia in 2011, a plethora of studies have investigated clinical, pathological and mechanistic aspects of the disease. Loss of function is supported by reduced levels of *C9orf72* in patient brain and functional work, revealing a role of the *C9orf72* protein in endocytic and autophagic pathways and motor function. Gain of function is supported by the presence in patient brain of both repeat RNA and protein aggregates. Repeat RNA aggregates termed RNA foci, a hallmark of noncoding repeat expansion diseases, have been shown to sequester proteins involved in RNA splicing, editing, nuclear export and nucleolar function. Repeat-associated non-ATG dependent translation gives rise to toxic dipeptide repeat proteins that form inclusions in patient tissue. Antisense oligonucleotides targeting *C9orf72* have shown promise for combating gain-of-function toxicity.

## Summary

Rapid progress is being made towards understanding this common genetic cause of amyotrophic lateral sclerosis and frontotemporal dementia. Overall, the weight of data currently sits in favour of gain of function as the most important disease mechanism, which has important implications for the development of effective and targeted therapies.

## Keywords

amyotrophic lateral sclerosis, *C9orf72*, frontotemporal dementia, gain or loss of function

## INTRODUCTION

A noncoding repeat expansion in *C9orf72* is a common genetic cause of amyotrophic lateral sclerosis (ALS) and frontotemporal lobar dementia (FTD) (C9FTD/ALS) [1,2]. Disease may occur through loss of function of *C9orf72*, or two distinct gain-of-function mechanisms: first, the formation of repeat RNA aggregates, termed RNA foci, in neuronal nuclei that sequester important RNA-binding proteins and second, the generation of toxic, dipeptide repeat (DPR) proteins, due to the repeat RNA mediating its own translation (Fig. 1) [3]. In this review, we will cover recent findings concerning the role of each of these mechanisms in C9FTD/ALS and their relevance for developing therapeutics. Key findings relating to loss or gain of function are summarized in Table 1 [4,5<sup>a</sup>,7<sup>a</sup>,8<sup>aa</sup>,9<sup>a</sup>–11<sup>a</sup>,12–14,15<sup>a</sup>,16<sup>a</sup>,17<sup>aa</sup>,18,19<sup>a</sup>–26<sup>a</sup>,27<sup>aa</sup>,28<sup>aa</sup>,29<sup>a</sup>–31<sup>a</sup>].

## GAIN AND LOSS OF FUNCTION IN NONCODING REPEAT EXPANSION DISORDERS

Noncoding repeat expansions have been commonly assigned into two groups: those which cause loss of function of the protein in which the mutation resides (fragile X syndrome and Friedreich's ataxia),

Department of Neurodegenerative Disease, UCL Institute of Neurology, London, UK

Correspondence to Adrian M. Isaacs, Department of Neurodegenerative Disease, UCL Institute of Neurology, London, WC1N 3BG, UK. Tel: +44 2078375470; e-mail: a.isaacs@prion.ucl.ac.uk

**Curr Opin Neurol** 2014, 27:515–523

DOI:10.1097/WCO.0000000000000130

This is an open access article distributed under the Creative Commons Attribution License 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**KEY POINTS**


- Evidence exists supporting a role for both gain-of-function and loss-of-function mechanisms in C9FTD/ALS.
- The weight of evidence currently suggests gain of function is the primary disease mechanism.
- Loss of function is more likely to modulate the disease phenotype than be a primary cause of disease.
- Therapeutics that target gain-of-function mechanisms, such as ASOs, are a promising avenue for therapy.

and those in which a gain-of-function mechanism has been identified because of repeat RNA species [myotonic dystrophy, fragile X-associated tremor/ataxia syndrome (FXTAS), Huntington’s disease-like 2 and spinocerebellar ataxia types 8, 10, 12 and 31]. In the fragile X mental retardation protein (*FMRP*) gene, CGG repeat size determines the clinical syndrome; expansions of 55–200 repeats result in the gain-of-function disease FXTAS [32,33], whereas

expansions of more than 200 repeats lead to hypermethylation of the *FMRP* gene, which silences transcription, leading to a loss of FMRP function and fragile X syndrome [34]. There is no evidence for such a bimodal mechanism in C9FTD/ALS, indeed, a large study found no effect of repeat size on clinical presentation [35]. The first mechanism attributed to the gain-of-function noncoding repeat expansion diseases was toxic functions of the repeat RNA [36]. Recently, a novel mechanism was identified whereby expanded CAG repeats are translated in the absence of an ATG initiation codon, termed repeat-associated non-ATG dependent (RAN) translation [37]. RAN translation has now been found to be common to several noncoding repeat expansions, including C9FTD/ALS [38], increasing the spectrum of potential mechanisms in disease.

**LOSS OF FUNCTION OF C9ORF72 PROTEIN**

The main support for a loss-of-function mechanism in C9FTD/ALS is numerous reports of decreased transcript levels of all three *C9orf72* mRNA variants



**FIGURE 1.** Potential mechanisms of disease in C9FTD/ALS. AP, alanine-proline; DPR, dipeptide repeat; GA, glycine-alanine; GP, glycine-proline; GR, glycine-arginine; PR, proline-arginine.

**Table 1.** Summary of the key evidence for gain-and loss-of-function mechanisms in C9FTD/ALS

	For (+) or against (-) hypothesis	Study
<b>Loss-of-function</b>		
Reduced mRNA and protein expression		
All mRNA isoforms reduced in C9FTD/ALS patient tissue and iPSC neurons	+	[1,4,5 <sup>a</sup> -7 <sup>a</sup> ,8 <sup>ab</sup> ]
Reduced C9orf72 protein in C9FTD/ALS patient brain	+	[9 <sup>a</sup> ]
Hypermethylation of <i>C9orf72</i> promoter reduces transcript levels and correlates with shorter disease duration	+	[10 <sup>a</sup> ]
Hypermethylation of <i>C9orf72</i> promoter reduces transcript levels, increasing resistance to cellular stress and reducing RNA foci and DPR proteins	-	[11 <sup>a</sup> ]
Models of loss of protein		
Dysregulation of cellular trafficking associated with reduction of C9orf72 protein	+	[12-14]
<i>C9orf72</i> orthologue knockdown in zebrafish has a motor phenotype	+	[15 <sup>a</sup> ]
<i>C9orf72</i> orthologue knockout in <i>Caenorhabditis elegans</i> has a motor phenotype	+	[16 <sup>a</sup> ]
<i>C9orf72</i> orthologue knockdown in mice has no motor phenotype	-	[17 <sup>ab</sup> ]
Genetics		
Lack of coding mutations in C9orf72 protein in C9ALS	-	[18]
Homozygous C9FTD case not as severe clinically and pathologically as pure loss-of-function diseases	-	[5 <sup>a</sup> ]
<b>Gain-of-function: RNA</b>		
RNA foci		
Sense and antisense foci identified in C9FTD/ALS patient tissue and iPSC neurons	+	[1,6 <sup>a</sup> ,8 <sup>ab</sup> ,17 <sup>ab</sup> ,19 <sup>a</sup> -22 <sup>a</sup> ]
Sense foci burden correlates with age-at-onset in C9FTD	+	[19 <sup>a</sup> ]
Sequestration of RNA-binding proteins		
ADARB2 – siRNA reduces sense RNA foci in C9ALS iPSC neurons and exacerbates glutamate-induced toxicity in control iPSC neurons	+	[8 <sup>ab</sup> ]
hnRNP A1 and pur-alpha – colocalization with sense RNA foci in iPSC motor neurons	+	[22 <sup>a</sup> ]
hnRNP A3 – identified in neuronal cytoplasmic and intranuclear inclusions	+	[23 <sup>a</sup> ]
hnRNP H – colocalization with sense RNA foci in patient brain	+	[24 <sup>a</sup> ]
Nucleolin – colocalization with sense RNA foci in patient brain and cells, indications of nucleolar stress	+	[25 <sup>a</sup> ]
Pur-alpha – overexpression rescues GGGGCC-dependent degeneration in <i>Drosophila</i>	+	[26 <sup>a</sup> ]
Antisense oligonucleotide (ASO) treatment		
ASOs reduce sense RNA foci and reverse transcriptome changes and toxicity in patient fibroblasts and iPSC neurons	+	[8 <sup>ab</sup> ,17 <sup>ab</sup> ,22 <sup>a</sup> ]
<b>Gain-of-function: DPR proteins</b>		
Inclusion pathology		
All DPR proteins found in neuronal cytoplasmic inclusions in C9FTD/ALS patient brain and iPSC neurons	+	[6 <sup>a</sup> ,20 <sup>a</sup> ,21 <sup>a</sup> ,27 <sup>ab</sup> ,28 <sup>ab</sup> ,29 <sup>a</sup> ]
Poly-GA DPR protein inclusions found prior to TDP-43 inclusions in patient brain	+	[30 <sup>a</sup> ]
Toxicity		
GA DPR protein inclusion distribution does not correlate with neurodegeneration in C9FTD/ALS patient brain	-	[31 <sup>a</sup> ]
Poly-GP and poly-PR DPR proteins made from GGGGCC repeats are toxic to HEK293 cells (other DPR proteins not assessed)	+	[21 <sup>a</sup> ]

ALS, amyotrophic lateral sclerosis; DPR, dipeptide repeat; FTD, frontotemporal dementia; GA, glycine-alanine; GP, glycine-proline; hnRNP, heterogeneous nuclear ribonucleoprotein; iPSC, induced pluripotent stem cell; PR, proline-arginine.

in patient-derived cells and tissue [1,4,5<sup>■</sup>–7<sup>■</sup>,8<sup>■</sup>]. Reduced levels of *C9orf72* transcripts may be due to hypermethylation of the *C9orf72* promoter or increased histone methylation [7<sup>■</sup>,10<sup>■</sup>,11<sup>■</sup>]. Unexpectedly, lower levels of *C9orf72* transcripts are also identified in ALS/FTD cases without *C9orf72* repeat expansion [15<sup>■</sup>], suggesting that loss of *C9orf72* could be part of a common pathway affected in these diseases. Current studies on *C9orf72* protein are limited by a lack of specific antibodies. However, one study developed a new *C9orf72* antibody which detects a protein of 48 kDa in human cell lines that is specifically reduced following treatment with siRNA targeting *C9orf72* [9<sup>■</sup>]. This antibody was used to show reduced *C9orf72* protein in frontal cortex, but not cerebellum, of C9FTD/ALS patient brain compared to ALS cases without *C9orf72* repeat expansion [9<sup>■</sup>], consistent with findings of reduced transcript levels.

When the mutation was discovered in 2011, *C9orf72* protein was of unknown function. However, recent studies show that *C9orf72* has a high homology to differentially expressed in normal and neoplasia proteins [13,14]. This family of proteins function as guanine nucleotide exchange factors (GEFs) that activate Rab guanosine 5'-triphosphate (GTP)ase and therefore regulate membrane trafficking [13,14]. Consistent with this, knockdown of *C9orf72* leads to reduced endocytosis and dysregulated autophagy in human neuroblastoma cells [12]. Impaired autophagy and endolysosomal degradation are implicated in neurodegenerative diseases [39], thus these data provide a basis for loss of function playing a role in C9FTD/ALS. However, better antibodies are urgently needed to definitively determine the cellular distribution and function of *C9orf72* protein; analysis of innate GEF activity will also provide important insight into *C9orf72* function.

Knockdown of the zebrafish orthologue of *C9orf72* (*zC9orf72*) with antisense morpholino oligonucleotides leads to axonopathy and motor deficits, which can be rescued by expression of human *C9orf72* [15<sup>■</sup>]. Homozygous knockout of the worm orthologue of *C9orf72* (alfa-1) also results in motor phenotypes [16<sup>■</sup>]. Together, these data suggest that the loss of *C9orf72* protein can lead to motor abnormalities, which argues for a role of loss of function in C9FTD/ALS. Conversely, intracerebroventricular delivery to adult mice of antisense oligonucleotides (ASOs) targeting *C9orf72* leads to knockdown of *C9orf72* throughout the central nervous system but does not result in any motor or behavioural phenotypes [17<sup>■</sup>]. It is possible that knockdown of *C9orf72* during development has differing effects to knockdown in adults which could explain the

discrepancy between the models. Also arguing against a loss-of-function mechanism, no mutations have been found in coding regions of the *C9orf72* gene [18]. Additionally, a rare homozygous *C9orf72* repeat expansion case did not exhibit clinical or pathological features outside the usual disease spectrum as would be expected from reports of homozygous cases in pure loss-of-function diseases [5<sup>■</sup>].

Reduced *C9orf72* transcription via cytosine-phosphate-guanine (CpG) hypermethylation correlated with shorter disease duration suggesting *C9orf72* reduction may play a role in disease [10<sup>■</sup>]. A second study showed *C9orf72* promoter CpG hypermethylation reduced *C9orf72* mRNA levels, but also decreased RNA foci and RAN protein formation in C9FTD/ALS lymphoblasts and brains. Furthermore, treatment with a demethylating agent increased the vulnerability of C9FTD/ALS lymphoblasts to external stressors, suggesting reduction of *C9orf72* levels may be a protective mechanism [11<sup>■</sup>].

On the basis of evidence described above, haploinsufficiency of *C9orf72* may cause defects in endosomal and autophagic processes and motor function. Therefore, the degree of reduction of *C9orf72* may well modulate the disease phenotype. However, clinical data suggest that loss of function is unlikely to be the predominant causative mechanism for neurodegeneration in C9FTD/ALS.

### GAIN OF FUNCTION: REPEAT RNA

RNA gain of function is a common mechanism in noncoding repeat expansion diseases [40]. The proposed mode of action in these diseases is via sequestration of essential RNA-binding proteins into aggregates of repeat-containing RNA foci, in the nucleus of affected cells. The most clearly defined molecular mechanism occurs in myotonic dystrophy (DM), a neuromuscular disease caused by CTG or CCTG repeat expansions in the dystrophin myotonia protein kinase (*DMPK*) or zinc finger protein 9 (*ZNF9*) genes (DM type 1 or 2, respectively) [36]. In DM type 1 or 2, RNA foci containing CUG repeat RNA sequester the splicing factor muscleblind-like protein 1, whose loss results in the mis-splicing of a muscle-specific chloride channel that is directly responsible for the myotonia observed in patients [41]. On the basis of this very clear example of RNA gain of function, great effort has focussed on investigating this mechanism in C9FTD/ALS. RNA foci composed of sense and antisense repeat RNA are present in frontal cortex, hippocampus, cerebellum and spinal cord of C9FTD/ALS patients [1,6<sup>■</sup>,8<sup>■</sup>,17<sup>■</sup>,19<sup>■</sup>–22<sup>■</sup>]. In support of a role in disease, increased burden of RNA foci in neurons correlates with lower age-at-onset of disease in

C9FTD cases, with the strongest correlation with sense foci in the frontal cortex, the region most affected in FTD [19<sup>■</sup>]. RNA foci were present in several types of glial cells (astrocytes, microglia and oligodendrocytes), but are predominantly a neuronal phenotype [17<sup>■</sup>,19<sup>■</sup>,20<sup>■</sup>], which is reflective of relative expression levels of *C9orf72* in these cell types in mouse [42]. This raises the possibility that toxicity could arise from non-cell-autonomous routes. Indeed, astrocytes derived from familial and sporadic ALS patients, including C9ALS cases, can exert toxicity to motor neurons [43].

Although RNA foci can be found in the cytoplasm, the vast majority are localized in the nucleus [17<sup>■</sup>,19<sup>■</sup>]. Several studies have employed biochemical techniques to identify binding partners of the expanded sense repeat *in vitro* using differing methods and sources of protein, reviewed in [44]. Sequestration of some of these proteins into sense RNA foci has been assessed in patient-derived cells and tissue (Table 2). Some inconsistency has been observed between studies, but investigations in larger cohorts will be needed to clarify whether this is due to variation between brain regions or patients or differences due to detection protocols. Splicing factors constituted a large proportion of identified RNA-binding proteins. Of these, the splicing factors heterogeneous nuclear ribonucleoprotein (hnRNP) A1, hnRNP H and serine/arginine-rich splicing factor 2 (SRSF2; also known as SC35) are found to be sequestered into sense RNA foci [22<sup>■</sup>,24<sup>■</sup>,45<sup>■</sup>]. Loss of function of these proteins would be predicted to cause downstream changes in splicing in target mRNAs. All these splicing factors affect splicing of a wide variety of targets, so a major challenge will be to determine whether specific targets are responsible

for neurodegeneration and to confirm specific alteration of these targets in C9FTD/ALS patients.

Proteins involved in nuclear mRNA export also bind to GGGGCC repeats and are sequestered into foci containing sense RNA transcripts [22<sup>■</sup>,45<sup>■</sup>]. Aly/REF export factor (ALYREF) functions as an adaptor for mature RNA, transferring it through the nuclear RNA export factor 1 pathway to the nuclear pore for export [46]. Pur-alpha has also been implicated in nuclear export and trafficking in dendritic RNA granules for the regulation of local translation [47,48]. Another RNA-binding protein, hnRNP A3, which binds GGGGCC repeat RNA *in vitro*, but does not colocalize with RNA foci [23<sup>■</sup>], is also proposed to function in a similar manner [49]. These proteins may contribute to export of *C9orf72* repeat-containing RNA from the nucleus for *C9orf72* and RAN protein translation or degradation. Sequestration of these factors into RNA foci may also prevent export of their other target mRNAs and affect downstream cellular functions.

Other proteins found to interact with GGGGCC repeats were adenosine deaminase, RNA-specific, B2 (ADARB2) [8<sup>■</sup>] and nucleolin [25<sup>■</sup>]. ADARB2 colocalizes with sense RNA foci in C9ALS induced pluripotent stem cell (iPSC)-differentiated motor neurons and patient motor cortex, leading to significant nuclear accumulation of the protein compared with controls. Unexpectedly, knockdown of ADARB2 results in reduced RNA foci in iPSC motor neurons, suggesting a role in foci formation or stabilization. ADARB2 (also known as ADAR3 or RED2) is a member of the ADAR (adenosine deaminase, RNA specific) RNA editing family, but unlike the other members, lacks editing activity [50]. Decrease in another member of this family, ADAR2,

**Table 2.** Reported sequestration of RNA-binding proteins into sense RNA foci in patient cells and tissue

RNA-binding protein	C9ALS or FTD	Patient-derived cells	Patient tissue	Study
ADARB2	ALS	iPSC-differentiated neurons	Motor cortex	[8 <sup>■</sup> ]
ALYREF	ALS	ND	Cerebellum and spinal cord	[45 <sup>■</sup> ]
hnRNP A1	ALS and ALS/FTD	iPSC-differentiated motor neurons		[22 <sup>■</sup> ]
	ALS	ND	Cerebellum	[45 <sup>■</sup> ]
hnRNP H <sup>a</sup>	ALS	ND	Cerebellum	[24 <sup>■</sup> ]
	ALS	Cerebellum and cerebellum and spinal cord		[45 <sup>■</sup> ]
Nucleolin	ALS	ND	Motor cortex	[25 <sup>■</sup> ]
Pur-alpha	ALS and ALS/FTD	iPSC-differentiated motor neurons		[22 <sup>■</sup> ]
SRSF2 (SC35)	ALS	ND	Cerebellum	[24 <sup>■</sup> ]
	ALS	ND	Cerebellum and spinal cord	[45 <sup>■</sup> ]

ADARB2, adenosine deaminase, RNA-specific, B2; ALS, amyotrophic lateral sclerosis; ALYREF, Aly/REF export factor; FTD, frontotemporal dementia; hnRNP, heterogeneous nuclear ribonucleoprotein; iPSC, induced pluripotent stem cell; ND, not determined; SRSF2 (SC35), serine/arginine-rich splicing factor 2 (also known as SC35).

<sup>a</sup>Sequestration of hnRNP H into RNA foci was not observed in C9FTD iPSC-neurons [6<sup>■</sup>].

is thought to underlie excitotoxic loss of motor neurons in ALS via reduced editing of AMPA receptors subunits [51].

GGGGCC RNA and DNA repeats exhibit stable G-quadruplex formation *in vitro* [25<sup>■</sup>,52<sup>■</sup>,53]. These structures are involved in telomere stability, RNA splicing, transport and degradation and regulation of translation [54–56]. A recent study investigated binding of proteins to GGGGCC RNA hairpins, GGGGCC G-quadruplexes and antisense GGCCCC hairpins [25<sup>■</sup>]. The nucleolar protein nucleolin was shown to have specificity for sense G-quadruplex structures, and dispersal of nucleolin staining in the nucleus was observed in C9ALS patient-derived cells and tissue, with sequestration into sense RNA foci visible in the motor cortex. Consequences of impaired nucleolar function were also observed, including decreased RNA processing and an increase in the number of P bodies, which are ribonucleoprotein complexes involved in the degradation of untranslated RNAs. These changes could be recapitulated when GGGGCC repeats were expressed in a cell line, implicating nucleolar stress as a gain-of-function RNA mechanism in C9FTD/ALS.

It is intriguing to speculate whether the heterogeneity seen in C9FTD/ALS could be attributed to differential sequestration of RNA-binding proteins between brain regions and patients, because of protein abundance or availability or RNA foci burden. No protein sequestration into antisense RNA foci has yet been identified, but this could be important as ASOs targeting sense RNA did not reverse all transcriptome changes in C9FTD/ALS-derived fibroblasts [17<sup>■</sup>]. This suggests antisense transcripts may also cause transcriptional changes, potentially through sequestration of RNA-binding proteins. Repeat RNA may also exert neurotoxic effects through processes other than RNA foci generation. In a fly model of myotonic dystrophy and a cell model of Huntington's disease, expression of expanded repeats led to double stranded RNAs that were processed by the dicer complex into short siRNAs that exert toxicity by silencing complementary CAG or CTG repeat-containing transcripts, respectively [57,58]. It will be interesting to determine whether this pathway contributes to C9FTD/ALS. In a similar manner, antisense transcripts may regulate the levels of sense transcripts, and vice versa, via antisense-mediated RNA degradation. This may explain why sense and antisense RNA foci are rarely found within the same cell [19<sup>■</sup>,21<sup>■</sup>]. In Huntington's disease, antisense transcripts reduce huntingtin protein expression, partially via dicer [59]. This mechanism could also contribute to the reduced *C9orf72* transcript levels observed in C9FTD/ALS patients.

In summary, the greatest evidence in support for a role of RNA gain of function in C9FTD/ALS are the abundant sense and antisense RNA foci in patient tissue that correlate with clinical phenotypes and can sequester RNA-binding proteins. However, a clear mechanism definitively linking sequestration of specific RNA-binding proteins to disease pathogenesis (as is the case for myotonic dystrophy) is currently lacking.

### GAIN OF FUNCTION: DIPEPTIDE REPEAT PROTEINS

The other novel and potentially toxic species in C9FTD/ALS are the DPR proteins formed by RAN translation of the expanded repeat [20<sup>■</sup>,21<sup>■</sup>,27<sup>■</sup>,28<sup>■</sup>,29<sup>■</sup>]. DPR proteins are translated from all frames of the GGGGCC repeat resulting in polymers of glycine-alanine (GA), glycine-proline (GP) and glycine-arginine (GR) in the sense frames, and glycine-proline (GP), alanine-proline (AP) and proline-arginine (PR) in the antisense frames. Although poly-GP is translated from both sense and antisense RNA, translation has been found to continue after the repeat expansion (using antibodies against downstream regions), and thus these poly-GP proteins have different carboxy terminal tails that may affect their function [21<sup>■</sup>]. All DPR proteins form widespread neuronal cytoplasmic inclusions in patient brain [20<sup>■</sup>,21<sup>■</sup>,27<sup>■</sup>,28<sup>■</sup>,29<sup>■</sup>] that frequently colocalize with p62-positive [but not TAR DNA-binding protein 43 (TDP-43)-positive] inclusions [28<sup>■</sup>,29<sup>■</sup>,31<sup>■</sup>,60]. Poly-GP, and poly-GA DPR proteins additionally display dot-like neuronal intranuclear inclusions [27<sup>■</sup>,28<sup>■</sup>,31<sup>■</sup>]. Unlike RNA foci, DPR inclusions appear to be an exclusively neuronal phenotype [27<sup>■</sup>,31<sup>■</sup>,60], possibly reflective of the clearance ability of mitotic cells.

A detailed pathological analysis of C9FTD/ALS cases found that TDP-43, but not poly-GA pathology, correlates with neurodegeneration [31<sup>■</sup>], suggesting a lack of pathogenicity of poly-GA inclusions. However, this does not rule out toxicity of soluble GA polymers or other DPR proteins. Interestingly, rare *C9orf72* cases have been reported with DPR but not TDP-43 pathology [28<sup>■</sup>,30<sup>■</sup>]. This suggests DPR proteins can be toxic without invoking TDP-43 dysfunction.

### DISSECTION OF GAIN-OF-FUNCTION MECHANISMS

We have reviewed evidence for gain-of-function and loss-of-function mechanisms in C9FTD/ALS. However, it still remains to be determined which species and pathways are responsible for neurodegeneration

and clinical phenotypes in disease. Co-occurrence of sense and antisense RNA foci with DPR, p62 or TDP-43 protein inclusions is only as frequent as expected by chance [8<sup>11</sup>,19<sup>11</sup>–21<sup>11</sup>,45<sup>11</sup>], suggesting a lack of interdependence between species and distinct toxic mechanisms. Overexpression of expanded GGGGCC repeats (outside the context of the *C9orf72* gene) can induce RNA foci formation [20<sup>11</sup>,24<sup>11</sup>] and DPR protein production [20<sup>11</sup>,21<sup>11</sup>,28<sup>11</sup>]. Repeats can also exert toxicity in cell lines [21<sup>11</sup>,24<sup>11</sup>,25<sup>11</sup>], flies [26<sup>11</sup>] and zebrafish [24<sup>11</sup>], suggesting that gain-of-function mechanisms are sufficient for neurodegeneration. Studies have yet to clearly attribute observed toxicity to repeat RNA or DPR protein species, but one study showed that increasing expression of *C9orf72* repeats specifically in the GP (sense) and PR (antisense) frames can exacerbate toxicity in human HEK293T cells [21<sup>11</sup>], showing that these DPR proteins can affect cell viability. Impaired degradation through the autophagic system is consistent with the accumulation of p62 and ubiquitin pathology that is abundant in C9FTD/ALS cases [61], and the sensitivity of C9FTD iPSCs specifically to autophagic stressors [6<sup>11</sup>]. However, it is not clear whether these effects are due to loss of the normal cellular function of *C9orf72* or gain of function due to the accumulation of protein aggregates.

### THERAPEUTIC TARGETING OF GAIN-OF-FUNCTION MECHANISMS

There is considerable excitement about the possibility of ASOs for the treatment of C9FTD/ALS. One reason for this is that ASOs should ameliorate both repeat RNA and DPR protein toxicity, and therefore do not need to wait for a better understanding of the contribution of each of these mechanisms to disease pathogenesis. ASOs will not alleviate loss of *C9orf72* function, but as discussed above, the weight of evidence currently suggests gain of function is likely to be the primary mechanism to address therapeutically. ASOs targeting sense transcripts reduce sense RNA foci and ameliorate transcriptome changes and toxicity in C9FTD/ALS-derived cells [8<sup>11</sup>,17<sup>11</sup>,22<sup>11</sup>]. ASOs targeting antisense transcripts may also be required as it was proposed that antisense RNA-mediated mechanisms were responsible for the proportion of transcriptional changes that remain dysregulated after treatment of patient cells with ASOs targeting sense repeats [17<sup>11</sup>]. In addition, antisense DPR proteins have been shown to be toxic to cells [21<sup>11</sup>]. Given the precedent of a clinical trial for ASOs targeting superoxide dismutase 1 (*SOD1*) for ALS patients with *SOD1* mutations [62], ASO treatment is currently the most promising prospect for treating C9FTD/ALS. The development of small

molecules that specifically bind the secondary structure formed by the GGGGCC repeats is another promising area for therapeutic intervention [52<sup>11</sup>,63<sup>11</sup>]. Such small molecules would also be predicted to prevent both repeat RNA and DPR protein mechanisms, with the potential advantage of simpler delivery.

### CONCLUSION

In the 3 years since the discovery of the *C9orf72* mutation in ALS and FTD, intense investigations have begun to unfold the mechanisms at play in these diseases. Haploinsufficiency of *C9orf72* may cause defects in endosomal and autophagic processes that lead to dysfunction of the motor system, but clinical data do not support causation of disease. Cellular and animal models inform us that gain-of-function mechanisms from expanded *C9orf72* repeats are sufficient to cause neurodegeneration. As has been found in other diseases classically thought of as either loss-of-function or gain-of-function diseases, it is likely that both mechanisms contribute to different aspects of the heterogeneous phenotypes in these diseases. As the same *C9orf72* mutation appears to result in a spectrum of disease phenotypes, differential regulation of toxic species or the degree of loss of function may play a role in differing disease presentations. Finally, current evidence suggests gain-of-function-based therapies such as ASOs hold promise for C9FTD/ALS.

### Acknowledgements

AMI receives funding from Alzheimer's Research UK, The Motor Neurone Disease Association, the MHMS General Charitable Trust and the UK Medical Research Council.

### Conflicts of interest

There are no conflicts of interest.

### REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. DeJesus-Hernandez M, Mackenzie IR, Boeve BF, *et al.* Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* 2011; 72:245–256.
2. Renton AE, Majounie E, Waite A, *et al.* A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 2011; 72:257–268.
3. Lashley T, Hardy J, Isaacs AM. RANTing about C9orf72. *Neuron* 2013; 77:597–598.
4. Gijssels I, Van Langenhove T, van der ZJ, *et al.* A C9orf72 promoter repeat expansion in a Flanders-Belgian cohort with disorders of the frontotemporal lobar degeneration-amyotrophic lateral sclerosis spectrum: a gene identification study. *Lancet Neurol* 2012; 11:54–65.

5. Fratta P, Poulter M, Lashley T, *et al*. Homozygosity for the C9orf72 GGGGCC repeat expansion in frontotemporal dementia. *Acta Neuropathol* 2013; 126:401–409.
- First description of homozygous C9orf72 repeat expansion carrier.
6. Almeida S, Gascon E, Tran H, *et al*. Modeling key pathological features of frontotemporal dementia with C9ORF72 repeat expansion in iPSC-derived human neurons. *Acta Neuropathol* 2013; 126:385–399.
- First description of RNA foci and DPR protein in C9FTD patient-derived iPSC neurons.
7. Belzil VV, Bauer PO, Prudencio M, *et al*. Reduced C9orf72 gene expression in c9FTD/ALS is caused by histone trimethylation, an epigenetic event detectable in blood. *Acta Neuropathol* 2013; 126:895–905.
- This article identifies histone methylation as a mechanism for reduction of C9orf72 transcription in C9FTD/ALS.
8. Donnelly CJ, Zhang PW, Pham JT, *et al*. RNA toxicity from the ALS/FTD C9ORF72 expansion is mitigated by antisense intervention. *Neuron* 2013; 80:415–428.
- One of three articles showing therapeutic potential of ASOs in C9ALS/FTD patient-derived cells.
9. Waite AJ, Baumer D, East S, *et al*. Reduced C9orf72 protein levels in frontal cortex of amyotrophic lateral sclerosis and frontotemporal degeneration brain with the C9ORF72 hexanucleotide repeat expansion. *Neurobiol Aging* 2014; 35:e5–13; 1779.
- First description of reduced C9orf72 protein in C9ALS/FTD patient tissue.
10. Xi Z, Zinman L, Moreno D, *et al*. Hypermethylation of the CpG island near the G4C2 repeat in ALS with a C9orf72 expansion. *Am J Hum Genet* 2013; 92:981–989.
- First description of CpG island methylation as a mechanism for reduced C9orf72 transcription.
11. Liu EY, Russ J, Wu K, *et al*. C9orf72 hypermethylation protects against repeat expansion-associated pathology in ALS/FTD. *Acta Neuropathol* 2014; Epub 2014 May 8.
- This study found a correlation between methylation status and clinical phenotype, suggesting that hypermethylation and thus decreased C9orf72 is protective.
12. Farg MA, Sundaramoorthy V, Sultana JM, *et al*. C9ORF72, implicated in amyotrophic lateral sclerosis and frontotemporal dementia, regulates endosomal trafficking. *Hum Mol Genet* 2014; 23:3579–3595.
13. Levine TP, Daniels RD, Gatta AT, *et al*. The product of C9orf72, a gene strongly implicated in neurodegeneration, is structurally related to DENN RabGEFs. *Bioinformatics* 2013; 29:499–503.
14. Zhang D, Iyer LM, He F, *et al*. Discovery of novel DENN proteins: implications for the evolution of eukaryotic intracellular membrane structures and human disease. *Front Genet* 2012; 3:1–10; Article 283. doi:10.3389/fgene.2012.00283.
15. Ciura S, Lattante S, Le B I. Loss of function of C9orf72 causes motor deficits in a zebrafish model of amyotrophic lateral sclerosis. *Ann Neurol* 2013; 74:180–187.
- First loss of C9orf72 function in zebrafish model.
16. Therrien M, Rouleau GA, Dion PA, *et al*. Deletion of C9ORF72 results in motor neuron degeneration and stress sensitivity in *C. elegans*. *PLoS One* 2013; 8:e83450.
- First loss of C9orf72 in *Caenorhabditis elegans* model.
17. Lagier-Tourenne C, Baughn M, Rigo F, *et al*. Targeted degradation of sense and antisense C9orf72 RNA foci as therapy for ALS and frontotemporal degeneration. *Proc Natl Acad Sci U S A* 2013; 110:E4530–E4539.
- One of three articles showing therapeutic potential of ASOs in C9FTD/ALS patient-derived cells, and one of the first descriptions of antisense RNA foci in C9FTD/ALS patient tissue.
18. Harms MB, Cady J, Zaidman C, *et al*. Lack of C9ORF72 coding mutations supports a gain of function for repeat expansions in amyotrophic lateral sclerosis. *Neurobiol Aging* 2013; 34:2234–2239.
19. Mizielinska S, Lashley T, Norona FE, *et al*. C9orf72 frontotemporal lobar degeneration is characterised by frequent neuronal sense and antisense RNA foci. *Acta Neuropathol* 2013; 126:845–857.
- One of the first descriptions of antisense RNA foci in C9FTD/ALS patient tissue and also reports a correlation between RNA foci burden and clinical phenotypes, suggesting a role for RNA foci in C9FTD/ALS.
20. Gendron TF, Bieniek KF, Zhang YJ, *et al*. Antisense transcripts of the expanded C9ORF72 hexanucleotide repeat form nuclear RNA foci and undergo repeat-associated non-ATG translation in c9FTD/ALS. *Acta Neuropathol* 2013; 126:829–844.
- One of the first descriptions of antisense RNA foci and DPR protein in C9FTD/ALS patient tissue.
21. Zu T, Liu Y, Banez-Coronel M, *et al*. RAN proteins and RNA foci from antisense transcripts in C9ORF72 ALS and frontotemporal dementia. *Proc Natl Acad Sci U S A* 2013; 110:E4968–4977.
- One of the first descriptions of antisense RNA foci and DPR protein in C9FTD/ALS patient tissue and provides evidence for cellular toxicity of DPR proteins.
22. Sareen D, O'Rourke JG, Meera P, *et al*. Targeting RNA foci in iPSC-derived motor neurons from ALS patients with a C9ORF72 repeat expansion. *Sci Transl Med* 2013; 5:208ra149.
- One of three articles showing therapeutic potential of ASOs in C9FTD/ALS patient-derived cells, and also reports sequestration of RNA-binding proteins into RNA foci.
23. Mori K, Lammich S, Mackenzie IR, *et al*. hnRNP A3 binds to GGGGCC repeats and is a constituent of p62-positive/TDP43-negative inclusions in the hippocampus of patients with C9orf72 mutations. *Acta Neuropathol* 2013; 125:413–423.
- Identification of hnRNP A3 as a binding protein of expanded C9orf72 repeat RNA and its presence in a subset of neuronal inclusions.
24. Lee YB, Chen HJ, Peres JN, *et al*. Hexanucleotide repeats in ALS/FTD form length-dependent RNA foci, sequester RNA binding proteins, and are neurotoxic. *Cell Rep* 2013; 5:1178–1186.
- Identification of hnRNP H as a protein sequestered by expanded C9orf72 repeat RNA.
25. Haeusler AR, Donnelly CJ, Periz G, *et al*. C9orf72 nucleotide repeat structures initiate molecular cascades of disease. *Nature* 2014; 507:195–200.
- Identification of nucleolin as a binding protein of G-quadruplexes formed by expanded C9orf72 repeat RNA and evidence suggesting a role for nucleolar stress in C9FTD/ALS.
26. Xu Z, Poidevin M, Li X, *et al*. Expanded GGGGCC repeat RNA associated with amyotrophic lateral sclerosis and frontotemporal dementia causes neurodegeneration. *Proc Natl Acad Sci U S A* 2013; 110:7778–7783.
- First description of C9orf72 repeats causing neurodegeneration in *Drosophila* and identification of repeat binding proteins.
27. Ash PE, Bieniek KF, Gendron TF, *et al*. Unconventional translation of C9ORF72 GGGGCC expansion generates insoluble polypeptides specific to c9FTD/ALS. *Neuron* 2013; 77:639–646.
- The first description, with reference [28], of RAN translation of the C9orf72 repeat in C9FTD/ALS.
28. Mori K, Weng SM, Arzberger T, *et al*. The C9orf72 GGGGCC repeat is translated into aggregating dipeptide-repeat proteins in FTL/ALS. *Science* 2013; 339:1335–1338.
- The first description, with reference [27], of RAN translation of the C9orf72 repeat in C9FTD/ALS.
29. Mori K, Arzberger T, Grasser FA, *et al*. Bidirectional transcripts of the expanded C9orf72 hexanucleotide repeat are translated into aggregating dipeptide repeat proteins. *Acta Neuropathol* 2013; 126:881–893.
- One of the first descriptions of antisense DPR protein in C9FTD/ALS patient tissue.
30. Proudfoot M, Gutowski NJ, Edbauer D, *et al*. Early dipeptide repeat pathology in a frontotemporal dementia kindred with C9ORF72 mutation and intellectual disability. *Acta Neuropathol* 2014; 127:451–458.
- Description of two C9orf72 repeat expansion carriers with predominant and early poly-GA pathology and limited TDP-43 pathology.
31. Mackenzie IR, Arzberger T, Kremmer E, *et al*. Dipeptide repeat protein pathology in C9ORF72 mutation cases: clinico-pathological correlations. *Acta Neuropathol* 2013; 126:859–879.
- An in-depth clinicopathological study showing that TDP-43 rather than poly-GA inclusions correlate with neurodegeneration in C9FTD/ALS.
32. Hagerman P. Fragile X-associated tremor/ataxia syndrome (FXTAS): pathology and mechanisms. *Acta Neuropathol* 2013; 126:1–19.
33. Hagerman RJ, Leehay M, Heinrichs W, *et al*. Intention tremor, parkinsonism, and generalized brain atrophy in male carriers of fragile X. *Neurology* 2001; 57:127–130.
34. Wang T, Bray SM, Warren ST. New perspectives on the biology of fragile X syndrome. *Curr Opin Genet Dev* 2012; 22:256–263.
35. van Blitterswijk M, DeJesus-Hernandez M, Niemantsverdriet E, *et al*. Association between repeat sizes and clinical and pathological characteristics in carriers of C9ORF72 repeat expansions (Xpansize-72): a cross-sectional cohort study. *Lancet Neurol* 2013; 12:978–988.
- First large study assessing repeat size and clinicopathological correlations in ALS, FTD and ALS/FTD C9orf72 repeat expansion carriers.
36. Cooper TA, Wan L, Dreyfuss G. R.N.A and disease. *Cell* 2009; 136:777–793.
37. Zu T, Gibbens B, Doty NS, *et al*. Non-ATG-initiated translation directed by microsatellite expansions. *Proc Natl Acad Sci U S A* 2011; 108:260–265.
38. Cleary JD, Ranum LP. Repeat associated non-ATG (RAN) translation: new starts in microsatellite expansion disorders. *Curr Opin Genet Dev* 2014; 26C:6–15.
39. Nixon RA. The role of autophagy in neurodegenerative disease. *Nat Med* 2013; 19:983–997.
40. Wojciechowska M, Krzyzosiak WJ. Cellular toxicity of expanded RNA repeats: focus on RNA foci. *Hum Mol Genet* 2011; 20:3811–3821.
41. Mankodi A, Takahashi MP, Jiang H, *et al*. Expanded CUG repeats trigger aberrant splicing of CIC-1 chloride channel premRNA and hyperexcitability of skeletal muscle in myotonic dystrophy. *Mol Cell* 2002; 10:35–44.
42. Suzuki N, Maroof AM, Merkle FT, *et al*. The mouse C9ORF72 ortholog is enriched in neurons known to degenerate in ALS and FTD. *Nat Neurosci* 2013; 16:1725–1727.
43. Meyer K, Ferraiuolo L, Miranda CJ, *et al*. Direct conversion of patient fibroblasts demonstrates noncell autonomous toxicity of astrocytes to motor neurons in familial and sporadic ALS. *Proc Natl Acad Sci U S A* 2014; 111:829–832.
44. Vatovec S, Kovanda A, Rogelj B. Unconventional features of C9ORF72 expanded repeat in amyotrophic lateral sclerosis and frontotemporal lobar degeneration. *Neurobiol Aging* 2014; 35:e1–12; 2421.



45. Cooper-Knock J, Walsh MJ, Higginbottom A, *et al.* Sequestration of multiple RNA recognition motif-containing proteins by C9orf72 repeat expansions. *Brain* 2014; 137:2040–2051.
- Identification of multiple RNA-binding proteins sequestered by expanded C9orf72 repeat RNA.
46. Stutz F, Bachi A, Doerks T, *et al.* REF, an evolutionary conserved family of hnRNP-like proteins, interacts with TAP/Mex67p and participates in mRNA nuclear export. *RNA* 2000; 6:638–650.
47. Kanai Y, Dohmae N, Hirokawa N. Kinesin transports RNA: isolation and characterization of an RNA-transporting granule. *Neuron* 2004; 43:513–525.
48. Johnson EM, Kinoshita Y, Weinreb DB, *et al.* Role of Pur alpha in targeting mRNA to sites of translation in hippocampal neuronal dendrites. *J Neurosci Res* 2006; 83:929–943.
49. Ma AS, Moran-Jones K, Shan J, *et al.* Heterogeneous nuclear ribonucleoprotein A3, a novel RNA trafficking response element-binding protein. *J Biol Chem* 2002; 277:18010–18020.
50. Chen CX, Cho DS, Wang Q, *et al.* A third member of the RNA-specific adenosine deaminase gene family, ADAR3, contains both single- and double-stranded RNA binding domains. *RNA* 2000; 6:755–767.
51. Kwak S, Hideyama T, Yamashita T, *et al.* AMPA receptor-mediated neuronal death in sporadic ALS. *Neuropathology* 2010; 30:182–188.
52. Fratta P, Mizielinska S, Nicoll AJ, *et al.* C9orf72 hexanucleotide repeat associated with amyotrophic lateral sclerosis and frontotemporal dementia forms RNA G-quadruplexes. *Sci Rep* 2012; 2:1–6; Article 1016. doi:10.1038/srep01016.
- The first description of the tertiary structures formed by C9orf72 repeat RNA.
53. Reddy K, Zamiri B, Stanley SY, *et al.* The disease-associated r(GGGGCC)<sub>n</sub> repeat from the C9orf72 gene forms tract length-dependent uni- and multi-molecular RNA G-quadruplex structures. *J Biol Chem* 2013; 288:9860–9866.
54. Ji X, Sun H, Zhou H, *et al.* Research progress of RNA quadruplex. *Nucleic Acid Ther* 2011; 21:185–200.
55. Subramanian M, Rage F, Tabet R, *et al.* G-quadruplex RNA structure as a signal for neurite mRNA targeting. *EMBO Rep* 2011; 12:697–704.
56. Collie GW, Parkinson GN. The application of DNA and RNA G-quadruplexes to therapeutic medicines. *Chem Soc Rev* 2011; 40:5867–5892.
57. Banez-Coronel M, Porta S, Kagerbauer B, *et al.* A pathogenic mechanism in Huntington's disease involves small CAG-repeated RNAs with neurotoxic activity. *PLoS Genet* 2012; 8:e1002481.
58. Yu Z, Teng X, Bonini NM. Triplet repeat-derived siRNAs enhance RNA-mediated toxicity in a *Drosophila* model for myotonic dystrophy. *PLoS Genet* 2011; 7:e1001340.
59. Chung DW, Rudnicki DD, Yu L, *et al.* A natural antisense transcript at the Huntington's disease repeat locus regulates HTT expression. *Hum Mol Genet* 2011; 20:3467–3477.
60. Mann DM, Rollinson S, Robinson A, *et al.* Dipeptide repeat proteins are present in the p62 positive inclusions in patients with frontotemporal lobar degeneration and motor neurone disease associated with expansions in C9ORF72. *Acta Neuropathol Commun* 2013; 1:1–13; Article 68. doi:10.1186/2051-5960-1-68.
61. Al Sarraj S, King A, Troakes C, *et al.* p62 positive, TDP-43 negative, neuronal cytoplasmic and intranuclear inclusions in the cerebellum and hippocampus define the pathology of C9orf72-linked FTL and MND/ALS. *Acta Neuropathol* 2011; 122:691–702.
62. Miller TM, Pestronk A, David W, *et al.* An antisense oligonucleotide against SOD1 delivered intrathecally for patients with SOD1 familial amyotrophic lateral sclerosis: a phase 1, randomised, first-in-man study. *Lancet Neurol* 2013; 12:435–442.
63. Zamiri B, Reddy K, Macgregor RB Jr, *et al.* TMPyP4 porphyrin distorts RNA G-quadruplex structures of the disease-associated r(GGGGCC)<sub>n</sub> repeat of the C9orf72 gene and blocks interaction of RNA-binding proteins. *J Biol Chem* 2014; 289:4653–4659.
- The study provides proof of principle for RNA G-quadruplex binding small molecules as a therapeutic strategy in C9FTD/ALS.