

ENVIRONMENTAL Science & Technology

Supporting information for *Environ. Sci. Technol.*, 1994, 28(12), 2014 – 2024, DOI: [10.1021/es00061a009](https://doi.org/10.1021/es00061a009)

HANNIGAN 2014-2024

Terms & Conditions

Electronic Supporting Information files are available without a subscription to ACS Web Editions. The American Chemical Society holds a copyright ownership interest in any copyrightable Supporting Information. Files available from the ACS website may be downloaded for personal use only. Users are not otherwise permitted to reproduce, republish, redistribute, or sell any Supporting Information from the ACS website, either in whole or in part, in either machine-readable form or any other form without permission from the American Chemical Society. For permission to reproduce, republish and redistribute this material, requesters must process their own requests via the RightsLink permission system. Information about how to use the RightsLink permission system can be found at <http://pubs.acs.org/page/copyright/permissions.html>.

Table A: Source Samples Tested for Mutagenicity in the Bacterial Assay

equivalent organic carbon(μg) ^b	-PMS ^a				+PMS ^a			
	mutant fraction ^c ($\times 10^5$)	relative survival ^d	stat signif ^e	CC HC	mutant fraction ^c ($\times 10^5$)	relative survival ^d	stat signif ^e	CC HC
catalyst-equipped automobile exhaust								
control	7.8 \pm 1.8	1.00			9.5 \pm 1.3	1.00		
1.96	12.4 \pm 2.0	0.72	+	-	12.3 \pm 1.5	0.91	-	-
5.87	27.7 \pm 4.0	0.50	+	+	16.0 \pm 2.0	0.79	+	+
17.6	111.2 \pm 18.3	0.19	+	+	32.4 \pm 4.0	0.49	+	+
overall				+				+
noncatalyst automobile exhaust								
control	9.7 \pm 1.5	1.00			7.4 \pm 1.2	1.00		
5.26	27.2 \pm 3.2	0.78	+	+	16.8 \pm 2.0	0.95	+	+
15.8	62.3 \pm 9.2	0.31	+	+	33.6 \pm 3.9	0.63	+	+
47.3					53.8 \pm 7.3	0.36	+	+
overall				+				+
heavy-duty diesel truck exhaust								
control	8.2 \pm 1.1	1.00			7.4 \pm 1.1	1.00		
3.22	23.7 \pm 2.9	0.54	+	+	9.2 \pm 1.5	0.66	-	-
9.67	85.3 \pm 11.1	0.25	+	+	20.4 \pm 2.8	0.53	+	+
29.0	320.5 \pm 65.3	0.07	NC		43.4 \pm 5.9	0.33	+	+
overall				+				+
fuel oil-fired boiler								
control	6.4 \pm 1.1	1.00			9.5 \pm 1.3	1.00		
1.20	27.5 \pm 3.1	0.82	+	+				
2.50					14.6 \pm 1.7	0.90	+	+
3.50	22.4 \pm 2.8	0.76	+	+				
5.00					14.2 \pm 1.8	0.77	+	+
7.50					15.4 \pm 2.0	0.71	+	+
10.0					21.4 \pm 2.7	0.62	+	+
10.5	63.7 \pm 8.5	0.37	+	+				
overall				+				+
natural gas home appliances								
control	13.3 \pm 1.9	1.00			9.3 \pm 1.3	1.00		
1.58	79.1 \pm 11.1	0.34	+	+	34.9 \pm 3.6	0.70	+	+
4.73	155.1 \pm 31.4	0.13	+	+	105.3 \pm 10.4	0.45	+	+
14.2	429.5 \pm 157.0	0.03	NC		250.3 \pm 33.0	0.20	+	+
overall				+				+
fireplace, pine								
control	9.7 \pm 1.8	1.00			12.3 \pm 1.6	1.00		
3.75	30.4 \pm 3.6	0.54	+	+	21.6 \pm 2.7	0.77	+	+
11.25	62.2 \pm 9.8	0.20	+	+	22.7 \pm 3.0	0.65	+	+
33.75	37.5 \pm 30.9	0.01	NC		45.5 \pm 7.3	0.28	+	+
overall				+				+

E 2024-32

Table A: continued

equivalent organic carbon(μg) ^b	-PMS ^a				+PMS ^a			
	mutant fraction ^c ($\times 10^5$)	relative survival ^d	stat signif ^e CC	stat signif ^e HC	mutant fraction ^c ($\times 10^5$)	relative survival ^d	stat signif ^e CC	stat signif ^e HC
fireplace, pine - nonpolar fraction ^{f,g}								
control	12.4 \pm 1.6	1.00			9.4 \pm 1.2	1.00		
3.75	8.0 \pm 1.2	1.20	-	-	10.4 \pm 1.3	0.84	-	-
11.25	12.5 \pm 1.5	1.16	-	-	9.4 \pm 1.3	0.83	-	-
33.75	12.9 \pm 2.0	0.67	-	-	10.3 \pm 1.5	0.63	-	-
overall			-				-	
fireplace, oak ^g								
control	10.5 \pm 1.4	1.00			9.4 \pm 1.2	1.00		
2.35	28.5 \pm 3.9	0.45	+	+	9.4 \pm 1.3	0.79	-	-
7.04	126.7 \pm 26.9	0.09	NC		12.1 \pm 1.8	0.59	-	-
21.13					15.9 \pm 4.3	0.14	+	+
overall			+				+	
fireplace, oak - nonpolar fraction								
control	10.5 \pm 1.4	1.00			9.0 \pm 1.9	1.00		
2.35	16.1 \pm 2.0	0.82	+	+	8.7 \pm 1.5	1.63	-	-
7.04	27.3 \pm 3.1	0.70	+	+	~1.5	1.55	-	-
21.13	108.7 \pm 15.6	0.22	+	+	13.5 \pm 2.4	1.09	+	-
overall			+				-	
fireplace, synthetic log								
control	9.7 \pm 1.8	1.00			12.3 \pm 1.6	1.00		
1.49	24.7 \pm 3.2	0.51	+	+	17.0 \pm 2.3	0.75	+	+
4.46	27.7 \pm 5.4	0.42	+	+	19.5 \pm 2.7	0.65	+	+
13.38	50.6 \pm 6.7	0.32	+	+	21.1 \pm 2.7	0.73	+	+
overall			+				+	
fireplace, synthetic log - nonpolar fraction								
control	10.5 \pm 1.4	1.00			8.0 \pm 1.5	1.00		
1.49	12.2 \pm 1.4	1.15	-	-	9.4 \pm 1.6	1.18	-	-
4.46	15.5 \pm 1.8	0.97	+	+	10.8 \pm 1.6	1.34	-	-
13.38	25.3 \pm 3.0	0.64	+	+	10.7 \pm 1.6	1.26	-	-
overall			+				-	
cigarette smoke								
control	13.0 \pm 1.8	1.00			11.0 \pm 1.6	1.00		
2.73	18.2 \pm 2.3	0.91	+	+	13.4 \pm 1.9	0.92	-	-
8.20	23.4 \pm 3.7	0.50	+	+	14.9 \pm 1.8	1.17	+	+
24.6	31.0 \pm 5.1	0.37	+	+	20.1 \pm 2.8	0.79	+	+
overall			+				+	
roofing tar pot								
control	8.2 \pm 1.1	1.00			7.4 \pm 1.1	1.00		
1.98	11.6 \pm 1.6	0.74	+	-	8.6 \pm 1.4	0.70	-	-
5.93	9.2 \pm 2.0	0.53	-	-	8.4 \pm 1.4	0.76	-	-
17.8	15.0 \pm 1.8	0.63	+	+	7.9 \pm 1.5	0.58	-	-
overall			+				-	

Table A: continued

equivalent organic carbon(μg) ^b	-PMS ^a				+PMS ^a			
	mutant fraction ^c ($\times 10^5$)	relative survival ^d	stat signif ^e		mutant fraction ^c ($\times 10^5$)	relative survival ^d	stat signif ^e	
			CC	HC			CC	HC
charbroiled burger smoke ^f								
control	12.4±1.6	1.00			11.0±1.6	1.00		
3.18	11.2±1.5	1.02	-	-	10.3±1.4	1.29	-	-
9.53	8.9±1.3	1.09	-	-	13.2±1.8	1.04	-	-
28.6	8.1±1.3	0.98	-	-	14.4±2.2	0.75	-	+
overall		-				-		
paved road dust ^g								
control	5.5±1.1	1.00			9.4±1.2	1.00		
0.50	6.4±1.2	0.95	-	-				
1.00	5.7±1.1	0.97	-	-				
2.00	7.3±1.2	1.04	-	-	9.4±1.2	0.88	-	-
6.00	7.7±1.3	0.95	-	-	9.0±1.2	0.83	-	-
18.0	8.6±1.5	0.87	+	-	13.8±1.5	0.92	+	-
overall		-				-		
brake lining wear								
control	8.2±1.1	1.00			7.4±1.1	1.00		
1.44	10.4±1.4	0.83	-	-	6.8±1.1	0.86	-	-
4.33	11.8±1.8	0.57	+	-	7.7±1.2	0.92	-	-
13.0	10.8±2.3	0.65	-	-	6.9±1.2	0.77	-	-
overall		-				-		
tire wear debris								
control	5.8±1.0	1.00			7.1±1.1	1.00		
4.20	6.0±1.1	0.96	-	-	6.7±1.0	0.96	-	-
12.6	10.4±3.4	0.39	+	-	8.4±1.1	1.03	-	-
37.8	13.0±3.7	0.21	+	-	8.6±1.5	0.68	-	-
overall		-				-		
urban vegetative detritus								
control	13.3±1.9	1.00			9.3±1.3	1.00		
1.64	17.9±2.4 ^h	0.87	-	+	9.0±1.4	0.81	-	-
4.93	13.3±2.0	0.86	-	-	9.8±1.5	0.76	-	-
14.8	14.9±1.9	1.12	-	+	10.7±1.7	0.69	-	-
overall		-				-		

^a Tested in the absence of postmitochondrial supernatant (-PMS) or tested in the presence of postmitochondrial supernatant (+PMS). ^b Equivalent organic carbon (μg) per 100 μL bacterial suspension. ^c Mutant cells per 10^5 original cells \pm the 99% confidence limit based on Poisson statistics. ^d Survival is relative to the concurrent negative control. ^e For an extract to be mutagenic, a mutant fraction must be greater than the concurrent negative control (CC) such that the 99% confidence limits on the sample mean value and on the concurrent negative control do not overlap, and that mutant fraction must exceed the 95% upper confidence limit of the historical negative controls (HC). For relative survival less than 0.10, mutagenicity is not interpreted and the result is designated no call (NC). ^f Sample was extremely toxic -PMS and was tested using a long-growth variation of the bioassay. ^g Sample was extremely toxic +PMS and was tested using a long-growth variation of the bioassay. ^h Expected numerical variation has 99% confidence limits at least as great as ± 2.7 .

E 2021- NY

Table B: Ambient Samples Tested for Mutagenicity in the Bacterial Assay

equivalent organic carbon(μg) ^b	-PMS ^a				+PMS ^a			
	mutant fraction ^c ($\times 10^5$)	relative survival ^d	stat signif ^e	CC HC	mutant fraction ^c ($\times 10^5$)	relative survival ^d	stat signif ^e	CC HC
Azusa; Jan, Feb, Mar								
control	5.6±1.0	1.00			6.4±1.1	1.00		
0.57	11.2±1.7	0.72	+	-	6.7±1.1	1.06	-	-
1.71	16.1±2.2	0.69	+	+	7.2±1.2	1.02	-	-
5.12	109.4±24.5	0.08	NC		15.6±2.2	0.71	+	+
overall			+				+	
Azusa; Apr, May, Jun								
control	7.8±1.8	1.00			6.4±1.1	1.00		
0.64	8.2±1.3	1.07	-	-	7.6±1.1	1.14	-	-
1.93	10.3±1.5	0.93	-	-	6.6±1.1	1.10	-	-
5.80	60.6±11.5	0.19	+	+	15.4±2.5	0.81	+	+
overall			+				+	
Azusa; Jul, Aug, Sep								
control	5.9±1.0	1.00			5.7±1.1	1.00		
0.81	11.5±1.6	0.82	+	-	5.6±1.1	0.97	-	-
2.43	13.8±2.2	0.55	+	-	6.1±1.2	0.89	-	-
7.28	33.9±4.8	0.40	+	+	7.7±1.6	0.72	-	-
overall			+				-	
Azusa; Oct, Nov, Dec								
control	5.9±1.0	1.00			5.7±1.1	1.00		
0.73	7.5±1.1	1.05	-	-	6.8±1.4	0.79	-	-
2.19	14.6±2.7	0.40	+	+	7.1±1.4	0.79	-	-
6.57	28.0±4.0	0.44	+	+	6.4±1.5	0.64	-	-
overall			+				-	

Table B: continued

equivalent organic carbon(μg) ^b	-PMS ^a				+PMS ^a			
	mutant fraction ^c ($\times 10^5$)	relative survival ^d	stat signif ^e	CC HC	mutant fraction ^c ($\times 10^5$)	relative survival ^d	stat signif ^e	CC HC
Long Beach; Jan, Feb, Mar								
control	11.2±1.8	1.00			9.1±1.4	1.00		
0.69	14.3±1.9	1.21	-	+	8.9±1.3	1.10	-	-
2.07	26.6±3.9	0.64	+	+	12.0±1.7	0.96	-	-
6.22	103.6±16.4	0.28	+	+	18.9±2.5	0.77	+	+
overall			+				+	
Long Beach; Apr, May, Jun								
control	11.2±1.8	1.00			9.1±1.4	1.00		
0.43	14.9±2.2	0.91	-	+	9.9±1.6	0.90	-	-
1.30	18.5±2.4	1.01	+	+	8.2±1.3	1.00	-	-
3.90	77.8±11.7	0.34	+	+	10.4±1.7	0.84	-	-
overall			+				-	
Long Beach; Jul, Aug, Sep ^{f,g}								
control	8.6±1.4	1.00			10.5±1.3	1.00		
0.24	8.4±1.4	0.96	-	-	12.1±1.4	0.97	-	-
0.48	10.1±1.8	0.74	-	-	8.9±1.2	1.00	-	-
0.96	32.1±25.7	0.01	NC		17.5±3.4	0.28	+	+
overall			NC				+	
Long Beach; Oct, Nov, Dec								
control	5.6±1.0	1.00			7.7±1.0	1.00		
1.28	7.9±1.2	0.94	-	-	8.6±1.0	1.05	-	-
3.83	15.9±2.3	0.58	+	+	9.5±1.1	1.07	-	-
11.5	70.6±11.0	0.21	+	+	15.9±1.8	0.73	+	+
overall			+				+	

^a Tested in the absence of postmitochondrial supernatant (-PMS) or tested in the presence of postmitochondrial supernatant (+PMS). ^b Equivalent organic carbon(μg) per 100 μL bacterial suspension. ^c Mutant cells per 10^5 original cells \pm the 99% confidence limit based on Poisson statistics. ^d Survival is relative to the concurrent negative control. ^e For an extract to be mutagenic, a mutant fraction must be greater than the concurrent negative control (CC) such that the 99% confidence limits on the sample mean value and on the concurrent negative control do not overlap, and that mutant fraction must exceed the 95% upper confidence limit of the historical negative controls (HC). For relative survival less than 0.10, mutagenicity is not interpreted and the result is designated no call (NC). ^f Sample was extremely toxic -PMS and was tested using a long-growth variation of the bioassay. ^g Sample was extremely toxic +PMS and was tested using a long-growth variation of the bioassay.