

Supporting Information for *J. Am. Chem. Soc.*, 1991, 113(19), 7417-7418, DOI: [10.1021/ja00019a046](https://doi.org/10.1021/ja00019a046)

# DHAL

# 7417-7418

## Terms & Conditions

Electronic Supporting Information files are available without a subscription to ACS Web Editions. The American Chemical Society holds a copyright ownership interest in any copyrightable Supporting Information. Files available from the ACS website may be downloaded for personal use only. Users are not otherwise permitted to reproduce, republish, redistribute, or sell any Supporting Information from the ACS website, either in whole or in part, in either machine-readable form or any other form without permission from the American Chemical Society. For permission to reproduce, republish and redistribute this material, requesters must process their own requests via the RightsLink permission system. Information about how to use the RightsLink permission system can be found at <http://pubs.acs.org/page/copyright/permissions.html>.



ACS Publications

MOST TRUSTED. MOST CITED. MOST READ.

### Supplementary material

Bisimidazole substrates **2a** and **3a** were synthesized from the corresponding dibromides and imidazole. In a typical reaction, 2.85 g imidazole in 15 ml dry THF was slowly added to 1.9 g sodium hydride (60% in mineral oil) under nitrogen and stirred for 30 minutes. 5.3 g  $\alpha,\alpha'$ -dibromo-p-xylene in 20 ml THF was slowly added, and the temperature was raised to 50°C for 4 hrs. After cooling to room temperature, 25 ml of ice cold water was added, and the mixture was stirred for 20 minutes. The organic phase was extracted with chloroform (3 x 50 ml), and the combined chloroform phase was dried over anhydrous sodium sulfate. The solvent was removed under vacuum, and the residue was recrystallized twice from ethyl acetate to give 2.6 g (55% yield) of **2a**. Mp 148-150°C;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : 5.2 (s, 4H), 6.85 (s, 2H), 7.05 (s, 2H), 7.20 (m, 4H), 7.55 (s, 2H).