

1 RAD51 AND MITOTIC FUNCTION OF MUS81 ARE ESSENTIAL FOR RECOVERY FROM  
2 LOW-DOSE OF CAMPTOTHECIN IN THE ABSENCE OF THE WRN EXONUCLEASE

3

4

5 <sup>1</sup> Francesca Antonella Aiello, <sup>1</sup> Anita Palma, <sup>1</sup> Eva Malacaria, <sup>2</sup> Li Zheng, <sup>3</sup> Judith L. Campbell, <sup>2</sup>  
6 Binghui Shen, <sup>1</sup> Annapaola Franchitto and <sup>1,4</sup> § Pietro Pichierri

7

8 <sup>1</sup> Mechanisms, Biomarkers and Models Unit, Department of Environment and Health, Istituto  
9 Superiore di Sanità, Roma (Italy)

10 <sup>2</sup> Department of Cancer Genetics and Epigenetics, Beckman Research Institute, City of Hope, 1500  
11 East Duarte Road, Duarte, CA 91010-3000, USA

12 <sup>3</sup> Braun Laboratories, California Institute of Technology, Pasadena, CA 91125, USA

13 <sup>4</sup> Istituto Nazionale Biostrutture e Biosistemi, Roma (Italy)

14

15 § Author to whom correspondence should be addressed:

16

17 Pietro Pichierri

18 Tel. +39 0649902994

19 Fax +39 0660513138

20 [pietro.pichierri@iss.it](mailto:pietro.pichierri@iss.it)

21

22 Running title: WRN exonuclease and fork protection

23

24 Key words: WRN, Genome stability, DNA repair, Replication stress

25

## 27 ABSTRACT

28

29 Stabilisation of the stalled replication fork is crucial to prevent excessive fork reversal or degradation,  
30 which can undermine genome integrity. The WRN protein is a human RecQ helicase that participates  
31 in the processing and recovery of perturbed replication forks. WRN is unique among the other human  
32 RecQ family members to possess exonuclease activity. However, the biological role of the WRN  
33 exonuclease is poorly defined, and little is known about an involvement in the response to perturbed  
34 replication. Recently, the WRN exonuclease has been linked to protection of stalled forks from  
35 MRE11-dependent degradation in response to clinically-relevant nanomolar doses of the  
36 Topoisomerase I inhibitor camptothecin. Alternative processing of perturbed forks has been  
37 associated to chemoresistance of BRCA-deficient cancer cells, thus, we used WRN exonuclease-  
38 deficiency as a model to investigate the fate of perturbed replication forks undergoing degradation,  
39 but in a BRCA wild-type condition. We find that, upon nanomolar doses of camptothecin, loss of  
40 WRN exonuclease stimulates fork inactivation and accumulation of parental gaps, which engages  
41 RAD51. Such alternative mechanism affects reinforcement of CHK1 phosphorylation and causes  
42 persistence of RAD51 during recovery from treatment. Notably, in WRN exonuclease-deficient cells,  
43 persistence of RAD51 correlates with elevated mitotic phosphorylation of MUS81 at Serine 87, which  
44 is essential to avoid accumulation of mitotic abnormalities. Altogether, these findings indicate that  
45 aberrant fork degradation, in the presence of a wild-type RAD51 axis, stimulates RAD51-mediated  
46 post-replicative repair and engagement of the MUS81 complex to limit genome instability and cell  
47 death.

48

## 49 AUTHOR SUMMARY

50

51 Correct progression of the molecular machine copying the chromosomes is threatened by multiple  
52 causes that induce its delay or arrest. Once the replication machinery is arrested, the cell needs to  
53 stabilise it to prevent DNA damage. Many proteins contribute to this task and the Werner's syndrome  
54 protein, WRN, is one of them.

55 Defining what happens to replication machineries when they are blocked is highly relevant. Indeed,  
56 destabilised replication machineries may form upon treatment with anticancer drugs and influence  
57 the efficacy of some of them in specific genetic backgrounds. We used cells that lack one of the two  
58 enzymatic functions of WRN, the exonuclease activity, to investigate the fate of destabilised  
59 replication machineries. Our data show that they are handled by a repair pathway normally involved  
60 in fixing DNA breaks but, in this case, recruited to deal with regions of the genome that are left  
61 unreplicated after their destabilisation. This alternative mechanism involves a protein, RAD51, which  
62 tries to copy DNA from the sister chromosome. In so doing, however, RAD51 produces a lot of DNA  
63 interlinking that requires upregulation of a complex, called MUS81/EME1, which resolves this  
64 interlinking prior cell division and prevents accumulation of mitotic defects and cell death.

65

66

67

68

69

## 70 INTRODUCTION

71

72 The response to perturbed replication is crucial for the maintenance of genome integrity [1–5]. In  
73 humans, the proper handling of perturbed replication forks is also linked to cancer avoidance and  
74 many proteins involved in this process act also as onco-suppressors [3–6]. The importance of dealing  
75 correctly with perturbed replication forks is also demonstrated by the existence of several human  
76 genetic diseases caused by mutations in factors involved in sensing, processing and recovering  
77 replication forks [7].

78 The Werner's syndrome protein (WRN) is one of the key factors of the response to perturbed  
79 replication and participates directly to the checkpoint in S-phase [8,9]. *WRN* is mutated in the genetic  
80 disease Werner's syndrome (WS), which is characterized by cancer predisposition and premature  
81 aging [9], and its loss confers sensitivity to several DNA-damaging agents inducing replication stress  
82 [8,10]. From an enzymatic point of view, WRN is both a DNA helicase and exonuclease; however,  
83 while its helicase activity has been linked to processing of reversed or collapsed replication forks  
84 [2,9], little is known about the biological relevance of the exonuclease activity. We recently reported  
85 that the exonuclease activity of WRN is involved in protecting replication forks that are perturbed by  
86 treatment with the Topoisomerase I poison Camptothecin (CPT) in the nanomolar range of  
87 concentration [11]. Exposure to low doses of CPT, as opposed to high doses, does not induce DSBs  
88 in the short-term but stimulates greatly formation of reversed forks [12,13]. Reversed replication forks  
89 are versatile yet vulnerable structures and several proteins participate in their stabilisation [14–16].  
90 Two proteins, BRCA2 and RAD51, are the most crucial for the stabilisation of reversed forks  
91 [14,16,17]. Thus, cells depleted of each of these two proteins have been used as a prototypical model  
92 to assess the consequences of inaccurate handling of reversed forks. However, BRCA2 and RAD51  
93 may also participate in DNA repair, which may be used to fix damage generated by fork instability  
94 [17–19]. Loss of WRN exonuclease determines a rapid MRE11-dependent degradation of the nascent  
95 strand, most likely after fork reversal, and affects correct replication recovery [11]. However, cells  
96 expressing the exonuclease-dead WRN retain ability to restart replication and are not overtly sensitive  
97 to low doses of CPT, suggesting that alternative mechanisms can be activated as a back-up. Since  
98 nanomolar doses of CPT are clinically-relevant in cancer therapy, cells expressing a catalytically-  
99 inactive WRN exonuclease can be used as a model to investigate the fate of CPT-perturbed replication  
100 forks undergoing pathological degradation but in a BRCA2-RAD51 wild-type background.

101 Here, we report that, upon prolonged exposure to nanomolar dose of CPT, loss of WRN exonuclease  
102 channels replication forks through a pathological RAD51-dependent mechanism that makes  
103 perturbed replication forks resistant to breakage. However, engagement of RAD51 and persisting

104 RAD51 foci make WRN exonuclease-deficient cells reliant on the mitotic function of the MUS81  
105 complex, which mitigates mitotic abnormalities deriving from accumulation of RAD51-dependent  
106 intermediates. Furthermore, our data suggest that enhanced accumulation of ssDNA and recruitment  
107 of RAD51 interfere with correct activation of CHK1, which provides a positive feedback to the  
108 formation of nascent ssDNA.  
109

## 110 RESULTS

**111 Loss of WRN exonuclease activity leads to a persisting and unusual formation of nascent ssDNA**  
**112 which compromises formation of DSBs in response to a low-dose of camptothecin**

113 Treatment with nanomolar concentrations of CPT does not induce DSBs immediately but initially  
114 stimulates fork reversal [12]. Under such conditions, loss of WRN exonuclease activity results in the  
115 rapid (< 2h) degradation of both nascent strands by the MRE11-EXO1 nucleases [11]. In this regard,  
116 WRN exonuclease deficiency is a useful model to determine what happens to nuclease-targeted forks  
117 after prolonged treatment with low dose of CPT. This prompted us to analyse the processing of  
118 perturbed replication forks beyond 2h of treatment.

119

120 We first examined the presence of nascent ssDNA as a sign of fork degradation by the native IdU  
121 assay. As expected, WS cells expressing the exo-dead WRN protein (WS<sup>E84A</sup>) showed less nascent  
122 ssDNA than the corrected wild-type counterpart (WS<sup>WT</sup>) at 1h of treatment (Fig. 1A). Surprisingly,  
123 the amount of nascent ssDNA in WRN exonuclease-deficient cells increased greatly over time and  
124 largely exceeded the wild-type level at 4h, while it increased only slightly in cells expressing the  
125 wild-type WRN (Fig. 1A). Since nascent strand degradation is MRE11-dependent in the absence of  
126 the WRN exonuclease while it is DNA2-dependent in wild-type cells (refs), we next examined if  
127 chemical inhibition of those nucleases might reduce the idiosyncratic accumulation of ssDNA  
128 detected at 4h of treatment with nanomolar CPT in WS<sup>E84A</sup> cells. Mirin treatment, which inhibits  
129 MRE11, barely reduced ssDNA detected at 4h of treatment in wild-type cells (Fig. 1B). Surprisingly,  
130 Mirin did not decrease the formation of ssDNA in WRN exonuclease-deficient cells but rather  
131 increased it even further (Fig. 1B). Inhibition of DNA2 by the small-molecule inhibitor C5 [20]  
132 increased formation of nascent ssDNA in wild-type but not in WRN exonuclease-deficient cells (Fig.  
133 1B). In contrast, concomitant inhibition of DNA2 and MRE11 was ineffective in modulating nascent  
134 ssDNA formation in wild-type cells while decreased its level in cells expressing the exo-dead form  
135 of WRN (Fig. 1B). Of note, ssDNA derived from end-resection of DSBs induced by a micromolar  
136 dose of CPT was efficiently reduced by DNA2 inhibitor C5 (Fig. S1A, B), providing a functional  
137 proof of the inactivation of the nuclease activity by the C5 compound in our cell model. These results  
138 indicate that different sets of nucleases are involved in the degradation of nascent ssDNA in WRN  
139 exonuclease-deficient cells when treatment is prolonged.

140

141 DNA breakage can occur even in response to low-doses of CPT if treatment is sufficiently prolonged  
142 [12,13]. Thus, to further investigate the origin of the late nascent ssDNA in the WRN exonuclease-  
143 deficient cells and the role of the different nucleases, we analysed the presence of DSBs after

144 treatment with nanomolar CPT by neutral Comet assay. As shown in Figure 1C, treatment with 50nM  
145 CPT for 4h is able to induce some DSBs in wild-type cells, although they are very low compared with  
146 those generated by the 5 $\mu$ M reference dose. In contrast, no DSBs were detected in WRN exonuclease-  
147 deficient cells after treatment with the low-dose of CPT, even if they were readily seen in response  
148 to the high-dose of the drug (Fig. 1C). Interestingly, pre-treatment with Mirin, which enhances  
149 ssDNA in WS<sup>E84A</sup> cells (Fig. 1B), resulted in DSBs (Fig. 1C). In contrast, formation of DSBs was  
150 unaffected in wild-type cells by Mirin while it was reduced by the DNA2 inhibitor (Fig. 1C).  
151 Interestingly, inhibition of both MRE11 and DNA2, which decreases ssDNA formation in WS<sup>E84A</sup>  
152 cells (Fig. 1B), reduced significantly the DSBs generated by Mirin (Fig. 1C). Concomitant inhibition  
153 of MRE11 and DNA2 also reduced DSBs in wild-type cells (Fig. 1C). The analyses of DSBs and  
154 ssDNA suggest that the increase in nascent ssDNA induced by MRE11 inhibition in WRN  
155 exonuclease-deficient cells derives from end-resection of DSBs. In contrast, the reduction of nascent  
156 ssDNA induced by concomitant treatment with C5 and Mirin would indicate that DNA2 and MRE11  
157 are involved in the formation of the late nascent ssDNA observed in WS<sup>E84A</sup> cells independently of  
158 detectable DSBs. Finally, we tested whether MUS81 was required for late DSB formation [21,22].  
159 As shown in Fig. S1C, DSBs derived from prolonged treatment with a low-dose of CPT in wild-type  
160 cells are not affected by depletion of MUS81.

161 Collectively, our results indicate that loss of the WRN exonuclease leads to accumulation of nascent  
162 ssDNA when treatment with nanomolar CPT is prolonged beyond 2h and that it makes cells refractory  
163 to formation of DSBs by CPT. Our data also suggest that the late accumulation of nascent ssDNA in  
164 WS<sup>E84A</sup> cells is related to the activity of multiple nucleases.

165

### 166 **Loss of WRN exonuclease stimulates engagement of RAD51 after CPT-induced fork** 167 **perturbation**

168 In WRN exonuclease-deficient cells, late reappearance and accumulation of nascent ssDNA together  
169 with absence of DSBs after CPT might correlate with engagement of an alternative or aberrant fork  
170 processing mode over time. Extended regions of ssDNA are a substrate for RAD51 during  
171 recombination, thus, we investigated whether loss of WRN exonuclease could affect recruitment of  
172 RAD51 after treatment with CPT.

173

174 To evaluate recruitment of RAD51, we first prepared chromatin from cells treated with nanomolar  
175 CPT and determined the amount of RAD51 present by Western blotting. In wild-type cells, RAD51  
176 barely increased its chromatin association after CPT treatment (Fig. 2A). Expression of the exo-dead  
177 WRN, however, greatly increased the amount of RAD51 in chromatin both in untreated cells and

178 CPT-treated cells. Treatment with CPT led to a minimal increase over untreated (about 20%; Fig.  
179 2A). As a further control, we also measured the level of RPA32, a subunit of the RPA heterotrimer  
180 that binds to ssDNA, in chromatin. RPA32 increased after CPT in wild-type cells (Fig. 2A). In  
181 contrast, the amount of RPA32 in chromatin was low in the absence of the WRN exonuclease and  
182 did not show any increase after treatment (Fig. 2A). RAD51 participates in multiple pathways [23,24]  
183 and chromatin recruitment may reflect such pleiotropy of roles. To further verify the increased  
184 recruitment of RAD51 in cells expressing the exo-dead WRN, we performed a quantitative  
185 immunofluorescence analysis of the RAD51 foci (Fig. 2B). We evaluated the number of foci by  
186 analysing the intensity of fluorescence in the RAD51 foci-positive nuclei. Consistent with  
187 biochemical data, loss of WRN exonuclease increased recruitment of RAD51 (Fig. 2B). Moreover,  
188 CPT treatment led to a higher number of RAD51 foci in WRN exonuclease-deficient cells compared  
189 with the wild-type (Fig. 2B)

190 These results suggest that, in the absence of the WRN exonuclease, RAD51 takes over the normal  
191 mechanism handling CPT-perturbed replication forks, perhaps to provide a backup mechanism for  
192 recovery.

193

194 To test if RAD51 participated in replication fork restart, we analysed whether, in cells expressing the  
195 exo-dead WRN, RAD51 chromatin levels remained elevated also during recovery from CPT. To this  
196 end, we treated cells with CPT and allowed them to recover for 2 and 4h prior to preparing chromatin  
197 fractions. As shown in Figure 3A, while RAD51 levels, as determined by Western blotting, tended to  
198 remain low during recovery in wild-type cells, they remained elevated in WRN exonuclease-deficient  
199 cells. Of note, the amount of chromatin-bound RPA32 and RPA70, two subunits of trimeric RPA,  
200 remained low in WS<sup>E84A</sup> cells while it increased greatly during recovery in cells expressing the wild-  
201 type WRN (Fig. 3A). To confirm this result, we performed quantitative immunofluorescence analysis  
202 of the formation of RAD51 foci during recovery from CPT. A reduction of RAD51 foci was observed  
203 during recovery from CPT in wild-type cells (Fig. 3B). In contrast, the number of RAD51 foci  
204 increased in WRN exonuclease-deficient cells (Fig. 3B), confirming that a RAD51-dependent  
205 pathway is over-activated in response to prolonged treatment with low-dose of CPT in these cells.

206

207 The persistently-high levels of RAD51 observed in the absence of the WRN exonuclease even during  
208 recovery prompted use to determine if they were correlated with the increased formation of nascent  
209 ssDNA. Having demonstrated that concomitant inhibition of MRE11 and DNA2 restores wild-type  
210 levels of nascent ssDNA in WRN exonuclease-deficient cells (Fig. 1C), we analysed RAD51 focus-  
211 forming activity after pre-treatment of cells with C5 and Mirin. As expected, RAD51 recruitment in



212 foci was elevated in WRN exonuclease-deficient cells after treatment and even more during recovery  
213 (Fig. 3C). Interestingly, concomitant pre-treatment with C5 and Mirin significantly reduced the  
214 formation of RAD51 foci in WS<sup>E84A</sup> cells (Fig. 3C), suggesting that it depends on late accumulation  
215 of nascent ssDNA observed in the absence of WRN exonuclease.

216

217 The elevated levels of RAD51 during recovery might be related to the presence of under-replicated  
218 DNA that requires recombination to be replicated or repaired [25]. Thus, we analysed whether cells  
219 recovering from CPT treatment presented under-replicated regions of DNA. Since under-replication  
220 is expected to leave regions of ssDNA in the parental strand (parental gaps) behind perturbed forks,  
221 we performed native IdU assay after a 24h treatment with IdU to label all parental DNA prior to add  
222 CPT and perform recovery (Fig. 3D). As shown in Figure 3E, little parental ssDNA was detectable  
223 in wild-type cells after treatment and its amount decreased substantially during recovery. In WS<sup>E84A</sup>  
224 cells, however, parental ssDNA was higher at the end of treatment and remained elevated during  
225 recovery (Fig. 3E). Interestingly, the increased amount of parental ssDNA paralleled that of RAD51  
226 foci, suggesting that RAD51 may be recruited to deal with under-replicated regions.

227

228 The presence of under-replicated DNA and the increased recruitment of RAD51 might indicate that  
229 perturbed replication forks get inactivated upon prolonged exposure to CPT in the absence of the  
230 WRN exonuclease. Thus, we carried out a single-molecule analysis of DNA replication by the DNA  
231 fiber assay (Fig. 4A). As observed previously [12,13], treatment with nanomolar CPT did not induce  
232 fork arrest but rather a delay in fork progression (Fig. S2). Indeed, the length of the CldU tract is  
233 increased during extended periods of treatment. Of note, length of the CldU tract was similar between  
234 WS<sup>WT</sup> and WS<sup>E84A</sup> cells at 1h and 4h of treatment (Fig. S2). When we analysed the ability to replicate  
235 after treatment, we found that most of perturbed replication forks remained active after 1h of CPT in  
236 wild-type cells, as indicated by the low level of stalled forks (red-only tracks) (Fig. 4B-D). After  
237 prolonged treatment, the number of stalled forks increased by around 2-fold, but perturbed forks  
238 remained mostly active (Fig. 4B). Notably, the number of stalled forks was higher in WRN  
239 exonuclease-deficient cells and around the 50% of the forks got inactivated at 4h of treatment (Fig.  
240 4B-D). Interestingly, in the absence of the WRN exonuclease, treatment with CPT stimulated firing  
241 of new origins (green-only tracks), which increased with time of treatment (Fig. 4C, D). In contrast,  
242 limited new origin firing was detectable in wild-type cells (Fig. 4C, D). Surprisingly, and in spite of  
243 the elevated levels of RAD51 in WS<sup>E84A</sup> cells, neither the percentage of inactive forks nor that of new  
244 origins was affected by inhibition of RAD51 during recovery (Fig. 4B-D). In contrast, we observed

245 that both fork inactivation and firing of new origins were increased by inhibition of RAD51 in wild-  
246 type cells (Fig. 4B-D).

247 These results suggest that RAD51 is not required to promote replication recovery but rather to  
248 promote repair of parental gaps left behind inactive forks after resumption of synthesis when the  
249 exonuclease activity of WRN is absent.

250

### 251 **Loss of WRN exonuclease reduces activation of CHK1**

252 Our data indicate that loss of the WRN exonuclease results in accumulation of ssDNA and RAD51  
253 accompanied by a concomitant decrease of RPA. Since RPA-coated ssDNA is required for  
254 checkpoint signalling upon replication fork perturbation, we investigated whether the functionality  
255 of the WRN exonuclease might also affect activation of the replication checkpoint in response to  
256 nanomolar concentration of CPT. As a readout of the activation of the ATR-dependent checkpoint  
257 response, we analysed phosphorylation of CHK1 at S345 by Western blotting. In wild-type cells,  
258 treatment with a nanomolar dose of CPT induced a time-dependent phosphorylation of CHK1 which  
259 is also readily observed in cells expressing the helicase-dead form of WRN (WS<sup>K577M</sup>) (Fig. 5A). In  
260 contrast, CPT-induced phosphorylation of CHK1 was reduced in cells expressing the exonuclease-  
261 dead mutant of WRN, and this phenotype was more evident at 4 and 6h of treatment (Fig. 5A). The  
262 requirement of the WRN exonuclease for correct CHK1 phosphorylation was specific for the low-  
263 dose CPT treatment as it was not observed after 5 $\mu$ M of CPT (Fig. S3A).

264 To confirm that loss of WRN exonuclease affected CHK1 phosphorylation by an independent assay,  
265 we monitored the status of S345 of CHK1 by immunofluorescence. As shown in Fig. 5B, a reduced  
266 phosphorylation of CHK1 at S345 was readily detected also by immunofluorescence in WRN  
267 exonuclease-deficient cells.

268

269 Next, we wanted to analyse whether decreased activation of CHK1 correlated with reduced activation  
270 of ATR-dependent signalling. To assess activation of ATR, we monitored phosphorylation of the  
271 activating site T1989 by immunofluorescence. Despite the defective phosphorylation of CHK1, ATR  
272 was phosphorylated similarly in wild-type cells and in cells expressing the exonuclease-dead form of  
273 WRN (Fig. 5C). Since loss of WRN exonuclease affects recruitment of RPA but did not affect  
274 activation of ATR, we analysed whether it could influence recruitment of other factors modulating  
275 ATR-checkpoint function. As loss of WRN exonuclease leads to accumulation of nascent ssDNA at  
276 4h of CPT (Fig. 1A), we analysed the presence of TopBP1 and RAD9, which associates with TopBP1  
277 [26,27], specifically at nascent ssDNA by our recently described IdU-PLA assay [11]. In parallel, we  
278 evaluated the presence of the total amount of ssDNA by IdU assay. Association of TopBP1 or RAD9

279 with nascent ssDNA was not detected under untreated conditions (data not shown), however,  
280 treatment with 50nM CPT for 4h resulted in recruitment of both factors at nascent ssDNA in wild-  
281 type cells (Fig. 5D, E). Of note, loss of WRN exonuclease reduced the recruitment of TopBP1 but  
282 not that of RAD9 at the nascent ssDNA (Fig. 5D, E) although in these conditions the IdU assay  
283 detected 2-times more ssDNA (Fig. 5F).

284

285 Loss of the WRN exonuclease leads to accumulation of nascent ssDNA, which is targeted by RAD51  
286 and not by checkpoint factors, possibly resulting in reduced CHK1 phosphorylation. Thus, we  
287 investigated whether pre-treatment with the RAD51 inhibitor B02 or treatment before sampling could  
288 re-establish a normal CHK1 activation in WRN exonuclease-deficient cells (Fig. S3B). Of note,  
289 inhibition of RAD51 further decreased the phosphorylation of CHK1 regardless the way it was added  
290 (Fig. S3B). This was surprising but prompted us to evaluate whether reduced CHK1 activation could  
291 be implicated in the enhanced accumulation of nascent ssDNA. To test this potential feedback effect,  
292 we expressed the S317/S345D CHK1 phosphomimetic mutant [28] in WS<sup>E84A</sup> cells (Fig. 6A) before  
293 evaluating the formation of nascent ssDNA by the IdU assay. As shown in Figure 6B, the  
294 phosphomimetic CHK1 was efficiently expressed in the cells and its expression was able to  
295 substantially reduce the amount of nascent ssDNA in WRN exonuclease-deficient cells restoring the  
296 wild-type levels.

297 Altogether, our results show that loss of WRN exonuclease activity affects proper activation of CHK1  
298 in response to a low-dose of CPT and that reduced phosphorylation of CHK1 probably correlates with  
299 reduced recruitment of checkpoint factors at ssDNA. They also suggest that reduced CHK1  
300 phosphorylation contributes to the accumulation of ssDNA in the nascent strand, possibly as part of  
301 a positive feedback loop.

302

### 303 **WRN exonuclease-deficient cells need the mitotic function of MUS81 to counteract mitotic** 304 **aberration and mis-segregation**

305 Inaccurate processing of perturbed replication forks, elevated under-replicated DNA and RAD51  
306 levels observed in the absence of WRN exonuclease could threaten mitosis because of DNA  
307 interlinking as shown in BRCA2-deficient cells [29]. Mitotic resolution of DNA interlinked  
308 intermediates involves the MUS81/EME1 complex [30,31]. Thus, we evaluated whether WRN  
309 exonuclease-deficient cells accumulated active MUS81 in mitosis by performing  
310 immunofluorescence with an antibody directed against the pS87-MUS81, which we have  
311 demonstrated to be a readout of the active MUS81/SLX4 complex [32]. In wild-type cells, little  
312 MUS81 phosphorylation at S87 was detectable either under unperturbed cell growth or in response

313 to low-dose of CPT (Fig. 7A). In contrast, many pS87-MUS81-positive nuclei were detectable in  
314 cells expressing the exonuclease-dead WRN already in unperturbed conditions (Fig. 7A). Notably, in  
315 WRN exonuclease-deficient cells, pS87-MUS81 levels were further enhanced by CPT and remained  
316 elevated also after recovery (Fig. 7A). Interestingly, and consistent with our previous results [32],  
317 phosphorylation of MUS81 never occurred in S-phase cells labelled with EdU (Fig. 7A). To  
318 determine whether elevated MUS81 activation correlated with under-replication and enhanced  
319 RAD51 recruitment, we asked if inhibition of RAD51 by B02 (ref) reverted the pS87-MUS81 levels  
320 in WRN exonuclease-deficient cells. As shown in Figure 7B, inhibition of RAD51 greatly decreased  
321 activation of the MUS81 complex after recovery from CPT, as indicated by reduction in pS87-  
322 MUS81-positive nuclei.

323  
324 Enhanced activation of MUS81 in mitosis in WS<sup>E84A</sup> cells might be indicative of persistence of  
325 unresolved DNA intermediates, which could induce mitotic abnormalities or segregation defects. To  
326 assess if loss of WRN exonuclease during the response to low-dose of CPT could result in segregation  
327 defects, we first analysed the presence of bulky anaphase bridges in DAPI-stained cells (Fig. 7C).  
328 Interestingly, anaphase cells were highly enriched in WRN exonuclease-deficient cells (Fig. 7C; see  
329 numbers above the bars), suggesting that these cells may have a delayed exit from anaphase. Of note,  
330 the number of anaphases with bridges was very low in WRN exonuclease-deficient cells as compared  
331 to cells expressing WRN wild-type (Fig. 7C). Since delayed exit from anaphase might derive from  
332 mitotic defects and could result in post-mitotic abnormalities, we decided to evaluate the presence of  
333 aberrant mitoses, multinucleated cells and micronuclei. As shown in Figure 7D, we found that both  
334 aberrant mitosis and cells with micronuclei were increased by loss of WRN exonuclease activity. In  
335 contrast, no difference in multinucleated cells was found between cells expressing wild-type or exo-  
336 dead WRN (Fig. 7D).

337  
338 The presence of under-replicated DNA or the persistence of unresolved DNA intermediates in G2/M  
339 triggers the formation of 53BP1 NBs in the subsequent G1 phase [33]. Since WRN exonuclease-  
340 deficient cells showed persistence of under-replicated DNA, we investigated whether they  
341 accumulated 53BP1 NBs. To this end, we performed immunofluorescence against 53BP1 and  
342 CyclinA in cells after recovery from 4h treatment with CPT and scored the number of 53BP1 NBs-  
343 positive cells in the CyclinA-negative population (i.e. G1 cells). As shown in Figure 7E, the number  
344 of 53BP1 NBs in WS<sup>E84A</sup> cells was higher than in wild-type cells even in untreated conditions.  
345 Treatment with CPT enhanced the number of 53BP1 NBs in wild-type and in WRN exonuclease-

346 deficient cells; however, the increase was substantially higher in cells expressing the exo-dead WRN  
347 (Fig. 7E and Fig. S5).

348 Notably, inhibition of RAD51 in WRN exonuclease-deficient cells resulted in persistent  
349 accumulation of mitotic cells in pro-metaphase (Fig. S4). This result suggested that loss of RAD51  
350 undermines correct mitotic progression in WRN-exonuclease-deficient cells but prevented  
351 assessment of any correlation between enhanced engagement of RAD51 and mitotic defects or  
352 formation of 53BP1 NBs.

353

354 As in WRN exonuclease-deficient cells engagement of RAD51 is functionally related to elevated  
355 levels of S87-MUS81 phosphorylation (Fig. 7B), we next analysed whether inactivation of the mitotic  
356 function of MUS81 by overexpression of the unphosphorylatable S87A-MUS81 mutant (ref) in these  
357 cells could aggravate the mitotic defects. Interestingly, over-expression of the S87A-MUS81 mutant  
358 in wild-type cells did not affect the percentage of anaphase bridges and micronuclei, while it increased  
359 the number of multinucleated cells (Fig. 8A). In sharp contrast, expression of the S87A-MUS81  
360 mutant substantially aggravated the mitotic defects in WRN exonuclease-deficient cells (Fig. 7A).  
361 Indeed, expression of the S87A-MUS81 protein increased the number of anaphase bridges and  
362 micronucleated cells. Similarly, expression of the S87A-MUS81 mutant enhanced the presence of  
363 53BP1 NBs in WS<sup>E84A</sup> cells (Fig. 8B).

364

365 To further assess the biological significance of the MUS81 hyperactivation observed in mitosis in the  
366 absence of the WRN exonuclease, we evaluated the sensitivity to nanomolar doses of CPT by  
367 clonogenic survival. As shown in Figure 8C, WRN exonuclease-deficient cells were slightly more  
368 sensitive to CPT than wild-type cells. Overexpression of the wild-type MUS81 resulted in a mild  
369 increase in sensitivity in wild-type cells but not in cells expressing the exo-dead WRN (Figure 7C).  
370 In contrast, overexpression of the S87A-MUS81 resulted in a substantial increase in the sensitivity of  
371 WRN exonuclease-deficient cells to CPT (Fig. 8C).

372

373 Altogether, our data indicate that the enhanced engagement of RAD51 observed in the absence of the  
374 WRN exonuclease requires the increased activation of the MUS81 complex in mitosis. Therefore,  
375 expression of a MUS81 mutant that disables mitotic activation of the MUS81/EME1 complex  
376 increases mitotic abnormalities and sensitivity to CPT of WRN exonuclease-deficient cells. Thus, in  
377 the absence of the WRN exonuclease, hyperactivation of the MUS81 complex functions as a fail-safe  
378 system that maintains mitotic abnormalities at low levels, allowing survival.

379

## 380 DISCUSSION

381

382 In recent years, an increasing interest arose around alternative mechanisms of fork processing and  
383 fork degradation since they correlate with response to chemotherapeutics in cells that are deficient  
384 for the primary pathway(s) as described in the absence of BRCA1/2 [15,17]. Most of these studies  
385 focused on the early events occurring in the absence of BRCA1 or BRCA2, but few of them  
386 investigated mechanisms involved during recovery from replication stress [29,34]. Furthermore, loss  
387 of BRCA1 or BRCA2 affects recombination as well as fork protection and this prevents the  
388 investigation of the role of recombination for the recovery of replication forks undergoing  
389 degradation. Recently, we reported that the WRN exonuclease activity protects against fork  
390 degradation when cells are treated with clinically-relevant doses of CPT [11]. Here, we used WRN  
391 exonuclease-deficient cells as a model to assess what happens at destabilised perturbed forks when  
392 treatment with nanomolar doses of CPT is prolonged. We find that, in the absence of the WRN  
393 exonuclease, nascent strands undergo continuous degradation that produces accumulation of ssDNA.  
394 This late accumulation of ssDNA follows its disappearance at early time points because of the  
395 activities of MRE11 and EXO1 [11]. Interestingly, the late wave of ssDNA at perturbed forks is only  
396 minimally affected by inactivation of each single exonuclease acting at perturbed forks, and is only  
397 reduced when MRE11 and DNA2 are both inhibited. This suggests that multiple nucleases take over  
398 with time at forks destabilised by the absence of WRN exonuclease while most of the degradation  
399 observed in cells deficient of BRCA1/2, or other factors assisting RAD51, seems to involve only  
400 MRE11-EXO1 [35–38]. Treatment with nanomolar doses of CPT does not induce DSBs unless  
401 treatment is prolonged [12,13]. Interestingly, loss of the WRN exonuclease makes cells resistant to  
402 the induction of DSBs after prolonged treatment with nanomolar CPT. Induction of DSBs in response  
403 to nanomolar doses of CPT has been correlated with activation of RECQ1 possibly to promote restart  
404 of those forks that failed to be processed otherwise [13]. Loss of the ability to induce DSBs at forks  
405 would be consistent with engagement of a distinct fork recovery mechanism in cells expressing the  
406 exo-dead WRN protein. Indeed, WRN exonuclease-deficient cells do not show RECQ1 PARylation  
407 [11], which is required to avoid unscheduled RECQ1 activation [13], which, together with the absence  
408 of DSBs support a pathway switch at CPT-perturbed forks.

409

410 Consistent with the pathway switch, inhibition of MRE11 is sufficient to restore DSBs after prolonged  
411 treatment with a low-dose of CPT in WRN exonuclease-deficient cells, suggesting that formation of  
412 DSBs does not necessarily occur downstream of pathological fork processing. Interestingly, RAD51  
413 is strongly accumulated in the absence of the WRN exonuclease and persists during recovery from

414 treatment. An elevated engagement of RAD51 in the absence of the WRN exonuclease has been  
415 reported in *Drosophila* [39], suggesting that the role of WRN exonuclease at perturbed forks is  
416 conserved. Similarly, unscheduled exonuclease-mediated processing of perturbed forks in yeast has  
417 been recently shown to engage a RAD51-mediated pathway [40]. Furthermore, parental ssDNA, a  
418 readout of template gaps, also accumulates in the absence of WRN exonuclease. RAD51 binds to  
419 ssDNA and initiates recombination [19,25]. The concomitant accumulation of ssDNA and elevated  
420 recruitment of RAD51 in the absence of the WRN exonuclease would be consistent with the  
421 engagement of gaps left behind inactivated forks in a template-switch mode of replication recovery,  
422 as shown after DNA damage in *Xenopus* egg extracts [23]. Consistent with this possibility, WRN  
423 exonuclease-deficient cells show enhanced inactivation of CPT-perturbed forks and new origin firing.  
424 Moreover, although RAD51 has been implicated in fork restart [24,41], in WRN exonuclease-  
425 deficient cells, RAD51 is not involved in fork reactivation after CPT treatment. Indeed, its inhibition  
426 does not increase fork inactivation. This result is in agreement with the participation of RAD51 in  
427 “gap repair” and supports the notion that prolonged treatment with nanomolar CPT channels  
428 perturbed forks into alternative fork processing pathways if the function of WRN exonuclease is  
429 absent. Consistent with this, in wild-type cells, inhibition of RAD51 reduces fork reactivation. This  
430 is not unexpected since RAD51 plays crucial roles in both fork remodelling and stability [14,16]. In  
431 addition, as RAD51 and WRN have been proposed to cooperate during recovery from fork arrest  
432 [42], it is reasonable to speculate that loss of WRN function also compromises the normal activity of  
433 RAD51 at fork.

434  
435 Loss of WRN exonuclease results in a mild defect in the activation of CHK1. Activation of the ATR-  
436 dependent checkpoint requires formation of ssDNA [43–45]. In the absence of the WRN exonuclease  
437 ssDNA accumulates but is hijacked by RAD51 and it is not completely free for the binding of  
438 checkpoint factors. Indeed, in WRN exonuclease-deficient cells, TopBP1 and its binding factor  
439 RAD9 are not more highly associated with ssDNA as compared with the wild-type. Notably,  
440 phosphorylation of ATR, a readout of its activation, is indistinguishable from the wild-type. It  
441 suggests that ATR also gets activated independently from ssDNA. Alternatively, a hyperactivation  
442 of ATR is also prevented by sequestering of ssDNA by RAD51. Indeed, overexpression of RAD51  
443 has been shown to affect checkpoint activation [46]. Notably, bypassing of the CHK1 activation  
444 defect by expression of a phosphomimetic CHK1 mutant in WRN exonuclease-deficient cells restores  
445 normal levels of ssDNA. This suggests that accumulation of ssDNA is also unleashed by reduced  
446 CHK1 activation through a positive feedback loop.

447

448 The observed elevated recruitment of RAD51, which is used during recovery in the absence of the  
449 WRN exonuclease to deal with under-replicated DNA, also leads to elevated phosphorylation of  
450 MUS81 at S87. Phosphorylation of MUS81 at S87 occurs in G2/M and is related to resolution of  
451 recombination intermediates [32]. Consistently, inhibition of RAD51 reduces S87 phosphorylation  
452 in WRN exonuclease-deficient cells. Thus, engagement of RAD51-dependent fork recovery, possibly  
453 by template switch since DSBs do not form, results in an increased number of interlinked  
454 intermediates calling for resolution by the MUS81 complex. Our data indicate that activation of  
455 MUS81 complex in G2/M is essential to overcome segregation defects arising from excessive  
456 RAD51-dependent recombination and support proliferation upon treatment with CPT. Indeed,  
457 expression of the unphosphorylatable S87A-MUS81 mutant increases abnormal mitosis and sensitizes  
458 WRN exonuclease-deficient.

459

460 Loss of the WRN exonuclease although resulting in fork degradation does not induce MUS81  
461 activation in S-phase, which is observed in the absence of BRCA2 [37]. However, the persistence of  
462 under-replicated DNA and requirement of MUS81 complex activity in G2/M shown by WRN  
463 exonuclease-deficient cells are also characteristic of BRCA2-deficient cells [34]. Thus, it is tempting  
464 to speculate that elevated fork degradation correlates with inability to replicate all the genome.  
465 Notably, BRCA2-deficient cells show much more severe mitotic defects [29,34]. In WRN  
466 exonuclease-deficient cells, mitotic abnormalities are increased by disabling MUS81 function in  
467 mitosis but are likely increased also by impairing RAD51 function during recovery, since RAD51  
468 inhibition results in increased accumulation of parental ssDNA and induces a significant mitotic  
469 block. As BRCA2 deficiency also interferes with the post-replicative function of RAD51 [23] it is  
470 tempting to speculate that the elevated mitotic defects might be the end-result of combined fork  
471 deprotection and recombination defects. Indeed, it is recently demonstrated that mitotic abnormalities  
472 in BRCA2-deficient cells are primarily linked to loss of the recombination function of RAD51 using  
473 separation-of-function mutants [29].

474

475 Collectively, our data show that WRN exonuclease-deficient cells can be a useful model to investigate  
476 the fate of deprotected or destabilised replication forks under a clinically-relevant, specific type of  
477 replication stress; and, together with published data, they can be summarised in the model shown in  
478 Figure 9. In response to nanomolar CPT, perturbed replication forks rapidly undergo fork reversal  
479 [12]. The WRN exonuclease is required somehow at this stage to prevent MRE11-dependent  
480 degradation. In wild-type cells, with time, reversed forks degenerate into DSBs, possibly because of  
481 unscheduled RECQ1-mediated fork restoration [13]. In the absence of WRN exonuclease, perturbed



482 replication forks undergo a further cycle of degradation of nascent strand by MRE11 and/or DNA2,  
483 which leads to ssDNA accumulation and engagement of RAD51. The accumulation of ssDNA and  
484 possibly engagement of RAD51 make perturbed replication forks resistant to DSBs and interfere with  
485 checkpoint signalling, resulting in a mild defect in CHK1 activation. In the absence of WRN  
486 exonuclease, more perturbed forks become inactivated over-time and RAD51 is required also during  
487 recovery from CPT to support repair at template gaps left behind the inactive forks. Engagement of  
488 RAD51 during recovery results in elevated activation of the MUS81 complex in G2/M to deal with  
489 intermediates, and to limit mitotic defects and cell death. As CPT is a chemotherapeutic, our data also  
490 indicate that tumors with impaired function of the WRN exonuclease can be sensitized to treatment  
491 by genetic or chemical interference with the MUS81 complex in mitosis, which is less relevant for  
492 survival in cells expressing the WRN wild-type.

493

494

495

## 496 MATERIALS AND METHODS

497

### 498 **Cell lines and culture conditions**

499 The SV40-transformed WRN-deficient fibroblast cell line (AG11395) was obtained from Coriell Cell  
500 Repositories (Camden, NJ, USA). To produce stable cell lines, AG11395 (WS) fibroblasts were  
501 transduced with retroviruses expressing the full-length cDNA encoding wild-type WRN (WS<sup>WT</sup>),  
502 exonuclease-dead (WS<sup>E84A</sup>), or helicase-dead (WS<sup>K577M</sup>)[47]. All the cell lines were maintained in  
503 Dulbecco's modified Eagle's medium (DMEM; Life Technologies) supplemented with 10% FBS  
504 (Boehringer Mannheim) and incubated at 37 °C in a humidified 5% CO<sub>2</sub> atmosphere.

505

### 506 **Chemicals**

507 Camptothecin (ENZO Lifesciences) was dissolved in DMSO and a stock solution (10 mM) was  
508 prepared and stored at -20°C. Mirin (Calbiochem), an inhibitor of MRE11 exonuclease activity, was  
509 used at 50 μM; the B02 compound (Selleck), an inhibitor of RAD51 activity, was used at 27 μM. C5  
510 (ref), DNA2 inhibitor C5 was dissolved in DMSO and used at final concentration of 300 μM [20].  
511 IdU and CldU (Sigma-Aldrich) were dissolved in sterile DMEM at 2.5mM and 200mM respectively  
512 and stored at -20°C.

513

### 514 **Plasmids transfection**

515 Plasmid expressing the phospho - mimic (Flag-CHK1<sup>317/345D</sup>) mutant form of CHK1, a kind gift from  
516 Professor K.K. Khanna (Queensland Institute of Medical Research, Australia) was generated as  
517 described [28]. To express the plasmids, cells were transfected using the Neon™ Transfection System  
518 Kit (Invitrogen), according to the manufacturer's instructions.

519

### 520 **Immunofluorescence assays**

521 Cells were grown on 35-mm coverslips and harvested at the indicated times after treatments. For  
522 RAD51 IF, after further washing with PBS, cells were pre-extracted with 0,5% TritonX-100 and fixed  
523 with 3% PFA / 2% sucrose at RT for 10min. After blocking in 3% BSA for 15 min, staining was  
524 performed with rabbit monoclonal anti-RAD51 (Bioss, 1:100) diluted in a 1% BSA / 0,1% saponin  
525 in PBS solution, for 1h at 37° in a humidified chamber. For 53BP1, pS87MUS81 and α-tubulin  
526 staining, cells were fixed with 4% PFA at RT for 10 min. Cells were subsequently permeabilized with  
527 0,4% Triton-X100. Staining with primary antibodies diluted in a 1% BSA / 0,1% saponin in PBS  
528 solution was carried out for 1h at RT. After extensive washing with PBS, specie-specific fluorophore-  
529 conjugated antibody (Invitrogen) was applied for 1h at RT followed by counterstaining with 0.5

530 mg/ml DAPI. Secondary antibody was used at 1:200 dilution. Images were acquired as greyscale files  
531 using Metaview software (MDS Analytical Technologies) and processed using Adobe Photoshop  
532 CS3 (Adobe). For each time point, at least 200 nuclei were examined, and foci were scored at 40×.  
533 Only nuclei with > 5 foci were considered as positive and were quantified using ImageJ.

534

### 535 **EdU incorporation assay**

536 To label replicated DNA, cells were incubated with 10 μM EdU for 30 minutes. Samples were fixed  
537 with 4% PFA at RT for 10 min and cells were subsequently permeabilized with 0,5% Triton-X100.  
538 EdU incorporation was detected using the Click-It Edu Alexa Fluor 488 Imaging Kit (Invitrogen)  
539 according to the manufacturer's instructions.

540

### 541 **Antibodies**

542 The primary antibodies used were: anti-pS10H3 (1:1000, Santa Cruz Biotechnologies), anti-Cyclin  
543 A (IF: 1:100, Santa Cruz Biotechnologies), anti-53BP1 (1:300, Millipore), anti-BrdU (1:80, Abcam;  
544 CldU detection), anti-BrdU (1:50, Becton Dickinson; anti-IdU detection), anti-pS87MUS81 (ref  
545 1:200), anti-RAD51 (1:1000, Bioss Antibodies), anti-αTubulin (1:50, Sigma-Aldrich) and anti-Lamin  
546 B1 (1:10000, Abcam).

547

### 548 **Chromatin fractionation and Western blot analysis**

549 Chromatin fractionation experiments were performed as previously described [48]. Western blotting  
550 was performed using standard methods. Blots were incubated with primary antibodies against: rabbit  
551 anti-pCHK1(S345) (Cell Signalling Technology), mouse anti-CHK1 (Santa Cruz Biotechnology),  
552 rabbit anti-RAD51 (Bioss Antibodies), mouse anti-RPA32 (Calbiochem), rabbit anti-RPA70  
553 (GeneTex), mouse anti-GAPDH (Santa Cruz Biotechnology) and rabbit anti-Lamin B1 (Abcam).  
554 After incubations with horseradish peroxidase-linked secondary antibodies (1:20000, Jackson  
555 Immunosciences), the blots were developed using the chemiluminescence detection kit  
556 WesternBright ECL HRP substrate (Advansta) according to the manufacturer's instructions.  
557 Quantification was performed on scanned images of blots using the Image Lab software, and the  
558 values shown on the graphs represent normalization of the protein content evaluated through  
559 LaminB1 or GAPDH immunoblotting.

560

561

562

563

564 **Clonogenic survival**

565 Cells were plated onto 35mm dishes, after 24h they were treated with different doses of CPT. After  
566 18, cells were washed, trypsinized and seeded in 60mm dishes. After 14-21 days, plates were stained  
567 with crystal violet and colonies counted.

568

569 **DNA fibres analysis**

570 DNA fibres were prepared, spread out and immunodecorated as previously described [11]. Images  
571 were acquired randomly from fields with untangled fibres using Eclipse 80i Nikon Fluorescence  
572 Microscope, equipped with a VideoConfocal (ViCo) system. The length of labeled tracks were  
573 measured using the Image-Pro-Plus 6.0 software. A minimum of 100 individual fibres were analysed  
574 for each experiment and the mean of at least three independent experiments presented.

575

576 **Detection of nascent single-stranded DNA**

577 To detect nascent single-stranded DNA (ssDNA), cells were plated onto 22x22 coverslips in 35mm  
578 dishes. After 24h, the cells were labelled for 15 min before the treatment with 250 $\mu$ M IdU (Sigma-  
579 Aldrich), cells were then treated with CPT 5 $\mu$ M for different time points. Next, cells were washed  
580 with PBS, permeabilized with 0.5% Triton X-100 for 10 min at 4°C and fixed with 2% sucrose, 3%  
581 PFA. For ssDNA detection, cells were incubated with primary mouse anti-BrdU antibody (Becton  
582 Dickinson) for 1h at 37°C in 1%BSA/PBS, followed by Alexa Fluor488-conjugated goat-anti-Mouse  
583 (Invitrogen), and counterstained with 0.5 $\mu$ g/ml DAPI. Slides were analysed with Eclipse 80i Nikon  
584 Fluorescence Microscope, equipped with a VideoConfocal (ViCo) system. For each time point, at  
585 least 100 nuclei were scored at 60 $\times$ . Parallel samples either incubated with the appropriate normal  
586 serum or only with the secondary antibody confirmed that the observed fluorescence pattern was not  
587 attributable to artefacts. Fluorescence intensity for each sample was then analysed using ImageJ  
588 software.

589

590 **Statistical analysis**

591 All the data are presented as means of at least two independent experiments. Statistical comparisons  
592 of WS<sup>WT</sup> or WRN-mutant cells to their relevant control were analysed by ANOVA or Mann-Whitney  
593 test.  $P < 0.5$  was considered as significant.

594

595

596

597

## 598 ACKNOWLEDGMENTS

599 We are grateful to Prof. Massimo Lopes (IMCR, University of Zurich) for scientific discussion. We  
600 thank all members of our laboratories for discussion. This work was supported by Associazione  
601 Italiana per la Ricerca sul Cancro (AIRC) to PP (IG17383) and to AF (IG119971), and in part by NIH  
602 R01CA085344 to BHS, R50CA211397 to LZ and GM123554 to JLC.

603

## 604 AUTHOR CONTRIBUTIONS

605 F.A.A. performed the analysis of CHK1 phosphorylation, fork recruitment by PLA and chromatin  
606 fractionation, and performed experiments to determine DNA damage. A.P. performed the analysis of  
607 MUS81 phosphorylation and experiments to evaluate mitotic abnormalities. E.M. analysed the  
608 persistence of RAD51 and parental ssDNA. F.A.A., A.P., E.M. analysed data, contributed to  
609 designing the experiments and writing the manuscript. A.F. and P.P. designed experiments, analysed  
610 data and wrote the paper. L.Z., J.L.C. and B.H.S. provided the DNA2 inhibitor C5, advised the  
611 relevant experiments, and revised the manuscript. All authors approved the paper.

612

## 613 CONFLICT OF INTEREST

614 The authors declare that they do not have any conflict of interest.

615

616

## 617 REFERENCES

618

- 619 1. Técher H, Koundrioukoff S, Nicolas A, Debatisse M. The impact of replication stress on  
620 replication dynamics and DNA damage in vertebrate cells. *Nat Rev Genet*. Nature Publishing  
621 Group; 2017; doi:10.1038/nrg.2017.46
- 622 2. Franchitto A, Pichierri P. Replication fork recovery and regulation of common fragile sites  
623 stability. *Cell Mol Life Sci*. 2014;71: 4507–4517. doi:10.1007/s00018-014-1718-9
- 624 3. Zeman MK, Cimprich KA. Causes and consequences of replication stress. *Nat Cell Biol*.  
625 2014;16: 2–9. doi:10.1038/ncb2897
- 626 4. Macheret M, Halazonetis TD. DNA Replication Stress as a Hallmark of Cancer. *Annu Rev*  
627 *Pathol Mech Dis*. 2015;10. doi:10.1146/annurev-pathol-012414-040424
- 628 5. Magdalou I, Lopez BS, Pasero P, Lambert S a E. The causes of replication stress and their  
629 consequences on genome stability and cell fate. *Semin Cell Dev Biol*. Elsevier Ltd; 2014;30:  
630 154–164. doi:10.1016/j.semcdb.2014.04.035
- 631 6. Hills SA, Diffley JFX. DNA Replication and Oncogene-Induced Replicative Stress. *Curr*  
632 *Biol*. Elsevier Ltd; 2014;24: R435–R444. doi:10.1016/j.cub.2014.04.012
- 633 7. Ciccia A, Elledge SJ. The DNA Damage Response: Making It Safe to Play with Knives. *Mol*  
634 *Cell*. Elsevier Inc.; 2010;40: 179–204. doi:10.1016/j.molcel.2010.09.019
- 635 8. Pichierri P, Ammazalorso F, Bignami M, Franchitto A. The Werner Syndrome protein:  
636 Linking the replication checkpoint response to genome stability. *Aging (Albany NY)*.  
637 2011;3: 311–318. doi:100293 [pii]
- 638 9. Rossi ML, Ghosh AK, Bohr V a. Roles of Werner syndrome protein in protection of genome  
639 integrity. *DNA Repair (Amst)*. Elsevier B.V.; 2010;9: 331–44.  
640 doi:10.1016/j.dnarep.2009.12.011
- 641 10. Franchitto A, Pichierri P. Understanding the molecular basis of common fragile sites  
642 instability: Role of the proteins involved in the recovery of stalled replication forks. *Cell*  
643 *Cycle*. 2011;10: 4039–4046. doi:10.4161/cc.10.23.18409
- 644 11. Iannascoli C, Palermo V, Murfunì I, Franchitto A, Pichierri P. The WRN exonuclease  
645 domain protects nascent strands from pathological MRE11/EXO1-dependent degradation.  
646 *Nucleic Acids Res*. 2015;43: 9788–803. doi:10.1093/nar/gkv836
- 647 12. Ray Chaudhuri A, Hashimoto Y, Herrador R, Neelsen KJ, Fachinetti D, Bermejo R, et al.  
648 Topoisomerase I poisoning results in PARP-mediated replication fork reversal. *Nature*  
649 *Structural & Molecular Biology*. Nature Publishing Group; 2012. pp. 417–423.  
650 doi:10.1038/nsmb.2258

- 651 13. Berti M, Ray Chaudhuri A, Thangavel S, Gomathinayagam S, Kenig S, Vujanovic M, et al.  
652 Human RECQ1 promotes restart of replication forks reversed by DNA topoisomerase I  
653 inhibition. *Nat Struct Mol Biol.* Nature Publishing Group; 2013;20: 347–54.  
654 doi:10.1038/nsmb.2501
- 655 14. Bhat KP, Cortez D. RPA and RAD51: fork reversal, fork protection, and genome stability.  
656 *Nat Struct Mol Biol.* 2018;25: 446–453. doi:10.1038/s41594-018-0075-z
- 657 15. Quinet A, Lemaçon D, Vindigni A. Replication Fork Reversal: Players and Guardians. *Mol*  
658 *Cell.* 2017;68: 830–833. doi:10.1016/j.molcel.2017.11.022
- 659 16. Kolinjivadi AM, Sannino V, de Antoni A, Técher H, Baldi G, Costanzo V. Moonlighting at  
660 replication forks - a new life for homologous recombination proteins BRCA1, BRCA2 and  
661 RAD51. *FEBS Lett.* 2017; 1–18. doi:10.1002/1873-3468.12556
- 662 17. Feng W, Jasin M. Homologous Recombination and Replication Fork Protection: BRCA2 and  
663 More! *Cold Spring Harb Symp Quant Biol.* 2017;82: 329–338.  
664 doi:10.1101/sqb.2017.82.035006
- 665 18. Costanzo V. Brca2, Rad51 and Mre11: Performing balancing acts on replication forks. *DNA*  
666 *Repair (Amst).* Elsevier B.V.; 2011;10: 1060–1065. doi:10.1016/j.dnarep.2011.07.009
- 667 19. Pellegrini L, Venkitaraman A. Emerging functions of BRCA2 in DNA recombination.  
668 *Trends Biochem Sci.* 2004;29: 310–316. doi:10.1016/j.tibs.2004.04.009
- 669 20. Liu W, Zhou M, Li Z, Li H, Polaczek P, Dai H, et al. A Selective Small Molecule DNA2  
670 Inhibitor for Sensitization of Human Cancer Cells to Chemotherapy. *EBioMedicine.* The  
671 Authors; 2016;6: 73–86. doi:10.1016/j.ebiom.2016.02.043
- 672 21. Murfun I, Nicolai S, Baldari S, Crescenzi M, Bignami M, Franchitto a, et al. The WRN and  
673 MUS81 proteins limit cell death and genome instability following oncogene activation.  
674 *Oncogene.* Nature Publishing Group; 2012;32: 610–20. doi:10.1038/onc.2012.80
- 675 22. Hanada K, Budzowska M, Davies SL, van Drunen E, Onizawa H, Beverloo HB, et al. The  
676 structure-specific endonuclease Mus81 contributes to replication restart by generating  
677 double-strand DNA breaks. *Nat Struct Mol Biol.* 2007;14: 1096–1104.  
678 doi:10.1038/nsmb1313
- 679 23. Hashimoto Y, Chaudhuri AR, Lopes M, Costanzo V. Rad51 protects nascent DNA from  
680 Mre11-dependent degradation and promotes continuous DNA synthesis. *Nat Struct Mol Biol.*  
681 2010;17: 1305–1311. doi:10.1038/nsmb.1927
- 682 24. Petermann E, Orta ML, Issaeva N, Schultz N, Helleday T. Hydroxyurea-Stalled Replication  
683 Forks Become Progressively Inactivated and Require Two Different RAD51-Mediated  
684 Pathways for Restart and Repair. *Mol Cell.* 2010;37: 492–502.

- 685 doi:10.1016/j.molcel.2010.01.021
- 686 25. Carr AM, Lambert S. Replication stress-induced genome instability: The dark side of  
687 replication maintenance by homologous recombination. *J Mol Biol. Elsevier Ltd*; 2013;425:  
688 4733–4744. doi:10.1016/j.jmb.2013.04.023
- 689 26. Lee J, Kumagai A, Dunphy WG. The Rad9-Hus1-Rad1 checkpoint clamp regulates  
690 interaction of TopBP1 with ATR. *J Biol Chem.* 2007;282: 28036–28044.  
691 doi:10.1074/jbc.M704635200
- 692 27. Delacroix S, Wagner JM, Kobayashi M, Yamamoto KI, Karnitz LM. The Rad9-Hus1-Rad1  
693 (9-1-1) clamp activates checkpoint signaling via TopBP1. *Genes Dev.* 2007;21: 1472–1477.  
694 doi:10.1101/gad.1547007
- 695 28. Gatei M, Sloper K, Sørensen C, Syljuäsen R, Falck J, Hobson K, et al. Ataxia-telangiectasia-  
696 mutated (ATM) and NBS1-dependent phosphorylation of Chk1 on Ser-317 in response to  
697 ionizing radiation. *J Biol Chem.* 2003;278: 14806–14811. doi:10.1074/jbc.M210862200
- 698 29. Feng W, Jasin M. BRCA2 suppresses replication stress-induced mitotic and G1  
699 abnormalities through homologous recombination. *Nat Commun.* 2017;8: 525.  
700 doi:10.1038/s41467-017-00634-0
- 701 30. Ying S, Minocherhomji S, Chan KL, Palmai-Pallag T, Chu WK, Wass T, et al. MUS81  
702 promotes common fragile site expression. *Nat Cell Biol. Nature Publishing Group*; 2013;15:  
703 1001–7. doi:10.1038/ncb2773
- 704 31. Naim V, Wilhelm T, Debatisse M, Rosselli F. ERCC1 and MUS81-EME1 promote sister  
705 chromatid separation by processing late replication intermediates at common fragile sites  
706 during mitosis. *Nat Cell Biol. Nature Publishing Group*; 2013;15: 1008–15.  
707 doi:10.1038/ncb2793
- 708 32. Palma A, Pugliese GM, Murfunì I, Marabitti V, Malacaria E, Rinalducci S, et al.  
709 Phosphorylation by CK2 regulates MUS81/EME1 in mitosis and after replication stress.  
710 *Nucleic Acids Res. Oxford University Press*; 2018;46: 5109–5124. doi:10.1093/nar/gky280
- 711 33. Lukas C, Savic V, Bekker-Jensen S, Doil C, Neumann B, Pedersen RS, et al. 53BP1 nuclear  
712 bodies form around DNA lesions generated by mitotic transmission of chromosomes under  
713 replication stress. *Nat Cell Biol. Nature Publishing Group*; 2011;13: 243–253.  
714 doi:10.1038/ncb2201
- 715 34. Lai X, Broderick R, Bergoglio V, Zimmer J, Badie S, Niedzwiedz W, et al. MUS81 nuclease  
716 activity is essential for replication stress tolerance and chromosome segregation in BRCA2-  
717 deficient cells. *Nat Commun.* 2017;8: 15983. doi:10.1038/ncomms15983
- 718 35. Schlacher K, Wu H, Jasin M. A Distinct Replication Fork Protection Pathway Connects



- 719 Fanconi Anemia Tumor Suppressors to RAD51-BRCA1/2. *Cancer Cell*. Elsevier Inc.;  
720 2012;22: 106–116. doi:10.1016/j.ccr.2012.05.015
- 721 36. Schlacher K, Christ N, Siaud N, Egashira A, Wu H, Jasin M. Double-Strand Break Repair-  
722 Independent Role for BRCA2 in Blocking Stalled Replication Fork Degradation by MRE11.  
723 *Cell*. Elsevier Inc.; 2011;145: 529–542. doi:10.1016/j.cell.2011.03.041
- 724 37. Lemaçon D, Jackson J, Quinet A, Brickner JR, Li S, Yazinski S, et al. MRE11 and EXO1  
725 nucleases degrade reversed forks and elicit MUS81-dependent fork rescue in BRCA2-  
726 deficient cells. *Nat Commun*. 2017;8: 860. doi:10.1038/s41467-017-01180-5
- 727 38. Kolinjivadi AM, Sannino V, De Antoni A, Zadorozhny K, Kilkenny M, Técher H, et al.  
728 Smarcal1-Mediated Fork Reversal Triggers Mre11-Dependent Degradation of Nascent DNA  
729 in the Absence of Brca2 and Stable Rad51 Nucleofilaments. *Mol Cell*. 2017;67: 867–881.e7.  
730 doi:10.1016/j.molcel.2017.07.001
- 731 39. Bolterstein E, Rivero R, Marquez M, McVey M. The *Drosophila* Werner exonuclease  
732 participates in an exonuclease-independent response to replication stress. *Genetics*.  
733 2014;197: 643–52. doi:10.1534/genetics.114.164228
- 734 40. García-Rodríguez N, Morawska M, Wong RP, Daigaku Y, Ulrich HD. Spatial separation  
735 between replisome- and template-induced replication stress signaling. *EMBO J*. 2018;  
736 e98369. doi:10.15252/embj.201798369
- 737 41. Hashimoto Y, Puddu F, Costanzo V. RAD51- and MRE11-dependent reassembly of  
738 uncoupled CMG helicase complex at collapsed replication forks. *Nat Struct Mol Biol*. Nature  
739 Publishing Group; 2011;19: 17–24. doi:10.1038/nsmb.2177
- 740 42. Sidorova JM, Kehrli K, Mao F, Monnat R. Distinct functions of human RECQ helicases  
741 WRN and BLM in replication fork recovery and progression after hydroxyurea-induced  
742 stalling. *DNA Repair (Amst)*. 2013;12: 128–139. doi:10.1016/j.dnarep.2012.11.005
- 743 43. Shechter D, Costanzo V, Gautier J. Regulation of DNA replication by ATR: signaling in  
744 response to DNA intermediates. *DNA Repair (Amst)*. 2004;3: 901–8.  
745 doi:10.1016/j.dnarep.2004.03.020
- 746 44. Friedel AM, Pike BL, Gasser SM. ATR/Mec1: coordinating fork stability and repair. *Curr*  
747 *Opin Cell Biol*. 2009;21: 237–244. doi:10.1016/j.ceb.2009.01.017
- 748 45. Flynn RL, Zou L. ATR: A master conductor of cellular responses to DNA replication stress.  
749 *Trends Biochem Sci*. 2011;36: 133–140. doi:10.1016/j.tibs.2010.09.005
- 750 46. Parpys AC, Seelbach JI, Becker S, Behr M, Wrona A, Jend C, et al. High levels of RAD51  
751 perturb DNA replication elongation and cause unscheduled origin firing due to impaired  
752 CHK1 activation. *Cell Cycle*. 2015;14: 3190–202. doi:10.1080/15384101.2015.1055996

- 753 47. Pirzio LM, Pichierri P, Bignami M, Franchitto A. Werner syndrome helicase activity is  
754 essential in maintaining fragile site stability. *J Cell Biol.* 2008;180: 305–314.  
755 doi:10.1083/jcb.200705126
- 756 48. Murfuni I, Basile G, Subramanyam S, Malacaria E, Bignami M, Spies M, et al. Survival of  
757 the Replication Checkpoint Deficient Cells Requires MUS81-RAD52 Function. Maizels N,  
758 editor. *PLoS Genet.* 2013;9: e1003910. doi:10.1371/journal.pgen.1003910  
759  
760

## 761 LEGENDS TO FIGURES

762

763 **Figure 1. Loss of WRN exonuclease activity leads to formation of nascent ssDNA which**  
 764 **compromises formation of DSBs in response to a low-dose of camptothecin** (A) Evaluation of  
 765 ssDNA by anti-IdU immunofluorescence under non-denaturing condition. Nascent DNA was pre-  
 766 labelled for 15 min with IdU before treatment and labelling remained during treatment with CPT. Dot  
 767 plots show the mean intensity of ssDNA staining for single nuclei from cells expressing the wild-type  
 768 ( $WS^{WT}$ ) or the *exo*-dead form of WRN ( $WS^{E84A}$ ). Cells were either left untreated or challenged with  
 769 50 nM CPT for increasing periods, as indicated. The intensity of the anti-IdU immunofluorescence  
 770 was measured in at least 200 nuclei from three independent experiments. Values are represented as  
 771 means  $\pm$ SE. Representative images of ssDNA labelling are shown. (B) Evaluation of nascent ssDNA  
 772 in cells treated with nuclease inhibitors. Cells were treated with Mirin, C5 or both 30 min before IdU  
 773 labelling and 45 min before CPT treatment for 4 h, and then subjected to the ssDNA assay. The graph  
 774 shows the mean intensity of IdU fluorescence measured from two independent experiments ( $n=200$ ),  
 775 data are presented as mean  $\pm$ SE. Statistical analysis in A and B was performed by the Mann–Whitney  
 776 test ( $**P < 0.1$ ;  $***P < 0.01$ ;  $****P < 0.001$ ) (C) Analysis of DSB accumulation by the neutral  
 777 Comet assay. Cells were treated or not with CPT 50 nM for the indicated time, or with 5  $\mu$ M CPT  
 778 (high-dose) for 1 h, and then subjected to the neutral Comet assay. Where indicated, cells were pre-  
 779 treated with Mirin, C5 or both. In the graph, data are presented as mean tail moment  $\pm$  SE from two  
 780 independent experiments (ns = not significant;  $**P < 0.1$ ;  $***P < 0.01$ ;  $****P < 0.001$ ; ANOVA  
 781 test). Representative images from the neutral Comet assay are shown.

782

783 **Figure 2. Loss of WRN exonuclease stimulates engagement of RAD51 after CPT.** (A) WB  
 784 analysis of chromatin association of RAD51 and RPA32 in wild-type ( $WS^{WT}$ ) and in cells expressing  
 785 the *exo*-dead mutant form of WRN ( $WS^{E84A}$ ). Cells were treated or not with CPT for 4 h. LaminB1  
 786 was used as loading control. The blot is representative of three replicates. The graphs show the  
 787 quantification of the amount of RAD51 or RPA32 normalised against LaminB1 (mean $\pm$ SE). (B)  
 788 Quantitative immunofluorescence analysis of RAD51 foci in  $WS^{WT}$  and  $WS^{E84A}$  cells. Cells were  
 789 treated with CPT 50nM for 4h, triton-extracted and subjected to RAD51 immunostaining. Graph  
 790 shows the intensity of RAD51 immunostaining for each cell with scorable foci ( $n>3$ ). Values are  
 791 presented as means  $\pm$  SE ( $*** P < 0.01$ ;  $**** P < 0.001$ ; Mann–Whitney test). Representative images  
 792 are shown.

793

794 **Figure 3. RAD51 recruitment persisted during recovery from CPT in the absence of the WRN**  
 795 **exonuclease.** (A) WB analysis of chromatin association of RAD51, RPA70 and RPA32 in wild-type  
 796 ( $WS^{WT}$ ) and in cells expressing the exo-dead mutant form of WRN ( $WS^{E84A}$ ). Cells were treated or  
 797 not with 50nM CPT for 4h followed by recovery as indicated. LaminB1 was used as loading control.  
 798 The blot is representative of three replicates. The graphs show the quantification of the amount of  
 799 RAD51 or RPA32 normalised against LaminB1 (mean $\pm$ SE). (B) Quantitative immunofluorescence  
 800 analysis of RAD51 foci in  $WS^{WT}$  and  $WS^{E84A}$  cells. Cells were treated with CPT 50nM for 4h and  
 801 recovered or not as indicated. Graph shows the intensity of RAD51 immunostaining for each cell  
 802 with scorable foci ( $n>3$ ). Values are presented as means  $\pm$  SE (\*\* $P < 0.01$ ; \*\*\*\* $P < 0.001$ ; Mann–  
 803 Whitney test). Representative images are shown. (C) Quantitative immunofluorescence analysis of  
 804 RAD51 foci in  $WS^{WT}$  and  $WS^{E84A}$  cells pre-treated with the nuclease inhibitors. Cells were pre-treated  
 805 with the indicated inhibitors prior to be challenged with CPT 50nM for 4h and recovered or not as  
 806 indicated. Graph shows the intensity of RAD51 immunostaining for each cell with scorable foci  
 807 ( $n>3$ ). Values are presented as means  $\pm$  SE (\*\* $P < 0.1$ ; \*\*\*\* $P < 0.001$ ; Mann–Whitney test). (D-E)  
 808 analysis of parental ssDNA. Parental DNA was labelled with IdU as indicated in the experimental  
 809 scheme (D). (E) The graph shows the amount of parental ssDNA calculated as mean intensity of IdU  
 810 fluorescence measured from two independent experiments ( $n=200$ ), data are presented as mean  $\pm$ SE.  
 811 Statistical analysis was performed by the Mann–Whitney test (\*\* $P < 0.1$ ; \*\*\*\* $P < 0.001$ ).  
 812 Representative images are shown.

813

814 **Figure 4. RAD51 inhibition does not impair replication fork recovery following treatment with**  
 815 **low dose of CPT.** (A) Experimental scheme of dual-labelling replication assay with DNA fibres. Red  
 816 tract: CldU; Green tract: IdU. (B) The graph shows the average number of stalled forks (red only  
 817 tracts opposed to the active forks marked as red+green tracks) after recovery from 50nM CPT  
 818 treatment. Where indicated, RAD51 inhibitor (B02; RAD51i) was added to cultures together with  
 819 CPT and during the IdU pulse. Data are presented as mean  $\pm$  SE. One-hundred IdU-positive tracts  
 820 were analysed in each experimental point ( $n=2$ ). (C) The graph shows the average number of new  
 821 origins (green only tracts) after recovery from 50nM CPT treatment. Where indicated, RAD51  
 822 inhibitor (B02; RAD51i) was added to the cultures together with CPT and during the IdU pulse.  
 823 Values are presented as mean  $\pm$  SE. In B and C statistical analysis was performed by Anova test. (D)  
 824 Representative DNA fibres fields are shown in the images.

825

826 **Figure 5. Loss of WRN exonuclease activity affects phosphorylation of CHK1.** (A) WB analysis  
 827 of CHK1 phosphorylation at S345 in wild-type ( $WS^{WT}$ ) and in cells expressing the exo-dead mutant

828 form of WRN ( $WS^{E84A}$ ) or the helicase-dead form ( $WS^{K577M}$ ). Cells were treated or not with 50nM  
 829 CPT as indicated. Total CHK1 and GAPDH were used as loading controls. The blot is representative  
 830 of three replicates. Below is reported quantification of p345CHK1 phosphorylation normalised  
 831 against total CHK1. (B) Immunofluorescence analysis of pS345CHK1 in  $WS^{WT}$  and  $WS^{E84A}$  cells  
 832 treated with CPT 50nM for 4h. Numbers in insets represent the mean percentage of pS345CHK1-  
 833 positive nuclei ( $n=2$ ; errors are not shown but are  $< 15\%$  of the mean). (C) Immunofluorescence  
 834 analysis of pT1989ATR in  $WS^{WT}$  and  $WS^{E84A}$  treated with CPT 50nM for 4h. Numbers in insets  
 835 represent the mean percentage of pS345CHK1-positive nuclei ( $n=2$ ; errors are not shown but are  $<$   
 836  $15\%$  of the mean). (D-E) Analysis of TopBP1 or RAD9 recruitment at nascent ssDNA by PLA.  
 837 Nascent strand was labelled with IdU for 15min before cells were treated with 50nM CPT for 4h.  
 838 PLA was performed under native conditions using anti-IdU to detect nascent ssDNA and anti-  
 839 TopBP1 or RAD9 to detect the protein. Negative controls are from samples processed with anti-IdU  
 840 only. The graphs show the number of PLA spots in each nucleus ( $n=300$  from 3 independent  
 841 replicates). Statistical analysis was performed by the Mann–Whitney test (ns = not significant; \*\*\*\* $P$   
 842  $< 0.001$ ). Representative images are shown. (F) Duplicated samples from D-E were analysed for the  
 843 presence of nascent ssDNA by native anti-IdU detection only. The graph shows the mean intensity  
 844 of IdU fluorescence measured from two independent experiments ( $n=200$ ), data are presented as mean  
 845  $\pm$ SE. Statistical analysis was performed by the Mann–Whitney test (\*\* $P < 0.01$ ).

846

847 **Figure 6. Expression of a phosphomimic CHK1 mutant restores wild-type levels of nascent**  
 848 **ssDNA in WRN exonuclease-deficient cells.** (A) WB analysis of FLAG-CHK1<sup>317/345D</sup> expression in  
 849  $WS^{E84A}$  cells. (B) Evaluation of nascent ssDNA formation. Cells treated with 50nM CPT for 4h were  
 850 analysed for the presence of nascent ssDNA by native anti-IdU detection. The graph shows the mean  
 851 intensity of IdU fluorescence measured from two independent experiments ( $n=200$ ), data are  
 852 presented as mean  $\pm$ SE. Statistical analysis was performed by the Mann–Whitney test ( $*P < 0.5$ ;  
 853 \*\*\* $P < 0.01$ ; \*\*\*\* $P < 0.001$ ). Representative images are shown.

854

855 **Figure 7. WRN exonuclease-deficient cells show enhanced MUS81 phosphorylation on S87 and**  
 856 **mitotic defects.** (A) Anti-pS87MUS81 immunofluorescence staining (red) was performed in wild-  
 857 type and WRN exonuclease-dead expressing cells. The S-phase cells (green) were revealed with short  
 858 EdU pulse followed by Click-IT reaction. Nuclei were depicted with DAPI staining (blue). The mean  
 859 frequency ( $\pm$ SE;  $n=3$ ) of pS87-MUS81-positive nuclei are indicated in the representative images. (B)  
 860  $WS$  cells expressing the  $WS^{E84A}$  mutant were treated with CPT for 4h and then released in fresh  
 861 medium for 18 hours. The RAD51 inhibitor B02 was added with CPT and during the recovery. The

862 frequency ( $\pm$  SE;  $n=2$ ) of pS87-MUS81-positive nuclei are indicated as percentage in the  
863 representative images. (C) The graph shows the mean percentage  $\pm$  SE of bulky anaphase bridges  
864 analysed in untreated and CPT-treated cells expressing WRN wild-type and the exonuclease-deficient  
865 mutant. The number of anaphases counted for each experimental point are indicated above as  $n$ .  
866 Randomly-selected representative anaphases with bridges are shown. (D) Representative images of  
867 mitotic aberrations analysed in  $\alpha$ -Tubulin (green) and DAPI-stained cells are shown above the graph  
868 indicating the frequency of each event after treatment with 50nM of CPT for 4h followed by a 18h  
869 recovery. Data are presented as mean $\pm$ SE. Statistical analysis was performed by the ANOVA test ( $*P$   
870  $< 0.5$ ). (E) Analysis of 53BP1 NBs. Cells were either untreated or treated with 50nM CPT as  
871 indicated. Samples were subjected to immunofluorescence using anti-53BP1 and anti-Cyclin A to  
872 evaluate 53BP1 fluorescence only in G1 cells (Cyclin A-negative). Graphs show the frequency of  
873 each class of nuclei in two independent replicates. Representative images from CPT-treated samples  
874 are shown.

875

876 **Figure 8. MUS81<sup>S87A</sup> mutant overexpression aggravates the mitotic phenotypes of WRN**  
877 **exonuclease-deficient cells.** (A) Bulky bridges, multinucleated and micro-nucleated cells were  
878 analysed with or without MUS81<sup>S87A</sup> mutant overexpression, in WRN wild-type and exonuclease-  
879 deficient cells. The graph represents the frequency of the aberration analysed in two independent  
880 experiments  $\pm$  SE. (B) Analysis of 53BP1 NBs formation in Cyclin A-negative cells. Representative  
881 images of fluorescence cells stained with anti-53BP1 (green) and Cyclin A (red). Nuclear DNA was  
882 counterstained with DAPI (blue). For each point at least 300 nuclei were counted and cells with  $> 5$   
883 53BP1 NBs were considered as positive. The graph shows the quantification of 53BP1-positive G1  
884 cells. (C) Clonogenic assay in cells treated with low-doses of CPT. Cells were exposed to different  
885 doses of CPT for 18h, re-plated at low density and survival evaluated as percentages of colonises  
886 normalised against the untreated. Statistical analyses in A-C were performed by ANOVA test ( $* P <$   
887  $0.5$ ;  $** P < 0.1$ ).

888

889 **Figure 9. Proposed model of the effect of prolonged treatment with nanomolar CPT doses in**  
890 **absence of WRN exonuclease** (see text for details).

891

892 SUPPORTING INFORMATION LEGENDS

893

894 **Supplementary Figures and Legends.** The file contains five supplementary figures and their  
895 legends.





Figure 1



A



B




Figure 2








Figure 3

D



E







Figure 3 cont'd


A



B



C



D




Figure 4








Figure 5

D



E



F




Figure 5 cont'd






Figure 6






Figure 7









Figure 7 cont'd

A



B




Figure 8

C




Figure 8 cont'd




Figure 9