

Measurements of quantum yields of bromine atoms in the photolysis of bromoform from 266 to 324 nm

Kyle D. Bayes

Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA

Darin W. Toohey

Program in Atmospheric and Oceanic Sciences, University of Colorado, Boulder, Colorado, USA

Randall R. Friedl and Stanley P. Sander

Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA

Received 23 August 2002; revised 4 November 2002; accepted 4 December 2002; published 5 February 2003.

[1] The quantum yield for the formation of bromine atoms in the photolysis of bromoform, CHBr_3 , has been measured between 266 and 324 nm. For 303 to 306 nm the quantum yields are unity within the experimental uncertainty of the measurements. At longer wavelengths, where the bromoform cross sections decrease rapidly, an apparent trend to slightly lower quantum yields is probably the result of systematic and random errors or incorrect CHBr_3 absorption cross sections. Support for a unit quantum yield for all wavelengths longer than 300 nm comes from the recent theoretical calculations of Peterson and Francisco. At 266 nm the bromine atom quantum yield is $0.76 (\pm 0.03)$, indicating that at least one additional dissociation channel becomes important at shorter wavelengths. For modeling of the troposphere, it is recommended that a quantum yield of unity be used for wavelengths of 300 nm and longer. *INDEX TERMS*: 0317 Atmospheric Composition and Structure: Chemical kinetic and photochemical properties; 0340 Atmospheric Composition and Structure: Middle atmosphere—composition and chemistry; 0365 Atmospheric Composition and Structure: Troposphere—composition and chemistry; *KEYWORDS*: bromoform, bromine atoms, quantum yield, photolysis

Citation: Bayes, K. D., D. W. Toohey, R. R. Friedl, and S. P. Sander, Measurements of quantum yields of bromine atoms in the photolysis of bromoform from 266 to 324 nm, *J. Geophys. Res.*, 108(D3), 4095, doi:10.1029/2002JD002877, 2003.

1. Introduction

[2] Bromoform, CHBr_3 , is present in the Earth's atmosphere in the parts per trillion (ppt) range [Berg *et al.*, 1984; Cicerone *et al.*, 1988; Fabian *et al.*, 1994; Sturges *et al.*, 2000]. As with other brominated methanes, the oceans appear to be the major primary source, with secondary contributions from polar microalgae and chlorination of domestic water supplies. Concentrations of bromoform are highest near macroalgae seaweeds, both dissolved in seawater and in the atmosphere [Moore *et al.*, 1993; Carpenter *et al.*, 1999]. A recent estimate of the emissions of organic bromine compounds concludes that the known sources are sufficient to account for the observed atmospheric concentrations [Carpenter and Liss, 2000].

[3] Although the concentration of bromine in the stratosphere is much less than that of chlorine, its presence can have a significant effect on ozone destruction because of its longer chain length and the interaction of the bromine and chlorine cycles [Wofsy *et al.*, 1975; Yung *et al.*, 1980;

McElroy *et al.*, 1986]. Atmospheric bromine chemistry has been reviewed recently [Lary, 1996; Lary *et al.*, 1996].

[4] There has been controversy about the importance of bromoform as a source of bromine in the lower stratosphere. In a survey of the total bromine budget, Wamsley *et al.* [1998] list bromoform under "negligible sources of stratospheric bromine." In contrast, Dvortsov *et al.* [1999, p. 1] concluded that for the midlatitude stratosphere close to the tropopause, "bromoform (CHBr_3) alone likely contributes more inorganic bromine than all the conventional long-lived sources (halons and methyl bromide) combined." This view was disputed by the measurements of Pfeilsticker *et al.* [2000] and Sturges *et al.* [2000] and also by the modeling study of Nielsen and Douglass [2001]; these studies conclude that the short-lived bromoalkanes (CH_2Br_2 , CHBr_3 , CHBr_2Cl) contribute no more than 10 to 30% of the bromine observed in the lower stratosphere.

[5] The atmospheric lifetime of bromoform in the troposphere is approximately a month [Singh, 1995; Butler and Rodriguez, 1996; Kurylo *et al.*, 1999]. Bromoform is destroyed by photolysis in the near ultraviolet ($\sim 75\%$) and by reaction with OH radicals ($\sim 25\%$). Both processes are thought to eventually release bromine atoms, but the mechanism and the efficiency of these releases have not

been determined. This paper reports laboratory measurements of bromine atom formation for CHBr_3 photolysis in the near ultraviolet.

2. Description of Experiments

[6] The photolysis of CHBr_3 was studied in a square (5×5 cm) aluminum tube 26 cm long, equipped with Brewster angle windows at both ends. The gas flow entered near the front window and exited near the rear window. A YAG laser (Continuum PL8000) provided pulsed radiation at 266 nm with a beam diameter of 5 mm. The YAG was also used at 532 nm to pump a dye laser (PDL-3) using various dyes (Rhodamine 640, Sulforhodamine 640, DCM). The red output from the dye laser was doubled to give radiation in the range 303 to 324 nm. The doubled dye laser pulses were expanded by a factor of three before entering the photolysis cell, giving them a diameter of about 5 mm. Both beams traversed the center of the photolysis cell. A joule meter (Molelectron EPM 1000) measured the average pulse energy at the front window just before and just after each photolysis run, and the average of the two measurements was used in the quantum yield determinations. While the doubled dye laser pulses passed through the Brewster window with no reflection loss, the 266 nm radiation had the wrong polarization, and so a correction was needed for the reflected radiation (measured as 4.5%).

[7] Bromine atoms were measured by vacuum ultraviolet resonance fluorescence with a combination of emission lines between 130 and 163 nm. A sealed lamp with MgF_2 window was excited by a ~ 10 watt, 180 MHz radio-frequency discharge and the emission was stabilized by an active feedback loop that maintained a constant IR output at ~ 840 nm [Toohey *et al.*, 1987; Brune *et al.*, 1989; McKinney *et al.*, 1997]. A solar blind photomultiplier (PM) with potassium bromide photochathode (EMR Model 541J) observed the scattered resonance lines at right angles to both the gas flow and the Br lamp radiation. Razor blade light traps on the walls opposite both lamp and detector served to reduce scatter off surfaces. TTL pulses (< 60 ns full width, half maximum) from an amplifier discriminator (Research Support Instruments) were counted as a function of time on a multichannel scaler (Ortec MCS-pci), stored and analyzed by weighted least squares using the PSI-Plot program.

[8] Bromoform was introduced by bubbling nitrogen or air at atmospheric pressure through liquid CHBr_3 at a temperature of 13 to 17°C. Most of the flow entered the vacuum system through a flow controller, while the excess mixture flowed through a 10 cm cell in a spectrophotometer (Cary model 4E). By continuously monitoring the absorbance at 266 nm, the concentration of CHBr_3 could be averaged over the course of a photolysis run. For the experiments using a mixture of molecular chlorine and vinyl bromide instead of bromoform, a similar procedure was used for monitoring Cl_2 using its absorbance at 330 nm. A separate spectrum of pure vinyl bromide showed that its cross section at 330 nm was 10^3 times less than that of Cl_2 and so for the concentrations used, vinyl bromide did not interfere with the Cl_2 measurements. Typical concentrations of the active ingredients are given in the caption to Figure 1.

[9] The pressure in the photolysis cell was varied from 1.5 to 20 Torr with no significant effect on the yield of




Figure 1. Examples of signals using method I (A) and method II (B). For trace A: $[\text{CHBr}_3] = 3.94 \cdot 10^{13}$ molecules cm^{-3} ; balance N_2 ; total pressure 2.04 Torr; laser at 266 nm, $0.96 \text{ mJ pulse}^{-1}$; 0.2 ms bin^{-1} , 3000 sweeps added. For trace B: $[\text{Cl}_2] = 2.6 \cdot 10^{13}$ molecules cm^{-3} ; $[\text{C}_2\text{H}_3\text{Br}] = 6.0 \cdot 10^{12}$ molecules cm^{-3} ; balance N_2 ; total pressure 2.04 Torr; laser at 305 nm, $1.18 \text{ mJ pulse}^{-1}$; 0.1 ms bin^{-1} , 2000 sweeps added.

bromine atoms. The quantum yield measurements were done at total pressures of 2 and 10 Torr and with a pulse repetition rate of 10 Hz. For the flow rates used and this pulse rate, each parcel of gas was exposed to three laser pulses. No difference in bromine atom yield was observed when the pulse repetition rate was varied between 10 Hz and 0.1 Hz. For the conditions used in the quantum yield experiments, the two spin states of the bromine atoms ($^2\text{P}_{3/2}$ and $^2\text{P}_{1/2}$) should have reached their equilibrium ratio within a millisecond. To be sure that this was not a problem, some runs used a mixture of 5% H_2 in N_2 as the carrier gas to increase the relaxation rate by a factor of ten [Johnson *et al.*, 1996]; there were no detectable differences in the yields or time dependence of Br signals when H_2 was present. The bromine atom signals increased linearly with pulse energy up to well above 1 mJ per pulse; quantum yield measurements were made with pulse energies of 0.8 to 1.2 mJ per pulse.

[10] The temperature of the gas in the photolysis cell was slightly above room temperature due to heating of the walls of the flow tube by the resonance lamp. A thermocouple at the center of the tube just downstream from the detection region registered 31°C. This temperature was used to calculate the bromoform and chlorine cross sections [DeMore *et al.*, 1997]. The bromoform (Aldrich Chemical 99 + %) was analyzed by gas chromatography; no CBr_4 was evident and only a trace of CH_2Br_2 ($< 0.1\%$) could be detected. The chlorine/vinyl bromide mixture was made

from neat vinyl bromide (Aldrich Chemical 98%) and a 1% Cl_2 in helium mixture (Matheson) and was stored in a 5 liter bulb in the dark.

3. Experimental Results

[11] Two methods were used to measure the quantum yields of Br atoms from CHBr_3 . Method I compared the amplitude of the Br signal when using the doubled radiation from the dye laser (303 to 324 nm) to that when using 266 nm radiation directly from the YAG laser. This method had the advantage of being able to switch the wavelength easily without disturbing the gas flow; the bromoform concentration was relatively stable throughout the day, affected only by a slight drift to higher values as the bath surrounding the bubbler warmed slightly. Since the identical gas mixture was used during both measurements, no correction for absorption of the bromine resonance lines by the CHBr_3 within the photolysis cell was needed.

[12] In method I, the bromine atom resonance fluorescence signals as a function of time, t , were fit to the equation,

$$\text{Signal} = S_0(\exp(-K_1 t)) + S_1(\exp(-K_2 t) - \exp(-K_1 t)) + Bk \quad (1)$$

This double exponential decay was necessary for runs at all pressures. The decay constants K_1 and K_2 were inversely proportional to total pressure, indicating diffusion controlled losses. The faster loss process ($K_1 = 250 \text{ s}^{-1}$ at 2 Torr) was interpreted as diffusion of bromine atoms out of the most sensitive detection region, and the slower process ($K_2 = 40 \text{ s}^{-1}$ at 2 Torr) was probably loss of bromine atoms on the walls of the flow tube. S_0 is the amplitude of the bromine atom signal at $t = 0$, and this was used as the measure of the number of Br atoms formed by the laser. S_1 was typically 45% as large as S_0 . The term Bk represents the background count, which contained contributions from scattered light, as well as a time-independent signal from the presence of bromoform within the flow tube. Trace A in Figure 1 shows a typical signal observed using method I together with the least squares fit to equation (1).

[13] For experiments at 320 and 324 nm, the Br signals were quite weak. For these experiments, values for K_1 , K_2 , and the ratio S_1/S_0 from 266 nm runs, which had strong signals, were inserted into equation (1), and then the least squares fits were used to determine S_0 and Bk . Normally values for K_1 , K_2 and S_1/S_0 were constant to within a few percent on any given day, as long as the flow conditions remained constant.

[14] After normalizing the signal amplitudes, S_0 , for slight differences in bromoform concentration (as measured by absorbance in the Cary), pressure within the photolysis cell, and laser pulse energy multiplied by wavelength (to convert energy to number of photons), the ratio of normalized signals gave the ratio of products of cross section (σ) times quantum yield (ϕ) at the two wavelengths:

$$\frac{S_0(\lambda)}{S_0(266)} = \frac{\sigma_\lambda \phi_\lambda}{\sigma_{266} \phi_{266}}$$

Literature values for the absorption cross sections of bromoform were then used to calculate the ratio of quantum


Table 1. Experimental Results Using Method I^a

λ , nm	Carrier	σ_λ	ϕ_λ/ϕ_{266}	95% C.L.	N	ϕ_λ	Estimated Error
303.0	N_2	0.501	1.323	0.65%	5	1.009	6.3%
306.0	N_2	0.329	1.216	2.7%	6	0.927	6.9%
306.0	air	0.329	1.183	3.7%	6	0.902	7.3%
310.0	N_2	0.188	1.181	1.0%	5	0.901	6.4%
314.0	air	0.1073	1.188	5.1%	6	0.906	8.2%
314.0	N_2	0.1073	1.158	10.0%	4	0.883	11.9%
320.0	air	0.0463	1.060	12.6%	6	0.809	14.2%
320.0	N_2	0.0463	1.086	8.7%	6	0.828	10.9%
324.4	N_2	0.0250	1.000	33%	3	0.762	34%

^aThe cross sections σ_λ are calculated for bromoform at 31°C; units are 10^{-20} cm^2 . N is the number of measurements that have been averaged for each condition. Values of ϕ_λ are calculated from the ratios ϕ_λ/ϕ_{266} by using $\phi_{266} = 0.763$ from method II (see Table 2).

yields at the two wavelengths [Gillotay *et al.*, 1989; Moortgat *et al.*, 1993]. The results obtained using method I are collected in Table 1.

[15] Method II compared the bromine atom signals for two different gas mixtures at the same photolysis wavelength. One gas mixture was the same CHBr_3/N_2 mixture that was used in method I. The other gas mixture contained molecular chlorine and vinyl bromide diluted with nitrogen. The laser pulse dissociated the molecular chlorine and the chlorine atoms were rapidly converted to bromine atoms by reaction with vinyl bromide [Park *et al.*, 1983]:



There was no evidence of photolysis of the vinyl bromide; no time-dependent Br signals appeared unless both Cl_2 and CH_2CHBr were present.

[16] The time dependence of the resonance fluorescence signals for the $\text{Cl}_2/\text{C}_2\text{H}_3\text{Br}$ mixtures was different from that observed with bromoform. As shown in trace B in Figure 1, the signal increases rapidly after the flash, as Cl atoms are converted into Br atoms, and then decays with the same double exponential decay observed for bromoform. For these mixtures, the signals could be fit to an equation of the form,

$$\text{Signal} = S_0(\exp(-K_1 t) - \exp(-K_0 t)) + S_1(\exp(-K_2 t) - \exp(-K_1 t)) + Bk \quad (2)$$

where K_0 is the rise-time constant and K_1 and K_2 are again the two decay constants.

[17] The mechanism used to model this time dependence was as follows. The chlorine atoms are formed by the laser pulse with a concentration $[\text{Cl}]_0$. Chlorine atoms are removed by reaction with vinyl bromide, with a rate constant k_1 , or they can diffuse out of the viewing region with a decay constant k_2 ; for simplicity, only a single diffusional decay will be used here. The bromine atoms are removed only by diffusion, with a rate constant k_3 . Solving the resulting differential equations, with the boundary condition that $[\text{Br}] = 0$ at $t = 0$, gives the following expression for the bromine atom concentration,

$$[\text{Br}] = [\text{Cl}]_0 \{ k_1 [\text{VB}] / (k_1 [\text{VB}] + k_2 - k_3) \} \cdot \{ \exp(-k_3 t) - \exp(-(k_1 [\text{VB}] + k_2)t) \}$$




Figure 2. The rise time constant, K_0 , as a function of the vinyl bromide concentration. The point marked B is from trace B in Figure 1. Except for the varying $[C_2H_3Br]$, the conditions are similar to those described for trace B.

where [VB] is the vinyl bromide concentration and the rate constants are defined above. Comparing this expression to the first term in the empirical fitted equation (2), it can be seen that the time constants are,

$$K_0 = k_1[VB] + k_2$$

$$K_1 = k_3$$

and the fitted amplitude S_0 can be written as,

$$S_0 = [Cl]_0 \{ (K_0 - k_2) / (K_0 - k_3) \}$$

[18] As required by the above model, the rise time constant K_0 was observed to depend linearly on the vinyl bromide concentration, as shown in Figure 2. The slope of the least squares line in Figure 2 gives a value for k_1 , $1.36 \times 10^{-10} \text{ cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$, that agrees very well with the only previous measurement (1.43×10^{-10}) by *Park et al.* [1983]. The intercept in Figure 2 gives a value for k_2 of 295 s^{-1} .

[19] Several corrections were needed when calculating quantum yields by method II. The most important correction was an allowance for the attenuation of the Br resonance radiation during its approximately 5 cm path through the two different gas mixtures. In separate experiments, measurements of the amplitude of the normalized signal as a function of $CHBr_3$ or Cl_2/VB mix concentrations within the flow cell resulted in reasonable Beer's Law plots. The slopes of these plots were used to calculate cross sections of $\sim 1.1 \times 10^{-16} \text{ cm}^2 \text{ molecule}^{-1}$ for $CHBr_3$ and $\sim 2.1 \times 10^{-16}$ for the Cl_2/VB mixture. Since the bromine resonance

lines occur at several wavelengths, these are only effective cross sections, but they are consistent with what is known about the absorption spectra of $CHBr_3$ and Cl_2 near 160 nm [*Henrici*, 1932; *Lee and Walsh*, 1959]. These Beer's Law slopes were then used to correct the measured S_0 's in the quantum yield measurements; these corrections were +2 to 3% for $CHBr_3$ and +11 to 12% for Cl_2/VB .

[20] A second adjustment was needed for the Cl_2/VB experiments to correct the amplitude of the signal so that it was proportional to the original chlorine atom concentration. This involved multiplying S_0 for each run by the factor $(K_0 - k_3)/(K_0 - k_2)$, where k_2 is the intercept in Figure 2 and k_3 , the diffusional loss rate constant for Br atoms, was taken to be the fitted value of K_1 in equation (2). Since values of K_0 were 1000 to 4000 s^{-1} and the fitted k_3 ($\sim 240 \text{ s}^{-1}$) were only slightly smaller than k_2 (295 s^{-1}), these corrections were in the range of 2 to 8%.

[21] With the amplitudes normalized for slight drifts in concentration, pressure and laser intensity as in method I, the ratio of corrected S_0 were taken to be the ratio of the product of the quantum yield and cross section for the two absorbers, $CHBr_3$ and Cl_2 . The quantum yield of Cl_2 is assumed to be 2.0 and so the quantum yield of $CHBr_3$ could be calculated. The results of method II are summarized in Table 2. The $CHBr_3$ quantum yields from both methods I and II are plotted as a function of wavelength in Figure 3.

[22] When N duplicate runs were averaged, the sample standard deviation was used with Student's t distribution and $(N-1)$ degrees of freedom to set 95% confidence limits on the reported mean values. These confidence limits, based only on the scatter of the data, are given in Tables 1 and 2. There were additional errors that could only be estimated: using the dye laser, there was an estimated wavelength uncertainty of 0.3 nm, which resulted in an uncertainty of about 4% in the cross sections; a temperature uncertainty of 2 K gave uncertainties in the cross sections of 1.5 to 2.5%; and for method I, the estimated uncertainty of 4.4% in the quantum yield of $CHBr_3$ at 266 nm contributed to the uncertainty reported at other wavelengths. These estimated uncertainties were combined with the 95% confidence limits to give the total uncertainties listed in the last columns of Tables 1 and 2.

4. Discussion

[23] It is not surprising that the quantum yield of bromine atoms in Figure 3 is unity or close to unity in the 300 nm region. The bromoform spectrum is an absorption continuum with a maximum near 220 nm and a long tail extending out to at least 360 nm [*Moortgat et al.*, 1993]. Such extended continua are common features of halogenated alkanes and they are usually interpreted as transitions to a

Table 2. Experimental Results Using Method II^a

λ , nm	σ_{CHBr_3}	σ_{Cl_2}	$\varphi_{CHBr_3}/\varphi_{Cl_2}$	95% C.L.	N	φ_{CHBr_3}	Estimated Error
266.0	48.3	0.496	0.381	2.8%	3	0.763	4.4%
303.0	0.501	13.9	0.470	3.6%	5	0.939	6.9%
305.0	0.379	15.2	0.512	11.5%	7	1.025	12.9%
306.0	0.329	15.9	0.513	5.9%	3	1.026	8.0%

^aThe cross sections, σ , are calculated for 31°C; units are 10^{-20} cm^2 . The quantum yield of chlorine atoms from Cl_2 , φ_{Cl_2} , is assumed to be 2.




Figure 3. The quantum yields of bromine atoms from bromoform as a function of wavelength. Values for method I come from Table 1. Values for method II come from Table 2. Trace B of Figure 1 is one of the seven runs contributing to the point at 305 nm. The error limits come from the last columns of Tables 1 and 2.

repulsive state that correlates with the separated ground state halogen atom and an alkyl radical [Herzberg, 1966]. The smooth decrease in absorption cross section to longer wavelengths is interpreted as a decreasing Franck-Condon overlap between the ground vibrational state and the repulsive upper state potential energy curve [Gordus and Bernstein, 1954].

[24] However, there are other dissociation processes that are energetically accessible in this wavelength region. In addition to the expected single bond rupture, A, three other channels are energetically possible, B, C and D:

Channel A	$\text{CHBr}_3 + h\nu \longrightarrow \text{Br} + \text{CHBr}_2$	$\lambda < 435 \text{ nm}$
Channel B	$\text{CHBr}_3 + h\nu \longrightarrow \text{HBr} + \text{CBr}_2$	$\lambda < 410 \text{ nm}$
Channel C	$\text{CHBr}_3 + h\nu \longrightarrow \text{Br}_2 + \text{CHBr}$	$\lambda < 337 \text{ nm}$
Channel D	$\text{CHBr}_3 + h\nu \longrightarrow \text{H} + \text{CBr}_3$	$\lambda < 304 \text{ nm}$
Channel E	$\text{CHBr}_3 + h\nu \longrightarrow 2 \text{ Br} + \text{CHBr}$	$\lambda < 207 \text{ nm}$

The formation of two bromine atoms from the absorption of one photon, channel E, is clearly not possible for the current experiments. The above wavelength limits were calculated using standard heats of formation together with calculated values for CHBr and CBr₂ [Dixon et al., 2002; Xu et al., 2002].

[25] Both method II and method I (using the method II quantum yield at 266 nm) agree that the bromine atom quantum yields at 303 to 306 nm are unity within the experimental error. Thus only channel A can be active in

this region. This finding is supported by the theoretical calculations of Peterson and Francisco [2002], who find that the only electronically excited state of CHBr₃ accessible at wavelengths longer than 270 nm is an unstable triplet state that is dissociative along the C-Br coordinate.

[26] Is the apparent decrease in quantum yield at wavelengths longer than 300 nm (Figure 3) the result of channels B and C becoming important? Mechanistically this seems unlikely. With a repulsive upper state, the longer the wavelength the longer is the carbon-bromine bond in the newly formed electronically excited molecule. This greater distance would make it harder for the departing Br atom to capture an H or another Br, channels B or C. Fluorescence of the electronically excited CHBr₃ would also decrease the bromine atom quantum yield, but this is a very inefficient process for a repulsive state.

[27] It is more likely that the bromine atom quantum yield remains at unity, and the apparent decrease at longer wavelengths observed in Table 1 and Figure 3 is due to random or systematic errors that become more prominent as the cross sections decrease. Certainly at 324 nm, the random errors dominate; for an integration time three times longer than that used at shorter wavelengths, the amplitude of the Br signal was only 350 counts/bin, while the background count was 35,000 counts/bin. The longer counting times also increased the error due to lamp drift. In addition, the quantum yields are fully dependent on the bromoform cross sections reported by Moortgat et al. [1993]. The cross sections at the longer wavelengths would have to be too large by 10 to 20% to return the long wavelength experimental points in Figure 3 to unity. However, Moortgat et al. [1993] claim that their cross section errors are only about $\pm 3\%$ in this region; this 3% uncertainty has not been included in the error bars shown in Figure 3.

[28] More surprising is the bromine atom quantum yield of 0.76 at 266 nm. This suggests that at least one of the channels B, C or D becomes important at this shorter wavelength. The molecular beam time-of-flight studies of Xu et al. [2002] report evidence that both channels A and C are important in the 200 nm region. They estimate that the Br₂ yield, resulting from channel C, is 0.16 at 267 nm and 0.26 at 234 nm. At these higher energies there is more chance of mixing with other electronic states having different properties and so additional dissociation processes occur. We believe that the more quantitative determination of the bromine atom quantum yield of 0.76 at 266 nm is to be preferred over the estimate of 0.84 of Xu et al. [2002].

[29] For modeling the bromine budget in the troposphere, where the radiation is mostly at wavelengths longer than 300 nm, it is recommended that a quantum yield of unity be used for channel A. In the atmosphere, the dibromomethyl radical that is also formed in channel A should rapidly form a peroxy radical, CHBr₂O₂. The fate of this peroxy radical has been investigated by Orlando et al. [1996].

[30] In the stratosphere, as more radiation shorter than 300 nm becomes available, the photochemistry of bromoform will become more complicated. However, measurements of CHBr₃ in the stratosphere [Sturges et al., 2000] show that its concentrations plunge rapidly with height above the tropopause, so that the importance of bromoform as a direct source of bromine atoms in the stratosphere is probably minor.

[31] **Acknowledgments.** The authors would like to thank Joseph Francisco and William Jackson for prepublication copies of their manuscripts and David Natzic for expert assistance in constructing the flow tube. The research described in this paper was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under contract to the National Aeronautics and Space Administration.

References

- Berg, W. W., L. E. Heidt, W. Pollock, P. D. Sperry, R. J. Cicerone, and E. S. Gladney, Brominated organic species in the Arctic atmosphere, *Geophys. Res. Lett.*, *11*, 429–432, 1984.
- Brune, W. H., J. G. Anderson, and K. R. Chan, In situ observations of BrO over Antarctica: ER-2 aircraft results from 54°S to 72°S latitude, *J. Geophys. Res.*, *94*, 16,639–16,647, 1989.
- Butler, J. H., and J. M. Rodriguez, Methyl bromide in the atmosphere, in *The Methyl Bromide Issue*, edited by C. H. Bell, N. Price, and B. Chakrabarti, pp. 28–90, John Wiley, New York, 1996.
- Carpenter, L. J., and P. S. Liss, On temperate sources of bromoform and other reactive organic bromine gases, *J. Geophys. Res.*, *105*, 20,539–20,547, 2000.
- Carpenter, L. J., W. T. Sturges, S. A. Penkett, and P. S. Liss, Short-lived alkyl iodides and bromides at Mace Head, Ireland: Links to biogenic sources and halogen oxide production, *J. Geophys. Res.*, *104*, 1679–1689, 1999.
- Cicerone, R. J., L. E. Heidt, and W. H. Pollock, Measurements of atmospheric methyl bromide and bromoform, *J. Geophys. Res.*, *93*, 3745–3749, 1988.
- DeMore, W. B., S. P. Sander, D. M. Golden, R. F. Hampson, M. J. Kurylo, C. J. Howard, A. R. Ravishankara, C. E. Kolb, and M. J. Molina, Chemical kinetics and photochemical data for use in stratospheric modeling: Evaluation number 12, *JPL Publ.*, *97-4*, 170–202, 1997.
- Dixon, D. A., W. B. de Jong, K. A. Peterson, and J. S. Francisco, Heats of formation of CBr, CHBr and CBr₂ from ab initio quantum chemistry, *J. Phys. Chem. A*, *106*, 4725–4728, 2002.
- Dvortsov, V. L., M. A. Geller, S. Solomon, S. M. Schauffler, E. L. Atlas, and D. R. Blake, Rethinking reactive halogen budgets in the midlatitude lower stratosphere, *Geophys. Res. Lett.*, *26*, 1699–1702, 1999.
- Fabian, P., R. Borchers, and K. Kourtidis, Bromine-containing source gases during EASOE, *Geophys. Res. Lett.*, *21*, 1219–1222, 1994.
- Gillotay, D., A. Jenouvrier, B. Coquart, M. F. Merienne, and P. C. Simon, Ultraviolet absorption cross-sections of bromoform in the temperature range 295–240K, *Planet. Space Sci.*, *37*, 1127–1140, 1989.
- Gordus, A. A., and R. B. Bernstein, Isotope effect in continuous ultraviolet absorption spectra: Methyl bromide-d₃ and chloroform-d, *J. Chem. Phys.*, *22*, 790–795, 1954.
- Henrici, A., Die ultraviolettabsorption der halogenderivate des methans, *Z. Phys.*, *77*, 35–51, 1932.
- Herzberg, G., *Molecular Spectra and Molecular Structure III: Electronic Spectra and Electronic Structure of Polyatomic Molecules*, pp. 451–453, Van Nostrand Reinhold, New York, 1966.
- Johnson, R. O., G. P. Perram, and W. B. Roh, Spin-orbit relaxation kinetics of Br(4²P_{1/2}), *J. Chem. Phys.*, *104*, 7052–7058, 1996.
- Kurylo, M. J., et al., Short-lived ozone-related compounds, in *Scientific Assessment of Ozone Depletion: 1998*, edited by C. A. Ennis, chap. 2, pp. 2.21–2.36, World Meteorol. Organ., Geneva, 1999.
- Lary, D. J., Gas phase atmospheric bromine photochemistry, *J. Geophys. Res.*, *101*, 1505–1516, 1996.
- Lary, D. J., M. P. Chipperfield, R. Toumi, and T. Lenton, Heterogeneous atmospheric bromine chemistry, *J. Geophys. Res.*, *101*, 1489–1504, 1996.
- Lee, J., and A. D. Walsh, The vacuum ultra-violet absorption spectra of the halogen molecules, part I, Chlorine, *Trans. Faraday Soc.*, *55*, 1281–1292, 1959.
- McElroy, M. B., R. J. Salawitch, S. C. Wofsy, and J. A. Logan, Reductions of antarctic ozone due to synergistic interactions of chlorine and bromine, *Nature*, *321*, 759–762, 1986.
- McKinney, K. A., J. M. Pierson, and D. W. Toohey, A wintertime in situ profile of BrO between 17 and 27 km in the Arctic vortex, *Geophys. Res. Lett.*, *24*, 853–856, 1997.
- Moore, R., R. Tokarczyk, and C. Geen, Sources of organobromines to the Arctic atmosphere, in *The Tropospheric Chemistry of Ozone in the Polar Regions*, edited by H. Niki and K. H. Becker, pp. 235–250, Springer-Verlag, New York, 1993.
- Moortgat, G. K., R. Meller, and W. Schneider, Temperature dependence (256–296K) of the absorption cross-sections of bromoform in the wavelength range 285–360 nm, in *The Tropospheric Chemistry of Ozone in the Polar Regions*, edited by H. Niki and K. H. Becker, pp. 359–369, Springer-Verlag, New York, 1993.
- Nielsen, J. E., and A. R. Douglass, A simulation of bromoform's contribution to stratospheric bromine, *J. Geophys. Res.*, *106*, 8089–8100, 2001.
- Orlando, J. J., G. S. Tyndall, T. J. Wallington, and M. Dill, Atmospheric chemistry of CH₂Br₂: Rate coefficients for its reaction with Cl atoms and OH and the chemistry of the CHBr₂O radical, *Int. J. Chem. Kinet.*, *28*, 433–442, 1996.
- Park, J.-Y., I. R. Slagle, and D. Gutman, Kinetics of the reaction of chlorine atoms with vinyl bromide and its use for measuring chlorine-atom concentrations, *J. Phys. Chem.*, *87*, 1812–1818, 1983.
- Peterson, K. A., and J. S. Francisco, Should bromoform absorb at wavelengths longer than 300 nm?, *J. Chem. Phys.*, *117*, 6103–6107, 2002.
- Pfeilsticker, K., W. T. Sturges, H. Bosch, C. Camy-Peyret, M. P. Chipperfield, A. Engel, R. Fitzenberger, M. Muller, S. Payan, and B. M. Sinnhuber, Lower stratospheric organic and inorganic bromine budget for the arctic winter 1998/99, *Geophys. Res. Lett.*, *27*, 3305–3308, 2000.
- Singh, H. G., Halogens in the atmospheric environment, in *Composition, Chemistry, and Climate of the Atmosphere*, edited by H. B. Singh, pp. 216–250, Van Nostrand Reinhold, New York, 1995.
- Sturges, W. T., D. E. Oram, L. J. Carpenter, and S. A. Penkett, Bromoform as a source of stratospheric bromine, *Geophys. Res. Lett.*, *27*, 281–284, 2000.
- Toohey, D. W., W. H. Brune, and J. G. Anderson, Mechanism and kinetics of Br + HO₂ → HBr + H₂O₂ → products over the temperature range 260–390 K, *J. Phys. Chem.*, *91*, 1215–1222, 1987.
- Wamsley, P. R., et al., Distribution of halon-1211 in the upper troposphere and lower stratosphere and the 1994 total bromine budget, *J. Geophys. Res.*, *104*, 1513–1526, 1998.
- Wofsy, S. C., M. B. McElroy, and Y. L. Yung, The chemistry of atmospheric bromine, *Geophys. Res. Lett.*, *2*, 215–219, 1975.
- Xu, D., J. S. Francisco, J. Huang, and W. M. Jackson, Ultraviolet photodissociation of bromoform at 234 nm and 267 nm by means of ion velocity imaging, *J. Chem. Phys.*, *117*, 2578–2585, 2002.
- Yung, Y. L., J. P. Pinto, R. T. Watson, and S. P. Sander, Atmospheric bromine and ozone perturbations in the lower stratosphere, *J. Atmos. Sci.*, *37*, 339–353, 1980.

K. D. Bayes, R. R. Friedl, and S. P. Sander, Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109, USA. (bayes@chem.ucla.edu)

D. W. Toohey, Program in Atmospheric and Oceanic Sciences, University of Colorado, Boulder, CO 80309, USA.