

## RESEARCH ARTICLE

# The polarity protein VANG-1 antagonizes Wnt signaling by facilitating Frizzled endocytosis

Chun-Wei He<sup>1</sup>, Chien-Po Liao<sup>1,§</sup>, Chung-Kuan Chen<sup>1,§,\*</sup>, Jérôme Teulière<sup>2</sup>, Chun-Hao Chen<sup>1,‡</sup> and Chun-Liang Pan<sup>1,¶</sup>

## ABSTRACT

Signaling that instructs the migration of neurons needs to be tightly regulated to ensure precise positioning of neurons and subsequent wiring of the neuronal circuits. Wnt-Frizzled signaling controls neuronal migration in metazoans, in addition to many other aspects of neural development. We show that *Caenorhabditis elegans* VANG-1, a membrane protein that acts in the planar cell polarity (PCP) pathway, antagonizes Wnt signaling by facilitating endocytosis of the Frizzled receptors. Mutations of *vang-1* suppress migration defects of multiple classes of neurons in the Frizzled mutants, and overexpression of *vang-1* causes neuronal migration defects similar to those of the Frizzled mutants. Our genetic experiments suggest that VANG-1 facilitates Frizzled endocytosis through  $\beta$ -arrestin2. Co-immunoprecipitation experiments indicate that Frizzled proteins and VANG-1 form a complex, and this physical interaction requires the Frizzled cysteine-rich domain. Our work reveals a novel mechanism mediated by the PCP protein VANG-1 that downregulates Wnt signaling through Frizzled endocytosis.

**KEY WORDS:** *C. elegans*, Wnt signaling, Planar polarity, Neuronal migration, Endocytosis, Frizzled

## INTRODUCTION

Wiring of the nervous system during development is established first by neuronal migration, followed by navigation of axon growth cones towards their innervation targets and subsequent axon branching and synapse formation. The migration of neurons and axon growth cones is regulated by highly dynamic signals in the environment that function as attractive or repulsive cues (Tessier-Lavigne and Goodman, 1996). Over the past three decades, several classes of conserved neuronal guidance cues have been uncovered, including Netrin, Slit, semaphorins, ephrins, Sonic Hedgehog and Wnt glycoproteins (Tessier-Lavigne and Goodman, 1996; Blockus and Chédotal, 2016; Dudanova and Klein, 2013; Jongbloets and Pasterkamp, 2014; Sun et al., 2011; Yam and Charron, 2013). Although substantial efforts have been dedicated to deciphering the

signaling pathways through which these factors instruct neuronal and growth cone migration, little is known about the regulatory mechanisms that downregulate them to prevent prolonged, aberrant signaling that could lead to guidance errors.

The availability of receptors on the plasma membrane is a well-established regulatory mechanism that modulates the magnitude and duration of signaling. In *Drosophila* embryos, trafficking of the Robo receptor to the growth cone membrane of pre-crossing commissural axons is prevented by the Golgi protein Commissureless, which traps Robo in the Golgi (Keleman et al., 2002; Tear et al., 1996; Kidd et al., 1998). In the *commissureless* mutant, constitutive Robo availability on the axonal membrane inadvertently transmits repulsive signaling by binding the Slit ligand, preventing the commissural axons from crossing the midline and hence causing the ‘commissureless’ axonal phenotype (Keleman et al., 2002; Kidd et al., 1998; Tear et al., 1996). Recent studies suggest that internalization of Frizzled proteins, seven-pass transmembrane proteins that function as Wnt receptors, modulates Wnt signaling and regulates axon guidance in *C. elegans* and rodents. Depending on the cellular contexts, Frizzled internalization may activate or terminate Wnt signaling. For example, internalization of Frizzled proteins triggered by PLR-1 E3 ligase-mediated ubiquitylation downregulates Wnt signaling to ensure proper axon development in *C. elegans* (Moffat et al., 2014), consistent with the vertebrate PLR-1 homologs ZNRF3 and RNF43 downregulating Frizzled proteins and inhibiting Wnt signaling in the mouse intestinal epithelium (Koo et al., 2012). Frizzled internalization could also activate Wnt signaling. In the mouse spinal cord, Vangl2, a four-pass transmembrane protein of the planar cell polarity (PCP) pathway, facilitates Frizzled endocytosis by antagonizing hyperphosphorylation of Frizzled proteins mediated by Dvl1/Dishevelled. In this context, Frizzled internalization activates Wnt signaling and promotes proper axon guidance (Shafer et al., 2011). Consistent with this, we recently showed that *C. elegans* VANG-1 promotes Wnt signaling by facilitating Frizzled internalization to instruct neurite branching patterns in the PLM mechanosensory neuron (Chen et al., 2017).

The aforementioned findings reveal an important, cell-autonomous function of Vangl2 in promoting Wnt signaling, in contrast to its complex non-autonomous requirement in *Drosophila* embryonic epithelia that involves interaction with Frizzled proteins in the neighboring cells (Bailly et al., 2017; Devenport, 2014). To explore how Vangl2 tunes Wnt signaling to regulate neuronal development, and whether Vangl2-dependent Frizzled internalization occurs in broader, more diverse cellular contexts, we systemically analyze the functions of *C. elegans vang-1* gene in multiple classes of migrating *C. elegans* neurons. In contrast to the regulation of Frizzled proteins in the non-migrating PLM neuron, we find that VANG-1 antagonizes Wnt signaling by facilitating Frizzled endocytosis in all these migrating neuronal classes. These

<sup>1</sup>Institute of Molecular Medicine, College of Medicine, National Taiwan University, Taipei 10002, Taiwan. <sup>2</sup>Department of Molecular Cell Biology, University of California, Berkeley, CA 94720-3204, USA.

\*Present address: Program in Biomedical Sciences, Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tsukuba, Ibaraki 305-8575, Japan. †Present address: Division of Biology and Bioengineering, California Institute of Technology, Pasadena, CA 91125, USA.

§These authors contributed equally to this work

¶Author for correspondence (chunliangpan@ntu.edu.tw)

© C.-K.C., 0000-0002-6973-2563; J.T., 0000-0002-0965-4399; C.-L.P., 0000-0003-0108-3138

observations expand the phenotypic repertoire through which VANG-1/Vangl2 modulates Wnt signaling to pattern neuronal migration precisely.

## RESULTS

### VANG-1 antagonizes Wnt signaling in QL descendant migration in *C. elegans*

The *C. elegans* Q neuroblasts are born in early first larval (L1) stage at equivalent positions on the right (hereafter referred to as QR) and the left side (hereafter referred to as QL) of the worm body. The posterior migration of QL descendants (QL.d) during early L1 requires the Abdominal-A family Hox gene *mab-5*, expression of which in the QL lineage is controlled by a  $\beta$ -catenin-dependent Wnt signaling pathway (Fig. 1A,B) (Clark et al., 1993; Wang et al., 1993; Harris et al., 1996; Whangbo and Kenyon, 1999; Korswagen, 2002). Genes in this pathway that are essential for *mab-5* activation include *egl-20/Wnt*, *mig-1/Frizzled*, *lin-17/Frizzled*, *mig-5/Dishevelled*, *bar-1/ $\beta$ -catenin*, and the TCF transcription factor *pop-1* (Harris et al., 1996; Eisenmann and Kim, 2000; Korswagen et al., 2000). Loss-of-function mutations of these genes diminish *mab-5* expression and result in reversed, anterior migration of the QL.d, including the PVM (QL.paa) (Fig. 1C,D).

Interestingly, we found that the *tm1422* deletion mutation of *vang-1*, the *C. elegans* homolog of the PCP gene *Van Gogh/Strabismus/Vangl2*, suppressed PVM migration defects of the *mig-1* mutant, but it did not affect PVM migration in the otherwise wild-type background (Fig. 1C,D). Similar suppression of PVM defects was seen with another *vang-1* deletion allele, *ok1142*, and also with *vang-1* feeding RNAi (Fig. S1A). In contrast to its significant suppression of the *mig-1* phenotypes, the *vang-1(tm1422)* mutation failed to suppress PVM migration defects in the *egl-20*, *mig-5* and *bar-1* mutants (Fig. 1D). *egl-20*, *mig-5* and *bar-1* each represent the single homolog for *Wnt*, *Dishevelled* and  $\beta$ -*catenin*, respectively, and are required for the posterior migration of QL.d. We hypothesize that *vang-1* antagonizes Wnt signaling specifically at the level of Frizzled receptors. Therefore, PVM defects caused by the absence of Wnt ligand (*egl-20*) or signals downstream of Frizzled receptors (*mig-5* and *bar-1*) are refractory to the effects of *vang-1* mutations. By contrast, LIN-17, another Frizzled protein, is also required to promote QL.d migration (Harris et al., 1996; Zinovyeva et al., 2008), and we speculate that restoration of PVM migration in the *mig-1*; *vang-1* double mutant could be mediated by Wnt signaling through the remaining LIN-17 receptor. This hypothesis predicts that the *vang-1* mutation is not able to suppress QL.d defects in the *mig-1 lin-17* double mutant, which lacks all relevant Frizzled receptors for QL.d migration. Consistent with this hypothesis, the *vang-1* mutation improved PVM defects of the *lin-17* single mutant but failed to do so in the *mig-1 lin-17* double Frizzled mutant (Fig. 1D; Fig. S2).

Expressing *vang-1* in the Q lineages of the *mig-1*; *vang-1* mutant, using the *egl-17* promoter, completely reversed the suppression of PVM mismigration conferred by the *vang-1* mutation (Fig. 1E). Furthermore, overexpressing *vang-1* in the QL.d resulted in PVM migration defects in the wild type and enhanced the PVM defects of the *mig-1* mutant (Fig. 1E). These results suggest that *vang-1* antagonizes Wnt signaling cell-autonomously in the QL.d neuroblasts. We confirmed this result by using somatic clustered regularly interspaced short palindromic repeats (CRISPR)-Cas9 gene editing (Fig. S1B) (Shen et al., 2014). In brief, we expressed Cas9 under the *egl-17* promoter together with ubiquitously expressed *vang-1* sgRNAs. Mutations of *vang-1* in the Q lineages significantly suppressed PVM defects of the *mig-1* mutant, confirming that *vang-1* acts in the Q lineage (Fig. S1B).


On the basis of the hypothesis that *vang-1* antagonizes *mig-1* functions, we predicted that the expression of *mab-5*, the transcriptional target of Wnt signaling in QL.d migration, would be restored in the *mig-1*; *vang-1* double mutant. This is indeed the case: *mab-5* expression in the QL.d was significantly increased in *mig-1*; *vang-1* compared with that in the *mig-1* mutant, which was barely detectable (Fig. 1F,G). By contrast, the *vang-1* mutation failed to restore *mab-5* expression in the *egl-20* mutant (Fig. 1F,G). Of note, the *vang-1* mutant showed significantly higher level of *mab-5* expression compared with the wild type, consistent with *vang-1* being a negative regulator of Wnt signaling (Fig. 1F,G). Taken together, we conclude that VANG-1 antagonizes Wnt signaling cell-autonomously at the level of Frizzled receptors during QL.d migration.

We hypothesize that the antagonistic function of *vang-1* in Wnt signaling is broadly conserved in early *C. elegans* neuronal development. Below we test this idea in three additional neuronal contexts: QR descendant (QR.d) migration, hermaphrodite-specific neuron (HSN) migration, and ALM touch neuron polarization.

### VANG-1 antagonizes Wnt signaling in QR descendant migration in *C. elegans*

In contrast to QL.d, QR.d migrate to the anterior, and these anterior migration events depend on Wnt signaling genes that are largely distinct from those involved in QL.d migration (Harris et al., 1996; Forrester et al., 2004; Zinovyeva et al., 2008). Specifically, in addition to *egl-20*, *cwn-1/Wnt*, *mom-5/Frizzled* and *dsh-2/Dishevelled* promote QR.d anterior migration, whereas *mig-1*, *mig-5* and *bar-1* are not required (Maloof et al., 1999; Zinovyeva et al., 2008). Additional Wnt receptors, such as LIN-17 and CFZ-2, are hypothesized to function with MOM-5 in QR.d migration, as QR.d migration defects in the *mom-5* mutant are less severe than those of the *cwn-1*; *egl-20* double mutant (Forrester et al., 2004; Zinovyeva et al., 2008). Consistent with previous studies, we found that AVM, one of the QR.d, terminated their anterior migration prematurely in the *cwn-1*, *egl-20* and *mom-5* mutants (Fig. 2A,B). The *vang-1* mutation ameliorated AVM migration defects in the *egl-20*, *cwn-1* and *mom-5* mutants, and caused the AVM to be more anterior in an otherwise wild-type background (Fig. 2A,B). Improvement of QR.d migration by the *vang-1* mutation likely resulted from Wnt signaling mediated by remaining Wnt ligands or Frizzled receptors in respective single mutants, similar to the suppression of QL.d defects in the *mig-1* mutant (see above). By contrast, the *vang-1* mutation failed to suppress AVM defects caused by the mutation of *mig-14*, the *C. elegans* homolog of the Wnt secretory factor *Wntless* (Fig. 2A,B) (Bänziger et al., 2006; Bartscherer et al., 2006; Goodman et al., 2006). In the *mig-14* mutant, Wnt secretion in general is profoundly depleted and very little extracellular Wnt protein is available for signaling. These results are consistent with the model that phenotypic improvement conferred by the *vang-1* mutation depends on the Wnt ligands. Similar to its effects on QL.d, overexpression of VANG-1 in the QR.d caused modest but significant AVM migration defects (Fig. 2A,B). This suggests that *vang-1* antagonizes Wnt signaling in QR.d migration cell-autonomously.

It was previously shown that *mom-5* mutations compromise anterior migration of the QR.p neuroblast, whereas *vang-1* mutations move QR.pap (future AVM) further anterior (Mentink et al., 2014). To test whether *vang-1* also plays a role during QR.p migration, we examined the positions of QR.p at a defined developmental time points during its anterior migration in L1 larvae. The V5R seam cell divides after the QR.a and QR.P start




**Fig. 1. *vang-1* antagonizes Wnt-Frizzled signaling in the migration of QL.d in *C. elegans*.** (A) Schematic of the Wnt signaling pathway that promotes posterior migration of QL.d. (B) Schematic of the Q neuroblast lineages. (C) Representative epifluorescent images of PVM neurons in the wild type and the mutants. (D,E) Quantification of PVM positions along the anterior-posterior axis of the worm body. Each dot represents a single PVM neuron, with mean±s.e.m. indicated. Number of neurons scored is indicated in parentheses. \**P*<0.05, \*\**P*<0.01, \*\*\**P*<0.001 (one-way ANOVA followed by Bonferroni's multiple comparison test); #*P*<0.01 (one-way ANOVA followed by Bonferroni's multiple comparison test; see Fig. S2). (F) Epifluorescent images of *Pmab-5::GFP* expression in QL.d of the wild type and the mutants. Dashed lines indicate QL.d in the GFP channel. (G) Quantification of *Pmab-5::GFP* expression. Number of QL.d scored is indicated in parentheses. \*\**P*<0.01 (one-way ANOVA followed by Bonferroni's multiple comparison test). n.s., not significant. Scale bars: 100 μm (C); 10 μm (F).

their anterior migration. We scored QR.p positions in L1 animals in which the V5R seam cell has divided but the two V5R daughter cells remain in contact (Fig. 2C). This experiment confirmed that the anterior migration of QR.p was compromised in the *mom-5* mutant, and the *vang-1* mutation significantly suppressed the QR.p defects (Fig. 2C,D). These data indicate that *vang-1* antagonizes Wnt signaling during QR.p neuroblast migration.

**VANG-1 antagonizes Wnt signaling in the early development of multiple classes of *C. elegans* neurons**

Because QL and QR neuroblasts share similar developmental history and functional property, we investigated whether *vang-1* antagonizes Wnt signaling in a completely unrelated neuronal type. The HSNs are bilaterally symmetric serotonergic motor neurons that control egg-laying muscles (Desai et al., 1988). HSNs migrate from



**Fig. 2. *vang-1* antagonizes Wnt-Frizzled signaling in the migration of QR.d in *C. elegans*.** (A) Representative epifluorescent images of AVM neurons in the wild type and the mutants. The touch neurons are marked by *zdl5(Pmec-4::GFP)*. (B) Quantification of AVM positions along the anterior-posterior axis of the worm body. Each dot represents a single AVM neuron, with mean $\pm$ s.e.m. indicated. Number of neurons scored is indicated in parentheses. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  (one-way ANOVA followed by Bonferroni's multiple comparison test). n.s., not significant. (C) Epifluorescent/DIC (top) or epifluorescent images of QR.d in L1 larvae. The Q lineages are labeled by *rdvls1(Pegl-17::Myri::mCherry)*. (D) Distribution of QR.p in L1 at the time of V5R division. \*\*\* $P < 0.001$  (Fisher exact test). Scale bars: 100  $\mu$ m (A); 10  $\mu$ m (C).


the posterior towards the future vulval primordium during embryogenesis under the control of multiple Wnt and Frizzled genes (Fig. 3A) (Garriga et al., 1993; Pan et al., 2006). Various Wnt pathway mutants, including *egl-20*, *mig-14*, *mig-1*, *cwn-1*; *egl-20* or *mig-1 mom-5* double mutants, showed migration defects of HSNs, which fall short of their wild-type positions near the vulval primordium (Fig. 3B-D). Similar to its effects on AVM migration, the *vang-1* mutation suppressed HSN migration defects in the *egl-20*, *mig-1* or *cwn-1*; *egl-20* double mutants (Fig. 3B-D). The suppression effect of the *vang-1* mutation was not seen in the *mig-1 mom-5* double mutant or when secreted Wnts were largely eliminated in the *mig-14* mutant (Fig. 3C,D). Restoration of *vang-1* in the HSNs, using the *unc-86* promoter, completely reversed the suppression of HSN migration defects in the *egl-20; vang-1* double mutant, making the HSN defects similar to those of the *egl-20* mutant (Fig. 3D). This rescue effect was abolished by *vang-1* RNAi, suggesting that the reversal of suppression is the action of functional *vang-1* transcripts rather than impairment of HSN migration by use of the *unc-86* promoter (Fig. 3E). These results confirm that *vang-1* antagonizes Wnt signaling cell-autonomously in HSN migration.

Could *vang-1* antagonize Wnt signaling in developmental contexts other than neuronal migration? To explore this

possibility, we examined the ALM, a unipolar touch neuron morphology of which is regulated by Wnts (Hilliard and Bargmann, 2006; Prasad and Clark, 2006; Pan et al., 2008; Zheng et al., 2015). The bilaterally symmetric ALM neurons extend a long neurite to the anterior. In the *mig-14* and *cwn-1*; *egl-20* double mutants, many ALM neurons grew a single posterior neurite that could reach as far as the lumbar ganglia, and some ALMs showed bipolar morphology with both anterior and posterior neurites (Fig. S3A,B). The *vang-1* mutation partially suppressed ALM defects in the *cwn-1*; *egl-20*, but not in the *mig-14* mutants (Fig. S3B). Moreover, *vang-1* overexpression in the touch neurons caused ALM defects similar to those seen in the *cwn-1*; *egl-20* mutants in a dose-dependent manner (Fig. S3C). Considering all data from the QL.d, QR.d, HSN and ALM experiments, we conclude that *vang-1* broadly antagonizes Wnt signaling in multiple events of early neuronal development in *C. elegans*.

#### ***vang-1* acts through *arr-1* to antagonize Wnt signaling in neuronal development**

Our genetic analyses of neuronal migration raise the possibility that *vang-1* negatively regulates Wnt signaling at the level of Frizzled receptors. A previous study of growth cone steering suggests that




**Fig. 3. *vang-1* antagonizes Wnt-Frizzled signaling in the migration of HSN in *C. elegans*.** (A) Schematic of HSN migration during embryogenesis and projection of its migratory path onto a late L4 larvae. (B) Representative epifluorescent images of HSN neurons in the wild type and the mutants. Asterisks indicate the position of the vulva. Scale bar: 100  $\mu$ m. (C-E) Quantification of HSN positions along the anterior-posterior axis of the posterior worm body. Each dot represents a single HSN neuron, with mean  $\pm$  s.e.m. indicated. Number of neurons scored is indicated in parentheses. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  (one-way ANOVA followed by Bonferroni's multiple comparison test). n.s., not significant.

Vangl2 facilitates Frizzled endocytosis by antagonizing Dvl1/Dishevelled-mediated Frizzled hyperphosphorylation (Shafer et al., 2011). We previously found that the localization of MIG-1 to RAB-5(+) early endosomes was decreased in the non-migratory *C. elegans* PLM neuron in the *vang-1* mutant, and that VANG-1 overexpression increased MIG-1 distribution to the early endosomes (Chen et al., 2017). On the basis of these observations, we hypothesize that VANG-1 antagonizes Wnt signaling by promoting Frizzled endocytosis, thereby terminating signal transduction induced by binding of Wnt ligands to the Frizzled receptors. This model makes three predictions. First, blocking endocytosis should suppress neuronal migration defects caused by a partial deficiency in Frizzled receptors. Second, suppression of neuronal defects conferred by the *vang-1* mutation is not further enhanced by blocking Frizzled endocytosis. Third, loss of VANG-1 impairs Frizzled endocytosis in the migrating neurons and results in higher Frizzled levels at the plasma membrane.

We tested the first prediction by examining animals that carried a deletion in the *arr-1* gene, which encodes the adaptor protein  $\beta$ -arrestin2 essential for Frizzled endocytosis in mammalian cells (Chen et al., 2003). We recently confirmed that *arr-1* also mediated

MIG-1 endocytosis in the non-migratory *C. elegans* PLM neuron (Chen et al., 2017). Although the *arr-1* mutation did not affect PVM migration in the wild type, similar to the *vang-1* mutation, it significantly suppressed PVM defects in the *mig-1* mutant (Fig. 4A). Likewise, defective AVM migration in the *mom-5* mutant and HSN undermigration of the *egl-20* mutant were also suppressed by the *arr-1* mutation (Fig. 4B,C). These results suggest that in the migrating *C. elegans* neurons, Frizzled endocytosis negatively regulates Wnt signaling. To test whether *vang-1* acts in the same genetic pathway as *arr-1*, we constructed multiple mutants that harbored both the *vang-1* and *arr-1* mutations, together with the *mig-1* or the *egl-20* mutation. PVM migration defects in the *mig-1; arr-1 vang-1* triple mutant were comparable to those of the *mig-1; vang-1* or the *mig-1; arr-1* mutants (Fig. 4A). This observation suggests that *arr-1* and *vang-1* act in a shared genetic pathway to temper Wnt-Frizzled signaling in QLd migration. Consistent with this, HSN undermigration was not further worsened in the *egl-20; arr-1 vang-1* triple mutant, compared with that in the *egl-20; arr-1* or *egl-20; vang-1* double mutant (Fig. 4C).

We further investigated the order of *vang-1* and *arr-1* function by genetic epistasis, taking advantage of the neuronal defects caused



**Fig. 4. *vang-1* acts with *arr-1* in the endocytic pathway to antagonize Wnt-Frizzled signaling in neuronal migration.** (A-C) Quantification of the positions of PVM (A), AVM (B) and HSN (C) in the wild type and mutant or transgenic strains. Each dot represents a single neuron, with mean  $\pm$  s.e.m. indicated. Number of neurons scored is indicated in parentheses. \* $P$ <0.05, \*\* $P$ <0.01, \*\*\* $P$ <0.001 (one-way ANOVA followed by Bonferroni's multiple comparison test). n.s., not significant.

by *vang-1* overexpression. PVM and AVM migration defects caused by *vang-1* overexpression were ameliorated in the *arr-1* mutant (Fig. 4A,B). Moreover, the *arr-1* mutation also significantly reduced the ALM polarity defects induced by excess *vang-1* activity (Fig. S3C). These results imply that *arr-1* acts downstream of *vang-1* to antagonize Wnt signaling.


### VANG-1 promotes endocytosis of the Frizzled receptors

Our previous work indicates that VANG-1 promotes the endocytosis of MIG-1 in the PLM touch neuron (Chen et al., 2017). Taken with results from the genetic experiments that VANG-1 attenuates Frizzled-dependent Wnt signaling and acts upstream of ARR-1, we hypothesized that VANG-1 facilitates endocytosis of the Frizzled receptors in the Q.d and other neuronal types. To test this, we investigated whether VANG-1 regulates Frizzled localization in the Q.d. For unknown reasons, fluorescence signals from our MIG-1::GFP transgene that fully rescued the PVM defects of the *mig-1* mutant, could not be detected in the migrating QL.d (data not shown). Therefore, we used the previously characterized *mom-5* transgene, *zuls145*(*Pnmy-2::MOM-5::GFP*) (Park et al., 2004), and observed MOM-5::GFP distribution with or without *vang-1* in QR.d. Because *zuls145* seemed to be integrated to the sex chromosome and failed to recombine with the sex chromosome-located *vang-1*(*tm1422*) allele, we used the CRISPR-Cas9 method to generate the *vang-1*(*twn3*) null mutation in the *zuls145* background (Fig. 5A; Fig. S1C). We first confirmed that the *vang-1*(*twn3*) mutation suppressed PVM migration defects of the *mig-1* mutant to levels comparable to other *vang-1* alleles (Fig. S1C). In the QR.d of the wild type, MOM-5::GFP

signals were distributed at the plasma membrane and intracellularly (Fig. 5B). By contrast, in the *vang-1*(*twn3*) mutant, intracellular MOM-5::GFP signals in QR.d were significantly reduced with reciprocal accumulation of MOM-5::GFP at the cell membrane (Fig. 5B,C). We further confirmed this result by performing *vang-1* RNAi, which resulted in the enrichment of MOM-5::GFP signals at the plasma membrane of QR.d (Fig. S4A,B). Moreover, overexpression of *vang-1* diminished membrane MOM-5::GFP, with reciprocal increase of MOM-5::GFP signals in the cytosol (Fig. 5B,C). An *arr-1* mutation engineered by CRISPR-Cas9 resulted in MOM-5::GFP accumulation at the cell membrane, confirming that these changes in MOM-5::GFP distribution indicate endocytosis blockade (Fig. 5B,C). We verified that the GFP-fused MOM-5 proteins are functional, as *zuls145* significantly rescued AVM undermigration of the *mom-5* mutants (Fig. S4C). Together, these results indicate that VANG-1 mediates MOM-5 endocytosis cell-autonomously.

### VANG-1 forms protein complexes with Frizzled receptors

We previously showed that VANG-1 complexed with MIG-1 when expressed in mammalian cells (Chen et al., 2017), and this was again confirmed in the current study (Fig. 6A,B; Fig. S5A). To investigate whether VANG-1 also forms complexes with MOM-5, we expressed HA-tagged VANG-1 and FLAG-tagged MOM-5 in HEK293 cells. Pull-down of MOM-5 by FLAG antibodies co-immunoprecipitated VANG-1, suggesting that VANG-1 and MOM-5 form protein complexes (Fig. 6A,C; Fig. S5B). To gain further insight into the interaction between VANG-1 and Frizzled



**Fig. 5. VANG-1 regulates the distribution of MOM-5 Frizzled receptors on the neuronal membrane.** (A) Schematic of the *vang-1(twn3)* and *arr-1(twn10)* mutations engineered by CRISPR-Cas9 into the *zuls145(Pnmy-2::MOM-5::GFP)* strain. Yellow highlights the changes in the *vang-1(twn3)* and *arr-1(twn10)* alleles. Black boxes indicate exons. (B, C) Confocal projection images (B) and quantification of MOM-5::GFP distribution (C) in QR.d. MOM-5::GFP is from *zuls145(Pnmy-2::MOM-5::GFP)* and the QR.d membrane marker is *rdvls1(Pegl-17::Myri::mCherry)* with mCherry pseudocolored in magenta. The V5R seam cell is indicated. Scale bar: 10  $\mu$ m. Each dot represents a single neuron, with mean  $\pm$ s.e.m. indicated. Number of neurons scored is indicated in parentheses. \* $P$ <0.05, \*\* $P$ <0.01, \*\*\* $P$ <0.001 (one-way ANOVA followed by Bonferroni's multiple comparison test).

proteins, we generated truncated MIG-1 and MOM-5 proteins that lacked the Wnt-binding, cysteine-rich domain ( $\Delta$ CRD). Removal of the CRD profoundly weakened the ability of either MIG-1 or MOM-5 to co-immunoprecipitate VANG-1, whereas deletion of the cytoplasmic domain ( $\Delta$ C) of MIG-1 had no effect (Fig. 6B,C; Fig. S5). We were not able to express MOM-5 lacking the cytoplasmic domain in mammalian cells for unknown reason. These results are consistent with the observation from a prior *Drosophila* study reporting that the Frizzled CRD binds Van Gogh (Wu and Mlodzik, 2008), and indicate that the CRD of Frizzled proteins is important for VANG-1-Frizzled interaction.


## DISCUSSION

In addition to its well-established function in the development of *Drosophila* eye and wing discs, PCP signaling also plays important roles in various aspects of vertebrate neuronal development, such as neural tube closure, polarization of inner hair cells, neuronal migration and axon guidance (Goodrich, 2008; Tissir and Goffinet, 2013; Zou, 2012). In neural tube closure, PCP signaling generates molecular asymmetries within and between neuronal precursors that polarize these cells uniformly along the tissue plane to enable convergent extension, a crucial process that elongates the vertebrate

embryo. Compared with these early morphogenetic events, our understanding of mechanisms by which PCP signaling controls long-range neuronal migration and axon guidance is incomplete. The current study establishes that VANG-1 acts cell-autonomously in migrating *C. elegans* neurons and tempers Wnt signaling by facilitating Frizzled endocytosis. These findings extend current understanding of how PCP signaling controls neuronal development and shapes the connectivity of the nervous system.

## VANG-1 shapes Wnt signaling depending on the cellular contexts of the neurons

*vang-1*, the *C. elegans* Vangl2 homolog, is involved in several morphogenetic events that bear some similarity to *Drosophila* or vertebrate PCP, such as the assembly of ventral nerve cord during which immature neurons display transient rosette formation similar to that in vertebrate convergent extension movement (Shah et al., 2017), or the repositioning of endodermal descendants during intestinal morphogenesis (Asan et al., 2016). *vang-1* also controls neuronal polarization (Sanchez-Alvarez et al., 2011), neurite outgrowth (Zheng et al., 2015) and neurite branching (Chen et al., 2017). Mentink et al. reported that *vang-1* controls the final positions of two QR.d, QR.paa (future AVM) and QR.pap (Mentink



**Fig. 6. VANG-1 forms protein complexes with Frizzled proteins that require the cysteine-rich domains.** (A) Schematic of various MIG-1 and MOM-5 deletion mutants used in the co-immunoprecipitation experiments. Boxes represent transmembrane domains. FL, full-length. (B,C) Co-immunoprecipitation experiments of MIG-1 (B) or MOM-5 (C). Arrows indicate MIG-1 and MOM-5 signals in respective immunoblots.

et al., 2014). In the *vang-1* mutant, QR.paa and QR.pap are positioned more anteriorly compared with those in the wild type, a phenotype that is opposite to that caused by loss of *egl-20*, *cwn-1* or *mom-5* (Mentink et al., 2014). This observation is consistent with our model that *vang-1* antagonizes a Wnt signaling cascade composed of *egl-20*, *cwn-1* and *mom-5* in QR.d. These results contrast with those in our prior study, which showed that *vang-1* promotes Wnt signaling to pattern neurite branching in the non-migratory PLM neuron (Chen et al., 2017). Because in both studies evidence suggests that *vang-1* facilitates Frizzled endocytosis, we speculate that cellular contexts determine whether VANG-1-mediated Frizzled internalization upregulates or diminishes Wnt signaling. In immature neurons that undergo long-range migration or cellular polarization, Frizzled internalization terminates Wnt signaling. By contrast, when neurons mature and begin to grow collateral branches in their neurites, as in the case of PLM, Frizzled endocytosis transduces Wnt signaling (Chen et al., 2017). These observations expand the spectrum of Wnt-PCP pathways, implying that VANG-1 directly tunes a Wnt signaling cascade previously thought to be distinct from the PCP pathway, thus blurring the boundaries between the β-catenin and PCP branches of Wnt signaling.

Anterior displacement of QL.d in the *egl-20* mutants has led to the hypothesis that these mutant QL.d have acquired properties of QR.d, which migrate anteriorly. This hypothesis gains some support

from the observation that a *cwn-1* mutation shifts the positions of QL.d in the *egl-20* mutants towards the posterior, which is explained by the fact that *cwn-1* drives QR.d anterior migration (Zinovyeva et al., 2008). It is therefore intriguing why the migration defects of QR.d, but not those of QL.d, are suppressed by *vang-1* mutations in the *egl-20* mutants. However, QL.d in the *cwn-1*; *egl-20* sometimes occupies a position that is even more posterior than QL.d in the wild type. The distribution of QL.d in the *cwn-1*; *egl-20* double mutant is also more posterior compared with that of QR.d in the same mutant (Zinovyeva et al., 2008). These observations suggest that QL.d in the *egl-20* mutants bears some phenotypic similarity to the wild-type QR.d, but their molecular properties could still be different. One of these differences is likely to be the distinct subsets of Frizzled receptors expressed or functioning in QL.d or QR.d. *mig-1* or *lin-17* are required in QL.d, but mutations of these Frizzled genes cause no undermigration or even slight overmigration in QR.d (Zinovyeva et al., 2008; Mentink et al., 2014). On the other hand, *mom-5* is important for QR.d migration yet dispensable for QL.d migration. We speculate that QL.d and QR.d may express different Frizzled receptors, and these Frizzled receptors engage different signaling cascades. All these potentially contribute to the observed difference in QL.d and QR.d regarding their sensitivity to the *vang-1* mutation in the *egl-20* mutant background.

We noted that the distribution of PVM becomes somewhat bimodal when *vang-1* mutations or RNAi are added to the *mig-1*



mutant (Fig. 1D; Fig. S1A,B). It has been shown that EGL-20 signaling regulates QL lineage migration independent of its distribution (Whangbo and Kenyon, 1999). We speculate that *mab-5* is activated and normal posterior QL.d migration is initiated when Wnt signaling exceeds a threshold in QL.d. Variability in *mab-5* expression increases in the *mig-1* and *lin-17* mutants, and it has been proposed that Wnt signaling acts to reduce variability of *mab-5* expression (Ji et al., 2013). We speculate that the *vang-1* mutations restore Wnt signaling intensity in general but do not improve all the regulatory steps within the Wnt signaling network, which in part explains low or no *mab-5* expression in some *mig-1*; *vang-1* mutant animals and a bimodal distribution of PVM positions.

### Genetic and molecular mechanisms by which VANG-1 promotes Frizzled endocytosis

In vertebrate cells, endocytosis of the Frizzled receptors could be mediated by  $\beta$ -arrestin2 or by ZNRF3-dependent ubiquitylation (Chen et al., 2003; Koo et al., 2012). Although Frizzled endocytosis by ubiquitylation clearly downregulates Wnt signaling in both mammalian intestinal epithelium and *C. elegans* neurons (Koo et al., 2012; Moffat et al., 2014), Frizzled internalization by  $\beta$ -arrestin2 could promote Wnt signaling both in vertebrate models of tumorigenesis and *C. elegans* neuronal development (Bonnans et al., 2012; Fereshteh et al., 2012; Seitz et al., 2014; Chen et al., 2017). In the present study, we show that VANG-1 and ARR-1 act in a common genetic pathway to internalize Frizzled receptors, and this downregulates Wnt signaling. It remains to be determined whether ZNRF3 and Vangl2- $\beta$ -arrestin2 cooperate to internalize Frizzled proteins, or if they represent distinct pathways to recycle Frizzled proteins into separate intracellular compartments. Physical interaction between VANG-1 and Frizzled ligands requires the Frizzled CRD but not the cytoplasmic domain. One possibility is that binding of VANG-1 to the Frizzled-CRD triggers Frizzled internalization mediated by the  $\beta$ -arrestin2 machinery. A prior study showed that Van Gogh, the *Drosophila* Vangl2, binds Frizzled-CRD, and when addressed in the context of epithelial cell polarization, this represents an interaction *in trans* between Frizzled proteins and Van Gogh from neighboring cells (Wu and Mlodzik, 2008). By contrast, our genetic and biochemical data suggest that in *C. elegans* migrating neurons, the interaction between VANG-1 and Frizzled-CRD can occur *in cis*, i.e. in the same cell. Although the crystal structure of the Frizzled-8-CRD complexed with the *Xenopus* Wnt-8 ligand has been determined recently (Janda et al., 2012), the structure of Vangl2 or its homologous proteins remains unknown. In the future, it will be important to decipher the structure of Vangl2 by itself and that of Vangl2 complexed with the Frizzled CRD. This will offer important insight into how binding of diffusible (such as Wnts) or membrane-tethered ligands (such as Vangl2) to the Frizzled-CRD initiates Frizzled endocytosis and tunes Wnt signaling.

### Modulation of Wnt signaling that terminates neuronal migration

Wnt signaling has been shown to promote the migration of neurons and growth cones by acting as either attractive or repulsive cues (Yoshikawa et al., 2003; Lyuksyutova et al., 2003; Pan et al., 2006). A recent report in *C. elegans* suggests that signaling through the Frizzled receptor MIG-1 and LIN-17 terminates the anterior migration of QR.paa and QR.pap neurons through the  $\beta$ -catenin BAR-1 (Mentink et al., 2014). Adding to the complexity of Wnt signaling in controlling neuronal migration, Mentink et al. further showed that *vang-1* also acts as a brake to prevent QR.pax from

overmigrating. As the migration of QR.p was not affected in the *vang-1* mutant, they concluded that *vang-1* functions independently of Wnt- $\beta$ -catenin signaling, although no genetic experiments were performed to test the interaction between *vang-1* and the Wnt pathway genes (Mentink et al., 2014). By contrast, we show here that *vang-1* antagonizes Wnt signaling by removing Frizzled receptors from the cell membrane. As Frizzled proteins transduce Wnt signaling through both  $\beta$ -catenin-dependent and  $\beta$ -catenin-independent pathways, our results suggest that *vang-1* has a broad, negative impact on Wnt signaling in early neural development. This conclusion is supported by observations made in four different neuronal classes (QL.d, QR.d, HSN and ALM). Unlike QL.d, for which the target of Wnt signaling for neuronal migration, the Hox gene *mab-5*, has been identified, genes that Wnt-Frizzled signaling targets to promote migration in QR.d and HSN or polarization in ALM remain unknown. Identification of such Wnt target genes in neuronal development helps to clarify the role of *vang-1* in fine-tuning Wnt signaling. Our current study, together with a prior report that PLR-1/ZNRF3 E3 ligase downregulates Wnt signaling by internalizing Frizzled proteins (Moffat et al., 2014), suggests that timely internalization of Frizzled proteins is a crucial step to prevent aberrant neuronal wiring caused by prolonged Wnt signaling. Understanding the molecular and structural detail of VANG-1-mediated endocytosis of Frizzled proteins will provide important insights into the modulation of Wnt signaling in early neuronal development.

## MATERIALS AND METHODS

### Key resources

Key experimental materials, including *C. elegans* and bacterial strains, cell lines and antibodies, are listed in Table S1. *C. elegans* strains were cultured and maintained as described (Brenner, 1974). The HEK293 cell line used in this study is from the American Type Culture Collection (ATCC) and was authenticated and tested for contamination. The antibodies used in this study were previously validated (Duval et al., 2014; Sanna et al., 2018; Schafer and Braun, 1995).

### Generation of the *vang-1(twn3)* allele by CRISPR-Cas9 gene editing

#### Germline CRISPR

We performed germline CRISPR as previously described with some modifications (Dickinson et al., 2013). In brief, we co-injected into worms a mixture of the following three plasmids: pDD162(*Peft-3::Cas9*), PU6::*vang-1* sgRNA(GACACGAGGAGTTGCGTT) or PU6::*arr-1* sgRNA(GGTGACTATGTCGACTTGA), and *unc-22* sgRNA (twitcher phenotypes, as a co-CRISPR marker) (Kim et al., 2014). Twitcher F1 transgenics were subjected to T7 endonuclease digestion to identify nucleotide mismatches caused by Cas9-mediated editing. Homozygotes of the *vang-1(twn3)* thus generated were confirmed by DNA sequencing.

#### Somatic CRISPR

Somatic CRISPR was performed as previously described (Shen et al., 2014). In brief, we co-injected the mixture of the following plasmids: *Pegl-17::Cas9* (Q/V5 lineage-specific), PU6::*vang-1* sgRNA(GACACGAGGAGTTGCGTT) and the co-injection markers *Pgcy-8::mCherry* (for isolation of transformants) and *Pmec-7::mCherry*. To optimize the expression of transgenes, animals were cultivated at 25°C, and PVM positions scored with *Pmec-7::mCherry*, which labels the six touch mechanosensory neurons.

### Plasmid construction and molecular biology

We used standard molecular biology techniques for cloning and plasmid construction. Constructs used for generating transgenes of the *twnEx* series used the pPD95.77 Fire vector as their backbone. The constructs for expression in the HEK293 cells were in the pcDNA, RK5F vector backbone.

Primer sequences for cloning *vang-1*, *mig-1* and *mom-5* are listed in Table S2. Germline transformation by microinjection was performed as described (Mello et al., 1991).

### RNA interference by feeding

Feeding RNAi was performed as described (Kamath et al., 2001). In brief, *vang-1* RNAi bacteria were induced by 1 mM isopropyl  $\beta$ -D-1-thiogalactopyranoside (IPTG) for 1.5 h before seeding on NGM plates supplemented with 1 mM IPTG. Gravid hermaphrodites were bleached on RNAi plates and newly hatched L1 larvae were allowed to develop to adulthood. This procedure was repeated one more time before gravid hermaphrodites were collected and bleached to obtain synchronized early L1 larvae arrested in M9 buffer. These arrested L1 larvae were then released on fresh *vang-1* RNAi plates and scored at appropriate stages for individual phenotypes (2–3 h post-hatching L1 for MOM-5::GFP quantification, and late L4 for PVM positions).

### Quantification of neuronal positions by epifluorescence microscopy

The transgene *zdis5(Pmec-4::gfp)* labels PVM and AVM, and integrated array *mgIs42(Ptph-1::gfp)* or extrachromosomal array *twmEx199(Punc-86::gfp)* mark HSN. Late L4 animals were anesthetized with 1% sodium azide. Neurons were imaged under the 10 $\times$  objective of the AxioImager M2 imaging system (Carl Zeiss). For quantification of AVM and PVM positions, we set the nose tip as zero and the tail tip as one. AVM and PVM positions were defined as fractions of the nose-to-tail length. For quantification of HSN positions, we set the vulva opening as zero and the tail tip as one, as normally the HSNs do not migrate anterior to the vulva. HSN positions are defined as fractions of the vulva-to-tail distance. Pixel-wise measurement of length was performed with ImageJ.

### Characterization of ALM polarity

The transgene *zdis5(Pmec-4::gfp)* also labels the ALM and PLM touch receptor neurons. L4 animals were anesthetized with 1% sodium azide. Neurons were imaged under the 10 $\times$  objective of the AxioImager M2 imaging system (Carl Zeiss). In the Wnt pathway mutant, a bipolar polarity defect was defined as ALM neurons with the normal anterior process and a posterior process longer than ten times the cell diameter; this definition is the same as previously published methods (Pan et al., 2008). For VANG-1 overexpression, a bipolar polarity defect was defined as ALM neurons with the normal anterior process and a posterior process longer than five times cell diameter. A reversed polarity defect was defined as ALM neurons having a long posterior process without the normal anterior process.

### Quantification of *mab-5* expression by fluorescence microscopy

The transgene *mulS16(Pmab-5::gfp)* was used to observe *mab-5* expression, with QL.a and QL.p marked by *rdvIs1(Pegl-17::Myri::mCherry)*. We synchronized early L1 larvae by hatching them in M9 buffer, and scored them after resuming feeding for 2–3 h on regular OP50 *Escherichia coli* plates. L1 larvae were observed using the 100 $\times$  objective of the AxioImager M2 imaging system (Carl Zeiss) and AxioCam MRm CCD camera (Carl Zeiss) under identical imaging parameters. GFP pixel density OF *mulS16* in QL.d, the boundary of which is defined by *rdvIs1*, was quantified using ImageJ.

### Characterization of MOM-5::GFP by confocal microscopy

We used *rdvIs1(Pegl-17::Myri::mCherry)*, which highlights the plasma membrane, to label the Q cell lineages, and *zuls145(Pnmy-2::mom-5::gfp)* to observe MOM-5::GFP signal (Ou et al., 2010; Park et al., 2004). Note that chromosomes are also labeled by *rdvIs1*. Early L1 larvae (post-hatching 1–2 h) were examined under the 100 $\times$  objective of the LSM 700 Confocal System (Carl Zeiss), and series of z-stack projection images were acquired using Zeiss Zen software under identical parameters. For this analysis, we focused on QR.a and QR.p. Pixel-wise quantification for GFP fluorescence intensity was quantified using ImageJ for individual optical slices, and summation of all optical sections from the same z-stack series was represented. To determine the subcellular distribution of MOM-5::GFP, cell membrane MOM-5::GFP signals were first quantified using *rdvIs1* as a

reference for QR.d membrane. Cytosolic MOM-5::GFP signals were derived by subtracting membrane MOM-5::GFP signals from total MOM-5::GFP signals.

### Co-immunoprecipitation and western blotting

Gene constructs were transfected into HEK293 cells with Lipofectamine (Invitrogen) and grown for 2 days before lysis with buffer (50 mM Tris, 150 mM NaCl, 2 mM EDTA pH 8.0, 0.5% sodium deoxycholate, 10 mM phenylmethylsulfonyl fluoride, 1 M dithiothreitol, 1% NP-40), protease inhibitor cocktail (Roche) and 20 mM sodium fluoride. For co-immunoprecipitation, we used anti-HA (Invitrogen) or anti-FLAG (Sigma) beads to pull down proteins from cell lysates. After proteins (1  $\mu$ g) had bound to the beads, we added 4 $\times$  sample buffer with 10%  $\beta$ -mercaptoethanol and incubated samples at 37 $^{\circ}$ C for 20 min to prevent aggregation of transmembrane proteins. Samples for western blot analysis were electrophoresed in 10% SDS-polyacrylamide gel and transferred onto the PVDF membrane. We used anti-HA, anti-FLAG or anti- $\beta$ -actin antibodies as described in the Table S1 to detect protein signals.

### Statistics

For all samples, *n* represents the number of cells scored. Cells were excluded only when they could not be unambiguously identified or visualized. One-way ANOVA followed by Bonferroni's multiple comparison test was used for multiple comparisons between conditions within a dataset. For data in Fig. 2D, Fisher exact test was used. Statistics was performed in Prism.

### Acknowledgements

We thank Gian Garriga and the *C. elegans* Genetics Center (CGC) for worm strains. CGC is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

### Competing interests

The authors declare no competing or financial interests.

### Author contributions

Conceptualization: C.-W.H., C.-L.P.; Methodology: C.-W.H., C.-L.P.; Validation: C.-L.P.; Formal analysis: C.-W.H., C.-L.P.; Investigation: C.-W.H., C.-P.L., C.-K.C., J.T., C.-H.C., C.-L.P.; Data curation: C.-W.H., C.-P.L., C.-K.C., J.T., C.-L.P.; Writing - original draft: C.-W.H., C.-L.P.; Writing - review & editing: C.-W.H., C.-P.L., C.-K.C., J.T., C.-H.C., C.-L.P.; Visualization: C.-W.H., C.-L.P.; Supervision: C.-L.P.; Project administration: C.-L.P.; Funding acquisition: C.-L.P.

### Funding

This study was funded by the Ministry of Science and Technology, Taiwan (MOST 103-2320-B-002-050-MY3, MOST 106-2320-B-002-051-MY3 to C.-L.P.).

### Supplementary information

Supplementary information available online at <http://dev.biologists.org/lookup/doi/10.1242/dev.168666.supplemental>

### References

- Asan, A., Raiders, S. A. and Priess, J. R. (2016). Morphogenesis of the *C. elegans* intestine involves axon guidance genes. *PLoS Genet.* **12**, e1005950.
- Bailly, E., Walton, A. and Borg, J. P. (2017). The planar cell polarity Vangl2 protein: from genetics to cellular and molecular functions. *Semin. Cell Dev. Biol.* **81**, 62–70.
- Bänziger, C., Soldini, D., Schütt, C., Zipperlen, P., Hausmann, G. and Basler, K. (2006). Wntless, a conserved membrane protein dedicated to the secretion of Wnt proteins from signaling cells. *Cell* **125**, 509–522.
- Bartscherer, K., Pelte, N., Ingelfinger, D. and Boutros, M. (2006). Secretion of Wnt ligands requires Evi, a conserved transmembrane protein. *Cell* **125**, 523–533.
- Blockus, H. and Chédotal, A. (2016). Slit-Robo signaling. *Development* **143**, 3037–3044.
- Bonnans, C., Flaceliere, M., Grillet, F., Dantec, C., Desvignes, J.-P., Pannequin, J., Severac, D., Dubois, E., Bibeau, F., Escρίου, V. et al. (2012). Essential requirement for beta-arrestin2 in mouse intestinal tumors with elevated Wnt signaling. *Proc. Natl. Acad. Sci. USA* **109**, 3047–3052.
- Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics* **77**, 71–94.
- Chen, W., Ten Berge, D., Brown, J., Ahn, S., Hu, L. A., Miller, W. E., Caron, M. G., Barak, L. S., Nusse, R. and Lefkowitz, R. J. (2003). Dishevelled 2 recruits beta-arrestin 2 to mediate Wnt5A-stimulated endocytosis of Frizzled 4. *Science* **301**, 1391–1394.

- Chen, C.-H., He, C.-W., Liao, C.-P. and Pan, C.-L. (2017). A Wnt-planar polarity pathway instructs neurite branching by restricting F-actin assembly through endosomal signaling. *PLoS Genet.* **13**, e1006720.
- Clark, S. G., Chisholm, A. D. and Horvitz, H. R. (1993). Control of cell fates in the central body region of *C. elegans* by the homeobox gene *lin-39*. *Cell* **74**, 43-55.
- Desai, C., Garriga, G., McIntire, S. L. and Horvitz, H. R. (1988). A genetic pathway for the development of the *Caenorhabditis elegans* HSN motor neurons. *Nature* **336**, 638-646.
- Devenport, D. (2014). The cell biology of planar cell polarity. *J. Cell Biol.* **207**, 171-179.
- Dickinson, D. J., Ward, J. D., Reiner, D. J. and Goldstein, B. (2013). Engineering the *Caenorhabditis elegans* genome using Cas9-triggered homologous recombination. *Nat. Methods* **10**, 1028-1034.
- Dudanova, I. and Klein, R. (2013). Integration of guidance cues: parallel signaling and crosstalk. *Trends Neurosci.* **36**, 295-304.
- Duval, N., Daubas, P., Bourcier de Carbon, C., St Clément, C., Tinevez, J. Y., Lopes, M., Ribes, V. and Robert, B. (2014). *Msx1* and *Msx2* act as essential activators of *Atoh1* expression in the murine spinal cord. *Development* **141**, 1726-1736.
- Eisenmann, D. M. and Kim, S. K. (2000). Protruding vulva mutants identify novel loci and Wnt signaling factors that function during *Caenorhabditis elegans* vulva development. *Genetics* **156**, 1097-1116.
- Fereshteh, M., Ito, T., Kovacs, J. J., Zhao, C., Kwon, H. Y., Tornini, V., Konuma, T., Chen, M., Lefkowitz, R. J. and Reya, T. (2012). *beta-Arrestin2* mediates the initiation and progression of myeloid leukemia. *Proc. Natl. Acad. Sci. USA* **109**, 12532-12537.
- Forrester, W. C., Kim, C. and Garriga, G. (2004). The *Caenorhabditis elegans* Ror RTK CAM-1 inhibits EGL-20/Wnt signaling in cell migration. *Genetics* **168**, 1951-1962.
- Garriga, G., Desai, C. and Horvitz, H. R. (1993). Cell interactions control the direction of outgrowth, branching and fasciculation of the HSN axons of *Caenorhabditis elegans*. *Development* **117**, 1071-1087.
- Goodman, R. M., Thombre, S., Firtina, Z., Gray, D., Betts, D., Roebuck, J., Spana, E. P. and Selva, E. M. (2006). *Sprinter*: a novel transmembrane protein required for Wg secretion and signaling. *Development* **133**, 4901-4911.
- Goodrich, L. V. (2008). The plane facts of PCP in the CNS. *Neuron* **60**, 9-16.
- Harris, J., Honigberg, L., Robinson, N. and Kenyon, C. (1996). Neuronal cell migration in *C. elegans*: regulation of Hox gene expression and cell position. *Development* **122**, 3117-3131.
- Hilliard, M. A. and Bargmann, C. I. (2006). Wnt signals and frizzled activity orient anterior-posterior axon outgrowth in *C. elegans*. *Dev. Cell* **10**, 379-390.
- Janda, C. Y., Waghray, D., Levin, A. M., Thomas, C. and Garcia, K. C. (2012). Structural basis of Wnt recognition by Frizzled. *Science* **337**, 59-64.
- Ji, N., Middelkoop, T. C., Mentink, R. A., Betist, M. C., Tnegawa, S., Mooijman, D., Korswagen, H. C. and van Oudenaarden, A. (2013). Feedback control of gene expression variability in the *Caenorhabditis elegans* Wnt pathway. *Cell* **155**, 869-880.
- Jongbloets, B. C. and Pasterkamp, R. J. (2014). Semaphorin signalling during development. *Development* **141**, 3292-3297.
- Kamath, R. S., Martinez-Campos, M., Zipperlen, P., Fraser, A. G. and Ahringer, J. (2001). Effectiveness of specific RNA-mediated interference through ingested double-stranded RNA in *Caenorhabditis elegans*. *Genome Biol.* **2**, RESEARCH0002.
- Keleman, K., Rajagopalan, S., Cleppien, D., Teis, D., Paiha, K., Huber, L. A., Technau, G. M. and Dickson, B. J. (2002). *Comm* sorts robo to control axon guidance at the *Drosophila* midline. *Cell* **110**, 415-427.
- Kidd, T., Russell, C., Goodman, C. S. and Tear, G. (1998). Dosage-sensitive and complementary functions of roundabout and commissureless control axon crossing of the CNS midline. *Neuron* **20**, 25-33.
- Kim, H., Ishidate, T., Ghanta, K. S., Seth, M., Conte, D., Jr, Shirayama, M. and Mello, C. C. (2014). A co-CRISPR strategy for efficient genome editing in *Caenorhabditis elegans*. *Genetics* **197**, 1069-1080.
- Koo, B.-K., Spit, M., Jordens, I., Low, T. Y., Stange, D. E., Van De Wetering, M., Van Es, J. H., Mohammed, S., Heck, A. J. R., Maurice, M. M. et al. (2012). Tumour suppressor RNF43 is a stem-cell E3 ligase that induces endocytosis of Wnt receptors. *Nature* **488**, 665-669.
- Korswagen, H. C. (2002). Canonical and non-canonical Wnt signaling pathways in *Caenorhabditis elegans*: variations on a common signaling theme. *Bioessays* **24**, 801-810.
- Korswagen, H. C., Herman, M. A. and Clevers, H. C. (2000). Distinct beta-catenins mediate adhesion and signalling functions in *C. elegans*. *Nature* **406**, 527-532.
- Lyuksyutova, A. I., Lu, C. C., Milanesio, N., King, L. A., Guo, N., Wang, Y., Nathans, J., Tessier-Lavigne, M. and Zou, Y. (2003). Anterior-posterior guidance of commissural axons by Wnt-frizzled signaling. *Science* **302**, 1984-1988.
- Maloof, J. N., Whangbo, J., Harris, J. M., Jongeward, G. D. and Kenyon, C. (1999). A Wnt signaling pathway controls hox gene expression and neuroblast migration in *C. elegans*. *Development* **126**, 37-49.
- Mello, C. C., Kramer, J. M., Stinchcomb, D. and Ambros, V. (1991). Efficient gene transfer in *C. elegans*: extrachromosomal maintenance and integration of transforming sequences. *EMBO J.* **10**, 3959-3970.
- Mentink, R. A., Middelkoop, T. C., Rella, L., Ji, N., Tang, C. Y., Betist, M. C., Van Oudenaarden, A. and Korswagen, H. C. (2014). Cell intrinsic modulation of Wnt signaling controls neuroblast migration in *C. elegans*. *Dev. Cell* **31**, 188-201.
- Moffat, L. L., Robinson, R. E., Bakoulis, A. and Clark, S. G. (2014). The conserved transmembrane RING finger protein PLR-1 downregulates Wnt signaling by reducing Frizzled, Ror and Ryk cell-surface levels in *C. elegans*. *Development* **141**, 617-628.
- Ou, G., Stuurman, N., D'Ambrosio, M. and Vale, R. D. (2010). Polarized myosin produces unequal-size daughters during asymmetric cell division. *Science* **330**, 677-680.
- Pan, C.-L., Howell, J. E., Clark, S. G., Hilliard, M., Cordes, S., Bargmann, C. I. and Garriga, G. (2006). Multiple Wnts and frizzled receptors regulate anteriorly directed cell and growth cone migrations in *Caenorhabditis elegans*. *Dev. Cell* **10**, 367-377.
- Pan, C. L., Baum, P. D., Gu, M., Jorgensen, E. M., Clark, S. G. and Garriga, G. (2008). *C. elegans* AP-2 and retromer control Wnt signaling by regulating mig-14/Wntless. *Dev. Cell* **14**, 132-139.
- Park, F. D., Tenlen, J. R. and Priess, J. R. (2004). *C. elegans* MOM-5/frizzled functions in MOM-2/Wnt-independent cell polarity and is localized asymmetrically prior to cell division. *Curr. Biol.* **14**, 2252-2258.
- Prasad, B. C. and Clark, S. G. (2006). Wnt signaling establishes anteroposterior neuronal polarity and requires retromer in *C. elegans*. *Development* **133**, 1757-1766.
- Sanchez-Alvarez, L., Visanuvimol, J., Mcewan, A., Su, A., Imai, J. H. and Colavita, A. (2011). VANG-1 and PRKL-1 cooperate to negatively regulate neurite formation in *Caenorhabditis elegans*. *PLoS Genet.* **7**, e1002257.
- Sanna, M. D., Quattrone, A. and Galeotti, N. (2018). Activation of ERK/CREB pathway in noradrenergic neurons contributes to hypernociceptive phenotype in H4 receptor knockout mice after nerve injury. *Neuropharmacology* **128**, 340-350.
- Schafer, K. and Braun, T. (1995). Monoclonal anti-FLAG antibodies react with a new isoform of rat Mg<sup>2+</sup> dependent protein phosphatase beta. *Biochem. Biophys. Res. Commun.* **207**, 708-714.
- Seitz, K., Dürsch, V., Harnoš, J., Bryja, V., Gentzel, M. and Schambony, A. (2014). *beta-Arrestin* interacts with the beta/gamma subunits of trimeric G-proteins and dishevelled in the Wnt/Ca<sup>2+</sup> pathway in xenopus gastrulation. *PLoS ONE* **9**, e87132.
- Shafer, B., Onishi, K., Lo, C., Colakoglu, G. and Zou, Y. (2011). Vangl2 promotes Wnt/planar cell polarity-like signaling by antagonizing Dvl1-mediated feedback inhibition in growth cone guidance. *Dev. Cell* **20**, 177-191.
- Shah, P. K., Tanner, M. R., Kovacevic, I., Rankin, A., Marshall, T. E., Noblett, N., Tran, N. N., Roenspies, T., Hung, J., Chen, Z. et al. (2017). PCP and SAX-3/Robo pathways cooperate to regulate convergent extension-based nerve cord assembly in *C. elegans*. *Dev. Cell* **41**, 195-203.e3.
- Shen, Z., Zhang, X., Chai, Y., Zhu, Z., Yi, P., Feng, G., Li, W. and Ou, G. (2014). Conditional knockouts generated by engineered CRISPR-Cas9 endonuclease reveal the roles of coronin in *C. elegans* neural development. *Dev. Cell* **30**, 625-636.
- Sun, K. L. W., Correia, J. P. and Kennedy, T. E. (2011). Netrins: versatile extracellular cues with diverse functions. *Development* **138**, 2153-2169.
- Tear, G., Harris, R., Sutaria, S., Kilomanski, K., Goodman, C. S. and Seeger, M. A. (1996). *Commissureless* controls growth cone guidance across the CNS midline in *Drosophila* and encodes a novel membrane protein. *Neuron* **16**, 501-514.
- Tessier-Lavigne, M. and Goodman, C. S. (1996). The molecular biology of axon guidance. *Science* **274**, 1123.
- Tissir, F. and Goffinet, A. M. (2013). Shaping the nervous system: role of the core planar cell polarity genes. *Nat. Rev. Neurosci.* **14**, 525-535.
- Wang, B. B., Müller-Immergluck, M. M., Austin, J., Robinson, N. T., Chisholm, A. and Kenyon, C. (1993). A homeotic gene cluster patterns the anteroposterior body axis of *C. elegans*. *Cell* **74**, 29-42.
- Whangbo, J. and Kenyon, C. (1999). A Wnt signaling system that specifies two patterns of cell migration in *C. elegans*. *Mol. Cell* **4**, 851-858.
- Wu, J. and Mlodzik, M. (2008). The frizzled extracellular domain is a ligand for Van Gogh/Stbm during nonautonomous planar cell polarity signaling. *Dev. Cell* **15**, 462-469.
- Yam, P. T. and Charron, F. (2013). Signaling mechanisms of non-conventional axon guidance cues: the Shh, BMP and Wnt morphogens. *Curr. Opin. Neurobiol.* **23**, 965-973.
- Yoshikawa, S., Mckinnon, R. D., Kokel, M. and Thomas, J. B. (2003). Wnt-mediated axon guidance via the *Drosophila* Derailed receptor. *Nature* **422**, 583-588.
- Zheng, C., Diaz-Cuadros, M. and Chalfie, M. (2015). Dishevelled attenuates the repelling activity of Wnt signaling during neurite outgrowth in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA* **112**, 13243-13248.
- Zinovyeva, A. Y., Yamamoto, Y., Sawa, H. and Forrester, W. C. (2008). Complex network of Wnt signaling regulates neuronal migrations during *Caenorhabditis elegans* development. *Genetics* **179**, 1357-1371.
- Zou, Y. (2012). Does planar cell polarity signaling steer growth cones? *Curr. Top. Dev. Biol.* **101**, 141-160.