

Counts and colours of faint galaxies in the U and R bands

David W. Hogg,¹ Michael A. Pahre,² James K. McCarthy,² Judith G. Cohen,²
Roger Blandford,¹ Ian Smail³ and B. T. Soifer²

¹Theoretical Astrophysics, California Institute of Technology, Mail Stop 130-33, Pasadena, CA 91125, USA

²Palomar Observatory, California Institute of Technology, Mail Stop 105-24, Pasadena, CA 91125, USA

³Department of Physics, University of Durham, South Road, Durham DH1 3LE

Accepted 1997 February 15. Received 1997 February 5; in original form 1996 April 15

ABSTRACT

Ground-based counts and colours of faint galaxies in the U and R bands in one field at high Galactic latitude are presented. Integrated over flux, a total of 1.2×10^5 source deg^{-2} are found to $U=25.5$ mag and 6.3×10^5 source deg^{-2} to $R=27$ mag, with $d \log N/dm \sim 0.5$ in the U band and $d \log N/dm \sim 0.3$ in the R band. Consistent with these number–magnitude curves, sources become bluer with increasing magnitude to median $U-R=0.6$ mag at $24 < U < 25$ mag and $U-R=1.2$ mag at $25 < R < 26$ mag. Because the Lyman break redshifts into the U band at $z \approx 3$, at least 1.2×10^5 source deg^{-2} must be at redshifts $z < 3$. Measurable U -band fluxes of 73 per cent of the 6.3×10^5 source deg^{-2} suggest that the majority of these also lie at $z < 3$. These results require an enormous space density of objects in any cosmological model.

Key words: galaxies: fundamental parameters – galaxies: photometry – cosmology: observations – ultraviolet: galaxies.

1 INTRODUCTION

The number of faint galaxies as a function of apparent magnitude is one of the fundamental observational constraints on cosmology and galactic evolution. The first such measurements made with CCDs and automated source-detection algorithms (Hall & MacKay 1984; Tyson 1988), showed signs of an excess over predictions based on naive extrapolations of local galaxy properties. Recent ground-based galaxy counts to $B=27.5$ (Metcalf et al. 1995), $V=27$, $R=27$, $I=25.5$ (Smail et al. 1995), and $K=24$ mag (Djorgovski et al. 1995) all reach integrated number densities around $6 \times 10^5 \text{ deg}^{-2}$ and show numbers increasing by a factor of $\sim 2 \text{ mag}^{-1}$, or $d \log N/dm \sim 0.3$ in the red and near-infrared. Given local luminosity function determinations (e.g., Loveday et al. 1992), these high numbers require either strong source evolution or extreme world models; it is certainly possible that a number of effects contribute. Furthermore, the variation of the slope $d \log N/dm$ with observed waveband, showing increased slope ~ 0.5 in the B band and ~ 0.6 in the U band and corresponding blueing of the objects with apparent magnitude (see Koo & Kron 1992, for a review) is an important clue to the physical processes generating the radiation in these objects.

The *Hubble Space Telescope* (*HST*) has proven very effective in the field of faint galaxy counts, benefiting from reduced sky brightness and the small angular sizes of faint galaxies (Smail et al. 1995), which are only marginally resolved even with the 0.1-arcsec resolution of the *HST* WFC. In a 3×10^4 s *HST* exposure, Cowie, Hu & Songaila (1995) count objects in the field to $I=26$, finding $\approx 8 \times 10^5 \text{ deg}^{-2}$. More recently, in the *Hubble Deep Field* (Williams et al. 1996), 1.5×10^5 s *HST* exposures were taken in each of four filters in a single field; counts by Williams et al. (1996) find roughly $10^6 \text{ object deg}^{-2}$ to $(F606W)_{AB} \approx 30$ mag.

Galaxy counts in the U band have not been pushed to numbers nearly as high as those in other optical bands. The subject is interesting because the U counts that do exist show the number count rises very rapidly, by a factor of 3 or 4 mag^{-1} or $d \log N/dm \sim 0.5$ or 0.6 (Koo 1986; Songaila, Cowie & Lilly 1990; Jones et al. 1991); i.e., U -band counts are much steeper than counts at longer wavelengths. We present the deepest published galaxy counts in the U band, reaching $U \approx 25.5$ mag, in an image taken under conditions of good seeing with the Hale Telescope, along with galaxy counts in the R band, in a deep image taken with the W. M. Keck Telescope. We also present $U-R$ colours to look for the colour trend implied by the difference in count slopes.

In terms of point-source sensitivity, the U -band image presented here will be surpassed by ultra-deep *HST* observations (such as those in the *Hubble Deep Field*) but this image has very good surface-brightness sensitivity and a much wider field of view.

At $z > 3$, the observed U band is at emitted wavelengths shortward of the Lyman limit, which is expected to be optically thick to absorption by neutral hydrogen in the intergalactic medium. This would be observed as anomalously low U -band flux, or anomalously red ($U - B$) colours for any population of $z > 3$ objects. Guhathakurta, Tyson & Majewski (1990) used the lack of such objects to demonstrate that to $R = 26$ mag, faint galaxy counts are not dominated by objects at $z > 3$. Steidel, Pettini & Hamilton (1995) have counted and, recently, spectroscopically confirmed (Steidel et al. 1996) a population of objects identified for anomalously red ($U - G$) colours and find that there are ~ 1500 objects deg^{-2} in the redshift range $3 < z < 3.4$ to $R \approx 25$ mag.

2 FIELD SELECTION AND OBSERVATIONS

The field, RA $00^{\text{h}}53^{\text{m}}23^{\text{s}}.20$, Dec $+12^{\circ}33'57''.5$ (J2000), was chosen for the purposes of deep K -band imaging (Djorgovski et al. 1995) and a faint object redshift survey (Cohen et al. 1996a) from among existing deep Medium Deep Survey (MDS, Griffiths et al. 1994) *HST* images which are taken in parallel mode in fields selected for low extinction and high Galactic latitude. The particular MDS field was chosen for its long *HST* exposure time and high Galactic latitude to minimize stellar contamination. Additional ground-based photometry on this field will be reported by Pahre et al. (in preparation) and the detailed results of the redshift survey will be reported by Cohen et al. (in preparation).

The U -band data were taken in 1995 September with the COSMIC instrument (Dressler 1993) at the prime focus of the 5-m Hale Telescope. Palomar CCD13, a Tektronix 2048×2048 array of 24- μm pixels (TK2048), was used in place of the standard COSMIC TK2048 CCD owing to high near-UV sensitivity of CCD13. Individual 600-s exposures were taken on a 5×5 grid with roughly 10-arcsec spacing. The seeing is 1.1 arcsec in the final stacked image. Patchy cloud cover necessitated independent photometric calibration (see below).

The R -band data were taken during periods of fair seeing (0.8 arcsec in the final stacked image) on spectroscopic runs in 1995 July and August with the Low Resolution Imaging Spectrograph (LRIS) (Oke et al. 1995) on the 10-m Keck Telescope. Individual 600-s exposures were taken at dithers of several arcsec with respect to one another. A temporary problem with the telescope caused the individual R -band exposures to be contaminated with transient, non-repeating streaks of scattered light, from bright stars outside the field reflecting from telescope structures. This limits the quality of the R image flat-field and background estimation, and reduces the completeness and quality of the photometry at very faint levels. In removing the streaks, small gradients on scales of ~ 20 arcsec were eliminated (taking with them any hypothetical population of large, low-surface-brightness galaxies; of course the detection algorithm is not optimized for such objects anyway). Again patchy cloud cover necessitated independent photometric calibration.

For both the U - and R -band data, before stacking, individual images were shifted and geometrically remapped to a Cartesian plane according to a distortion map determined from the dithered images themselves. The remapped, shifted images were stacked with the IRAF/IMCOMBINE task, making use of sigma-clipping to remove cosmic rays. Matched 1×1 arcmin² sections of the stacked Hale U - and Keck R -band images are shown in Figs 1 and 2, and observational details are summarized in Table 1.

The U - and R -band images were calibrated with U - and R_c -band CCD images taken under photometric conditions with CCD13 on the Palomar Sixty-Inch Telescope (P60). The details of the P60 data reduction are presented in Pahre et al. (in preparation). Calibration shows that the filter and CCD used on the Hale Telescope make a non-standard U bandpass, which we denote ' U_{13} '. We find $U = U_{13} + 0.006(U_{13} - R)$. In the following we use U_{13} but the magnitudes can be corrected to pseudo- U -band by adding 0.08 mag, correct for a typical faint object, which has $(U_{13} - R) = 1.3$ mag. This correction is on the same order as the estimated uncertainty in the calibration, between 0.05 and 0.1 mag.

3 ANALYSIS

3.1 Object detection

Objects are detected in both the U - and R -band images with the image analysis package SEXTRACTOR (Bertin & Arnouts 1996). The algorithm is: (2) convolve the image with a Gaussian filter with FWHM matched to the point spread function (PSF); (3) find objects above the threshold which corresponds to a point source with $U_{13} = 25.63$ mag or $R = 28.49$ mag. Often detected objects have multiple peaks; a peak is split off into its own object only if its part of the object contains > 10 per cent of the total flux. We did not use the package for star/galaxy separation or any 'cleaning' of spurious detections. Stars only make up a small fraction of the faint sources at high Galactic latitude (Smail et al. 1995) and cleaning spurious objects is handled by our noise object and completeness corrections described below.

3.2 Photometry

In order to match the seeing of our images (1.1-arcsec FWHM in the U -band and 0.8 arcsec in the R -band), aperture photometry is performed in 1.5-arcsec diameter apertures, which is between 1.3 and 2 times the seeing FWHM. To these aperture magnitudes aperture corrections are added to attempt, in a statistical way, to account for object flux coming from outside the aperture. The faintest objects are consistent with having stellar profiles, in these images and even in R -band images with superior seeing (Smail et al. 1995), so we added the stellar correction, -0.60 mag in the U -band image and -0.25 mag in the R -band. For all objects we also measure a 1σ isophotal magnitude to which is added an aperture correction varying linearly from 0 to -0.60 over the range $23.5 < U_{13} < 25.5$ mag and from 0 to -0.25 over $24.5 < R < 26.0$ mag. Every object is assigned the brighter of the two corrected magnitudes.




Figure 1. A 1×1 arcmin² section of the final, stacked *U*-band image. Sources with $25.0 < U_{13} < 25.5$ mag are circled.

The reddening in this field is $A_V = 0.13$ mag (Burstein & Heiles 1982) so we corrected the *U*-band fluxes by 0.20 mag and *R*-band by 0.11 mag.

3.3 Removal of noise objects

The transforming, shifting and stacking of the images introduces pixel-to-pixel correlations in the noise, making it difficult to estimate analytically the contamination of the counts by spurious detections of peaks in the noise. Spurious detections of this type are corrected for in a statistical way, by subtracting ‘noise counts’ from the positive counts. Noise counts are estimated by running the detection and photometry algorithms on the images but searching for negative rather than positive objects. The raw negative counts are subtracted from the raw positive counts before applying completeness corrections (see below) and all Poisson error bars include the uncertainty added by this procedure. In the *U*-band image, noise objects account for 10 and 29 per cent of the objects in the two faintest half-magnitude bins and in

the *R*-band image, only 0.6 and 8 per cent in the two faintest half-magnitude bins.

Spurious objects are sometimes produced by cosmic ray hits. However, so many individual exposures (47 in U_{13} , 14 in *R*) were stacked with the sigma-clipping algorithm that no significant-flux cosmic-ray events could plausibly remain.

3.4 Completeness correction

The counts are corrected for completeness in a manner similar to that of Smail et al. (1995). Detected objects are cut out of the final *U*- and *R*-band images, dimmed and added back into randomly selected subfields. The detection algorithm is run and the catalogues of objects in the subfields are compared before and after adding the additional object. By this procedure we generate a ‘completeness matrix’ P_{ij} which stores the probability that an added object of true magnitude m_i is in fact detected with magnitude m_j . Because some objects are not recovered at all and because others are blended into existing brighter objects, etc., the




Figure 2. A 1×1 arcmin² section of the final, stacked *R*-band image, matched to Fig. 1. Sources with $26.5 < R < 27.0$ mag are circled.

Table 1. Imaging data.

| band | instrument | field area (arcmin ²) | total t_{exp} (s) | pixel scale (arcsec) | seeing ^a (arcsec) | 3σ detection ^b (mag) |
|----------|-------------------|--------------------------------------|-------------------------------|-------------------------|---------------------------------|---|
| U_{13} | Hale/COSMIC/CCD13 | 81 | 28000 | 0.28 | 1.1 | 26.36 |
| U | P60/CCD13 | 160 | 5400 | 0.37 | 2.2 | 23.13 |
| R | Keck/LRIS | 39 | 8400 | 0.22 | 0.8 | 28.01 |
| R_C | P60/CCD13 | 160 | 600 | 0.37 | 1.3 | 23.25 |

^aSeeing FWHM in final, stacked images.

^bSignal-to-noise ratio of 3 through an aperture of diameter 1.5 times the seeing FWHM, with no aperture correction applied.

sum over j of P_{ij} will not in general be unity. The completeness information is a matrix because the statistical incompleteness in each magnitude bin depends on the true functional form of the number–magnitude plot; however,

under the assumption that the galaxy count slopes do not change dramatically over the magnitude range of interest, this matrix converts naturally into a fractional completeness as a function of magnitude.

To construct P_{ij} in practice, bright objects were cut out of the images, dimmed by a factor of 10, replaced at random locations, and then searched for and photometered by the detection algorithm. In the U -band image, 10^5 random replacements were performed, and in the R -band image, 6×10^4 , with the number increasing with magnitude in proportion to the counts. Each element of the matrix has an associated uncertainty from Poisson statistics. The detection fraction f_j in each bin was generated by assuming that the U counts follow a power law $d \log N/dm \approx 0.47$ and that the R counts follow the power law $d \log N/dm \approx 0.33$. Our noise-object-subtracted, completeness-corrected U - and R -band object counts are shown in Figs 3 and 4, along with the completeness-corrected counts of other authors. Note that because this completeness correction corrects not just for missing numbers but also for photometry errors, and because the counts increase with magnitude, it is possible for the completeness correction near (but not at) the detection threshold to be negative, as is seen for at least one point in Fig. 3.

Faint objects get smaller in angular size with increasing magnitude (Smail et al. 1995; Im et al. 1995), and since more compact objects are easier to detect at the same flux level, the detected fractions calculated by this technique would, in better data, be lower limits. A better procedure would involve changing the angular sizes of the objects as well as dimming them before replacing them. However, the seeing in our images is not good enough for the changing sizes to be a significant effect at faint levels.




Figure 3. The U_{13} -band number counts. Open squares are the positive counts with the ‘noise’ counts (negative counts, see text) subtracted, from the data presented here; error bars are Poisson noise for the difference. Filled squares are the completeness-corrected counts (see text), also from the data presented here. The solid line is a fit to the corrected counts; it has slope $d \log N/dm = 0.467$. The dotted line shows counts of Songaila et al. (1990), converted to this magnitude system assuming $U_{13} \approx U_{AB} - 0.79$ mag for a typical object.

3.5 Colour measurement

Magnitudes through 1.5-arcsec diameter apertures, measured with the NOAO APPHOT package, were subtracted to make $(U-R)$ colours for the entire R -selected sample to $R=27$ and that part of the U -selected sample which overlaps the R -band image to $U=25$. Median colours as a function of magnitude and colour histograms for several U - and R -selected subsamples are shown in Fig. 5. To $R=27$ mag, 73 per cent of the sources in the R -band image are detected with a confidence of 1σ or better in the U -band image. The median measured colour in the faintest U -band magnitude bin is $U-R=0.6$ mag. In the faintest R -band magnitude bin it is $U-R=1.3$ mag, although clearly there are large magnitude uncertainties in the U -band data for such faint objects.

4 RESULTS AND DISCUSSION

4.1 Colours and faint-end count slopes

The faint ends of the counts have different slopes in the U and R bands, which means, under fairly robust assumptions, the mean object colour must be a strong function of magnitude, a trend clearly visible in the colour–magnitude diagram. Since the night sky would be infinitely bright in the U band if this trend continued forever, at some faint magnitude the U -band count slope must break to match the R -




Figure 4. The R -band number counts. Open squares are the positive counts with the ‘noise’ counts (negative counts, see text) subtracted, from the data presented here; error bars are Poisson noise for the difference. Filled squares are the completeness-corrected counts (see text), also from the data presented here. The solid line is a fit to the corrected counts; it has slope $d \log N/dm = 0.334$. Dashed and dotted lines are the completeness-corrected counts of other authors. The Steidel & Hamilton (1993) counts have been converted to this magnitude system assuming $R \approx R_{AB} - 0.16$ for a typical object.




Figure 5. Histograms of $U-R$ colours (corrected to the standard U band) through 1.5-arcsec apertures for several U - (top) and R -selected (bottom) samples. The squares show median colours as a function of magnitude, medianed in groups of 61 in the U band and groups of 201 in the R band.

band slope and thereby end the blueing trend. One physical explanation for the blueing trend is that the fraction of the radiation owing to young stars may increase with apparent magnitude. In this scenario, the objects should get no bluer than about $f_v \propto v^0$, the ultraviolet spectral slope of a star-forming galaxy (Kinney et al. 1993), so a natural prediction is that the break in the U -band counts will appear when the median $(U-R)$ obtains the $f_v \propto v^0$ value, -0.4 mag. This should happen at $27 < U < 28$ mag. Very deep counts in the *Hubble Deep Field* (HDF) data (Williams et al. 1996), which reach point-source detection levels several magnitudes fainter than these observations, report a break at $U \sim 25.3$, although it is not robustly detected.

As is shown in Figs 3 and 4, the U -band counts found here are in good agreement with the previous study near this depth (Songaila et al. 1990), the R -band counts are also in very good agreement with previous studies (Tyson 1988; Steidel & Hamilton 1993; Smail et al. 1995; Metcalfe et al. 1995) as are the colours (Guhathakurta et al. 1990; Steidel & Hamilton 1993). The results here are similar to those of Guhathakurta et al. (1990) except that the latter have $< 2.7 \times 10^5$ source deg^{-2} and only identify sources to $R \approx 26$, although that limit is fuzzy because sources were identified by those authors not in a single band but in a summed UBR image.

4.2 The number problem

We count faint objects in the U and R bands to surface densities, integrated over flux, of $1.2 \times 10^5 \text{ deg}^{-2}$ to $U=25.5$ mag and $6.3 \times 10^5 \text{ deg}^{-2}$ to $R=27$ mag. These numbers corresponds to 0.5 and 2.5×10^{10} objects over the entire sky. These numbers are in excess of the total number of galaxies expected within the observable Universe in any

no-evolution or passive-evolution model (i.e., models in which the comoving number density of galaxies is conserved), for the following simple reason: if the local galaxy luminosity function is that given by Loveday et al. (1992) (or Mobasher, Sharples & Ellis 1993, or Lin et al. 1996) and galaxies exist down to luminosities 5 mag fainter than L^* , the space density of galaxies is $5.7 \times 10^{-2} h^3 \text{ Mpc}^{-3}$, where h is the Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$. In units of Hubble volume $V_H \equiv (c/H_0)^3$, this corresponds to a density of $1.5 \times 10^9 V_H^{-1}$. In an Einstein-de Sitter universe, $(\Omega_M, \Omega_\Lambda) = (1, 0)$, there is an all-sky comoving volume of only $4.2 V_H$ to $z=3$. The product of number density and volume is 6×10^9 galaxies of the entire sky, so there are four times too many sources observed in these data to be easily explained by naive models. The number of galaxies increases with apparent magnitude, so the problem is worse with deeper data: Cowie et al. (1995) show a factor of 10 times too many sources, and the counts in the HDF (Williams et al. 1996) show a factor of at least 15 times too many sources. We show that only a small fraction of the galaxies presented here lie beyond $z \approx 3$ because at this redshift the Lyman limit redshifts into the U band and the flux is cut-off by either self-absorption or absorption in intervening material (e.g., Guhathakurta et al. 1990; Steidel et al. 1995, 1996); 73 per cent of the sources to $R=27$ mag are detected in U_{13} . Of course these numbers are consistent with the population of galaxies at $z > 3$ found and confirmed spectroscopically by Steidel et al. (1995, 1996), because the high-redshift population is only a few per cent of the total source counts.

For $(\Omega_M, \Omega_\Lambda) = (0.05, 0)$ the comoving volume is $14 V_H$, for $(\Omega_M, \Omega_\Lambda) = (0.2, 0.8)$ it is $19 V_H$, so switching world models does not solve the problem unless one considers even more extreme world models that are almost certainly ruled out by gravitational lens statistics (Turner 1990). Several authors have found local luminosity function amplitudes higher than that of Loveday et al. (1992) by factors of a few (Marzke, Huchra & Geller 1994; Lilly et al. 1995; Small 1996) but factors large enough to solve the number problem would be surprising. There is also some disagreement over the faint-end slope α [in the parametrization $\phi(L) \propto L^\alpha$] of the local luminosity function (e.g., Lilly et al. 1995). If the slope is steeper than the standard flat value (i.e., if α is more negative than $\alpha = -1.0$), there could be significantly more galaxies in the local Universe than the above estimates suggest. For example, comparing with $\alpha = -1.0$, if $\alpha = -1.5$ but L^* and the bright-end amplitude are held fixed, there is a factor of ≈ 5 more galaxies to a luminosity limit of 5 mag fainter than L^* . This would go some way towards alleviating the number problem, although the slope is well-enough determined by local surveys that such a large discrepancy seems unlikely.

With current observational constraints, it is more natural to look to evolution in the sources to solve the number problem. It has been suggested that the galaxy merging rate is high, so large numbers of small galaxies at high redshift evolve into small numbers of large galaxies locally (Guidedone & Rocca-Volmerange 1991; Broadhurst, Ellis & Glazebrook 1992). It is possible that there is a large population of small galaxies which explode or evaporate after their first burst of star formation and supernovae (Babul & Rees 1992). Also, there is a less-explored possibility that galaxies

may form at high redshift with more-or-less their present-day masses but if star formation occurs in small, spatially isolated bursts (Katz 1992), each present-day galaxy would be observed as many different objects at large lookback time. There may be support for some of these models in the redshift distribution of faint galaxies; several authors have found a steepening of the faint-end slope of the luminosity function with redshift (Eales 1993; Lilly et al. 1995; Ellis et al. 1996), and enormous information will come from the current generation of super-deep redshift surveys (e.g. Cowie et al. 1996; Cohen et al. 1996a,b; Koo et al. 1996).

4.3 Stronger redshift limits with the *HST*

The ultraviolet sensitivities of the WFPC2 and STIS instruments on *HST* suggest the extension of this technique – limiting redshift distributions by looking for the Lyman limit break – to lower redshift. In fact, the broad-band UV filters of these instruments can actually be used to locate the Lyman break over a range of redshifts from 0.9 to 1.8, a range which is difficult to identify at present even with 10-m-class telescopes because there are very few lines in the useful window of ground-based visual spectroscopy. With a modest amount of observing time it will be possible to obtain at least statistical redshift distributions significantly deeper than the practical limits of ground-based spectroscopy even if there were spectroscopic features in this redshift range. Such observations are particularly crucial since many of the above-mentioned models make very different predictions for the fraction of galaxies at $z > 1$ and $z > 2$.

ACKNOWLEDGMENTS

We are grateful to the W. M. Keck Foundation for the vision to fund the construction of the W. M. Keck Observatory. We benefited from helpful conversations with Kurt Adelberger, John Gizis, Tomislav Kundić, Gerry Neugebauer, Sterl Phinney and Chuck Steidel, and from comments from an anonymous referee. We are grateful for financial support from NSF grant AST 92-23370 (DWH, RB), NSF grant AST-91-57412 and the Bressler Foundation (MAP), support through a PPARC Advanced Fellowship (IS), and grants from NSF and NASA (BTS). The calibrated images used for this investigation will be made available to the public; contact the authors for more information.

This work is based on observations obtained at the W. M. Keck Observatory, which is operated jointly by the California Institute of Technology and the University of California; and at the Palomar Observatory, which is owned and operated by the California Institute of Technology.

REFERENCES

- Babul A., Rees M. J., 1992, MNRAS, 255, 346
 Bertin E., Arnouts S., 1996, A&AS, 117, 393
 Broadhurst T. J., Ellis R. S., Glazebrook K., 1992, Nat, 355, 55
 Burstein D., Heiles C., 1982, AJ, 87, 1165
 Cohen J. G., Hogg D. W., Pahre M. A., Blandford R. D., 1996a, ApJ, 462, L9
 Cohen J. G., Cowie L. L., Hogg D. W., Songaila A., Blandford R. D., Hu E. M., Shopbell P., 1996b, ApJ, 471, L5
 Cowie L. L., Hu E. M., Songaila A., 1995, AJ, 110, 1576
 Cowie L. L., Songaila A., Hu E. M., Cohen J. G., 1996, AJ, 112, 839
 Djorgovski S. et al., 1995, ApJ, 438, L13
 Dressler A., 1993, in Neugebauer G., eds., Palomar Observatory Annual Report 1993, p. 2
 Eales S., 1993, ApJ, 404, 51
 Ellis R. S., Colless M. M., Broadhurst T. J., Heyl J. S., Glazebrook K., 1996, MNRAS, 280, 235
 Griffiths R. E. et al., 1994, ApJ, 435, L19
 Guhathakurta P., Tyson J. A., Majewski S. R., 1990, ApJ, 357, L9
 Guiderdone B., Rocca-Volmerange B., 1991, A&A, 252, 435
 Hall P., Mackay C. D., 1984, MNRAS, 210, 979
 Im M., Casertano S., Griffiths R. E., Ratnatunga K. U., Tyson J. A., 1995, ApJ, 441, 494
 Jones L. R., Fong R., Shanks T., Ellis R. S., Peterson B. A., 1991, MNRAS, 249, 481
 Katz N., 1992, ApJ, 391, 502
 Kinney A. L., Bohlin R. C., Calzetti D., Panagia N., Wyse R. F. G., 1993, ApJS, 86, 5
 Koo D. C., 1986, ApJ, 311, 651
 Koo D. C., Kron R. G., 1992, ARA&A, 30, 613
 Koo D. C. et al., 1996, ApJ, 469, 535
 Lilly S. J., Tresse L., Hammer F., Le Fevre O., Crampton D., 1995, ApJ, 455, 108
 Lin H., Kirshner R. P., Smetman S. A., Landy S. D., Oemler A., Tucker D. L., Schechter P. L., 1996, ApJ, 464, 60
 Loveday J., Peterson B. A., Efstathiou G., Maddox S. J., 1992, ApJ, 390, 338
 Marzke R. O., Huchra J. P., Geller M. J., 1994, AJ, 108, 437
 Metcalfe N., Shanks T., Fong R., Roche N., 1995, MNRAS, 273, 257
 Mobasher B., Sharples R. M., Ellis R. S., 1993, MNRAS, 263, 560
 Oke J. B. et al., 1995, PASP, 107, 375
 Smail I., Hogg D. W., Yan L., Cohen J. G., 1995, ApJ, 449, L105
 Small T. A., 1996, PhD thesis, California Inst. Technol.
 Songaila A., Cowie L. L., Lilly S. J., 1990, ApJ, 348, 371
 Steidel C. C., Hamilton D., 1993, AJ, 105, 2017
 Steidel C. C., Pettini M., Hamilton D., 1995, AJ, 110, 2017
 Steidel C. C., Giavalisco M., Pettini M., Dickinson M., Adelberger K., 1996, ApJ, 462, L17
 Turner E. L., 1990, ApJ, 365, L43
 Tyson J. A., 1988, AJ, 96, 1
 Williams R. et al., 1996, AJ, 112, 1335