

Convergence of the Discrete Ordinates  
Method for the Transport Equation

P. M. Anselone\* and A. G. Gibbs\*\*

1. Introduction

The transport equation (transfer equation, linear Boltzmann equation) governs the distribution of neutrons in a nuclear reactor or of radiation in a stellar or planetary atmosphere. It is an integro-differential equation of some complexity in its more general forms. Primary independent variables are position and direction, speed (alternatively energy, frequency or wavelength), plus possibly time. The differential operator in the equation describes attenuation due to both absorption and scattering away from a particular direction. The integral operator describes scattering into a particular direction from other directions and production by fission. Source terms may be present. On any boundary surface there are boundary conditions corresponding to incoming directions. For time-dependent problems there are initial conditions.

---

\* Oregon State University, Corvallis, Oregon 97331

\*\* Battelle, Pacific Northwest Laboratory, Richland, Washington 99352

Supported by the U. S. Atomic Energy Commission at the Pacific Northwest Laboratory, Richland, Washington, and the Union Carbide Nuclear Division, Oak Ridge, Tenn.

## DISCLAIMER

**This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.**

## **DISCLAIMER**

**Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.**

Since the transport equation can be solved explicitly only in a few very special cases, approximation methods are commonly used. The discrete ordinates method involves the replacement of the integral operator by a numerical integral operator in order to obtain differential equations for approximate solutions. Empirically, this method yields very satisfactory results for a large class of transport problems. However, a complete and rigorous convergence and error analysis is presently available only in some rather idealized cases.

The discrete ordinates method was first applied by Wick [18] and by Chandrasekhar [7, 8] to particular steady-state problems from astrophysics with homogeneous media bounded by single planes or parallel planes (half-spaces or slabs), usually under conditions of isotropic scattering and isotropic sources. Convergence theorems for the approximations were obtained for various problems of this type with isotropic scattering by Anselone [1-5], Kofink [13], Keller [10, 11] and Wendroff [17]. Subsequently, convergence theorems have been obtained for certain transport problems involving anisotropic scattering, other geometries, and time-dependence, by Keller [12], Nestell [16], Madsen [14] and Wilson [20].

We shall survey these contributions from an abstract and unifying point of view in order to isolate what is essential to the convergence of the discrete ordinates approximations. In the process, methods of analysis

will be developed which appear to be applicable to very general transport problems and to other integro-differential equations. The discussion is in terms of a sequence of model problems of increasing complexity. All functions considered will be real. For mathematical convenience and physical relevance specified functions are continuous and usually nonnegative. We remark that the problems surveyed here involve discrete ordinates approximations only for the angular integrals, and thus do not represent the most general case of discretization as practiced, say, in many nuclear reactor applications today, where the spatial variable is also discretized. However, as noted below, the powerful analytical tools developed in applications to angular discrete ordinates approximations also show promise for investigations of the more general case.

#### 2a. An Isotropic Transport Problem in a Finite Slab

Consider the problem for  $\phi(x, \mu)$  with  $a \leq x \leq b$ ,  $-1 \leq \mu \leq 1$ , given by

$$\mu \frac{\partial \phi(x, \mu)}{\partial x} + \phi(x, \mu) = \frac{c}{2} \int_{-1}^1 \phi(x, \mu') d\mu' + g(x), \quad (2.1)$$

$$\phi(a, \mu) = 0 \quad \text{for } \mu > 0, \quad \phi(b, \mu) = 0 \quad \text{for } \mu < 0,$$

where  $\phi(x, \mu)$  is the one-speed angular flux at depth  $x$  in a homogeneous slab in a direction making an angle  $\theta = \cos^{-1} \mu$  with the positive  $x$ -axis. The positive



number  $c$  is of order 1 and represents the mean number of secondary neutrons per collision. In the above equation, distance has been measured in units of neutron mean free paths, i.e.,  $x = \sigma\tau$ , where  $\sigma$  is the total cross section and  $\tau$  is the actual distance.

The boundary conditions correspond to a situation with no reflection of neutrons at the surface and no incident neutrons. Actually, a non-zero incident flux boundary condition could have been assumed. However, such a problem can be reduced to (2.1) for  $\phi_1 = \phi - \phi_0$ , where  $\phi_0$  satisfies the homogeneous equation

$$\mu \frac{\partial \phi_0(x, \mu)}{\partial x} + \phi_0(x, \mu) = 0$$

and the given boundary conditions. If this reduction is not made, then an unnecessary approximation is introduced for  $\phi_0$ , which can be found explicitly. For the same reason, zero boundary conditions can and should be assumed also in applications of the discrete ordinates method to more general transport problems.

To proceed, assume a numerical integration rule such that, for any continuous function  $h(\mu)$ ,  $-1 \leq \mu \leq 1$ ,

$$\sum_{j=\pm 1}^{\pm n} w_{nj} h(\mu_{nj}) \rightarrow \int_{-1}^1 h(\mu) d\mu, \quad n \rightarrow \infty,$$

$$w_{n,-j} = w_{n,j} > 0, \quad \mu_{n,-j} = -\mu_{n,j},$$

$$0 < \mu_{n1} < \mu_{n2} \cdots < \mu_{nn} \leq 1.$$

The Gauss quadrature formula is an example. The symmetry conditions on the  $w_{nj}$  and  $\mu_{nj}$  are a convenience rather than a necessity.

Discrete ordinates approximations  $\phi_n(x, \mu)$ ,  $n = 1, 2, \dots$ , satisfy

$$\mu \frac{\partial \phi_n(x, \mu)}{\partial x} + \phi_n(x, \mu) = \frac{c}{2} \sum_{j=\pm 1}^{\pm n} w_{nj} \phi_n(x, \mu_{nj}) + g(x), \quad (2.2)$$

$$\phi_n(a, \mu) = 0 \quad \text{for } \mu > 0, \quad \phi_n(b, \mu) = 0 \quad \text{for } \mu < 0.$$

For  $\mu = \mu_{ni}$ ,  $i = \pm 1, \dots, \pm n$ , this is a system of ordinary differential equations for  $\phi_n(x, \mu_{ni})$ . Then (2.2) yields  $\phi_n(x, \mu)$ .

The problems for  $\phi$  and  $\phi_n$  have equivalent integral equation formulations which are more convenient for the convergence analysis. Let

$$f(x) = \frac{c}{2} \int_{-1}^1 \phi(x, \mu) d\mu + g(x), \quad (2.3)$$

$$f_n(x) = \frac{c}{2} \sum_{j=\pm 1}^{\pm n} w_{nj} \phi(x, \mu_{nj}) + g(x). \quad (2.4)$$

From (2.1) and (2.3), it follows that

$$\phi(x, 0) = f(x),$$

(2.5)

$$\phi(x, \mu) = \frac{1}{\mu} \int_z^x e^{(y-x)/\mu} f(y) dy, \quad \begin{array}{l} z = a, \mu > 0, \\ z = b, \mu < 0. \end{array}$$

The equations for  $\phi_n$  in terms of  $f_n$  are identical.

From (2.3) and (2.5),

$$(2.6) \quad f(x) - \frac{c}{2} \int_a^b E_1(|x-y|) f(y) dy = g(x),$$

where  $E_1$  is the exponential integral function of order one,

$$(2.7) \quad E_1(s) = \int_0^1 e^{-s/\mu} \mu^{-1} d\mu, \quad s > 0,$$

which has a logarithmic singularity at  $s = 0$ . Similarly,

$$(2.8) \quad f_n(x) - \frac{c}{2} \int_a^b E_{nl}(|x-y|) f_n(y) dy = g(x),$$

where  $E_{nl}$  is a numerical integration approximation to  $E_1$ :

$$(2.9) \quad E_{nl}(s) = \sum_{j=1}^n w_{nj} \frac{e^{-s/\mu_{nj}}}{\mu_{nj}}, \quad s \geq 0,$$

$$(2.10) \quad E_{nl}(s) \rightarrow E_1(s) \quad \text{uniformly for } s \geq \epsilon \text{ as } n \rightarrow \infty$$

for each  $\epsilon > 0$ . Thus, the discrete ordinates method is seen to be equivalent to the approximation of the integral operator in (2.6) by the integral operator in (2.8).

Express (2.6) and (2.8) in operator form

$$(2.11) \quad (I - K) f = g, \quad (I - K_n) f_n = g$$

on the space  $C[a, b]$  with the max norm. Then  $K$  and  $K_n$  are Fredholm integral operators with kernels

$$k(x, y) = \frac{c}{2} E_1(|x-y|), \quad k_n(x, y) = \frac{c}{2} E_{nl}(|x-y|).$$



These are bounded linear operators on  $C[a, b]$  and for  $c$  not too large (the case of a subcritical medium, assumed here) we have

$$\|K\| = \max_x \int_a^b k(x, y) dy < 1.$$

Hence,  $(I - K)^{-1}$  exists,

$$\|(I - K)^{-1}\| \leq (1 - \|K\|)^{-1},$$

and  $f$  and  $\phi$  are uniquely determined.

From (2.10),

$$\|K_n - K\| \rightarrow 0, \quad \|K_n\| \rightarrow \|K\| \quad \text{as } n \rightarrow \infty.$$

Thus, we are in the realm of standard operator approximation theory. For  $n$  sufficiently large,  $\|K_n\| < 1$ ,  $(I - K_n)^{-1}$  exists and is bounded uniformly in  $n$ , and

$$(2.12) \quad (I - K_n)^{-1} - (I - K)^{-1} = (I - K_n)^{-1} (K_n - K) (I - K)^{-1},$$

$$(2.13) \quad \|(I - K_n)^{-1} - (I - K)^{-1}\| \leq \|(I - K_n)^{-1}\| \|K_n - K\| \|(I - K)^{-1}\|,$$

$$(2.14) \quad \|(I - K_n)^{-1} - (I - K)^{-1}\| \leq \frac{\|(I - K)^{-1}\|^2 \|K_n - K\|}{1 - \|(I - K)^{-1}\| \|K_n - K\|},$$

$$(2.15) \quad \|(I - K_n)^{-1} - (I - K)^{-1}\| \leq \frac{\|(I - K_n)^{-1}\|^2 \|K_n - K\|}{1 - \|(I - K_n)^{-1}\| \|K_n - K\|},$$

$$(2.16) \quad \|(I - K_n)^{-1} - (I - K)^{-1}\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

The error bound in (2.14) is "theoretical" in the sense

that it involves  $(I - K)^{-1}$ , whereas (2.15) is "practical" because it depends on  $(I - K_n)^{-1}$ , and can thus be computed. The bound (2.13) is of mixed type. It follows from (2.16) and (2.5)ff that  $f_n \rightarrow f$  and  $\phi_n \rightarrow \phi$  uniformly as  $n \rightarrow \infty$ .

This is essentially the path followed by Anselone [4, 5] and by Keller [11], although Keller's analysis was less abstract and he defined the discrete ordinates approximations only at the quadrature points. Previously, Keller [10] and Wendroff [17] treated the differential operators directly without inverting them. The results were less satisfactory:  $L^2$  convergence on quadrature points, and uniform convergence on quadrature points under a restrictive assumption. Kofink [13] established  $L^2$  convergence on quadrature points by exploiting an equivalence between the discrete ordinates method and the spherical harmonics method.

## 2b. An Isotropic Transport Problem in a Half Space

Anselone [1-4]—also derived uniform convergence theorems for isotropic transport problems in the case of semi-infinite slabs ( $0 \leq x \leq \infty$ ). In particular, the classical Milne problem leads to the homogeneous equation  $(I - K)f = 0$ , where  $K$  is the same integral operator as above. The Wiener-Hopf method, based on Fourier transforms, was devised originally to solve this equation.

A more direct method, which anticipated later theories of positive and monotone operators, was given by Hopf [9] for the case  $c = 1$ . It involves a change of variable  $f(x) = x + q(x)$ . Then  $q$  satisfies the inhomogeneous equation  $(I - K)q = E_3$ , where  $E_3$  is the exponential integral function of order 3,

$$E_3(s) = \int_0^1 e^{-s/\mu} \mu d\mu, \quad s \geq 0.$$

A nonnegative solution  $q$  is sought in the space of bounded continuous functions on  $[0, \infty)$ . Although now  $\|K\| = 1$ , monotonicity considerations yield a unique solution  $q$  given by the uniformly convergent Neumann series

$$q = \sum_{m=0}^{\infty} K^m E_3.$$

The discrete ordinates approximation problem can be recast in the form  $(I - K_n)f_n = 0$ . Let  $f_n(x) = x + q_n(x)$ . Then  $(I - K_n)q_n = E_{n3}$ , where  $E_{n3}$  is a numerical integration approximation to  $E_3$ , and  $q_n$  is given by

$$q_n = \sum_{m=0}^{\infty} K_n^m E_{n3}.$$

A detailed term by term analysis yields  $q_n \rightarrow q$  uniformly as  $n \rightarrow \infty$ . It follows that  $f_n \rightarrow f$  and  $\phi_n \rightarrow \phi$  uniformly.

### 3a. An Anisotropic Transport Problem in a Finite Slab

Consider the problem for  $\phi(x, \mu)$  with  $a \leq x \leq b$ ,  $-1 \leq \mu \leq 1$ , given by

$$\mu \frac{\partial \phi(x, \mu)}{\partial x} + \phi(x, \mu) = \frac{1}{2} \int_{-1}^1 p(x, \mu, \mu') \phi(x, \mu') d\mu' + g(x, \mu), \quad (3.1)$$

$$\phi(a, \mu) = 0 \quad \text{for } \mu > 0, \quad \phi(b, \mu) = 0 \quad \text{for } \mu < 0.$$

Here,  $x$  represents the optical depth measured from the plane of the origin, i.e.,

$$x(\tau) = \int_0^\tau \sigma(\tau') d\tau',$$

with  $\sigma(\tau)$  the total cross section at position  $\tau$ , and  $\tau$  the distance measured from 0. The differential kernel  $p(x, \mu, \mu') d\mu'$  represents the average number of neutrons emerging with direction cosines in  $d\mu'$ , following a collision by a neutron with direction cosine  $\mu'$  at position  $x$ . The function  $g$  represents sources within the slab. We assume here that both  $p$  and  $g$  are continuous functions of  $x$  and  $\mu$ .

Let

$$(3.2) \quad f(x, \mu) = \frac{1}{2} \int_{-1}^1 p(x, \mu, \mu') \phi(x, \mu') d\mu'.$$

Then the problem is expressed in operator form as

$$(3.3) \quad D\phi = f, \quad f = L\phi + g,$$

where  $f, g \in C(X)$  with  $X = [a, b] \times [-1, 1]$ . Under reasonable conditions on  $\phi$ ,  $M = D^{-1}$  exists as an operator on  $C(X)$ , and  $\phi = Mf = D^{-1}f$  is given by

$$\begin{aligned} \phi(x, 0) &= f(x, 0), \\ (3.4) \quad \phi(x, \mu) &= \frac{1}{\mu} \int_z^x e^{(x' - x)/\mu} f(x', \mu) dx', \quad \begin{array}{l} z = a \text{ for } \mu > 0, \\ z = b \text{ for } \mu < 0. \end{array} \end{aligned}$$

Therefore, an equivalent formulation of the problem is

$$(3.5) \quad \phi = Mf, \quad (I - K)f = g, \quad K = LM.$$

It can be shown that  $K$  is a bounded integral operator which maps  $C(X)$  into  $C(X)$ . Since  $K$  has a non-negative kernel

$$\|K\| = \|Ke\|, \quad e \in C(X), \quad e \equiv 1.$$

If  $p$  is not too large, then  $\|K\| < 1$ , which we assume in what follows. Then  $(I - K)^{-1}$  exists as a bounded operator on  $C(X)$ , and  $f, \phi$  are uniquely determined.

Assume a convergent, positive quadrature rule:

$$\sum_{j=1}^n w_{nj} h(\mu_{nj}) \rightarrow \int_0^1 h(\mu) d\mu \quad \text{as } n \rightarrow \infty, \quad h \in C[-1, 1],$$

$$w_{nj} > 0, \quad 1 \leq j \leq n, \quad n = 1, 2, \dots$$

In particular, for  $h = e \equiv 1$ ,

$$\sum_{j=1}^n w_{nj} \rightarrow 1 \quad \text{as } n \rightarrow \infty.$$

Hence, there exists  $B < \infty$  such that

$$\sum_{j=1}^n w_{nj} \leq B, \quad n = 1, 2, \dots$$

The discrete ordinates approximations  $\phi_n$  satisfy

$$\mu \frac{\partial \phi_n(x, \mu)}{\partial x} + \phi_n(x, \mu) = \frac{1}{2} \sum_{j=1}^n w_{nj} p(x, \mu, \mu_{nj}) \phi(x, \mu_{nj}) + g(x, \mu), \quad (3.6)$$

$$\phi_n(a, \mu) = 0 \quad \text{for } \mu > 0, \quad \phi_n(b, \mu) = 0 \quad \text{for } \mu < 0.$$

Equivalent formulations are

$$(3.7) \quad D\phi_n = f_n, \quad f_n = L_n \phi_n + g,$$

$$(3.8) \quad \phi_n = M f_n, \quad (I - K_n) f_n = g, \quad K_n = L_n M,$$

where  $K_n$  is a bounded linear operator on  $C(X)$ .

Since the "kernel" of  $K_n$  is positive,  $\|K_n\| = \|K_n e\|$ , where  $e \equiv 1$ .

It follows from (3.5) that  $K_n \rightarrow K$ , i.e.,  $\|K_n h - Kh\| \rightarrow 0$  as  $n \rightarrow \infty$  for each  $h \in C(X)$ . However,  $\|K_n - K\| \not\rightarrow 0$  in the anisotropic case, so the standard operator approximation theory used in Section 2 above is not applicable. An alternative course of action is pursued here.

It follows from  $K_n \rightarrow K$ ,  $\|K\| = \|Ke\|$  and  $\|K_n\| = \|K_n e\|$  that  $\|K_n\| \rightarrow \|K\|$ . Recall that  $\|K\| < 1$ . Hence, for  $n$  sufficiently large,  $\|K_n\| < 1$ ,  $(I - K_n)^{-1}$  exists and is bounded uniformly in  $n$ , and

$$(3.9) \quad (I - K_n)^{-1} - (I - K)^{-1} = (I - K_n)^{-1} (K_n - K) (I - K)^{-1},$$

$$(3.10) \quad f_n - f = (I - K_n)^{-1} (K_n f - Kf),$$

$$(3.11) \quad \|f_n - f\| \leq \| (I - K_n)^{-1} \| \|K_n f - Kf\| .$$

It follows that  $f_n \rightarrow f$  and  $\phi_n \rightarrow \phi$  uniformly as  $n \rightarrow \infty$ . The error bound in (3.11) is of mixed type, neither purely theoretical nor purely practical.

The fact that  $K$  and  $K_n$  are positive operators is essential to the foregoing analysis, which is an abstract version of that carried out by Keller [12] in a more classical spirit.

### 3b. An Anisotropic Transport Problem in a Half Space

We remark that the case of anisotropic scattering problems in a half-space was treated by Nestell [16] by an adaptation of the positive operator theory of Hopf. That work is an extension of the isotropic scattering problem discussed in Section 2b above.

## 4. Collectively Compact Operator Approximation Theory

We shall consider the same problem as in 3 from another point of view, which yields both theoretical and practical (computable) error bounds. The analysis is based on collectively compact operator approximation theory (cf. Anselone [6]).

It can be shown that the operator  $K$  in (3.4) is compact:  $\{Kh : \|h\| \leq 1\}$  is relatively compact or, equivalently, bounded and equicontinuous. The sequence  $\{K_n : n \geq 1\}$  is collectively compact:  $\{K_n h : \|h\| \leq 1, n \geq 1\}$  is relatively compact. Thus, we have



$K_n \rightarrow K$ ,  $K$  compact,  $K_n$  collectively compact.

These are the hypotheses for a general operator approximation theory in a Banach space setting. Some of the main conclusions of the theory are as follows.

The operator  $(I - K)^{-1}$  exists iff for  $n$  sufficiently large  $(I - K_n)^{-1}$  exists and is bounded uniformly in  $n$ , in which case

$$(I - K_n)^{-1} \rightarrow (I - K)^{-1}.$$

Let  $f = (I - K)^{-1}g$  and  $f_n = (I - K_n)^{-1}g$ . Then  $\|f_n - f\| \rightarrow 0$ . Let

$$\Delta_n = \|(I - K)^{-1}\| \|(K_n - K)K_n\|,$$

$$\Delta^n = \|(I - K_n)^{-1}\| \|(K_n - K)K\|.$$

Then

$$\Delta_n \rightarrow 0, \quad \Delta^n \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

$$\|f_n - f\| \leq \frac{\|(I - K)^{-1}\| \|K_n g - Kg\| + \Delta_n \|f\|}{1 - \Delta_n} \quad \text{for } \Delta_n < 1,$$

$$\|f_n - f\| \leq \frac{\|(I - K_n)^{-1}\| \|K_n g - Kg\| + \Delta^n \|f_n\|}{1 - \Delta^n} \quad \text{for } \Delta^n < 1.$$

Moreover,  $(I - K)^{-1}$  exists whenever  $(I - K_n)^{-1}$  exists and  $\Delta^n < 1$ . Thus the existence of  $f = (I - K)^{-1}g$  can be inferred from approximations.

The convergence of the discrete ordinates approximations was established by means of the collectively compact theory by Nestell [16] for an anisotropic transport problem in a finite slab with  $p = p(\mu, \mu')$  and by Nelson [15] in greater generality. Both authors worked directly with explicit representations of  $K$  and  $K_n$ . Since  $K$  is an integral operator on functions of two variables, its kernel involves four variables. The kernel has a weak singularity. The definition of  $K_n$  is similar, with the  $\mu$ -integral replaced by a sum. The complications of  $K$  and  $K_n$ , particularly the singularity, make it difficult to see what makes the analysis go through. We propose a different approach here, which is more transparent and extends more readily to a larger class of transport problems. By (3.5) and (3.7),

$$K = LM, \quad K_n = L_n M$$

where  $M$  is defined in (3.4) and

$$(L\phi)(x, \mu) = \frac{1}{2} \int_{-1}^1 p(x, \mu, \mu') \phi(x, \mu') d\mu',$$

$$(L_n \phi)(x, \mu) = \frac{1}{2} \sum_{j=1}^n w_{nj} p(x, \mu, \mu_{nj}) \phi(x, \mu_{nj}).$$

The basis of the analysis is to take advantage of the fact that only  $L$ , which has a continuous kernel, is approximated in order to define  $K_n = L_n M$ . The singularity comes from  $M$ , which can be treated separately.

Consider  $\phi = Mf$  for  $f \in C(X)$ . From (3.4),

$$\phi(x, 0) = f(x, 0),$$

$$\phi(x, \mu) = \int_0^{\frac{(x-z)}{\mu}} e^{-t} f(x - t\mu, \mu) dt, \quad \begin{array}{l} z = a \text{ for } \mu > 0, \\ z = b \text{ for } \mu < 0. \end{array}$$

It follows that  $\phi$  is continuous except perhaps at  $(a, 0)$  and  $(b, 0)$ . The boundary conditions

$$\phi(a, \mu) = 0 \text{ for } \mu > 0, \quad \phi(b, \mu) = 0 \text{ for } \mu < 0,$$

are satisfied and

$$\phi(x, \mu) \rightarrow \phi(a, 0) \quad \text{as } x \rightarrow a, \mu \rightarrow 0-,$$

$$\phi(x, \mu) \rightarrow \phi(b, 0) \quad \text{as } x \rightarrow b, \mu \rightarrow 0+,$$

$$\phi(x, \mu) - (1 - e^{(a-x)/\mu})\phi(a, 0) \rightarrow 0 \quad \text{as } x \rightarrow a, \mu \rightarrow 0+,$$

$$\phi(x, \mu) - (1 - e^{(b-x)/\mu})\phi(b, 0) \rightarrow 0 \quad \text{as } x \rightarrow b, \mu \rightarrow 0-.$$

Thus,  $R(M)$ , the range of  $M$ , consists of bounded functions which are continuous except perhaps at two points. By routine arguments,  $L$  and  $L_n$  map  $R(M)$  into  $C(X)$ . Define the domains of  $L$  and  $L_n$  by  $D(L) = D(L_n) = R(M)$ . Then the operators  $K = LM$  and  $K_n = L_n M$  map  $C(X)$  into  $C(X)$ .

It follows easily from  $K_n - K = (L_n - L)M$  and the properties of  $R(M)$  that  $K_n \rightarrow K$ . To prove that  $K$  is compact and  $\{K_n\}$  is collectively compact, consider  $Kf = LMf$  and  $K_n f = L_n Mf$  for  $\|f\| \leq 1$ . As above, let  $\phi = Mf$ . Then for any  $\epsilon > 0$  and any  $\nu > 0$  there

exists  $\delta = \delta(\epsilon, \nu)$  independent of  $f$  such that

$$|\phi(x, \mu) - \phi(x', \mu)| < \epsilon \quad \text{for } |x - x'| < \delta, \quad |\mu| > \nu.$$

This is an equicontinuity property in  $x$ , uniform for  $|\mu| > \nu$ . Essentially,  $M$  is a compact operator with respect to  $x$ , uniformly for each  $\nu > 0$ . Similarly, the operator  $L$  is compact with respect to  $\mu$  uniformly for  $a \leq x \leq b$ . Examination of  $(L\phi)(x, \mu) - (L\phi)(x, \mu')$  and  $(L\phi)(x, \mu) - (L\phi)(x', \mu)$  reveals that the sets

$$\{Kf = Lmf : \|f\| \leq 1\}, \quad \{K_n f = L_n Mf : \|f\| \leq 1, n \geq 1\}$$

are bounded and equicontinuous. Therefore,  $K$  is compact and  $K_n$  is collectively compact.

The error bounds in the collectively compact theory depend particularly on

$$\|(K_n - K)g\| = \|(L_n - L)Mg\|,$$

$$\|(K_n - K)K\| = \|(L_n - L)MLM\|,$$

$$\|(K_n - K)K_n\| = \|(L_n - L)ML_n M\|.$$

Calculations based on these equations should yield sharper bounds than would be obtained otherwise.

##### 5. A Three-Dimensional Anisotropic Transport Problem

Let  $\Gamma$  be a domain in  $R^3$  with boundary  $\partial\Gamma$  which, at least piece-wise, has a well-defined tangent plane at each point. Directions will be denoted by  $\Omega \in S$ , the

unit sphere in  $\mathbb{R}^3$ . Consider the problem for  $\phi(x, \Omega)$ ,  $x \in \Gamma$ ,  $\Omega \in \mathbb{S}$ :


$$\Omega \cdot \nabla \phi(x, \Omega) + \sigma(x, \Omega)\phi(x, \Omega) = f(x, \Omega), \quad (5.1)$$

$$f(x, \Omega) = \int_{\mathbb{S}} p(x, \Omega, \Omega')\phi(x, \Omega')d\Omega' + g(x, \Omega),$$

with  $\phi(x, \Omega) = 0$  for  $x \in \partial\Gamma$  and  $\Omega$  an incoming direction. The gradient operator  $\nabla$  acts with respect to  $x$ . Assume that  $f$ ,  $g$  and  $p$  are continuous and that  $p$  is nonnegative. In operator form (5.1) becomes

$$(5.2) \quad D\phi = f, \quad f = L\phi + g.$$

Motivated by the fact that  $\Omega \cdot \nabla$  is the directional derivative in the direction  $\Omega$ , we introduce additional notation as illustrated in the following diagram.



The ray through  $x$  in the direction  $-\Omega$  is given by  $x - s\Omega$ ,  $s \geq 0$ . Let  $d(x, \Omega) = \|x - y\|_2$ , the Euclidean distance from  $x$  to  $\partial\Gamma$  along the ray.

An equivalent formulation of  $D\phi = f$  is

$$(5.3) \quad -\frac{d}{ds} \phi(x - s\Omega, \Omega) + \sigma(x - s\Omega, \Omega) \phi(x - s\Omega, \Omega) = f(x - s\Omega, \Omega).$$

Solve (5.3) to obtain

$$(5.4) \quad \phi(x, \Omega) = \int_0^{d(x, \Omega)} f(x - s\Omega, \Omega) e^{-\int_0^s \sigma(x - r\Omega, \Omega) dr} ds.$$

Thus,  $M = D^{-1}$  exists as an operator on  $C(X)$ , where  $X = \bar{\Gamma} \times S$ , and  $\phi = Mf$  is given by (5.4). The original problem now becomes

$$(5.5) \quad \phi = Mf, \quad (I - K)f = g, \quad K = LM,$$

where  $L$  is the integral operator in (4.1).

Discrete ordinates approximations  $\phi_n(x, \Omega)$  satisfy

$$(5.6) \quad D\phi_n = f_n, \quad f_n = L_n \phi_n + g,$$

where  $L_n$  is a numerical integration approximation to  $L$  defined in terms of a positive convergent quadrature rule. Now-familiar reasoning yields the equivalent formulation of (5.6),

$$(5.7) \quad \phi_n = Mf_n, \quad (I - K_n)f_n = g, \quad K_n = L_n M.$$

The tasks are the same as before, namely to prove that  $(I - K)^{-1}$  exists,  $(I - K_n)^{-1}$  exists for  $n$  sufficiently large, and

$$(I - K_n)^{-1} \rightarrow (I - K)^{-1},$$

which implies that  $f_n \rightarrow f$  and  $\phi_n \rightarrow \phi$  uniformly. If  $\|K\| < 1$  and  $K_n \rightarrow K$ , then these conclusions follow as in Section 3. If also  $K$  is compact and  $\{K_n\}$  is collectively compact, then the error bounds of Section 4 are obtained. This program will be undertaken in a forthcoming sequel.

The collectively compact operator approximation theory may also be applicable to the case where the approximations are made with respect to the spatial variable, either by approximating the detailed variation of  $\sigma(x, \Omega)$  and  $p(x, \Omega, \Omega')$  with respect to position  $x$  by appropriate smoothed functions, and/or by carrying out the spatial integrations by some quadrature approximation. In this more general setting, we have  $M$  approximated by  $M_j$  as well as  $L$  approximated by  $L_n$ . Thus, it will be necessary to consider  $L_n M_j \rightarrow LM$  in an appropriate sense.

## 6. A Positive Operator Approach

Explicit results for the three dimensional problem can be obtained rather easily if an additional assumption is made. Consider the problem of Section 5 in the form

$$(6.1) \quad T\phi = g, \quad T = D - L$$

under the restriction

$$(6.2) \quad \sigma(x, \Omega) - \int_S p(x, \Omega, \Omega') d\Omega' > c > 0, \quad x \in \Gamma, \Omega \in S.$$



Physically, this requirement means that absorption dominates production by fission, i.e., that  $\sigma_a > (\nu - 1)\sigma_f$ , where  $\sigma_a$  and  $\sigma_f$  are the absorption and fission cross sections, and  $\nu$  is the mean number of neutrons produced produced per fission. This inequality is always satisfied in a non-multiplying medium ( $\sigma_f = 0$ ) as long as some absorption is present ( $\sigma_a > 0$ ); however, it may not be satisfied in some nuclear reactor applications.

Suppose that for each  $g \in C(X)$  the equation  $T\phi = g$  has a unique solution  $\phi = T^{-1}g$  such that  $\phi \geq 0$  if  $g \geq 0$ . Then  $T^{-1}$  is a positive operator from  $C(X)$  to an appropriate solution space. Let  $e(x, \Omega) \equiv 1$  on  $X$ . Then (6.2) is equivalent to

$$(6.3) \quad Te > c > 0 \quad \text{on } X.$$

By an elementary argument,  $\phi = T^{-1}g$  is bounded for each  $g \in C(X)$  and  $\|\phi\| < c^{-1} \|g\|$  in terms of the sup norm. Thus,  $T^{-1}$  maps  $C(X)$  onto a subspace of the bounded functions on  $X$ ,  $T^{-1}$  is bounded, and  $\|T^{-1}\| = \|T^{-1}e\| < 1/c$ .

The discrete ordinates approximations  $\phi_n$  satisfy

$$(6.4) \quad T_n \phi_n = g, \quad T_n = D - L_n.$$

As above, suppose that  $T_n^{-1}$  exists and is a positive operator on  $C(X)$ . Since  $L_n \rightarrow L$ , we have  $T_n \rightarrow T$  and, in particular,  $T_n e \rightarrow Te$ . Therefore, for  $n$  sufficiently large,

$$(6.5) \quad T_n e > c > 0 \quad \text{on } X,$$

$T_n^{-1}$  is bounded and  $\|T_n^{-1}\| < 1/c$ . Now,

$$T_n^{-1} - T^{-1} = T_n^{-1}(T - T_n)T^{-1}.$$

Let  $\phi = T^{-1}g$  and  $\phi_n = T_n^{-1}g$ . Then

$$\phi_n - \phi = T_n^{-1}(T - T_n)\phi,$$

$$\|\phi_n - \phi\| \leq \|T_n^{-1}\| \|T_n\phi - T\phi\|,$$

$$\|\phi_n - \phi\| \rightarrow 0.$$

The error bound for  $\|\phi_n - \phi\|$  is of mixed type, neither purely theoretical nor purely practical.

A special case of this scheme was carried out by Madsen [14] in a classical analysis setting.

## 7. A Time-Dependent Transport Problem

Wilson [20] investigated the discrete ordinates method for a generalized transport equation in several space dimensions:

$$(7.1) \quad \frac{\partial \phi(t, x, v)}{\partial t} + v \cdot \nabla \phi(t, x, v) + \|v\|_2 \sigma(t, x, v) \phi(t, x, v) \\ = \int_{\mathfrak{B}} p(t, x, v, v') \phi(t, x, v') dv' + g(t, x, v).$$

Independent variables are time  $t \in (0, T)$ , position  $x \in \Gamma$ , a rectangle in  $R^n$ , and velocity  $v \in \mathfrak{B}$ , a ball in  $R^n$ . The gradient acts with respect to  $x$ . An

initial condition specifies  $\phi(0, x, v)$ . Partial specular reflection on  $\partial\Gamma$ , ranging from full reflection to an exterior vacuum, was assumed by Wilson.

In operator form (7.1) becomes  $D\phi = L\phi + g$ . Discrete ordinates approximations  $\phi_n$  satisfy equations  $D\phi_n = L_n\phi_n + g$ , where  $L_n$  is a numerical integration approximation to  $L$  with a convergent positive quadrature rule.

Wilson [19] obtained existence and uniqueness theorems for weak and strong solutions of a general problem which includes both  $D\phi = L\phi + g$  and  $D\phi_n = L_n\phi_n + g$  as special cases. The analysis involves an equivalent formulation of the problem as an integral equation which is of Volterra type in  $t$ . Direct iteration yields the existence and uniqueness of  $\phi$  and  $\phi_n$ , plus inequalities of the form

$$(7.2) \quad \|\phi_n(t)\| \leq (a \|\phi_n(0)\| + b \|g\|) e^{(c \|\sigma\| + d \|L_n\|)t},$$

with max norms on the functions over undisplayed arguments.

Since  $D(\phi_n - \phi) = L_n(\phi_n - \phi) + L_n\phi - L\phi$  with zero initial and boundary conditions, (7.2) implies

$$(7.3) \quad \|\phi_n(t) - \phi(t)\| \leq b \|L_n\phi - L\phi\| e^{(c \|\sigma\| + d \|L_n\|)t},$$

$$(7.4) \quad \phi_n \rightarrow \phi \quad \text{uniformly as } n \rightarrow \infty.$$

The error bound in (7.3) is of mixed type.

## 8. Conclusions

In the preceding, we have attempted to survey developments in the investigation of convergence properties and error bounds for the discrete ordinates approximations to the transport equation. This summary has been carried out in an abstract setting which, we believe, serves to unify and clarify much previous work in this area. The direction of future investigations is also clearly indicated.

While the first studies of convergence properties of the discrete ordinates methods were carried out nearly fifteen years ago, in an astrophysical setting, they have generally not come to the attention of workers interested in practical applications of the method to nuclear reactor computational problems. In fact, as recently as 1968, a standard work on reactor computing methods refers to the convergence of the discrete ordinates method as an unsolved problem.

A number of practical implications follow from the results surveyed above. For example, the equivalence of the spherical harmonics ( $P_N$ ) and Gauss-quadrature methods for slab geometry transport problems (in the sense that the solutions agree at the quadrature points) establishes the convergence of, and provides error bounds for, the  $P_N$  solutions at the quadrature points.

Hopefully, future extensions of the results surveyed here to include three-dimensional systems and discretization of the spatial variable will provide convergence proofs and practical error estimates which will be of use even in the most complicated practical applications of discrete ordinates methods.

## References

1. Anselone, P. M. Integral Equations of the Schwarzschild-Milne Type. *J. Math. and Mech.*, 7 (1958), pp. 557-570.
2. Anselone, P. M. Convergence of the Wick-Chandrasekhar Approximation Technique in Radiative Transfer. *Astrophys. J.*, 128 (1958), pp. 124-129.
3. Anselone, P. M. Convergence of the Wick-Chandrasekhar Approximation Technique in Radiative Transfer, II. *Astrophys. J.*, 130 (1959), pp. 881-883.
4. Anselone, P. M. Convergence of Chandrasekhar's Method for Inhomogeneous Transfer Problems. *J. Math. and Mech.*, 10 (1961), pp. 537-546.
5. Anselone, P. M. Convergence of Chandrasekhar's Method for the Problem of Diffuse Reflection. *Royal Astr. Soc. Monthly Notices*, 120 (1960), pp. 498-503.
6. Anselone, P. M. Collectively Compact Operator Approximation Theory and Applications to Integral Equations. Prentice-Hall, Inc., Englewood Cliffs, N. J., 1971.
7. Chandrasekhar, S. On the Radiative Equilibrium of a Stellar Atmosphere, II. *Astrophys. J.*, 100 (1944), pp. 76-86.
8. Chandrasekhar, S. Radiative Transfer. Oxford University Press, 1950.
9. Hopf, E. Mathematical Problems of Radiative Equilibrium. Cambridge University Press, 1934.

10. Keller, H. B. Approximate Solutions of Transport Problems. SIAM J. Appl. Math., 8 (1960), pp. 43-73.
11. Keller, H. B. On the Pointwise Convergence of the Discrete Ordinate Method. SIAM J. Appl. Math., 8 (1960), pp. 560-567.
12. Keller, H. B. Convergence of the Discrete Ordinate Method for the Anisotropic Scattering Transport Equation. Provisional Intl. Comp. Center, Rome Symposium, Birkhauser-Verlag, Basel/Stuttgart, 1960, pp. 292-299.
13. Kofink, W. Studies of the Spherical Harmonics Method in Neutron Transport Theory, Nuovo Cimento 10, 9 (1958), Supplemento, pp. 497-541.
14. Madsen, N. K. Pointwise Convergence of the Three-Dimensional Discrete Ordinate Method. SIAM J. Num. Anal., 8 (1971), pp. 266-269.
15. Nelson, P. Convergence of the Discrete Ordinates Method for Anisotropically Scattering Multiplying Particles in a Subcritical Slab. SIAM J. Num. Anal., 10 (1973), pp. 175-181.
16. Nestell, M. K. The Convergence of the Discrete Ordinates Method for integral Equations of Anisotropic Radiative Transfer. Tech. Report 23, Math Dept., Oregon State University, 1965.
17. Wendroff, B. On the Convergence of the Discrete Ordinate Method. SIAM J. Appl. Math., 8 (1960), pp. 508-511.

18. Wick, G. C. Über Ebene Diffusions Problem. Z. für Physik 121 (1943), pp. 702-718.
19. Wilson, D. G. Time-Dependent Linear Transport, I., J. Math. Anal. Appl., to appear.
20. Wilson, D. G. Time Dependent Linear Transport, III; Convergence of Discrete Ordinate Method, submitted for publication.