

Trinity University Digital Commons @ Trinity

Biology Faculty Research

Biology Department

2011

AMPK Directly Inhibits NDPK Through a Phosphoserine Switch to Maintain Cellular Homeostasis

R. U. Onyenwoke

L.J. Forsberg

L. Liu

Tyisha Williams

Trinity University, twilliam@trinity.edu

O. Alzate

See next page for additional authors

Follow this and additional works at: https://digitalcommons.trinity.edu/bio_faculty

Part of the [Biology Commons](#)

Repository Citation

Onyenwoke, R.U., Forsberg, L.J., Liu, L., Williams, T., Alzate, O., & Brenman, J.E. (2011). AMPK directly inhibits NDPK through a phosphoserine switch to maintain cellular homeostasis. *Molecular Biology of the Cell*, 23(2), 381-389. doi:10.1091/mbc.E11-08-0699

This Article is brought to you for free and open access by the Biology Department at Digital Commons @ Trinity. It has been accepted for inclusion in Biology Faculty Research by an authorized administrator of Digital Commons @ Trinity. For more information, please contact jcostanz@trinity.edu.

Authors

R. U. Onyenwoke, L. J. Forsberg, L. Liu, Tyisha Williams, O. Alzate, and J. E. Brenman

AMPK directly inhibits NDPK through a phosphoserine switch to maintain cellular homeostasis

Rob U. Onyenwoke^a, Lawrence J. Forsberg^a, Lucy Liu^a, Tyisha Williams^a, Oscar Alzate^b, and Jay E. Brenman^{a,b}

^aNeuroscience Center and ^bDepartment of Cell and Developmental Biology, University of North Carolina School of Medicine, Chapel Hill, NC 27599

ABSTRACT AMP-activated protein kinase (AMPK) is a key energy sensor that regulates metabolism to maintain cellular energy balance. AMPK activation has also been proposed to mimic benefits of caloric restriction and exercise. Therefore, identifying downstream AMPK targets could elucidate new mechanisms for maintaining cellular energy homeostasis. We identified the phosphotransferase nucleoside diphosphate kinase (NDPK), which maintains pools of nucleotides, as a direct AMPK target through the use of two-dimensional differential in-gel electrophoresis. Furthermore, we mapped the AMPK/NDPK phosphorylation site (serine 120) as a functionally potent enzymatic “off switch” both in vivo and in vitro. Because ATP is usually the most abundant cellular nucleotide, NDPK would normally consume ATP, whereas AMPK would inhibit NDPK to conserve energy. It is intriguing that serine 120 is mutated in advanced neuroblastoma, which suggests a mechanism by which NDPK in neuroblastoma can no longer be inhibited by AMPK-mediated phosphorylation. This novel placement of AMPK upstream and directly regulating NDPK activity has widespread implications for cellular energy/nucleotide balance, and we demonstrate in vivo that increased NDPK activity leads to susceptibility to energy deprivation-induced death.

Monitoring Editor

Benjamin Margolis
University of Michigan
Medical School

Received: Aug 16, 2011

Revised: Oct 27, 2011

Accepted: Nov 16, 2011

INTRODUCTION

AMP-activated protein kinase (AMPK) functions as a cellular energy sensor activated by hypoxia, low glucose, and other stressors that lower ATP levels and raise AMP levels (Hardie *et al.*, 2006; Shaw, 2006). In response to AMP/ATP ratio-altering events, activated AMPK turns on ATP-generating pathways and inhibits ATP-consuming pathways, thereby restoring the AMP/ATP ratio (Williams and Brenman, 2008). AMPK was first discovered as a protein whose activity inhibited preparations of acetyl-CoA carboxylase (ACC1), a regulator of cellular fatty acid synthesis (Winder

et al., 1997). AMPK is a heterotrimeric protein with a 63-kDa catalytic α subunit and two regulatory β and γ subunits (38 and 36 kDa, respectively), each of which is encoded by distinct genes (α 1, α 2; β 1, β 2; γ 1, γ 2, γ 3; Davies *et al.*, 1994; Mitchelhill *et al.*, 1994; Gao *et al.*, 1996; Stapleton *et al.*, 1996; Nielsen *et al.*, 2003), and AMPK is implicated in a number of signaling pathways (Hardie, 2004, 2007; Shaw, 2009).

Upstream activation of AMPK is mediated by the tumor suppressor liver kinase B1 (LKB1; Shaw *et al.*, 2004) and Ca^{2+} /calmodulin-dependent kinase kinase β (Hawley *et al.*, 2005; Hurley *et al.*, 2005). Although LKB1 has clear roles in metabolism, LKB1 is also known as Par-4 in *Caenorhabditis elegans*, a key regulator of cell polarity (Watts *et al.*, 2000; Chartier *et al.*, 2011). Nonetheless, all known AMPK upstream kinases phosphorylate AMPK α threonine 172 (Thr-172) in the “activation loop” of the catalytic α subunit (both α 1 and α 2 isoforms), and this phosphorylation event causes >100-fold increase in kinase activity (Hawley *et al.*, 1996). Conversely, dephosphorylation of Thr-172 by phosphatases can turn AMPK activity off (Sanders *et al.*, 2007; Rubenstein *et al.*, 2008). In general, mammalian AMPK activity stimulates processes involved in ATP-producing, catabolic pathways (e.g., increasing the glucose transporter GLUT4 and mitochondrial biogenesis) and inhibits ATP-consuming anabolic

This article was published online ahead of print in MBoC in Press (<http://www.molbiolcell.org/cgi/doi/10.1091/mbc.E11-08-0699>) on November 23, 2011.

The authors declare no conflict of interest.

Address correspondence to: Jay E. Brenman (brenman@med.unc.edu); Rob U. Onyenwoke (onyenwok@email.unc.edu).

Abbreviations used: AMPK, AMP-activated protein kinase; hGFAP-Cre, human glial fibrillary acidic protein-Cre; LKB1, liver kinase B1; NDPK, nucleoside diphosphate kinase; 2D-DIGE, two-dimensional differential in-gel electrophoresis.

© 2012 Onyenwoke *et al.* This article is distributed by The American Society for Cell Biology under license from the author(s). Two months after publication it is available to the public under an Attribution-Noncommercial-Share Alike 3.0 Unported Creative Commons License (<http://creativecommons.org/licenses/by-nc-sa/3.0>).

“ASCB®,” “The American Society for Cell Biology®,” and “Molecular Biology of the Cell®” are registered trademarks of The American Society of Cell Biology.

pathways (e.g., gluconeogenesis, lipogenesis, and protein synthesis; Hardie and Hawley, 2001; Barnes *et al.*, 2002; Holmes and Dohm, 2004; Hardie, 2007). The best-defined direct target of AMPK is the fatty acid synthesis and rate-limiting enzyme ACC, which AMPK phosphorylates and inhibits to subsequently lower malonyl-CoA levels and increase fatty acid uptake into mitochondria (Merrill *et al.*, 1997; Winder *et al.*, 1997; Hardie and Hawley, 2001).

Beyond lipid synthesis, AMPK can also switch off protein synthesis by using two different pathways. These pathways include activation of elongation factor-2 kinase, which causes inhibition of the elongation step of translation (Winder *et al.*, 1997; Horman *et al.*, 2002), and inhibition of the target-of-rapamycin (TOR) pathway, which stimulates the initiation step of protein synthesis by the phosphorylation of multiple targets (Proud, 2004). TOR is directly activated by an upstream signaling pathway involving the TSC1–TSC2 (tuberous sclerosis complex) heterodimer. AMPK directly phosphorylates TSC2 and thereby activates the TSC (Inoki *et al.*, 2003). There is also evidence suggesting that AMPK might directly target and inhibit TOR (Cheng *et al.*, 2004). More recent studies identified MRLC, raptor, a clock-related gene, and ULK1 as direct targets of AMPK (Lee *et al.*, 2007; Gwinn *et al.*, 2008; Lamia *et al.*, 2009; Egan *et al.*, 2011).

In this study, we searched for potential new targets of AMPK activity. From these efforts, we identified nucleoside diphosphate kinase (NDPK) as a potential downstream target of AMPK α . NDPK is a ubiquitous enzyme that catalyzes the transfer of the γ -phosphate group from a nucleoside or deoxynucleoside triphosphate (NTP or dNTP) to a nucleoside or deoxynucleoside diphosphate (NDP or dNDP, respectively) involving a high-energy phosphoenzyme intermediate (Rosengard *et al.*, 1989; Engel *et al.*, 1995). Functionally, NDPK maintains pools of nucleoside and deoxynucleoside triphosphates for processes central to energy utilization, for example, DNA synthesis and translation, using diphosphate substrates (Engel *et al.*, 1995). In addition, the *Drosophila* homologue of NDPK is required in vivo during normal development for guided cell migration (Rosengard *et al.*, 1989; Randazzo *et al.*, 1991; Dammal *et al.*, 2003; Nallamotheu *et al.*, 2008). Here we suggest mechanisms by which AMPK normally inhibits NDPK activity through phosphorylation of a highly conserved serine within NDPK.

RESULTS

Two-dimensional differential in-gel electrophoresis analyses identify NDPK as a phosphoprotein

The overall goal of our study was the identification of potential new targets of AMPK—keys for understanding energy homeostasis and metabolic disease that might be mediated by AMPK signaling.

We performed two-dimensional differential in-gel electrophoresis (2D-DIGE) using cytosolic brain lysates derived from wild-type (WT) and AMPK α 1/2 double-knockout (KO) mice (Williams *et al.*, 2011) devoid of all AMPK catalytic activity to identify proteins altered in abundance or posttranslational modification between the WT and KO. One particular protein spot showed a potential shift in migration when comparing WT- and KO-derived lysates but a similar total abundance (Figure 1A).

Given that AMPK α functions as a kinase and the WT spot migrated higher on the gel, this spot might contain a modified protein that was no longer phosphorylated in the KO and thus would migrate differently. A second 2D-DIGE experiment was performed with only WT lysate. However, the WT sample was divided into two samples; the first sample was treated with phosphatase, and the second sample was untreated. A qualitatively similar result to the WT/KO 2D-DIGE comparison was observed (Figure 1B).




FIGURE 1: 2D-DIGE identifies NDPK as a phosphoprotein with reduced phosphorylation in AMPK α 1/2-targeted knockout brain. (A) 2D-DIGE identifies overlapping/adjacent protein spots with slower mobility in wild-type (red) compared with AMPK α -knockout (green) brain. (B) Phosphatase treatment of wild-type extracts (blue) results in greater mobility compared with untreated samples (magenta). All four protein spots were identified as NDPK. (C) Although total NDPK protein levels are similar, immobilized metal ion affinity chromatography purification of phosphoproteins demonstrates more phospho-NDPK in wild-type brain.

After performing the 2D-DIGE experiments, all four adjacent spots of the gel corresponding to the WT or KO spots were excised and identified. All protein spots were confirmed to be nucleoside diphosphate kinase A (NDPK-A; EC 2.7.4.6), also known as Nm23 (nonmetastatic 23). Western blotting indicated that there was no significant difference in NDPK protein expression levels in the total WT versus KO lysate fractions (Figure 1C, Load, and Supplemental Figure S1A). However, a difference was noted when phosphoprotein enrichment was performed prior to Western blot analysis for NDPK, which indicated more phospho-NDPK exists in WT compared with KO brain lysate (Figure 1C, Eluate, and Supplemental Figure S1A).

***Drosophila* NDPK activity is inhibited by AMPK function, and loss of NDPK function can compensate for loss of AMPK α function**

Although there are multiple genes that encode for NDPK activity in mammals (Boissan *et al.*, 2009), there is only a single *NDPK* gene in *Drosophila* with previously characterized genetic loss-of-function mutations available for studies. We therefore used *Drosophila* genetics and a well-established biochemical NDPK activity assay (Timmons *et al.*, 1995; Krishnan *et al.*, 2001) to study NDPK activity in *Drosophila*. Using two genetically defined null alleles, we demonstrated gene-dosage responsiveness for NDPK activity (Figure 2A). We subsequently confirmed that the decreased NDPK biochemical activity also roughly correlated with the NDPK protein levels detected in various genetic *NDPK* mutant genotypes by Western blot (Figure 2B and Supplemental Figure S1B).

We next investigated the seemingly inverse relationship between NDPK and AMPK activities by reducing AMPK α function through the use of a transgenic RNA interference (RNAi)-based expression system that phenocopies genetic loss of AMPK α function and only allows *Drosophila* development to reach the late pupal/pharate adult stage without producing eclosing adults (Johnson *et al.*, 2010). When larvae with decreased AMPK α function, through RNAi (Johnson *et al.*, 2010; Supplemental Figure S2, A and B), were assayed for NDPK activity, a significant increase was observed (Figure 2C).




FIGURE 2: NDPK assay activity in *Drosophila* directly correlates with NDPK but inversely with AMPK α protein expression levels. (A) Quantification of the specific activities and (B) protein levels of the *Drosophila* NDPK alleles compared with wild-type show NDPK activity closely tracks NDPK protein levels ($n = 5$). ** $p < 0.001$ and * $p < 0.005$ vs. the wild type. (C) The RNAi-mediated reduction of the AMPK α level in first-instar larvae leads to an increase in NDPK activity ($n = 3$). Actin-GAL4 was the driver for the expression of the AMPK α -RNAi transgenic element, and the RNAi-mediated reduction of AMPK α protein only occurs when the GAL4 is present. *** $p < 0.0005$. Data are shown as mean \pm SEM.

Although these results are suggestive that AMPK might inhibit NDPK activity, because no prior report described the regulation of NDPK activity by phosphorylation, it was unclear whether loss of NDPK would enhance or inhibit AMPK genetic function. Initially, we used the transgenic RNAi-based expression system to reduce AMPK α function and found that introducing a single NDPK loss-of-function mutation for either NDPK allele (Figure 2B and Supplemental Figure S1B) rescued otherwise lethal AMPK α knockdown animals to viability/eclosion (Table 1). In addition, heterozygous NDPK loss-of-function alleles were able to rescue previously published (Mirouse *et al.*, 2007) null AMPK α lethal loss-of-function alleles to viability as well, an effect otherwise only seen by dominant-negative S6 kinase, which is known to antagonize AMPK function (Montagne *et al.*, 1999). In support of these results, increased NDPK expression did not rescue AMPK α knockdown or loss-of-function animals but instead made the transgenic animals susceptible to energetic stress by accelerating starvation-induced death in a defined starvation paradigm (Johnson *et al.*, 2010; Figure 3).

AMPK directly inhibits NDPK activity through phosphorylation

The preceding results suggested that AMPK and NDPK genetically antagonize each other. Combination of these genetic results with the biochemical identification of decreased NDPK phosphorylation

Transgene or loss-of-function mutation	AMPK α RNAi rescue (% expected)	AMPK α loss-of-function allele 1 rescue (number rescued/total scored)	AMPK α loss-of-function allele 2 rescue (number rescued/total scored)
UAS-S6k (DN)	Yes (11)	Yes (14/130)	Yes (23/122)
UAS-TOR (DN)	Yes (13)	No (0/68)	No (0/42)
NDPK ¹	Yes (26)	Yes (7/91)	Yes (12/96)
NDPK ²	Yes (41)	Yes (5/101)	Yes (8/71)
UAS-AMPK α	Yes (100)	Yes (22/65)	Yes (36/75)
UAS-AMPK γ	No	No (0/63)	No (0/94)
UAS-GFP	No	No (0/52)	No (0/77)

TABLE 1: AMPK α RNAi and AMPK α loss-of-function rescue by NDPK.

in AMPK α mutant brain suggests a potential model by which AMPK directly phosphorylates NDPK to inhibit NDPK function. The relationship between NDPK and AMPK was subsequently investigated in vitro using AMPK purified from a cell expression system (Dyck *et al.*, 1996), which was subsequently added to NDPK protein. When these protein complexes were incubated together before performing NDPK assays, NDPK activity was significantly decreased and substantially further decreased upon additional AMPK activation (Figure 4A and Supplemental Figure S1C) with cobalt chloride (Lee *et al.*, 2003; Supplemental Figure S3; CoCl₂ treatment was the most effective AMPK activator of those tested), which demonstrates AMPK activity inhibits NDPK activity in vitro (Figure 4B).

After establishing that AMPK can directly phosphorylate NDPK in vitro, we identified potential NDPK phosphorylation sites in vivo. The protein spot corresponding to NDPK from WT mouse brain was excised for phosphopeptide mapping (Yale Mass Spectrometry




FIGURE 3: NDPK transgenic overexpression in *Drosophila* leads to decreased survival under starvation conditions. Male flies ($n = 30$) were starved in empty food vials containing filter paper saturated with deionized H₂O. Tubulin-GAL4 was the driver for overexpression of the NDPK transgenic element, and NDPK protein overexpression only occurs when the GAL4 element is present. Asterisk denotes statistically different values; for the 60-, 72-, 84-, and 96-h time points, $p < 0.0072$, < 0.0005 , < 0.0001 , and < 0.013 , respectively; $n = 3$. Data are shown as mean \pm SEM.




FIGURE 4: Activated AMPK activity decreases NDPK activity. Activated AMPK decreases NDPK activity. (A) CoCl₂ addition to HEK293 cells transiently transfected with GST-AMPKα1, HA-β1, and HA-γ1 increases purified activated phospho-AMPKα and (B) incubating purified NDPK with AMPK decreases NDPK activity. **p < 0.005 and *p < 0.05 vs. GST alone (n = 5).

Core; 73.7% coverage, including all NDPK serines and threonines; Supplemental Figure S4). A single peptide (Figure 5A, underlined red) with three candidate serines was found to contain a single phosphoserine. Of importance, this sequence flanked the catalytic NDPK histidine residue and contained three total serine residues, only two of which were conserved from *Drosophila* to human.

To identify which serine residues may be phosphorylated by AMPK, the two conserved serine residues were mutated singly for in vitro AMPK phosphorylation/NDPK activity assays. Single mutations of either serine (S120 or S125) to alanine in NDPK had no significant effect on NDPK activity. Similarly, mutating S125 to the phosphomimetic amino acid glutamate also had no effect on NDPK activity (Figure 5B). However, mutating S120 to glutamate (E) resulted in insoluble protein under native protein purification conditions. In addition, when solubilized under denaturing/renaturation conditions, the S120E mutant protein was inactive even when WT NDPK that underwent the same treatment maintained significant activity (Supplemental Figure S5). We therefore added purified AMPK to purified NDPK protein—either WT or with the serine-to-alanine mutations to measure AMPK-mediated inhibition of NDPK activity. Indeed, adding the AMPK complex to S125A decreased NDPK activity; however, adding AMPK to S120A had no inhibitory effect on NDPK activity, suggesting that S120 is a residue that can be phosphorylated to inhibit NDPK activity (Figure 5B).

To determine whether AMPK could phosphorylate NDPK, in vitro kinase assays were performed to monitor the incorporation of ³²P into a modified version of the NDPK phosphopeptide (the NDPK-tide, with all serines except S120 mutated to alanine; Figure 5A and Supplemental Figure S6) versus the SAMS peptide, a specific, well-established gold standard substrate for AMPK activity (Davies et al.,




FIGURE 5: An identified in vivo phospho-NDPK peptide and identification of an AMPK-dependent NDPK phosphoserine inhibitory site. (A) Sequence alignment of the active-site region of NDPK (purple shading) and identified NDPK phosphopeptide (red underline). The active-site histidine and conserved serines (Ser-120 and Ser-125) are highlighted in green, red, and blue, respectively. (B) The specific activities of wild-type and each NDPK mutant with (dark-colored columns) and without (light-colored columns) the addition of activated AMPK. The decrease in activity for the S120A mutant protein was not statistically significant (p = 0.36, n = 3). *p < 0.05, n = 3. Data are shown as mean ± SEM.

1989). The kinase assays revealed very similar calculated specific activities for the SAMS and the NDPKtide peptides, $V_{max} = 1.47$ and $1.29 \text{ nmol min}^{-1} \text{ mg}^{-1}$, respectively, with the SAMS peptide being only a slightly better—but statistically insignificant—substrate (Figure 6, A and B, and Supplemental Figure S7).

DISCUSSION

Elaborating targets for the cellular energy sensor AMPK is key for understanding the roles this molecule plays in energy homeostasis and metabolic disease. Through the use of proteomic techniques, we were able to identify proteins that are potentially regulated and/or phosphorylated by AMPK and further prioritize these proteins for study based on genetic evaluation in *Drosophila*. On the basis of these criteria and studies, the protein NDPK was identified as a good candidate for additional study.

Phosphopeptide mapping identified a peptide (Figure 5A, underlined in red) that contained a phosphorylated serine residue. Mutagenesis studies performed on the two conserved serine residues indicated that S120 is the critical residue for NDPK regulation (Figure 5B), which has been a speculated but, up to this point, not experimentally validated mechanism for NDPK regulation (Venerando et al., 2011). These results correlate well with previous studies indicating that NDPK is phosphorylated at serine residue(s)




FIGURE 6: In vitro AMPK kinase assays demonstrate similar kinetics and affinity of AMPK for NDPK and SAMS peptide. (A) Kinase assays were performed with purified AMPK from untransfected (untrans.) HEK cell lysates, kinase-dead (KD) myc-tagged AMPK, and wild-type (WT) myc-tagged AMPK. A synthesized NDPK peptide (containing only one serine, Ser-120) and the SAMS peptide were compared as AMPK substrates ($n = 3$). (B) Kinase assays were performed with varying amounts of the substrate peptide and WT-AMPK to measure the kinetics of the SAMS and NDPK peptides ($n = 3$).

(MacDonald *et al.*, 1993; Almaula *et al.*, 1995; Treharne *et al.*, 2009). In fact, two prior studies, one in *Escherichia coli* (Almaula *et al.*, 1995) and a proteome-wide phospho-mapping study in *Drosophila* (Zhai *et al.*, 2008), found this serine (S120) to be a candidate for phosphorylation. In addition, in vitro assays mixing protein kinase CK2 (CK2; formerly casein kinase 2) with NDPK demonstrated that serine phosphorylation of NDPK may be a mechanism to negatively regulate its activity (Biondi *et al.*, 1996) and indicate that the serine phosphorylation of NDPK may inhibit NDPK's phosphotransferase function (Treharne *et al.*, 2009), which are in agreement with our results. Therefore, NDPK activity, which normally has a high turnover rate, might be partially inhibited by serine phosphorylation (Biondi *et al.*, 1996). Conversely, the need for large amounts of NTPs could be satisfied rapidly via phosphatase-catalyzed dephosphorylation of NDPK, thus increasing NDPK activity (Biondi *et al.*, 1996). However, until now, the mechanistic details of NDPK inhibition due to a specific phosphorylation event in vivo have been unclear.

The critical serine identified in this work, S120, is strictly conserved in all prokaryotic and eukaryotic NDP kinases and has been linked to a direct involvement in the catalytic mechanism of stabilizing the NDPK catalytic site (His-118) through site-directed mutagenesis and crystal structure studies (Tepper *et al.*, 1994; Giraud *et al.*, 2006). A serine 120-to-glycine (S120G) mutation of *nm23-H1* (NDPK-A) was even identified in several aggressive neuroblastomas (Chang *et al.*, 1994). Of interest, biochemical studies indicated that this mutant S120G was still active, which is comparable to our results with our S120 mutants (Chang *et al.*, 1994; Freije *et al.*, 1997).

We verified serine 120 as a residue capable of being phosphorylated by AMPK through the use of in vitro kinase assays and comparisons to the SAMS peptide, which is a specific substrate for

AMPK that contains the optimal AMPK-binding sequence (Gwinn *et al.*, 2008). Although the NDPKtide does not contain the described optimal AMPK-binding sequence, several described AMPK substrates also do not contain this sequence, including histone H2B, myosin light chain, the tumor suppressor p53, and the cyclin-dependent kinase inhibitor p27 (Jones *et al.*, 2005; Lee *et al.*, 2007; Liang *et al.*, 2007; Bjorklund *et al.*, 2010; Bungard *et al.*, 2010; Kim *et al.*, 2011). Of importance, the NDPK peptide displayed kinetic characteristics similar to those of the SAMS peptide (Figure 6B), and these kinetic characteristics were completely abolished when S120 was individually mutated to an alanine (Supplementary Figure S7). Thus the NDPKtide sequence and serine 120 are a specific peptide region and residue targeted by AMPK for phosphorylation.

Although many studies have suggested phosphorylation may affect NDPK activity, a clear genetic link to a kinase in vivo has been missing to this point. A previous study (Treharne *et al.*, 2009) provided 1) biochemical evidence to indicate a direct interaction between AMPK and NDPK, although other groups have been unable to reproduce these results (Annesley *et al.*, 2011), and 2) data suggesting that AMPK activity decreases NDPK's phosphotransferase activity through the phosphorylation of NDPK serine residues, but was unable to mechanistically define the interaction and indicated that the AMPK/NDPK interaction may still be indirect. Other studies investigating the AMPK/NDPK interaction have also served to confound this interaction—see, for example, the retractions of several papers that had sought to directly address this interaction biochemically (Crawford *et al.*, 2005, 2006a, 2006b, 2007). Nevertheless, these retracted papers claimed that NDPK regulated and activated AMPK, which is not the regulatory interaction we describe here.

In our proposed model, active NDPK helps produce nucleotides for various anabolic pathways in the presence of inactive AMPK, that is, a high-energy and/or low-stress cellular state (Figure 7A). However, when cellular ATP falls and AMP levels rise, AMPK is activated and phosphorylates the critical serine 120 residue of NDPK to decrease its activity and, thereby, save ATP stores (Figure 7B). Thus AMPK is able to fulfill its primary energy gauge function and balance the formation of nucleotides with the conservation of ATP under energetic stress. Such a model might also explain why a mutation of this critical serine to a nonphosphorylatable glycine residue would lead to more aggressive neuroblastomas (Chang *et al.*, 1994); tumors could synthesize deoxynucleotides for DNA replication and use the increased biosynthetic products of NDPK for elevated levels of glycolysis (Green *et al.*, 2011) even during energetic stress if this "off switch" was absent. This paradigm illustrates a potential direct link between tumor suppression and control of cellular metabolism. In addition, the activation of NDPK would also trigger an increase in the ADP/ATP ratio, which would further enhance AMPK activity by protecting AMPK from dephosphorylation. Neuroblastomas with the NDPK S120G mutation would then have elevated AMPK activities, as well as constitutive NDPK activation, which might increase glucose uptake/oxidation and help with tumor survival. Thus this AMPK-mediated NDPK inhibitory function is consistent with numerous studies suggesting that AMPK, as a molecule downstream of the human tumor suppressor LKB1, has tumor-suppressive activities (Jones *et al.*, 2005; Shackelford and Shaw, 2009). Of course, it is unlikely that AMPK activity is the only activity capable of regulating NDPK activity; however, in brain lysates, it appears that AMPK activity is a major one.

MATERIALS AND METHODS

Materials

All chemicals were of an analytical grade and, unless otherwise noted, from Sigma-Chemical (St. Louis, MO) or Fisher Scientific (Fair




FIGURE 7: AMPK mediates inhibition of NDPK activity during nutrient stress. Model for AMPK-mediated NDPK inhibition. (A) Under nutrient-rich conditions, AMPK remains inactive, and NDPK is active and uses ATP as an energy source to produce nucleotides/deoxynucleotides for several anabolic pathways. (B) During nutrient-limiting conditions, AMPK is active (and phosphorylated at Thr-172) and inhibits NDPK activity through the phosphorylation of Ser-120, thereby conserving cellular ATP stores. Dashed lines indicate decreased biochemical activities.

Lawn, NJ). [γ - 32 P]ATP (specific activity 3000 Ci/mmol) was from PerkinElmer (Boston, MA). The SAMS (HMRSAMSGLHLVKRR) and NDPK (RNIIHGSDAVKAKRR) peptides were synthesized by Abgent (San Diego, CA) and the University of North Carolina Microprotein Sequencing and Peptide Synthesis Facility (Chapel Hill, NC), respectively. The hemagglutinin (HA)-tagged human AMPK γ 1 and rat AMPK β 1 constructs were gifts from R. Shaw. Glutathione S-transferase (GST)-tagged AMPK α 1 was a gift from L. Witters. The antibodies used were as follows: anti-NDPK (C-20, sc-343; Santa Cruz Biotechnology, Santa Cruz, CA), anti-phospho-AMPK α (40H9) and anti-AMPK α (23A3; Cell Signaling, Beverly, MA), and anti- α -tubulin (clone B-5-1-2; Sigma-Aldrich). Anti-dNDPK (reactive to protein of *Drosophila* origin, dNDPK) was a gift from T. Hsu.

Transgenic animals and brain sample/lysate preparation

Three-month-old male mice were killed to obtain brain tissue (kindly provided by T. Williams, University of North Carolina). The genotypes used were as follows: wild type, human glial fibrillary acidic protein-Cre (hGFAP-Cre) mice (K. McCarthy, University of North Carolina), and conditional AMPK α 1/2 knockout mice (Williams et al., 2011) to produce AMPK α 1 $^{-/-}$ α 2 $^{F/F}$ -hGFAP-Cre mice. The animals were handled under protocols approved by the Institutional Animal Care and Use Committee (Institutional Animal Care and Use Committee ID 09-149.0) of the University of North

Carolina—Chapel Hill and in accordance with National Institutes of Health guidelines.

For the preparation of brain lysate, a whole mouse brain (375–425 mg) was processed in 10 ml of ice-cold lysis buffer A (50 mM Tris-HCl, pH 7.5, protease [P2714; Sigma-Aldrich] and phosphatase [P5726; Sigma-Aldrich] inhibitor cocktails [note: for the preparation of the phosphatase-treated WT brain lysate for 2D-DIGE, the phosphatase inhibitor cocktail was omitted for 2.5 U/ml alkaline phosphatase (final concentration; P6774; Sigma) and the sample was incubated for 1 h with the addition of 1 mM MgCl₂ (37°C) before the high-speed centrifugation step], and benzonase nuclease) using a Tissuemiser homogenizer (Fisher Scientific). The lysate was centrifuged at 1000 \times g for 20 min (4°C). The supernatant was then centrifuged for a second time. The resultant clarified supernatant was centrifuged at 100,000 \times g to produce the cytosol. The pellet was discarded, and the supernatant (cytosol) was directly used for Western blotting and then for 2D-DIGE analyses after it was cleaned using a 2D Clean-Up Kit (GE Healthcare, Piscataway, NJ).

2D-DIGE protocols and protein identifications

The 2D-DIGE experiments were performed by the University of North Carolina Systems-Proteomics Center using previously described methodologies (Osorio et al., 2007). Protein spot identification was performed by the Yale Mass Spectrometry and Proteomics Resource Core (New Haven, CT) using peptide mass fingerprinting tandem mass spectrometry data, as previously described (Osorio et al., 2007; Pinaud et al., 2008).

Purification of phosphoproteins by immobilized metal affinity chromatography

Phosphoproteins were purified from mouse brain lysates using a PhosphoProtein Kit (Qiagen, Valencia, CA), as described by the manufacturer. The brain lysates were prepared as described; however, 0.25% 3-[(3-cholamidopropyl)dimethyl-ammonio]-1-propanesulfonate was included in the buffer and throughout the purification process, as recommended by the manufacturer's protocol.

Plasmid construction

A pET21b (Novagen, Gibbstown, NJ) construct containing the cDNA for wild-type NDP kinase A, inserted in the *Bam*HI-*Nde*I site, was obtained from M.-L. Lacombe. The NDPK gene was PCR amplified from this construct using the primer pair F, 5'-AAAGGATCCG-GCCAACTGTGAGCGTACCTTC-3', and R, 5'-AAAGGATCCGGC-CAACTGTGAGCGTACCTTC-3', digested with *Bam*HI and *Sall*, and ligated into a pET28b (Novagen) vector cut with the same restriction enzymes for expression as a histidine (His)-tagged protein.

Site-directed mutagenesis of NDPK was carried out using bridging PCR. The primer pairs for the construction of each NDPK variant are as follows: S120E (F, 5'-ATTATACATGGCGAGGATTCTGTGGAGAGTGC-3'; R, 5'-TCCACAGAATCCTCGCCATGTATAATGTTCTG-3'), S125E (F, 5'-TTCTGTGGAGGAGGCAGAGAAGGAGATCGG-3', R, 5'-CCTTCTCTGCCTCCTCCACAGAATCACTGCC-3'), S120A (F, 5'-ATTATACATGGCGCTGATTCTGTGGAGAGTGC-3', R, 5'-TCCACAGAATCAGCGCCATGTATAATGTTCTG-3'), and S125A (F, 5'-TTCTGTGGAGGCTGCAGAGAAGGAG-ATCGG-3', R, 5'-CCTTCTCTGCAGCCTCCACAGAATCACTGCC-3').

Purification of NDP kinase

Full-length His-tagged wild-type and mutant proteins were expressed in high yields using *E. coli* BL21-Gold (DE3) pLysS competent cells (Agilent Technologies, Palo Alto, CA) transformed with the appropriate construct using heat shock, as described by the

manufacturer, and purified using batch/gravity-flow column purification with Talon IMAC resin (Clontech, Mountain View, CA) under native conditions (denaturing conditions, i.e., 4 M urea, were used throughout the purification process for the S120E mutant) following the manufacturer's instructions.

Cell culture and AMPK α / β / γ coimmunoprecipitation

For mammalian cell expression of AMPK, the AMPK subunits were used as previously described (Dyck *et al.*, 1996). HEK293 cells were cultured in complete DMEM (Life Technologies, Carlsbad, CA) containing 10% fetal bovine serum (Atlanta Biologicals, Lawrenceville, GA) at 37°C in 5% CO₂. For the transient expression of AMPK protein, the cells were plated 24 h before the experiments in 15-cm dishes and then transfected with the three plasmids using Lipofectamine 2000 (1 μ g DNA per 2 μ l; Invitrogen, Carlsbad, CA) following the manufacturer's protocols. Note: For some kinase assays, GST-AMPK α 1 was replaced with WT or kinase-dead [KD] myc-AMPK (in a pCMV-myc vector; Clontech; Kazgan *et al.*, 2010).

After 24 h, fresh medium containing CoCl₂ (200 μ M) was added to the cells for 1 h in the incubator to activate AMPK, as previously described (Lee *et al.*, 2003). Cells were then harvested, lysed in 0.5 ml of lysis buffer A plus 1.0% Triton X-100 with shaking for 1 h (4°C), and centrifuged at 16,000 \times g for 10 min (4°C). GST- and myc-tagged AMPK were purified from the supernatants via GST pull-down, using glutathione Sepharose 4B (Amersham, GE Healthcare), and immunoprecipitation performed, using anti-c-Myc antibody (9E10; Developmental Studies Hybridoma Bank, University of Iowa, Iowa City, IA) at a 1:100 dilution for 1 mg/ml lysate and A/G agarose (Pierce Protein Research Products, Rockford, IL), respectively, according to the manufacturer's instructions and as previously described (Kazgan *et al.*, 2010). Washed, bead-adsorbed GST-AMPK was used for NDPK assays, as previously described (Dyck *et al.*, 1996), and both GST- and myc-tagged AMPK were used for kinase assays.

NDPK assays

A well-known procedure to assay NDP kinase activities was used (Timmons *et al.*, 1993; Krishnan *et al.*, 2001). In brief, 10 μ l of diluted enzyme (lysate or purified NDPK; see further discussion) was added to a 990-ml reaction mixture containing 100 mM Tris-HCl, pH 7.5, 10 mM MgCl₂, 100 mM KCl, 0.4 mM NADH, 6 mM ATP, 0.7 mM TDP, 4 mM phosphoenolpyruvate (PEP), and 10 U of pyruvate kinase and lactate dehydrogenase each. The absorbance of NADH at 340 nm was then recorded. A unit of activity is defined as the amount required to convert 1 μ mol of NADH to NAD⁺ in 1 min.

For NDPK assays with flies or fly larvae, 20 adult flies or 40–50 fly larvae were homogenized in 50 μ l of ice-cold buffer (100 mM Tris-HCl, pH 7.5, 10 mM MgCl₂, and 100 mM KCl). The lysates were centrifuged at 10,000 \times g for 10 min (4°C). The clarified supernatant was then diluted 1:10 for inclusion in the described assay (0.5–2 μ g of protein used). Purified NDPK was purchased from Sigma-Aldrich (N2635) and also produced as purified His-tagged versions and used at amounts of 10–50 and 50–200 ng, respectively, in NDPK assays. For the inhibition assays including AMPK, 3 μ g of bead-adsorbed GST-AMPK was incubated with purified NDPK or purified His-tagged NDPK at room temperature for 2 h before executing the NDPK assays.

Kinase assays

Kinase assays were performed according to previously described methods (Davies *et al.*, 1989). Briefly, AMPK activity assays with GST-AMPK were performed at room temperature (25°C) in a 25- μ l reac-

tion mixture containing 3–12 μ g of protein in kinase buffer (50 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid, pH 7.0, 75 mM NaCl, 5 mM sodium acetate, 5 mM magnesium chloride, 1 mM dithiothreitol, 8% glycerol, 0.1 mM EDTA, 200 μ M AMP and ATP, and 2 μ Ci of [γ -³²P]ATP) with or without the SAMS or NDPK peptide. After a 30-min incubation period, the reaction mixtures were counted in a scintillation counter. Kinase assays with myc-tagged KD-AMPK and myc-tagged WT-AMPK were performed in the same manner as described earlier, but 0.5 μ g of protein was added to the reaction mixture. AMPK activity is expressed as picomoles of ³²P incorporation into the peptide per minute per microgram of protein.

Western blotting

Fly protein lysates for immunoblotting were prepared by collecting equal numbers of male and female flies (50 total) of each genotype in a 1.5-ml microfuge tube. One milliliter of lysis buffer A was added to each sample. Flies were then ground to homogeneity, incubated for 1 h with shaking (4°C), and centrifuged at 16,000 \times g for 10 min (4°C). Supernatants were collected, and protein concentrations were determined using the Bio-Rad DC protein assay (Richmond, CA). Note: The preparation of brain lysate samples was described earlier.

Proteins (50 μ g) were then boiled in loading buffer and subjected to SDS-PAGE (Invitrogen), followed by Western analyses using 1:1000 dilutions of all primary antibodies, with the exception of anti- α tubulin (1:16,000). Secondary antibodies (IRDye infrared antibodies; LI-COR Biosciences, Lincoln, NE) were used at a dilution of 1:2000. Scanning, analyzing, and quantification of blots were performed via the Odyssey Infrared Imaging System (LI-COR Biosciences). Three or more independent experiments were performed for all immunoblotting data.

Fly stocks, crosses, rescue experiments, and lifespan measurements

Drosophila melanogaster strains obtained from the Bloomington Stock Center (Bloomington, IN) included the following: *Act-GAL4*, *tub-GAL4*, *awd*^{KRS6} (referred to as NDPK¹ in this study), *awd*^{MSM95} (NDPK²), *UAS-AMPK α ^{RNAi}*, the *SNF1A*¹ and *SNF1A*³ mutants, *UAS-SNF1A*, *UAS-S6k.KQ*, *UAS-Tor.TED*, and *UAS-GFP*. *UAS-SNF4* and *UAS-NDPK* were gifts from D. Kretschmar and T. Hsu, respectively. All flies were maintained at 25°C in yeast-cornmeal vials, and all crosses were also performed in cornmeal-yeasted vials.

For the AMPK α RNAi rescue experiments, males carrying a transgene or loss-of-function mutation on the second or third chromosome were mated to virgin females carrying a *GAL4* (either *tub-GAL4* or *Act-GAL4*, respectively). From these crosses, male progeny carrying the transgene or loss-of-function mutation and the *GAL4* were mated to virgin females carrying *UAS-AMPK α ^{RNAi}*. The progeny from this second cross were then scored for rescue, that is, viable adult flies in spite of AMPK α RNAi knockdown.

For the AMPK α loss-of-function rescue experiments, males carrying a transgene or loss-of-function mutation on the second or third chromosome were mated in parallel to virgin females from both the *SNF1A*¹ and *SNF1A*³ loss-of-function lines. The male progeny from both of these crosses were scored for rescue, that is, viable adult flies in spite of carrying the lethal loss-of-function mutation/phenotypically, non-bar-eyed males.

Measurements of lifespan have been widely used in *Drosophila* as a metric of stress sensitivity (Johnson *et al.*, 2010). Thirty 3- to 5-d-old male flies were starved in empty food vials that contained pieces of filter paper saturated with deionized H₂O. We assessed the percentage survival of at least three replicate vials three times daily.

Statistical analyses

Comparisons were made using the unpaired Student's *t* test with *p* < 0.05 considered significant. Values are presented as the mean ± the SE of the mean (SEM) and are represented as error bars. Indirect immunofluorescent detection of a secondary antibody (LI-COR Biosciences) was scanned and standardized to an internal standard to calculate and quantify arbitrary units using the Odyssey Infrared Imaging System, and a representative Western blot is shown in each figure.

ACKNOWLEDGMENTS

We thank B. Viollet (Université Paris Descartes, Paris, France) and K. McCarthy (University of North Carolina–Chapel Hill) for *AMPK α_1 ^{-/-} α_2 ^{F/F}*-hGFAP-Cre mice, R. Shaw (Salk Institute, La Jolla, CA) for HA-tagged *AMPK β* and *AMPK γ* constructs, L. Witters (Dartmouth Medical School, Hanover, NH) for the GST-tagged *AMPK α* construct, T. Hsu (Boston University, Boston, MA) for anti-NDPK antibody and *UAS-NDPK* flies, M.-L. Lacombe (Université Pierre et Marie Curie, Paris, France) for the plasmid containing wild-type *NDPK*, and D. Kretzschmar (Oregon Health and Science University, Portland, OR) for *UAS-AMPK γ* flies. R.U.O. is supported through T32 National Cancer Institute Training Grant 5-T32-CA09156-35 to the University of North Carolina–Chapel Hill Lineberger Comprehensive Cancer Center. J.E.B. is funded by National Institutes of Health Grant R01NS063858 and through University of North Carolina–Chapel Hill University Funds.

REFERENCES

- Almoula N, Lu Q, Delgado J, Belkin S, Inouye M (1995). Nucleoside diphosphate kinase from *Escherichia coli*. *J Bacteriol* 177, 2524–2529.
- Annesley SJ, Bago R, Mehta A, Fisher PR (2011). A genetic interaction between NDPK and AMPK in *Dictyostelium discoideum* that affects motility, growth and development. *Naunyn Schmiedeberg's Arch Pharmacol* 384, 341–349.
- Barnes K, Ingram JC, Porras OH, Barros LF, Hudson ER, Fryer LG, Foulfelle F, Carling D, Hardie DG, Baldwin SA (2002). Activation of GLUT1 by metabolic and osmotic stress: potential involvement of AMP-activated protein kinase (AMPK). *J Cell Sci* 115, 2433–2442.
- Biondi RM, Engel M, Sauane M, Welter C, Issinger OG, Jimenez de Asua L, Passeron S (1996). Inhibition of nucleoside diphosphate kinase activity by in vitro phosphorylation by protein kinase CK2. Differential phosphorylation of NDP kinases in HeLa cells in culture. *FEBS Lett* 399, 183–187.
- Bjorklund MA, Vaahntomeri K, Peltonen K, Viollet B, Makela TP, Band AM, Laiho M (2010). Non-CDK-bound p27 (p27^{INCKD}) is a marker for cell stress and is regulated through the Akt/PKB and AMPK-kinase pathways. *Exp Cell Res* 316, 762–774.
- Boissan M, Dabernat S, Peuchant E, Schlattner U, Lasco I, Lacombe ML (2009). The mammalian Nm23/NDPK family: from metastasis control to cilia movement. *Mol Cell Biochem* 329, 51–62.
- Bungard D, Fuerth BJ, Zeng PY, Faubert B, Maas NL, Viollet B, Carling D, Thompson CB, Jones RG, Berger SL (2010). Signaling kinase AMPK activates stress-promoted transcription via histone H2B phosphorylation. *Science* 329, 1201–1205.
- Chang CL, Zhu XX, Thoraval DH, Ungar D, Rawwas J, Hora N, Strahler JR, Hanash SM, Radany E (1994). Nm23-H1 mutation in neuroblastoma. *Nature* 370, 335–336.
- Chartier NT, Salazar Ospina DP, Benkemoun L, Mayer M, Grill SW, Maddox AS, Labbe JC (2011). PAR-4/LKB1 mobilizes nonmuscle myosin through anillin to regulate *C. elegans* embryonic polarization and cytokinesis. *Curr Biol* 21, 259–269.
- Cheng SW, Fryer LG, Carling D, Shepherd PR (2004). Thr2446 is a novel mammalian target of rapamycin (mTOR) phosphorylation site regulated by nutrient status. *J Biol Chem* 279, 15719–15722.
- Crawford RM, Treharne KJ, Arnaud-Dabernat S, Daniel JY, Foretz M, Viollet B, Mehta A (2006a). Understanding the molecular basis of the interaction between NDPK-A and AMPK alpha 1. *Mol Cell Biol* 26, 5921–5931.
- Crawford RM, Treharne KJ, Arnaud-Dabernat S, Daniel JY, Foretz M, Viollet B, Mehta A (2007). Protein kinase CK2 acts as a signal molecule switching between the NDPK-A/AMPK 1 complex and NDPK-B. *FASEB J* 21, 3398.
- Crawford RM, Treharne KJ, Best OG, Muimo R, Riemen CE, Mehta A (2005). A novel physical and functional association between nucleoside diphosphate kinase A and AMP-activated protein kinase alpha1 in liver and lung. *Biochem J* 392, 201–209.
- Crawford RM, Treharne KJ, Best OG, Riemen CE, Muimo R, Gruenert DC, Arnaud-Dabernat S, Daniel JY, Mehta A (2006b). NDPK-A (but not NDPK-B) and AMPK alpha1 (but not AMPK alpha2) bind the cystic fibrosis transmembrane conductance regulator in epithelial cell membranes. *Cell Signal* 18, 1595–1603.
- Dammai V, Adryan B, Lavenburg KR, Hsu T (2003). *Drosophila* awd, the homolog of human nm23, regulates FGF receptor levels and functions synergistically with shi/dynamin during tracheal development. *Genes Dev* 17, 2812–2824.
- Davies SP, Carling D, Hardie DG (1989). Tissue distribution of the AMP-activated protein kinase, and lack of activation by cyclic-AMP-dependent protein kinase, studied using a specific and sensitive peptide assay. *Eur J Biochem* 186, 123–128.
- Davies SP, Hawley SA, Woods A, Carling D, Haystead TA, Hardie DG (1994). Purification of the AMP-activated protein kinase on ATP-gamma-Sepharose and analysis of its subunit structure. *Eur J Biochem* 223, 351–357.
- Dyck JR, Gao G, Widmer J, Stapleton D, Fernandez CS, Kemp BE, Witters LA (1996). Regulation of 5'-AMP-activated protein kinase activity by the noncatalytic beta and gamma subunits. *J Biol Chem* 271, 17798–17803.
- Egan DF et al. (2011). Phosphorylation of ULK1 (hATG1) by AMP-activated protein kinase connects energy sensing to mitophagy. *Science* 331, 456–461.
- Engel M, Veron M, Theisinger B, Lacombe ML, Seib T, Dooley S, Welter C (1995). A novel serine/threonine-specific protein phosphotransferase activity of Nm23/nucleoside-diphosphate kinase. *Eur J Biochem* 234, 200–207.
- Freije JM, Blay P, MacDonald NJ, Manrow RE, Steeg PS (1997). Site-directed mutation of Nm23-H1. Mutations lacking motility suppressive capacity upon transfection are deficient in histidine-dependent protein phosphotransferase pathways in vitro. *J Biol Chem* 272, 5525–5532.
- Gao G, Fernandez CS, Stapleton D, Auster AS, Widmer J, Dyck JR, Kemp BE, Witters LA (1996). Non-catalytic beta- and gamma-subunit isoforms of the 5'-AMP-activated protein kinase. *J Biol Chem* 271, 8675–8681.
- Giraud MF, Georgescauld F, Lasco I, Dautant A (2006). Crystal structures of S120G mutant and wild type of human nucleoside diphosphate kinase A in complex with ADP. *J Bioenerg Biomembr* 38, 261–264.
- Green AS, Chapuis N, Lacombe C, Mayeux P, Bouscary D, Tamburini J (2011). LKB1/AMPK/mTOR signaling pathway in hematological malignancies: From metabolism to cancer cell biology. *Cell Cycle* 10, 2115–2120.
- Gwinn DM, Shackelford DB, Egan DF, Mihaylova MM, Mery A, Vasquez DS, Turk BE, Shaw RJ (2008). AMPK phosphorylation of raptor mediates a metabolic checkpoint. *Mol Cell* 30, 214–226.
- Hardie DG (2004). The AMP-activated protein kinase pathway—new players upstream and downstream. *J Cell Sci* 117, 5479–5487.
- Hardie DG (2007). AMP-activated/SNF1 protein kinases: conserved guardians of cellular energy. *Nat Rev Mol Cell Biol* 8, 774–785.
- Hardie DG, Hawley SA (2001). AMP-activated protein kinase: the energy charge hypothesis revisited. *Bioessays* 23, 1112–1119.
- Hardie DG, Hawley SA, Scott JW (2006). AMP-activated protein kinase—development of the energy sensor concept. *J Physiol* 574, 7–15.
- Hawley SA, Davison M, Woods A, Davies SP, Beri RK, Carling D, Hardie DG (1996). Characterization of the AMP-activated protein kinase kinase from rat liver and identification of threonine 172 as the major site at which it phosphorylates AMP-activated protein kinase. *J Biol Chem* 271, 27879–27887.
- Hawley SA, Pan DA, Mustard KJ, Ross L, Bain J, Edelman AM, Frenguelli BG, Hardie DG (2005). Calmodulin-dependent protein kinase kinase-beta is an alternative upstream kinase for AMP-activated protein kinase. *Cell Metab* 2, 9–19.
- Holmes B, Dohm GL (2004). Regulation of GLUT4 gene expression during exercise. *Med Sci Sports Exerc* 36, 1202–1206.
- Horman S, Browne G, Krause U, Patel J, Vertommen D, Bertrand L, Lavoine A, Hue L, Proud C, Rider M (2002). Activation of AMP-activated protein kinase leads to the phosphorylation of elongation factor 2 and an inhibition of protein synthesis. *Curr Biol* 12, 1419–1423.

- Hurley RL, Anderson KA, Franzone JM, Kemp BE, Means AR, Witters LA (2005). The Ca²⁺/calmodulin-dependent protein kinase kinases are AMP-activated protein kinase kinases. *J Biol Chem* 280, 29060–29066.
- Inoki K, Zhu T, Guan KL (2003). TSC2 mediates cellular energy response to control cell growth and survival. *Cell* 115, 577–590.
- Johnson EC, Kazgan N, Bretz CA, Forsberg LJ, Hector CE, Worthen RJ, Onyenwoke R, Brenman JE (2010). Altered metabolism and persistent starvation behaviors caused by reduced AMPK function in *Drosophila*. *PLoS One* 5, pii12799.
- Jones RG, Plas DR, Kubek S, Buzzai M, Mu J, Xu Y, Birnbaum MJ, Thompson CB (2005). AMP-activated protein kinase induces a p53-dependent metabolic checkpoint. *Mol Cell* 18, 283–293.
- Kazgan N, Williams T, Forsberg LJ, Brenman JE (2010). Identification of a nuclear export signal in the catalytic subunit of AMP-activated protein kinase. *Mol Biol Cell* 21, 3433–3442.
- Kim J, Kundu M, Viollet B, Guan KL (2011). AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. *Nat Cell Biol* 13, 132–141.
- Krishnan KS, Rikhy R, Rao S, Shivalkar M, Mosko M, Narayanan R, Etter P, Estes PS, Ramaswami M (2001). Nucleoside diphosphate kinase, a source of GTP, is required for dynamin-dependent synaptic vesicle recycling. *Neuron* 30, 197–210.
- Lamia KA et al. (2009). AMPK regulates the circadian clock by cryptochrome phosphorylation and degradation. *Science* 326, 437–440.
- Lee JH et al. (2007). Energy-dependent regulation of cell structure by AMP-activated protein kinase. *Nature* 447, 1017–1020.
- Lee M, Hwang JT, Lee HJ, Jung SN, Kang I, Chi SG, Kim SS, Ha J (2003). AMP-activated protein kinase activity is critical for hypoxia-inducible factor-1 transcriptional activity and its target gene expression under hypoxic conditions in DU145 cells. *J Biol Chem* 278, 39653–39661.
- Liang J et al. (2007). The energy sensing LKB1-AMPK pathway regulates p27(kip1) phosphorylation mediating the decision to enter autophagy or apoptosis. *Nat Cell Biol* 9, 218–224.
- MacDonald NJ, De la Rosa A, Benedict MA, Freije JM, Krutsch H, Steeg PS (1993). A serine phosphorylation of Nm23, and not its nucleoside diphosphate kinase activity, correlates with suppression of tumor metastatic potential. *J Biol Chem* 268, 25780–25789.
- Merrill GF, Kurth EJ, Hardie DG, Winder WW (1997). AICA riboside increases AMP-activated protein kinase, fatty acid oxidation, and glucose uptake in rat muscle. *Am J Physiol* 273, E1107–E1112.
- Mirouse V, Swick LL, Kazgan N, St Johnston D, Brenman JE (2007). LKB1 and AMPK maintain epithelial cell polarity under energetic stress. *J Cell Biol* 177, 387–392.
- Mitchellhill KI, Stapleton D, Gao G, House C, Michell B, Katsis F, Witters LA, Kemp BE (1994). Mammalian AMP-activated protein kinase shares structural and functional homology with the catalytic domain of yeast Snf1 protein kinase. *J Biol Chem* 269, 2361–2364.
- Montagne J, Stewart MJ, Stocker H, Hafen E, Kozma SC, Thomas G (1999). *Drosophila* S6 kinase: a regulator of cell size. *Science* 285, 2126–2129.
- Nallamotheu G, Woolworth JA, Dammai V, Hsu T (2008). Awd, the homolog of metastasis suppressor gene Nm23, regulates *Drosophila* epithelial cell invasion. *Mol Cell Biol* 28, 1964–1973.
- Nielsen JN, Mustard KJ, Graham DA, Yu H, MacDonald CS, Pilegaard H, Goodyear LJ, Hardie DG, Richter EA, Wojtaszewski JF (2003). 5'-AMP-activated protein kinase activity and subunit expression in exercise-trained human skeletal muscle. *J Appl Physiol* 94, 631–641.
- Osorio C, Sullivan PM, He DN, Mace BE, Ervin JF, Strittmatter WJ, Alzate O (2007). Mortalin is regulated by APOE in hippocampus of AD patients and by human APOE in TR mice. *Neurobiol Aging* 28, 1853–1862.
- Pinaud R, Osorio C, Alzate O, Jarvis ED (2008). Profiling of experience-regulated proteins in the songbird auditory forebrain using quantitative proteomics. *Eur J Neurosci* 27, 1409–1422.
- Proud CG (2004). Role of mTOR signalling in the control of translation initiation and elongation by nutrients. *Curr Top Microbiol Immunol* 279, 215–244.
- Randazzo PA, Northup JK, Kahn RA (1991). Activation of a small GTP-binding protein by nucleoside diphosphate kinase. *Science* 254, 850–853.
- Rosengard AM, Krutzsch HC, Shearn A, Biggs JR, Barker E, Margulies IM, King CR, Liotta LA, Steeg PS (1989). Reduced Nm23/Awd protein in tumour metastasis and aberrant *Drosophila* development. *Nature* 342, 177–180.
- Rubenstein EM, McCartney RR, Zhang C, Shokat KM, Shirra MK, Arndt KM, Schmidt MC (2008). Access denied: Snf1 activation loop phosphorylation is controlled by availability of the phosphorylated threonine 210 to the PP1 phosphatase. *J Biol Chem* 283, 222–230.
- Sanders MJ, Grondin PO, Hegarty BD, Snowden MA, Carling D (2007). Investigating the mechanism for AMP activation of the AMP-activated protein kinase cascade. *Biochem J* 403, 139–148.
- Shackelford DB, Shaw RJ (2009). The LKB1-AMPK pathway: metabolism and growth control in tumour suppression. *Nat Rev Cancer* 9, 563–575.
- Shaw RJ (2006). Glucose metabolism and cancer. *Curr Opin Cell Biol* 18, 598–608.
- Shaw RJ (2009). LKB1 and AMP-activated protein kinase control of mTOR signalling and growth. *Acta Physiol (Oxf)* 196, 65–80.
- Shaw RJ, Kosmatka M, Bardeesy N, Hurley RL, Witters LA, DePinho RA, Cantley LC (2004). The tumor suppressor LKB1 kinase directly activates AMP-activated kinase and regulates apoptosis in response to energy stress. *Proc Natl Acad Sci USA* 101, 3329–3335.
- Stapleton D et al. (1996). Mammalian AMP-activated protein kinase subfamily. *J Biol Chem* 271, 611–614.
- Tepper AD, Dammann H, Bominaar AA, Veron M (1994). Investigation of the active site and the conformational stability of nucleoside diphosphate kinase by site-directed mutagenesis. *J Biol Chem* 269, 32175–32180.
- Timmons L, Hersperger E, Woodhouse E, Xu J, Liu LZ, Shearn A (1993). The expression of the *Drosophila* awd gene during normal development and in neoplastic brain tumors caused by Igl mutations. *Dev Biol* 158, 364–379.
- Timmons L, Xu J, Hersperger G, Deng XF, Shearn A (1995). Point mutations in awdKpn which revert the prune/Killer of prune lethal interaction affect conserved residues that are involved in nucleoside diphosphate kinase substrate binding and catalysis. *J Biol Chem* 270, 23021–23030.
- Treharne KJ, Best OG, Mehta A (2009). The phosphorylation status of membrane-bound nucleoside diphosphate kinase in epithelia and the role of AMP. *Mol Cell Biochem* 329, 107–114.
- Venerando A, Pagano MA, Tosoni K, Meggio F, Cassidy D, Stobbart M, Pinna LA, Mehta A (2011). Understanding protein kinase CK2 mis-regulation upon F508del CFTR expression. *Naunyn Schmiedebergs Arch Pharmacol* 384, 473–488.
- Watts JL, Morton DG, Bestman J, Kempthues KJ (2000). The *C. elegans* par-4 gene encodes a putative serine-threonine kinase required for establishing embryonic asymmetry. *Development* 127, 1467–1475.
- Williams T, Brenman JE (2008). LKB1 and AMPK in cell polarity and division. *Trends Cell Biol* 18, 193–198.
- Williams T, Courchet J, Viollet B, Brenman JE, Polleux F (2011). AMP-activated protein kinase (AMPK) activity is not required for neuronal development but regulates axogenesis during metabolic stress. *Proc Natl Acad Sci USA* 108, 5849–5854.
- Winder WW, Wilson HA, Hardie DG, Rasmussen BB, Hutber CA, Call GB, Clayton RD, Conley LM, Yoon S, Zhou B (1997). Phosphorylation of rat muscle acetyl-CoA carboxylase by AMP-activated protein kinase and protein kinase A. *J Appl Physiol* 82, 219–225.
- Zhai B, Villen J, Beausoleil SA, Mintseris J, Gygi SP (2008). Phosphoproteome analysis of *Drosophila melanogaster* embryos. *J Proteome Res* 7, 1675–1682.