

DOE/ER/60100--1

DE83 017472

A Study of Polynuclear Aromatic Hydrocarbons
on an
Amino Bonded Phase Liquid Chromatographic Column
in the
Normal and Reversed Phase
by
Dori Karlesky, Dennis C. Shelly
and Isiah M. Warner

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE
PORTIONS OF THIS REPORT ARE ILLEGIBLE.

It has been reproduced from the best available copy to permit the broadest possible availability.

MASTER

HP
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

A STUDY OF POLYNUCLEAR AROMATIC HYDROCARBONS ON AN
AMINO-BONDED-PHASE LIQUID-CHROMATOGRAPHIC COLUMN
IN THE NORMAL AND REVERSED PHASE

Dori Karlesky, Dennis C. Shelly and Isiah M. Warner*
Department of Chemistry
Texas A&M University
College Station, TX. 77843

ABSTRACT

Experiments were run using an n-propyl amine polar bonded phase (Chromosorb LC-9) liquid chromatographic column in both the normal and reversed phase mode. Results confirm that the mechanism of separation in the normal phase is due mainly to a charge transfer interaction between the lone pair electrons on the stationary phase nitrogen and the π electron cloud of the solute PNAs. Elution order seems to depend upon a combination of π energy, and type of ring condensation of the solute. Plots of log I versus number of aromatic carbons for catacondensed PNAs suggest that while the specific interaction is different than that seen in silica chromatography, the overall adsorption effect is comparable. In the reversed phase there may be two types of separation mechanisms: 1) a pure partitioning effect in highly polar mobile phases (methanol/water), or 2) a mixture of liquid-solid adsorption and liquid-liquid partition in less polar solvent systems (acetonitrile/water).

INTRODUCTION

Polynuclear aromatic compounds (PNAs) constitute an important class of chemical pollutants, many of which are known or suspect carcinogens and mutagens (1, 2). Several industrial sources of these pollutants have been identified and include power plants, motor vehicles and oil refineries. Reliable analytical methods are needed to determine their presence since many of these compounds are on the EPA priority pollutant list. Due to the fact that PNAs are commonly found in complex matrices, it has been necessary in the past to combine several analytical techniques to provide complete characterization.

A typical analysis scheme for PNAs would include a pre-separation step such as solvent extraction or adsorption on silica or alumina, and then a subsequent analysis by GC and/or LC. However, solvent extraction and the use of "classical" adsorbents can be time consuming and tedious. Moreover, impurities such as water in the mobile phase can significantly modify the adsorbent, producing a non-reproducible separation (3). These adsorbents display a strong attraction for PNAs and often trace components will be strongly retained on the column. This results in decreased recovery and poor separation. As an alternative, gel filtration on Sephadex LH-20 has been incorporated into a separation procedure by Giger and Blumer (4). Novotiny et al. (5)

used Sephadex LH-20 for initial purification and then an amino bonded phase for further fractionation of PNAs. Studies by others (6,7) have demonstrated that complex mixtures can be resolved by using adsorption or gel chromatography followed by reversed phase HPLC.

Considerable interest has been focused on the class separation of PNAs on alkyl amine chemically bonded phase columns. Galya and Suatoni (8) have reported using a μ Bondapak NH_2 column to separate saturates, aromatics, resins, and asphaltenes (SARA) from crude oils. Wise and co-workers (9) have used the same column for the HPLC separation of crude oils and marine sediment samples. Chmielowiec and George (10) in a subsequent study of several polar stationary phases found superior PNA selectivity for a diamino bonded phase column. Similarly, a chemically bonded pyrrolidone substrate has been used to separate PNAs in shale oil samples (11). These three studies reported separations which were performed in the normal phase and which exhibited an elution order generally based on ring number and the type of ring condensation. The separation mechanism on the amino, diamino, and pyrrolidone column packings, in the normal phase mode, is postulated to be a charge-transfer interaction between the lone pair electrons of the stationary phase nitrogen and the π electron cloud of the PNAs. An interesting development in Mourey's work was the investigation of a reversed phase separation on pyrrolidone (11). These experiments demonstrated that normal and reversed phase

separations are complementary in terms of selectivity with the latter technique more efficient for solutes with less than six aromatic rings.

Separation via adsorption chromatography is based on the competition of the solute and solvent molecules for active sites on the stationary phase. In classical adsorption, the active sites on the adsorbent generally interact with polar functional groups on compounds to be separated. The elution order in this type of separation is usually predictable and follows the constituent polarity. In a charge transfer mechanism, the interaction between the adsorbent surface and solute is more specific. When the mobile phase is relatively non-polar, the lone pair electrons on the polar amine functionality of the stationary phase interact with the π cloud on the PNAs forming a weak bond. Elution order in this case may be dependent more on either the electron affinity or the total energy of the π electron system rather than its polarity. A modification of this explanation may be required, however, when considering the interactions involved when the mobile phase is more polar.

In order to explore further the separation mechanism of the n-propyl amine bonded phase column, this work reports the retention of PNAs on this stationary phase in both the normal and reversed phase modes. The selectivity and capacity factors are given for several examples of three, four, and five ring PNAs which can be considered as members of different ring classes.

This data as well as theoretical considerations are discussed in terms of previously described mechanisms.

EXPERIMENTAL

Apparatus

All data was obtained on a microprocessor controlled Altex model 312 MP liquid chromatograph with a 100 μ L sample loop. The PNAs were separated on a 30 cm X 10 mm i.d. Chromosorb LC-9 column (Supelco Inc., Bellefonte, PA) and monitored with a UV absorption detector at 254 nm.

Reagents and Standards

All PNAs used were purchased at 95% purity or better and were used without further purification. Mobile phase eluents were glass distilled cyclohexane for normal phase chromatography, and spectrograde methanol, glass distilled acetonitrile, and type three reagent grade water obtained using a Milli-Q water purification system, for reversed phase chromatography. Reagent grade isopropanol, and 99+% dichloromethane were solvents used for converting from the normal to the reversed phase modes. Standards for the normal phase mode were prepared by dissolving the appropriate amount of each standard in cyclohexane to prepare solutions with concentrations of approximately 10^{-4} M. For the reversed phase work, standards were prepared in both acetonitrile and methanol.

Procedure

The PNA standards were run in the normal phase (100% cyclohexane) and the retention times recorded. The column was then converted to reversed phase by flushing with 20 column volumes each of isopropanol, dichloromethane and methanol. In the final step, equilibrium was attained with an additional 100 column volumes of methanol. The retention times for the PNA standards were measured in mobile phases consisting of varying percentages of methanol/water or acetonitrile/water. In both modes of chromatography, the flow rate was 2 mL/min. After all reversed phase work was complete the column was converted back to normal phase by flushing with the above solvents in the reverse order. Comparison of the number of theoretical plates before and after conversion to reverse phase showed no significant changes indicating little or no irreversible effects.

RESULTS AND DISCUSSIONS

Chromosorb LC-9 packing consists of an n-propyl amino silane chain chemically bonded to 10 micron silica particles. Because the surface of this material is highly polar, it is useful for separating polar compounds. However, when used with a non-polar mobile phase, such as cyclohexane, non-polar aromatic hydrocarbon separation is possible, analogous to the separation obtained on silica or alumina. Generally, elution volumes seem to increase with the number of condensed rings.

Because the retention times of various solutes can change

from run to run due to minor experimental differences, it is necessary to relate retention time to a parameter independent of these differences. One possibility is to use the retention index (I). This is a parameter derived in a manner analogous to the Kovat's index for gas chromatography. The calculation is based on a linear relationship between $\log V_R$ and the number of aromatic rings. The basic assumption is that slight variations in chromatographic conditions which cause a shift in retention of one PNA will also affect the retention of all other PNAs although their position relative to each other will not change. In other words, the observed V_R will change but I values will not.

The retention indices for several PNAs were calculated as previously described by Popl et al. (12). To compare the retention characteristics of PNAs on several nitrogen containing stationary phases, the logarithms of the retention indices (I) for several PNAs on μ Bondapak NH_2 (13), diamine (10), pyrrolidone (11) and Chromosorb LC-9 are given in Table I. The hydrocarbon standards for the μ Bondapak NH_2 , pyrrolidone, and diamine packing material were assigned the following retention indices: benzene 10, naphthalene 100, phenanthrene 1000, 1,2-benzanthracene 10,000 and benzo (b) chrysene 100,000. For Chromosorb LC-9, the same hydrocarbon standards were used with the following exceptions: anthracene 1000, and 1,2,3,4-dibenzanthracene 100,000. As reported by Popl et al. (14), the selection of the index compounds for separation of PNAs on alumina was dictated by an observed pH

Table I

Comparison of the Log of Retention Index (log I) of PNAs on Several Nitrogen Containing Liquid Chromatographic Packing Materials

Compound	Chromosorb LC-9	μ Bondapak	Pyrrolidone	Diamine
	(Cyclohexane)	NH ₂ (Hexane)	(Heptane)	(n-Heptane)
<u>One Ring Aromatics</u>	log I ^a	log I ^b	log I ^c	log ^b
Benzene	1.00	1.00	1.00	1.00
<u>Two Ring Aromatics</u>				
Biphenyl	1.93	2.16	1.40	2.25
Naphthalene	2.00	2.00	2.00	2.00
Acenaphthylene	2.64	2.59	2.26	3.20
<u>Three Ring Aromatics</u>				
Fluorene	2.59	2.55	2.26	2.50
p-Terphenyl	3.03	3.28	2.88	3.49
Anthracene	3.00	2.94	2.96	2.92
Phenanthrene	3.12	3.00	3.00	3.00
9,10-Diphenyl- anthracene	2.57	—	3.16	3.47
Dibenzothiophene	2.77	2.75	—	—
<u>Four Ring Aromatics</u>				
Fluoranthene	3.60	3.51	3.46	3.59
Pyrene	3.43	3.37	3.47	3.43
Chrysene	3.87	3.90	3.83	—
p-Quaterphenyl	4.05	4.50	3.90	—
Triphenylene	4.23	4.07	3.91	—
1,2-Benzanthracene	4.00	4.00	4.00	4.00
<u>Five Ring and Larger Aromatics</u>				
Perylene	4.70	4.61	3.92	—
Benz(a)pyrene	4.44	4.38	4.11	—
Benz(e)pyrene	4.60	4.46	—	—
Benz(g,h,i)perylene	5.00	4.83	4.45	—
1,2,5,6-Dibenz- anthracene	4.68	4.94	4.60	—
1,2,3,4-Dibenz- anthracene	5.00	4.93	4.64	—
Picene	—	5.03	4.63	—
Benze(b)chrysene	—	5.00	—	5.00



^aIndex compounds were assigned retention indices as follows: benzene 10, naphthalene 100, anthracene 1000, 1,2-benzanthracene 10,000 and 1,2,3,4-dibenzanthracene 100,000.

^bIndex compounds were assigned retention indices as follows: benzene 10, naphthalene 100, phenanthrene 1000, 1,2-benzanthracene 10,000 and benzo(b)chrysene 100,000, and are taken directly from the references.

^cIndex compounds were assigned retention indices as follows: benzene 10, naphthalene 100, phenanthrene 1000, and 1,2-benzanthracene 10,000, and were calculated from the k' values given in reference 11. For compounds with higher elution volumes, the linear dependence of log V_r versus log I was extrapolated to the region of I \geq 10,000.

dependence. For PNAs with certain structural characteristics such as an acidic hydrogen (fluorene) or low ionization potentials (anthracene), it was found that retention increased on basic alumina and decreased on acidic alumina. Popl chose index compounds for which the $\log V_r$ vs $\log I$ relationship was linear because of this acid-base effect.

Although we are dealing with charge transfer interactions, the same criteria Popl used can be applied when choosing PNAs for index compounds on amine polar bonded phases. That is, those compounds for which the $\log V_r$ versus $\log I$ relationship is linear would be suitable index compounds. Even though the compounds chosen above are not necessarily charge transfer homologs, if a linear relationship is satisfied, then the compounds are satisfactory index compounds. Figure 1 shows a plot of $\log I$ vs $\log V_r$ on Chromosorb LC-9 for the index compounds chosen. The plot is linear with a coefficient of determination, r^2 , of .9984. Although this r^2 is based on a small data set (only five points), the probability of five unrelated data points giving an r value this high would be .001 (15). If the choice of index compounds is not critical, then one would suspect that the $\log I$ values of the same PNAs on similar columns should yield similar relationships, even if the index compounds chosen to calculate the $\log I$ values are slightly different. Figure 2 shows a plot of $\log I$ vs $\log V_r'$ (the corrected retention volume) for the same compounds on Chromosorb LC-9. The $\log I$ values were calculated using the



retention volumes of the index compounds on μ Bondapak NH_2 . Phenanthrene was used as a three ring standard and benzo (b) chrysene for a five ring index standard. This relationship is also linear with a coefficient of determination of .9908. The probability of obtaining this r value with 14 unrelated points is .001 (15). It should be noted that the new index standard will not produce identical retention indices for both amine columns; however, the relationships between individual PNAs is preserved. Anthracene has a log I value of 3.00 and 2.94 on the Chromosorb LC-9 and μ Bondapak NH_2 columns respectively. Likewise, phenanthrene has log I values of 3.12 and 3.00. Although these values are not identical, phenanthrene has a larger log I value than anthracene on both columns. Figure 2 shows that the relationship of each individual PNA to any other is the same using both sets of standards. Therefore, one may conclude that choosing anthracene instead of phenanthrene for a three ring index standard, and 1,2,3,4-dibenzanthracene in place of benzo (b) chrysene for a five ring index standard will not significantly affect the data comparison, even if the index compounds are not charge transfer homologs.

If a polar bonded phase amine packing material separates primarily via a charge transfer mechanism, then one would expect that the elution order of PNAs would be dictated by some parameter of the solute PNA which reflects the electron transfer interaction. Some possible choices would be total π energy,

electron affinity (EA), the reduction potential ($E_{1/2 \text{ red}}$), ionization energy (IE), the energy of the lowest unoccupied molecular orbital (LUMO) or the highest occupied molecular orbital (HOMO). Some values for these constants for several PNAs are given in Table II. Plots of $\log I$ versus each of these parameters were made for each different type of packing material. The coefficients of determination for these plots are given in Table III. The parameter r^2 is the coefficient of determination which was determined for each plot, n is the number of data points used and P is the probability of obtaining or exceeding the given r using completely unrelated data (15). Ranking of r^2 indicates that the plot of $\log I$ vs π energy is the best fit to the data, ionization energy is next and all others show a much poorer fit. This correlation supports the theory that the separation mechanism involves a charge transfer interaction between the nitrogen containing stationary phase and the solute PNA. This conclusion is also supported by the fact that both Wise (9) and Mourey (11) noticed that the addition of alkyl substituents does not significantly shift the retention of a PNA on either an amine or pyrrolidone bonded phase. This is because addition of an alkyl group does not appreciably change the total π energy.

Table IV gives the coefficient of determinations for $\log I$ vs π energy plots for different ring classes on different packing materials. It appears that the coefficient for catacondensed PNAs is generally greater than that for pericondensed solutes. This

Table II

Solute Parameters Used in Adsorption Mechanism Study

Compound	Π Energy [16]	EA(eV) [15]	IE(eV) [18]	$E_{1/2\text{red}}(\text{V})$ [17]	LUMO [16]	HOMO [16]
Benzene	5.87	—	9.22	-2.30	0.800	-0.800
Biphenyl	11.85	0.085	8.36	—	0.599	-0.599
Naphthalene	8.12	0.152	8.28	-1.54	0.535	-0.534
Acenaphthylene	11.87	—	8.21	—	0.550	-0.555
Fluorene	11.85	—	—	—	0.599	-0.596
p-Terphenyl	17.84	—	—	—	0.514	-0.519
Anthracene	13.81	0.552	7.76	-1.09	0.375	-0.370
Phenanthrene	13.94	0.308	8.18	-1.50	0.526	-0.520
Fluoranthene	16.04	—	8.14	-1.45	0.535	-0.403
Pyrene	16.01	0.579	7.85	-1.16	0.400	-0.404
Chrysene	17.99	0.419	8.17	-1.35	0.460	-0.460
p-Quaterphenyl	23.83	—	—	—	0.473	-0.470
Triphenylene	18.07	0.284	—	-1.55	0.584	-0.580
1,2-Benz- anthracene	17.90	0.696	7.79	-0.85	0.320	-0.326
Perylene	20.06	—	7.49	-0.85	0.320	-0.326
Benz(a)pyrene	20.04	0.829	—	—	0.340	-0.340
Benz(e)pyrene	20.15	0.486	7.85	-1.24	0.486	-0.483
Benz(ghi)perylene	22.25	—	7.76	-1.01	0.396	-0.396
1,2,5,6-Dibenz- anthracene	21.99	0.676	—	-1.19	0.423	-0.426
1,2,3,4-Dibenz- anthracene	22.05	—	—	-1.25	0.444	-0.444
Picene	22.03	.0490	7.23	-1.33	0.446	-0.446

may be due in part because the retention index standards are catacondensed structures, but it also may imply that a steric effect should be considered as well. Work done by Hammers, Spanjer, and DeLigny (19) on an amino butyl polar bonded phase column indicates the presence of a steric effect. These authors noticed a discrepancy in the behavior of flat fused arenes and

TABLE III

Comparison of Coefficients of Determination, r^2 , for Plots of Log I versus Solute Parameters^a

Solute Parameter ^b	Chromosorb LC-9			μ Bondapak NH ₂			Pyrrolidone			Diamine		
	r^2	n	P	r^2	n	P	r^2	n	P	r^2	n	P
π Energy	0.880	20	0.001	0.948	21	0.001	0.863	11	0.001	0.891	11	0.001
IE (eV)	0.702	13	0.001	0.757	14	0.001	0.712	13	0.001	0.726	9	0.01
EA (eV)	0.526	11	0.02	0.457	12	0.02	0.548	11	0.02	0.836	6	0.01
$E_{1/2\text{red}}(\text{V})$	0.590	16	0.001	0.535	15	0.01	0.531	14	0.01	0.708	7	0.01
LUMO	0.550	20	0.001	0.530	21	0.001	0.557	20	0.001	0.618	11	0.01
HOMO	0.381	20	0.01	0.347	21	0.01	0.369	20	0.01	0.669	11	0.01

^aSee Tables I and II for data^bAll π Energies, LUMO and HOMO numerical data were obtained using the Huckel molecular orbital theory. r^2 is the coefficient of determination, n is the number of data points and P is the probability of obtaining or exceeding the given r with unrelated data.

Table IV

Coefficients of Determination for Plots of Log I Versus Π Energy for Different N-Containing Packings^a

Ring Condensation Type	Pyrrolidone		Chromosorb LC-9		μ Bondapak NH ₂	
Non-Condensed	0.928	n=5	0.946	n=5	0.982	n=5
Peri-Condensed	0.954	n=6	0.985	n=7	0.984	n=7
Cata-Condensed	0.993	n=9	0.979	n=8	0.989	n=9

^aSee Table I for data

slightly twisted polyphenyls. Because the interaction is charge transfer, it follows that the greater the π conjugation the stronger the interaction. All else being equal, the amount of π conjugation depends upon the arrangement of the molecules. For example, interaction appears to be greater for flat molecules than for slightly twisted polyphenyls.

Popl et al. (12) found that on silica, there is a linear dependence of $\log I$ on the number of aromatic carbon atoms for parent catacondensed PNAs. They found that the following equation describes the observed effect,

$$\log I = 0.25 n_{C_{Ar}} - 0.5$$

where $n_{C_{Ar}}$ is the number of aromatic carbon atoms in the molecule. Table V gives the equation of the line which describes the relationship between carbon number and $\log I$ for catacondensed PNAs on μ Bondapak NH_2 , the diamine, and pyrrolidone packing

Table V

Equations Describing the Relationship Between $\log I$ and Number of Aromatic Carbons in Catacondensed PNAs

Pyrrolidone	$\log I = 0.230 n_{C_{Ar}} - 0.286$
μ Bondapak NH_2	$\log I = 0.249 n_{C_{Ar}} - 0.496$
Diamine	$\log I = 0.245 n_{C_{Ar}} - 0.497$
Silica	$\log I = 0.25 n_{C_{Ar}} - 0.5$

materials. In the monoamine, and especially the diamine systems, the equations are nearly identical to the silica relationship. This indicates that the interaction is similar to adsorption on silica. However, this is not to say that all PNAs behave the same on both the silica and the amine polar bonded phases. Alkyl substitution of PNAs results in greater chromatographic differences on silica than on the amine packing material as shown by Wise et al. (9). Therefore, the separation mechanisms or "adsorption" interactions are different, although the overall adsorption effects are similar.

After examining the performance of Chromosorb LC-9 in the normal phase, we converted the column to the reversed phase to examine its behavior under these conditions. The capacity factors for several PNAs in 100% CH₃CN and 100% MeOH are given in Table VI. The capacity factors are very small, indicating that this particular column does not retain PNAs very strongly under these conditions and would not be a practical way to chromatograph PNAs. However, the relative retention (or lack of retention) of these compounds in the reversed phase should reveal information about the mechanism of separation of the N-containing packing materials. With the more polar solvents used in reversed phase chromatography, adsorption effects are minimized and thus retention is shorter. Mourey et al. (11) reported much higher k' values for the same PNAs in an even more polar mobile phase (60%

methanol/water) on pyrrolidone, a cyclic amine bonded phase. The best explanation for this increased retention is a combination of two effects: 1) the bonded phase polarity and 2) the mobile phase polarity. Obviously, the pyrrolidone functionality is much less

Table VI

Capacity Factors, k' , for Selected PNAs on the Amino Bonded Column in the Reversed Phase

Compound	100% CH ₃ CN	100% MeOH
<u>Two Rings</u>		
Naphthalene	0.121	0.170
<u>Three Rings</u>		
Anthracene	0.165	0.188
<u>Four Rings</u>		
p-Quaterphenyl	0.123	0.136
Chrysene	0.170	0.180
Triphenylene	0.177	0.180
1,2-Benzanthracene	0.167	0.170
9,10-Dimethyl-1,2-benzanthracene	0.165	0.165
Fluoranthene	0.170	0.121
Pyrene	0.159	0.175
<u>Five Rings and More</u>		
Benz(a)pyrene	0.177	0.234
Benz(e)pyrene	0.180	0.244
1,2,3,4-Dibenzanthracene	0.167	0.229
Pentacene	0.175	0.159
9,10-Diphenylanthracene	0.144	0.141
Perylene	0.203	0.252
Benz(g,h,i)perylene	0.196	0.275
Rubrene	0.139	0.149

polar than the n-propyl amine in the reversed phase. Thus it is possible to chromatograph PNAs on pyrrolidone in the reversed phase but not on an amino bonded phase column.

Adding water to a silica column can greatly affect the activity of the column and cause irreproducible effects. However, when the silanol groups are chemically bonded with some polar functional group (such as these nitrogen containing moieties) the column becomes much more stable to water. Although it is likely that due to incomplete coverage there are some exposed silanol groups which can interact with water, the percentage of these is low, and reproducible results can be achieved if the column is allowed to come to equilibrium with the mobile phase.



Although the k' values are very small they seemed to be very reproducible. Table VII shows the retention of three PNAs (naphthalene, anthracene, and pyrene) on Chromosorb LC-9 using different mobile phases. The standard deviation of each point is listed along with the number of repetitions. This data is presented graphically in Figures 3 and 4. The limits of the standard deviation are marked on the last point of each plot since this is the only place the deviation was large enough to visually observe. The upper limit of the percent water in the mobile phase is determined by the solubility of the PNAs. In the methanol/water system (Figure 3), the retention of each PNA seems

Table VII

Retention of Three PNAs on Chromosorb LC-9 Using Varying Percentages of Methanol/Water and Acetonitrile/Water

Percent Methanol	Number of Repetitions	Pyrene	Anthracene	Naphthalene
100	8	20.60 \pm 0.01	19.92 \pm 0.02	10.40 \pm 0.02
60	8	26.14 \pm 0.11	23.50 \pm 0.05	21.33 \pm 0.05
40	5	54.72 \pm 0.68	37.14 \pm 0.42	25.62 \pm 0.14
Percent Acetonitrile				
100	5	19.25 \pm 0.02	18.98 \pm 0.03	18.83 \pm 0.05
80	8	16.92 \pm 0.03	16.72 \pm 0.02	16.83 \pm 0.05
60	8	16.10 \pm 0.07	15.97 \pm 0.17	15.81 \pm 0.25
40	8	21.60 \pm 0.09	20.55 \pm 0.06	19.47 \pm 0.05
20	5	56.51 \pm 0.67	49.70 \pm 1.00	26.04 \pm 0.72

to increase as the polarity of the solvent increases. For reversed phase chromatography, this is exactly what would be expected if the separation occurred according to a partition mechanism. In the acetonitrile/water system (Figure 4), however, results are a bit more interesting. The three compounds are strongly retained on the column in 80% water/acetonitrile. As the percent water is decreased the retention volumes decrease with a minimum occurring at about 40% water/acetonitrile. At 100% acetonitrile (0% water), however, the elution volumes again increase. Similar effects have been observed by Goldstein (20) on pyrrolidone in isopropanol, and have been compared to parallel behavior observed on silica. As the composition of the mobile



phase changes from 100% acetonitrile to acetonitrile/water, the stationary phase surface adsorbs an aqueous layer, prohibiting solute interaction with the stationary phase and consequently solute adsorption is decreased. As more water is added, the aqueous layer becomes more extensive and the solutes slowly begin to preferentially partition into the stationary phase. Thus, there are two effects in competition here. At high percentage of organic modifier, a liquid-solid adsorption effect seems to be predominant which becomes less important as a more aqueous solvent system is reached where a liquid-liquid partitioning effect seems to be more important.

CONCLUSIONS

The separation mechanism for PNAs on the amino, diamino, and pyrrolidone columns in the normal phase is postulated to be a charge transfer interaction between the lone pair electrons on the stationary phase and the π electron cloud. This possible mechanism is supported by the high correlation between the log I values and the total π energy of the PNAs studied. Ranking of coefficients of determination for these plots indicate that π energy has the best fit to the data and the ionization potential is the next best. These values were taken from Hückel molecular orbital calculations and possibly more exact calculations for π energy could give a better fit. According to the r^2 values, it would seem that pericondensed PNAs are retained differently than

catacondensed PNAs, suggesting that there could be a geometric effect in addition to π energy and ionization potential.

Plots of log I versus number of aromatic carbons for catacondensed PNAs on the monoamine, diamine, and pyrrolidone materials suggest that the retention is due to interactions resulting in adsorption effects which are similar in both silica and amine separations. However, the specific interactions are different than those seen in silica chromatography.

The reversed phase data show that the selectivity on Chromosorb LC-9, in general, for PNAs in 100% acetonitrile or 100% methanol is poor. It is difficult to classify bonded phase, reverse phase chromatography as either a strict partition or adsorption process. The data seems to point to both processes happening. With methanol/water systems the data seems to indicate a partitioning mechanism. On the other hand, in an acetonitrile water system the mechanism seems to be a mixture of liquid-solid adsorption and liquid-liquid partition, depending upon the polarity and water content of the mobile phase.

In general, our data support previously drawn conclusions concerning the separation mechanisms of PNAs on nitrogen-containing supports. However, to our knowledge, these conclusions have not been supported experimentally. We have formulated these conclusions from both normal and reversed phase studies using data from our experiments as well as the published results of other studies.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the technical assistance of Ms. Karen Timmons. This work was supported in part by contracts from the Department of Energy (Contract No. De-AS05-80EV10404), the Office of Naval Research and a grant from the Robert A. Welch Foundation.

CURRENT AUTHOR AFFILIATION

Dr. I. M. Warner, Chemistry Department, Emory University, Atlanta, Georgia 30322

REFERENCES

1. Møller, M. and Alfheim, I., Mutagenicity and PAH Analysis of Airborne Particulate Matter, *Atmos. Environ.*, 14, 83, 1980.
2. Nordén, B., Edlund, U. and Wold, S., Carcinogenicity of Polycyclic Aromatic Hydrocarbons Studied by SIMCA Pattern Recognition, *Acta Chem. Scand. B.*, 32, 602, 1978.
3. Snyder, L. R., Principles of Adsorption Chromatography, Marcel Dekker, New York, 1968.
4. Giger, W. and Blumer, M., Polycyclic Aromatic Hydrocarbons in the Environment; Isolation and Characterization by Chromatography, Visible, Ultraviolet, and Mass Spectrometry, *Anal. Chem.*, 46, 1663, 1974.
5. Novotny, M., Lee, M. L., and Bartle, K. D., The Methods for Fractionation, Analytical Separation and Identification of Polynuclear Aromatic Hydrocarbons in Complex Mixtures, *J. Chromatogr. Sci.*, 12, 606, 1974.
6. Schmit, J. A., Henry, R. A., Williams, R. C., and Dieckman, J. F., Applications of High Speed Reversed Phase Liquid Chromatography, *J. Chromatogr. Sci.* 9, 645, 1971.
7. Golden, C. and Sawicki, E., Ceramic Alumina in the HPLC Determination of Benzo(a)pyrene in Air Particulate Material, *Anal. Lett.* 9, 957, 1976.

8. Galya, L. C. and Suatoni, J. C., Rapid SARA Separation by High Performance Liquid Chromatography, 3, 229, 1980.
9. Wise, S. A., Chesler, S. N., Hertz, H. S., Hilpert, L. R. and May, W. E., Chemically-Bonded Aminosilane Stationary Phase for the High-Performance Liquid Chromatographic Separation of Polynuclear Aromatic Compounds, Anal. Chem., 49, 2306, 1977.
10. Chmielowiec, J. and George, A. E., Polar Bonded-Phase Sorbents for High Performance Liquid Chromatographic Separations of Polycyclic Aromatic Hydrocarbons, Anal. Chem., 52, 1152, 1980.
11. Mourey, T. H., Siggia, S., Uden, P. C. and Crowley, R. J., High Performance Liquid Chromatographic Separation of Polycyclic Aromatic Hydrocarbons on Microparticulate Pyrrolidone and Application to the Analysis of Shale Oil, Anal. Chem., 52, 885, 1980.
12. Popl, M., Dolansky, V., and Mostecky, J., Influence of the Molecular Structure of Aromatic Hydrocarbons on Their Adsorptivity on Silica Gel, J. Chromatogr., 117, 117, 1976.
13. Wise, S. A., Bonnett, W. J. and May, W. E., Polynuclear Aromatic Hydrocarbons: Chemistry and Biological Effects, Bjørseth, A. and Dennis, A., Eds., Battelle Press, Columbus, Ohio, 1980, p. 791.
14. Popl, M., Dolansky, V. and Mostecky, J., Influence of the Molecular Structure of Aromatic Hydrocarbons on Their Adsorptivity on Alumina, J. Chromatogr. 91, 649, 1974.
15. Franklin, J. L., The Handbook of Spectroscopy, CRC Press Inc., Cleveland, Ohio, (In Press).
16. Coulson, C. A., and Streitwieser, A., Dictionary of Electron Calculations, Pergamon Press Inc., San Francisco, California, 1966.
17. Pysh, E. S. and Yang, N. C., Polarographic Oxidation Potentials of Aromatic Compounds, J. Amer. Chem. Soc. 85, 2124, 1963.
18. Krishna, B. and Bahuguna, G. P., Revised IOC- ω Technique, Indian J. Pure Appl. Phys., 16, 766, 1978.
19. Hammers, W. E., Spanjer, M. C. and DeLigny, C. L., Selectivity of Nucleosil 10 NH₂ as an Adsorbent in High Performance Liquid Chromatography, J. Chromatogr., 174, 291, 1979.

20. Goldstein, G., Separation of Polycyclic Aromatic Hydrocarbons by Liquid Chromatography on Cross-Linked Polyvinylpyrrolidone, *J. Chromatogr.*, 129, 61, 1976.

FIGURE CAPTIONS

FIGURE 1. Calibration plot for PNAs on Chromosorb LC-9.

FIGURE 2. Comparison calibration plot.

FIGURE 3. Change in retention volumes of some polynuclear aromatic hydrocarbons with water content in a methanol/water solvent system. The retention volumes are not corrected for the column void volume (15.3mL).

FIGURE 4. Change in retention volumes of some polynuclear aromatic hydrocarbons with water content in an acetonitrile/water solvent system. The retention volumes are not corrected for the column void volume (15.3 mL).