

1  
OCTOBER 1981

14. 59  
PPPL-1834  
uc-20G

11-9-81

8233

**MASTER**

NONLINEAR GYROKINETIC EQUATIONS FOR  
LOW-FREQUENCY ELECTROMAGNETIC WAVES  
IN GENERAL PLASMA EQUILIBRIA

BY

E.A. FRIEMAN AND LIU CHEN

**PLASMA PHYSICS  
LABORATORY**



**PRINCETON UNIVERSITY  
PRINCETON, NEW JERSEY**

This work was supported by the U.S. Department of Energy  
Contract No. DE-AC02-76-CHO 3073. Reproduction, transla-  
tion, publication, use and disposal, in whole or in part,  
by or for the United States government is permitted.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. It is the property of the United States Government and is loaned to your organization; it and its contents are not to be distributed outside your organization. For copies of this report, contact the Office of Scientific and Technical Information, P.O. Box 616, Springfield, VA 22161-0616. For all other information, contact the Office of Primary Distribution, P.O. Box 909, Gaithersburg, MD 20878-0909. This report is available in microfiche and microfilm editions. For more information, contact the Office of Scientific and Technical Information, P.O. Box 616, Springfield, VA 22161-0616. This report is available in microfiche and microfilm editions. For more information, contact the Office of Scientific and Technical Information, P.O. Box 616, Springfield, VA 22161-0616.

Nonlinear Gyrokinetic Equations for Low-Frequency  
Electromagnetic Waves in General Plasma Equilibria

by

E. A. Frieman\* and Liu Chen

Plasma Physics Laboratory, Princeton University

Princeton, New Jersey 08544

ABSTRACT

A nonlinear gyrokinetic formalism for low-frequency (less than the cyclotron frequency) microscopic electromagnetic perturbations in general magnetic field configurations is developed. The nonlinear equations thus derived are valid in the strong-turbulence regime and contain effects due to finite Larmor radius, plasma inhomogeneities, and magnetic field geometries. The specific case of axisymmetric tokamaks is then considered, and a model nonlinear equation is derived for electrostatic drift waves. Also, applying the formalism to the shear Alfvén wave heating scheme, it is found that nonlinear ion Landau damping of kinetic shear-Alfvén waves is modified, both qualitatively and quantitatively, by the diamagnetic drift effects. In particular, wave energy is found to cascade in wavenumber instead of frequency.

\*Present address: Science Application Inc., La Jolla, CA 92038 and  
University of California, San Diego, CA, 92092.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

## I. INTRODUCTION

Electromagnetic instabilities with frequencies lower than the ion-cyclotron frequency and perpendicular (to the magnetic field) wavelengths comparable to the ion Larmor radius are believed to be important for the transport processes in magnetically confined plasmas. One complicating factor in analyzing these low-frequency microscopic instabilities, which are driven by plasma inhomogeneities and shall be loosely termed kinetic drift-Alfvén waves, is that the destabilizing mechanisms are sensitive to effects associated with magnetic field configurations such as magnetic shear, magnetic gradient and curvature drifts, and trapped particles. To overcome this difficulty, Rutherford and Frieman<sup>1</sup> as well as Taylor and Hastie<sup>2</sup> have developed, independently, a formalism, now known as the gyrokinetic formalism, to treat the linear aspects of kinetic drift waves in general magnetic configurations. Recently, the linear gyrokinetic formalism has been extended to include electromagnetic perturbations<sup>3,4</sup> associated with shear and compressional Alfvén waves. Since the transport induced by the instabilities is ultimately determined by the nonlinear processes, it is, therefore, desirable to further extend the gyrokinetic formalism into the nonlinear regime while retaining crucial features such as finite Larmor radius and arbitrary magnetic field configurations. This constitutes the principal motivation of the present research.

In this work we develop the nonlinear gyrokinetic formalism based on a multiple-scale expansion.<sup>5</sup> That is, microscopic fluctuations vary on the fast (linear) time scale (i.e., typically, the diamagnetic drift frequency time scale), while macroscopic quantities are assumed to vary on the slow transport time scale. Here the cyclotron frequency time scale is the fastest time scale. Furthermore, consistent with experimental

observations,<sup>6,7</sup> our formulation allows nonlinear time scales to be comparable to the linear ones and, hence, the results are valid in the strong-turbulence regime.

Section II contains the theoretical formulation and derivations of the nonlinear gyrokinetic equations. The specific case of axisymmetric tokamaks is further considered in Sec. III using the ballooning-mode representation<sup>8-10</sup> and a nonlinear drift-wave equation is derived in a limiting case. Noting that our results are also applicable to nonlinear heating via externally launched low-frequency waves, we also consider, in Sec. IV, nonlinear ion Landau damping (ion induced scattering) of the mode-converted kinetic (shear) Alfvén waves<sup>11</sup> and find that the inclusion of diamagnetic drift effects modifies, both qualitatively and quantitatively, the decay processes. Final conclusions and discussion are given in Sec. V.

## II. THEORETICAL ANALYSES

(1) Guiding-center transformation and the two spatial scales. As in the linear formalism,<sup>1,2,12</sup> it is more convenient to carry out the analysis in the guiding-center phase space  $(\tilde{x}, \tilde{v})$  which is related to the particle phase space  $(x, v)$  via the following guiding-center transformation

$$\tilde{x} = x + v \times \underline{e}_{\parallel} / \Omega \quad , \quad (1)$$

$$\tilde{v} = v(\varepsilon, \mu, \alpha) \quad , \quad (2)$$

where  $\varepsilon = v^2/2 + q\phi_0/m$ ,  $\mu = v_{\perp}^2/2B$ ,  $\phi_0$  and  $B$  are, respectively, the macroscopic (equilibrium) electric potential and magnetic field,  $\Omega = qB/mc$ ,  $\underline{e}_{\parallel} = B/B$ ,  $\alpha$  is the gyrophase, and

$$\underline{v}_\perp = v_\perp (\underline{e}_1 \cos \alpha + \underline{e}_2 \sin \alpha) , \quad (3)$$

with  $\underline{e}_1$ ,  $\underline{e}_2$ , and  $\underline{e}_\parallel$  being the local orthogonal unit vectors. Furthermore, noting that perturbations of interest here have perpendicular (to  $\underline{B}$ ) wavelengths of the order of the Larmor radius,  $\rho$ , which is much smaller than the macroscopic scale length,  $L_0$ , (i.e.,  $\lambda \equiv \rho/L_0$  is a small parameter), we may separate physical quantities into microscopic and macroscopic parts by averaging over the microscopic spatial variations. Thus

$$\underline{P} = \underline{P} + \delta \underline{P} , \quad (4)$$

where

$$\underline{P} = \int d^2 \underline{x}_{\perp 10} \underline{P} / \int d^2 \underline{x}_{\perp 10} \equiv \langle \underline{P} \rangle_{\underline{x}_{\perp 10}} , \quad (5)$$

$\langle \delta \underline{P} \rangle_{\underline{x}_{\perp 10}} = 0$  and  $\underline{x}_{\perp 10}$  corresponds to fast perpendicular spatial variations.

In the  $(\underline{x}_\perp, \underline{v})$  phase space, the Vlasov equation then becomes

$$L_g f = -(q/m)(\delta R f) , \quad (6)$$

where

$$L_g = \partial/\partial t + \partial \underline{v}/\partial t \cdot \nabla_{\underline{v}} + \partial \underline{x}/\partial t \cdot \nabla_{\underline{x}} + v_\parallel \underline{e}_\parallel \cdot \nabla_{\underline{x}} \\ + v_\perp (\lambda_{B1} + \lambda_{B2}) + (q/m)(\underline{E} - \underline{E}_0) \cdot \nabla_{\underline{v}} \partial/\partial \epsilon$$

$$\begin{aligned}
 & + (q/m) \underline{\tilde{E}} \cdot [ (\underline{v}_1/B) (\partial/\partial \mu) + (\underline{e}_\alpha/v_1) (\partial/\partial \alpha) ] \\
 & + \underline{v}_E \cdot \underline{\tilde{v}}_X - \Omega \partial/\partial \alpha \quad , \quad (7)
 \end{aligned}$$

$$\underline{\lambda}_{B1} = \underline{v} \times \underline{\tilde{v}}_X (\underline{e}_\alpha/\Omega) \cdot \underline{\tilde{v}}_X \quad , \quad (8)$$

$$\underline{\lambda}_{B2} = (\underline{\tilde{v}}_X \mu) (\partial/\partial \mu) + (\underline{\tilde{v}}_X \alpha) (\partial/\partial \alpha) \quad , \quad (9)$$

$$\underline{\tilde{v}}_X \mu = -[\mu \underline{\tilde{v}}_X B + v_{\parallel} \underline{\tilde{v}}_X \underline{e}_\parallel \cdot \underline{\tilde{v}}_1] / B \quad , \quad (10)$$

$$\underline{\tilde{v}}_X \alpha = (\underline{\tilde{v}}_X \underline{e}_2) \cdot \underline{e}_1 + (v_{\parallel}/v_1^2) \underline{\tilde{v}}_X \underline{e}_1 \cdot (\underline{v}_1 \times \underline{e}_1) \quad , \quad (11)$$

$$\delta R = \delta \underline{a} \cdot \underline{\tilde{v}}_V = \delta \underline{a} \cdot \underline{\tilde{v}}_V + \delta \underline{a} \times \underline{e}_\parallel / \Omega \cdot \underline{\tilde{v}}_X \quad , \quad (12)$$

$$\delta \underline{a} = \delta \underline{E} + \underline{v} \times \delta \underline{B} / c \quad , \quad (13)$$

$\underline{E}_0 = -\underline{\tilde{v}}_X \Phi_0$ ,  $\underline{e}_\alpha = -\underline{e}_1 \sin \alpha + \underline{e}_2 \cos \alpha$ , and  $\underline{v}_E = c \underline{E} \times \underline{e}_\parallel / B$ . Here  $L_g$  contains  $\partial \underline{v} / \partial t$  and  $\partial \underline{X} / \partial t$  because the macroscopic quantities are in general time-dependent. Performing spatial averaging on Eq. (6), we obtain

$$L_g F = -(q/m) \langle \delta R \delta F \rangle_{\underline{X}_{10}} \quad , \quad (14)$$

and

$$L_g \delta F = -(q/m) [ \delta R F + \delta R \delta F - \langle \delta R \delta F \rangle_{\underline{X}_{10}} ] \quad . \quad (15)$$

(11) Ordering. To proceed further with Eqs. (14) and (15), we adopt

the following orderings for the microscopic fluctuations

$$\begin{aligned}
 |\partial/\partial t|/|\rho| &\sim |\rho \underline{v}_E \cdot \underline{\nabla}_X| \sim |\delta F/F| \sim |q\delta\phi/T| \\
 &\sim |\delta B/B| \sim \{v_E/v_t\} \sim O(\lambda) \quad , \quad (16)
 \end{aligned}$$

$v_t$  is the characteristic (thermal) velocity, and

$$|\rho \underline{\nabla}_X| \sim O(1) \quad . \quad (17)$$

For macroscopic quantities, however, since they evolve on the transport (including that induced by turbulence) time scale, we take

$$|\partial/\partial t|/|\rho| \sim O(\lambda^3) \quad , \quad (18)$$

in addition to

$$|\rho \underline{\nabla}_X| \sim O(\lambda) \quad . \quad (19)$$

We remark that the ordering adopted here is consistent with experimental observations as well as most of the proposed phenomenological anomalous transport mechanisms.<sup>6,7</sup> Furthermore, since the nonlinear term,  $\delta R \delta F$ , is of  $O(\lambda^2)$ , i.e., comparable to the linear term,  $L_g \delta F$ , our ordering in principle contains strong-turbulence effects. Also, as will be shown later, in Eq. (47) the ordering of the transport time scale, Eq. (18), is consistent with that of the fluctuations, Eq. (16).

(iii) Solution of F. Using the small parameter  $\lambda$ , we have with  $F =$

$$F_0 + F_1 + \dots, 1, 2$$

$$F_0 = F_0(\epsilon, \mu, \tilde{x}_1) ; \quad (20)$$

that is,  $\tilde{e}_\parallel \cdot \tilde{\nabla}_x F_0 = 0$  ,

$$F_1 = (\beta/B)(\partial F_0/\partial \mu) , \quad (21)$$

$$\tilde{\beta} = -(\tilde{v}_1 \cdot \tilde{v}_D + \int^\alpha (d\alpha'/\Omega) \{ \tilde{v}_1 (\tilde{v}_1 \cdot \tilde{\nabla}_x \tilde{e}_\parallel \cdot \tilde{v}_1) - v_{1\perp}^2 \tilde{\nabla}_x \cdot \tilde{e}_\parallel / 2 \} ) , \quad (22)$$

$$\tilde{v}_D = \tilde{v}_d + \tilde{v}_{E0} , \quad (23)$$

$$\tilde{v}_d = \tilde{e}_\parallel \times [ (v_1^2/2) \tilde{\nabla}_x \ln B + v_{1\perp}^2 \tilde{\nabla}_x \tilde{e}_\parallel ] / \Omega , \quad (24)$$

and  $\tilde{F}_1$  is the  $\alpha$ -dependent part of  $F_1$ . To determine the  $\alpha$ -independent part of  $F_1$ , one needs to go to  $O(\lambda^2)$ , where turbulence effects,  $\langle \delta R \delta F \rangle_{\tilde{x}_{10}}$ , would also enter. In fact, following the procedures of neoclassical theory,<sup>13</sup> a formal transport theory including turbulence could be developed; this is, however, beyond the scope of the present work and will be left for future investigations. For the present purpose of obtaining a nonlinear gyrokinetic equation from Eq. (15), knowing  $F$  and  $\tilde{F}_1$  is sufficient.

(iv) Nonlinear gyrokinetic equations. We now concentrate on Eq. (15) for the fluctuating  $\delta F$ . Since only terms up to  $O(\lambda^2)$  are of interest here, the macroscopic background can be treated as frozen, and  $L_g$  becomes accurate to  $O(\lambda)$ ,

$$L_g = L_{gf} = \partial/\partial t + \tilde{v}_1 \tilde{e}_\parallel \cdot \tilde{\nabla}_x + \tilde{v} \cdot (\tilde{\lambda}_{B1} + \tilde{\lambda}_{B2})$$



$$\begin{aligned}
 & + (q/m) \mathbf{E}_0 \left[ (\mathbf{v}_\perp / B) (\partial / \partial \mu) + (\mathbf{e}_\perp / v_\perp) (\partial / \partial \alpha) \right] \\
 & + \mathbf{v}_{E0} \cdot \nabla_{\mathbf{x}} - \Omega \partial / \partial \alpha \quad . \quad (25)
 \end{aligned}$$

Following the linear formalism,<sup>3,4</sup> we let

$$\delta F = (q/m) \delta F_a + \delta G \quad , \quad (26)$$

where

$$\delta F_a = [\delta \Phi \partial / \partial \epsilon + (\delta \Phi - v_\parallel \delta A_\parallel / c) \partial / \partial \mu] F_0 \quad , \quad (27)$$

in order to remove the  $O(\lambda)$  term,  $\delta R F_0$ , in Eq. (15). We note that we have adopted  $\delta \Phi$  and  $\delta A_\parallel$  as field variables. Thus,  $\delta \mathbf{E} = \nabla_{\mathbf{x}} \times \delta \mathbf{A}$  and  $\delta \mathbf{E} = -(\nabla_{\mathbf{x}} \delta \Phi + \partial \delta A_\parallel / c \partial t)$ . Substituting Eqs. (25) and (26) into Eq. (15), we obtain

$$L_{gf} \delta G = -(q/m) (R_\ell + R_{nl}) \quad , \quad (28)$$

where  $R_\ell$  is the linear term<sup>3,4</sup> given by

$$R_\ell = R_{\ell 1} + R_{\ell 2} + R_{\ell 3} + R_{\ell 4} \quad , \quad (29)$$

$$R_{\ell 1} = (\partial \delta L / \partial t) (\partial F_0 / \partial \epsilon) - \nabla_0 \delta L \times \mathbf{e}_\parallel / \Omega \cdot \nabla_{\mathbf{x}} F_0 \quad , \quad (30)$$

$$\delta L \equiv \delta \Phi - \mathbf{v} \cdot \delta \mathbf{A} / c \quad , \quad (31)$$

$$\underline{\nabla}_0 \equiv \partial / \partial \underline{x}_{10} \quad , \quad (32)$$

$$R_{\lambda 2} = (\partial F_0 / \partial \mu) (\partial / \partial t + \underline{v}_{\parallel} \underline{e}_{\parallel} \cdot \underline{\nabla}_{\underline{x}}) \delta L \quad , \quad (33)$$

$$R_{\lambda 3} = \{ \delta \Phi (\underline{v}_{\parallel} \underline{e}_{\parallel} + \underline{x}_D) \cdot \underline{\nabla}_{\underline{x}} - \delta \Phi \underline{v}_{\perp} \cdot \underline{\nabla}_{\underline{x}} \ln B \\ - (\underline{\nabla}_{\underline{v}} \beta) \cdot \underline{\nabla}_{\underline{x}} \delta \Phi + \Omega (\delta \Phi \beta)' (\partial / \partial \epsilon + \partial / \partial \mu) \} (\partial F_0 / \partial \mu) \quad , \quad (34)$$

$$R_{\lambda 4} = -(1/c) \{ \underline{v}_{\parallel} (\underline{v} \cdot \underline{\nabla}_{\underline{x}} \underline{e}_{\parallel}) \cdot \delta \underline{A} + \delta \underline{A}_{\parallel} (\underline{v} \cdot \underline{\nabla}_{\underline{x}}) \cdot \underline{v} \\ + (q/m) \underline{e}_{\perp} \cdot \underline{E}_0 \delta \underline{A}_{\parallel} - \underline{v}_{\parallel} \delta \underline{A}_{\perp} \underline{v} \cdot \underline{\nabla}_{\underline{x}} \ln B \\ - (\underline{\nabla}_{\underline{v}} \tilde{\beta}) \cdot [ \underline{\nabla}_0 (\underline{v} \cdot \delta \underline{A}) - (\underline{v} \cdot \underline{\nabla}_0) \delta \underline{A} ] \\ + \Omega (\beta \underline{v}_{\parallel} \delta \underline{A}_{\parallel})' (\partial / \partial \mu) \} (\partial F_0 / \partial \mu) \quad , \quad (35)$$

$$R_{n\lambda} = \delta R_0 \delta F - \langle \delta R_0 \delta F \rangle_{\underline{x}_{10}} \quad , \quad (36)$$

$$\delta \underline{R}_0 = \delta \underline{a}_0 \cdot \underline{\nabla}_{\underline{v}} + \delta \underline{a}_0 \times \underline{e}_{\parallel} / \Omega \cdot \underline{\nabla}_0 \quad , \quad (37)$$

$$\delta \underline{a}_0 = -\underline{\nabla}_0 \delta \Phi + [ \underline{\nabla}_0 (\underline{v} \cdot \delta \underline{A}) - (\underline{v}_{\perp} \cdot \underline{\nabla}_0) \delta \underline{A} ] / c \quad , \quad (38)$$

and  $(a)'_{\alpha} \equiv \partial a / \partial \alpha$ . Expanding  $\delta G = \delta G_0 + \delta G_1 + \dots$  and noting that the right-hand side of Eq. (28) is of  $O(\lambda^2)$ , we have, for  $O(\lambda)$ ,

$$\partial \delta G_0 / \partial \alpha = 0 \quad . \quad (39)$$

Gyrophase averaging the  $O(\lambda^2)$  equation then yields

$$\langle L_{gf} \rangle_{\alpha} \delta G_o = -(q/m) \langle R_{\parallel} + R_{n\parallel} \rangle_{\alpha} , \quad (40)$$

where

$$\langle L_{gf} \rangle_{\alpha} = \partial/\partial t + v_{\parallel} \frac{e}{\hbar} \cdot \nabla_{\mathbf{x}} + v_{\perp} \cdot \nabla_{\perp} , \quad (41)$$

and  $\langle \dots \rangle_{\alpha} \equiv (1/2\pi) \int_0^{2\pi} d\alpha (\dots)$ . If we further let

$$\delta G_o = -(q/m) \langle \delta L \rangle_{\alpha} \partial F_o / B \partial \mu + \delta H_o , \quad (42)$$

Eq. (40) becomes

$$\langle L_{gf} \rangle_{\alpha} \delta H_o = (-q/m) (S_{\parallel 1} + S_{\parallel 2} + \langle R_{n\parallel} \rangle_{\alpha}) , \quad (43)$$

where

$$S_{\parallel 1} = (\partial \langle \delta L \rangle_{\alpha} / \partial t) (\partial F_o / \partial \mu) - \nabla_{\perp} \langle \delta L \rangle_{\alpha} \times \frac{e}{\hbar} \cdot \nabla_{\mathbf{x}} F_o , \quad (44)$$

$$\begin{aligned} S_{\parallel 2} = & (\partial F_o / B \partial \mu) \{ v_{\parallel} \langle \frac{e}{\hbar} \cdot \nabla_{\mathbf{x}} \delta L \rangle_{\alpha} - v_{\parallel} \frac{e}{\hbar} \cdot \nabla_{\mathbf{x}} \langle \delta L \rangle_{\alpha} \\ & - v_{\perp} \cdot \nabla_{\perp} \langle \delta L \rangle_{\alpha} - \langle v_{\perp} \cdot \nabla_{\perp} \delta \Phi \rangle_{\alpha} \cdot \nabla_{\perp} \ln B \\ & - \langle \nabla_{\perp} \tilde{\beta} \cdot \nabla_{\perp} \delta \Phi \rangle_{\alpha} - \langle \nabla_{\perp} \delta A / c \rangle_{\alpha} - \langle \nabla_{\perp} \delta A / c \rangle_{\alpha} v_{\perp} \frac{e}{\hbar} \cdot \nabla_{\perp} \ln B \\ & + v_{\parallel} \langle \nabla_{\perp} \delta A / c \rangle_{\alpha} \cdot \nabla_{\perp} \ln B + \langle \nabla_{\perp} \tilde{\beta} \rangle_{\alpha} \cdot [ \nabla_{\perp} (\nabla_{\perp} \delta A) ] \end{aligned}$$

$$\begin{aligned}
 & - \langle (\mathbf{v} \cdot \nabla_{\mathbf{x}}) \delta A / c \rangle_{\alpha} - v_{\parallel} \langle (\mathbf{v} \cdot \nabla_{\mathbf{x}}) \mathbf{e}_{\parallel} \cdot \delta A / c \rangle_{\alpha} \\
 & - \langle \delta A_{\parallel} (\mathbf{v} \cdot \nabla_{\mathbf{x}}) \mathbf{e}_{\parallel} \cdot \mathbf{v} / c \rangle_{\alpha} - (q/m) \mathbf{e}_{\parallel} \cdot \mathbf{E} \langle \delta A_{\parallel} / c \rangle_{\alpha} \quad , \quad (45)
 \end{aligned}$$

and, after some algebraic manipulations,

$$\begin{aligned}
 \langle R_{\parallel} \rangle_{\alpha} & = \langle \delta R_{\parallel} \delta F_{\parallel} \rangle_{\alpha} = \langle \delta R_{\parallel} [(q/m) \delta F_{\parallel} + \delta G_{\parallel}] \rangle_{\alpha} \\
 & = -\nabla_{\mathbf{x}} \langle \delta L \rangle_{\alpha} \times \mathbf{e}_{\parallel} / \Omega \cdot \nabla_{\mathbf{x}} \cdot \quad (46)
 \end{aligned}$$

Here we have noted that

$$\langle \langle \delta R_{\parallel} \delta F_{\parallel} \rangle_{\alpha} \rangle_{\mathbf{x}_{\perp 0}} = 0 \quad (47)$$

Equation (47) indicates that effects of turbulence on the  $\alpha$ -independent part of  $F$  is of  $O(\lambda^3)$  and, therefore, consistent with the ordering of the transport time scale, Eq. (18). Combining Eqs. (43) and (46) gives

$$\begin{aligned}
 \bar{L}_{\mathbf{g}} \delta H_{\parallel 0} & = \{ \partial / \partial t + v_{\parallel} \mathbf{e}_{\parallel} \cdot \nabla_{\mathbf{x}} + [\mathbf{v}_{\perp D} + (c/B) \mathbf{e}_{\parallel} \times \nabla_{\mathbf{x}} \langle \delta L \rangle_{\alpha}] \cdot \nabla_{\mathbf{x}} \} \delta H_{\parallel 0} \\
 & = -(q/m) (S_{\parallel 1} + S_{\parallel 2}) \quad (48)
 \end{aligned}$$

Equation (48) is the desired nonlinear gyrokinetic equation. Combining Maxwell's equations and  $\delta F$ , given by Eqs. (26), (42), and (48), the microscopic dynamics is then, in principle, completely determined.

Since the theoretical analyses employ two spatial scales, it is

instructive to proceed further with the following WKB ansatz

$$\begin{aligned} \delta F(\underline{x}, \underline{v}) &= \int_{\underline{k}_\perp} \delta \bar{F}(\underline{x}, \underline{v}; \underline{k}_\perp) \exp[i \int_{x_{10}}^x \underline{k}_\perp \cdot d\underline{x}_\perp] \\ &= \int_{\underline{k}_\perp} \delta \bar{F}(\underline{x}, \underline{v}; \underline{k}_\perp) \exp[i \int_{x_{10}}^x \underline{k}_\perp \cdot d\underline{x}_\perp - iL(\underline{k}_\perp)] \quad , \quad (49) \end{aligned}$$

where  $L(\underline{k}_\perp) = \underline{k}_\perp \cdot \underline{v} \times \underline{e}_\parallel / \Omega$ , and  $\delta \bar{F}$  as well as  $\underline{k}_\perp$  contain slow spatial variations. Equation (48) then reduces to

$$\begin{aligned} [\partial/\partial t + v_\parallel \underline{e}_\parallel \cdot \nabla_{\underline{X}} + i \underline{k}_\perp \cdot \underline{v}_\perp] \delta \bar{H}_0(\underline{k}_\perp) + (c/B) \int_{\underline{k}'_\perp + \underline{k}''_\perp = \underline{k}_\perp} [\underline{e}_\parallel \cdot (\underline{k}'_\perp \times \underline{k}''_\perp)] \\ \langle \delta \bar{L} \rangle_\alpha(\underline{k}'_\perp) \delta \bar{H}_0(\underline{k}''_\perp) = -(q/m) S_{\lambda 1}(\underline{k}_\perp) \quad , \quad (50) \end{aligned}$$

where

$$\langle \delta \bar{L} \rangle_\alpha(\underline{k}_\perp) = [\delta \bar{\Phi} - v_\parallel \delta \bar{A}_\parallel / c] J_0(\gamma) + v_\perp J_1(\gamma) \delta \bar{B}_\parallel / k_\perp c \quad , \quad (51)$$

$$S_{\lambda 1}(\underline{k}_\perp) = [(\partial F_0 / \partial \epsilon)(\partial / \partial t) + i \underline{e}_\parallel \cdot \underline{k}_\perp / \Omega \cdot \nabla_{\underline{X}} F_0] \langle \delta \bar{L} \rangle_\alpha(\underline{k}_\perp) \quad , \quad (52)$$

$\gamma = k_\perp v_\perp / \Omega$ , and  $S_{\lambda 2}(\underline{k}_\perp)$  is of higher order and ignorable here. Equation (50) and, hence, Eq. (48) show that the nonlinearity arises from a gyrophase-averaged effective  $\delta \underline{E}_{\text{eff}} \times \underline{B} \cdot \nabla_{\underline{X}}$  coupling. Here,

$$\delta \underline{E}_{\text{eff}} = -\nabla_0 (\delta \Phi - v_\parallel \delta \bar{A} / c) \quad . \quad (53)$$

It is interesting to note that the nonlinear polarization drift is contained, not very obviously, within the finite Larmor radius corrections

in  $J_0$  and  $J_1$ . In fact, the electrostatic nonlinear drift-wave equation first obtained by Hasegawa and Mima<sup>14</sup> can be readily derived from Eq. (50) in the appropriate limits, i.e., adiabatic electrons and cold fluid ions (Sec. III). Equation (48) [or Eq. (50)], of course, is much more general and also is not easy to solve. In the next two sections, we consider more specific applications of the general results obtained so far.

### III. AXISYMMETRIC TOKAMAKS

(1) General formulation. Here we consider the specific case of axisymmetric tokamaks and, using the ballooning-mode representation,<sup>8-10,15</sup> further explore the properties of the nonlinear gyrokinetic equation, Eq. (48). Thus, employing the  $\psi$  (poloidal flux),  $\xi$  (toroidal angle), and  $\chi$  (poloidal angle-like) coordinates, we have

$$\vec{B} = \nabla_{\vec{x}} \psi \times \nabla_{\vec{x}} \phi + I(\psi, \chi) \nabla_{\vec{x}} \xi, \quad (54)$$

$$F_0 = F_0(\psi, \mu, \varepsilon), \quad (55)$$

$$\delta \mathcal{H}_0 = \int_n \int_m \delta h_{n,m}(\psi, \underline{v}) \exp[i(n\xi - m\chi + n \int^\psi k d\psi)] , \quad (56)$$

$$\delta h_{n,m}(\psi, \underline{v}) = \int_{-\infty}^{\infty} d\hat{\theta}_n \delta \bar{h}_n(\psi, \hat{\theta}_n, \underline{v}) g_{n,m}(\hat{\theta}_n) . \quad (57)$$

$$g_{n,m}(\hat{\theta}_n) = \exp[i(m\hat{\theta}_n - n \int^{\hat{\theta}_n} v d\hat{\theta})] , \quad (58)$$

$v = IJ/R^2$ , and  $J = (\nabla_{\vec{x}} \psi \times \nabla_{\vec{x}} \xi \cdot \nabla_{\vec{x}} \chi)^{-1}$  is the Jacobian. Equation (48) or, equivalently, Eq. (43) then becomes

$$\int_{-\infty}^{\infty} d\hat{\theta}_n g_{n,m}(\hat{\theta}_n) \{ [\partial/\partial t + (v_{\parallel}/JB)(\partial/\partial \hat{\theta}_n) + ik_{\perp}^i v_{\perp D}^i] \delta \bar{h}_n + (q/m) \bar{S}_{\perp n} \} = -(q/m) \langle R_{n\ell} \rangle_{\alpha} (m, n) , \quad (59)$$

where

$$k_{\perp}^i = (nB/RB_{\chi}) (\underline{e}_{\parallel} \times \underline{e}_{\phi}) - nRB_{\chi} \int_0^{\hat{\theta}_n} (\partial v/\partial \psi) d\hat{\theta} - k_{\parallel} , \quad (60)$$

$$\bar{S}_{\perp n} = [(\partial F_0/\partial \epsilon)(\partial/\partial t) - i(nB/c)(\partial F_0/\partial \psi)] \delta \bar{L}_n , \quad (61)$$

$$\delta \bar{L}_n = J_0(\gamma)(\delta \bar{\Phi} - v_{\parallel} \delta \bar{A}_{\parallel}/c)_n + J_1(\gamma) v_{\perp} \delta \bar{b}_{\perp n} / k_{\perp} c , \quad (62)$$

$$\langle R_{n\ell} \rangle_{\alpha} (m, n) = -\frac{1}{2} \sum_{n'=n} \sum_{m'=m} \sum_{n''=n-k} \int_{-\infty}^{\infty} d\hat{\theta}_{n'} \int_{-\infty}^{\infty} d\hat{\theta}_{n''} g_{n',m'} g_{n'',m''} C_{n',n''} (\delta \bar{L}_{n'} \delta \bar{h}_{n''} - \delta \bar{L}_{n''} \delta \bar{h}_{n'}) , \quad (63)$$

$$C_{n',n''} = (\underline{e}_{\parallel}/\Omega) \cdot (k_{\perp}^i \times k_{\perp}^{i'}) = (Bn'/\Omega) [n'' \int_{\hat{\theta}_{n'}}^{\hat{\theta}_{n''}} (\partial v/\partial \psi) d\hat{\theta} + n(k' - k)] , \quad (64)$$

and, again,  $\gamma = k_{\perp} v_{\perp}/\Omega$ . After some manipulations, Eq. (63) can be shown to be

$$\langle R_{n\ell} \rangle_{\alpha} (m, n) = \int_{-\infty}^{\infty} d\hat{\theta}_n g_{n,m}(\hat{\theta}_n) (\bar{R}_{n\ell})_n , \quad (65)$$

where

$$\begin{aligned}
 (\bar{R}_{nl})_n &= -\pi \sum_{n'+n''} \sum_{n'k'+n''k''} \sum_p \exp(-in''2\pi p Q) \\
 \tilde{C}_{n',n''} &= \int_{-\infty}^{\infty} d\hat{\theta}_n \delta(\hat{\theta}_n - \hat{\theta}_{n'}) \int_{-\infty}^{\infty} d\hat{\theta}_{n''} \delta(\hat{\theta}_{n''} - \hat{\theta}_n - 2\pi p) \\
 & (\delta\bar{L}_{n'} \delta\bar{h}_{n''} - \delta\bar{L}_{n''} \delta\bar{h}_{n'}) \quad , \quad (66)
 \end{aligned}$$

$$Q \equiv (1/2\pi) \int_0^{2\pi} v d\hat{\theta} \quad , \quad (67)$$

and

$$\tilde{C}_{n',n''} = (Bn'/Q) [n''2\pi p (\partial Q / \partial \phi) + n(k' - k)] \quad . \quad (68)$$

Combining Eqs. (59) and (65), we finally derive the nonlinear gyrokinetic equation for axisymmetric tokamaks

$$[\partial / \partial t + (v_{||} / JB) (\partial / \partial \hat{\theta}_n)] + i k_{\perp} \cdot v_{\perp} [\delta\bar{h}_n] = -(q/m) [\bar{S}_{ln} + (\bar{R}_{nl})_n] \quad . \quad (69)$$

We now make some qualitative remarks. For simplicity, we shall ignore global amplitude modulations by letting  $k = k' = k'' = 0$  and, hence,  $\tilde{C}_{n',n''} \propto p$ . For flute-like modes, which are highly localized about the mode-rational surfaces,  $\exp(-in''2\pi p Q) \approx 1$  and  $\delta\bar{h}$  as well as  $\delta\bar{L}$  are weakly dependent on  $\hat{\theta}$ . Therefore, we have, roughly,  $(\bar{R}_{nl})_n \propto \sum_p p = 0$ ; i.e., nonlinear coupling is much reduced, which may be expected because there is little overlap between the modes in the  $\phi$  (radial) coordinate. On the other hand, for modes which are strongly ballooning, and hence,  $(\delta\bar{\phi}, \delta\bar{L})(\hat{\theta} + p2\pi) = 0$  for  $p \neq 0$ ; the nonlinear coupling, again, is small.



This is because, in this limit, radial structures are rather broad, and hence the  $\delta \vec{E} \times \vec{B} \cdot \nabla_{\vec{x}}$  coupling is rendered ineffective. Thus qualitatively speaking, Eq. (66) appears to indicate that the most effective nonlinear coupling occurs among modes which are moderately ballooning.

(11) Electrostatic drift waves with adiabatic electrons and cold fluid ions. We assume, for simplicity,  $F_0$  to be a local Maxwellian; i.e.,

$$F_0 = F_M(\psi, \epsilon) \quad , \quad (70)$$

and neglect any equilibrium electric potential  $\phi_0$ . Since electrons are adiabatic, their nonlinear contributions are negligible, and the quasi-neutrality condition becomes

$$(1 + \tau)\delta\tilde{n}_n = 2\pi \int B d\mu \int dv_{\parallel} \delta\tilde{n}_{in} J_0(\gamma_1) \equiv \delta\tilde{n}_{in} \quad . \quad (71)$$

Here,  $\delta\phi_n \equiv e\delta\tilde{\phi}_n/T_e$ ,  $\tau = T_e/T_i \gg 1$ , and  $J_0(\gamma_1) = 1 - \gamma_1^2/4$ . Meanwhile, neglecting the ion-sound term, we multiply the ion Eq. (69) by  $J_0(\gamma_1)$  and carry out the velocity integration to obtain

$$\begin{aligned} (\partial/\partial t + ik_{\perp} \cdot \vec{v}_{d1})\delta\tilde{n}_{in} &= \tau(1 - b_1)(\partial/\partial t - i\omega_{*in})\delta\phi_n \\ &- (e/m_i)2\pi \int B d\mu \int dv_{\parallel} J_0(\gamma_1) \langle \vec{R}_{n\ell} \rangle_{in} \quad , \end{aligned} \quad (72)$$

where, for  $j = e, i$ ,

$$\vec{v}_{dj} = \hat{e}_{\theta} \times (\nu_j^2/2)(\nabla_{\vec{x}} \ln B + \hat{e}_{\theta} \cdot \nabla \hat{e}_{\theta})/\Omega_j \quad , \quad (73)$$

$$\omega_{*j} = (v_j^2 n B / 2 \Omega_j) |\partial \ln N_{oj} / \partial \psi|, \quad (74)$$

and  $b_1 = k_1^2 \rho_1^2 / 2$ . In deriving Eq. (72), we have noted that in the zeroth-order approximation,  $\delta \bar{n}_{1n} = \delta \bar{n}_{1n} F_{M1}$ . Expanding the  $J_0$ 's in  $(\bar{R}_{n\ell})_{1n}$  and using the quasi-neutrality condition, Eq. (71), it is then straightforward to derive

$$\begin{aligned} & [(1 + k_1^2 \rho_s^2)(\partial/\partial t) - i k_1 \cdot \bar{v}_{de} + i \omega_{*e}] \delta \phi_n \\ &= \pi C_s^2 \sum_{n'=-n}^{n'+n} \sum_{n''=n-k}^{n'+n''k} \int_p \exp(-in'' 2\pi p Q) \hat{C}_{n', n''} \int_{-\infty}^{\infty} d\hat{\theta}_{n''} \delta(\hat{\theta}_n - \hat{\theta}_{n'}) \\ & \int_{-\infty}^{\infty} d\hat{\theta}_{n''} \delta(\hat{\theta}_{n''} - \hat{\theta}_n - 2\pi p) \rho_s^2 [(k'')^2 - (k_1')^2] \delta \phi_{n''} \delta \phi_{n''}, \quad (75) \end{aligned}$$

where  $C_s^2 = \tau v_1^2 / 2$  and  $\rho_s = C_s / \kappa_1$ . Equation (75) may be regarded as the tokamak version of the nonlinear drift-wave equation of Hasegawa and Mima.<sup>12</sup> It contains the poloidal mode-coupling effects due to the  $\nabla B$  and curvature drifts and employs the ballooning-mode representation. The assumption of adiabatic electrons and cold fluid ions thus results in a single nonlinear equation instead of coupled ones, as in more general cases. To make Eq. (75) more tractable, we further assume concentric, circular magnetic surfaces and ignore the global wavenumber ( $k$ ). Equation (75) then reduces to

$$\begin{aligned} & [(1 + \hat{k}_1^2 \rho_s^2)(\partial/\partial t) - i \bar{\omega}_{de} T_c(\hat{\theta}) + i \omega_{*e}] \delta \phi(n, \hat{\theta}) \\ &= \pi \Omega_i \hat{S} \sum_{n'=-n}^{n'+n} \int_p \exp(-in'' 2\pi p Q) \rho_s^4 (k_n, k_{n''} 2\pi p) \end{aligned}$$

$$i(\hat{k}_\perp'')^2 - (\hat{k}_\perp')^2 \gamma \delta\phi(n', \hat{\theta}) \delta\phi(n'', \hat{\theta} + 2\pi p) , \quad (76)$$

where  $\omega_{*e} = k_n \rho_s C_s / n$ ,  $r_n^{-1} = |d \ln N / dr|$ ,  $k_n = n0/r$ ,  $\bar{\omega}_{de} = 2r_n \omega_{*e} / R$ ,  
 $\hat{S} = r(\partial Q / \partial r) / Q$ ,

$$T_c = \cos \hat{\theta} + \hat{S} \hat{\theta} \sin \hat{\theta} , \quad (77)$$

$$\hat{k}_\perp'^2 = k_n^2 (1 + \hat{S}^2 \hat{\theta}^2) , \quad (78)$$

$$(\hat{k}_\perp'')^2 = k_n^2 (1 + \hat{S}^2 \hat{\theta}^2) , \quad (79)$$

and

$$(\hat{k}_\perp'')^2 = k_n^2 [1 + \hat{S}^2 (\hat{\theta} + 2\pi p)^2] . \quad (80)$$

We note that Eq. (76) is a two-dimensional  $(n, \hat{\theta})$  nonlinear equation and may be viewed as the simplest model equation for electrostatic drift waves in axisymmetric tokamaks. Detailed studies of Eq. (76) will be reported in a future publication.

#### IV. NONLINEAR ION LANDAU DAMPING OF KINETIC ALFVEN WAVES

As noted in Sec. I, the results obtained in Sec. II are rather general and, therefore, are also useful for describing nonlinear processes associated with low-frequency wave heating. To illustrate this potential application, we consider here nonlinear ion Landau damping of the mode-converted kinetic (shear) Alfvén waves.<sup>11</sup> The major difference between our

work and Ref. 16 is that we include the diamagnetic drift effects. Another minor difference is that no small ion-Larmor-radius expansion is taken here.

For the sake of simplicity, we adopt the WKB description, Eqs. (49) and (50), and ignore the geometrical effects. Furthermore,  $F_0$  is taken to be Maxwellian and for the present purpose we can assume a weak-turbulence subsidiary ordering as well as ignore the compressional Alfvén perturbations  $\delta A_{\perp} = 0$  (i.e.,  $\beta \equiv (\text{plasma pressure/magnetic pressure}) \ll 1$ ). Thus, the relevant equations are

$$\delta \bar{F}(\underline{k}) = -(q/T) \delta \bar{\Phi}(\underline{k}) F_0 + \delta \bar{H}_0(\underline{k}) \exp(i \underline{k} \cdot \underline{v} \times \underline{e}_{\parallel} / \Omega) , \quad (81)$$

and

$$\begin{aligned} (\omega_{\underline{k}} - k_{\parallel} v_{\parallel}) \delta H_0(\underline{k}) &= (q/T) (\omega + \omega_{\star})_{\underline{k}} F_0 \delta \bar{L}(\underline{k}) J_0(\gamma) \\ &\quad - i \langle c/B \rangle \int_{\substack{\underline{k}' + \underline{k}'' \\ = \underline{k}}} [\delta \bar{L}(\underline{k}') J_0(\gamma') \delta \bar{H}_0(\underline{k}'') (\underline{k} \times \underline{k}') \cdot \underline{e}_{\parallel}] , \end{aligned} \quad (82)$$

where  $\delta \bar{L}(\underline{k}) = \delta \bar{\Phi}(\underline{k}) - v_{\parallel} \delta \bar{A}_{\parallel}(\underline{k})/c$ ,  $\omega_{\star \underline{k}} = k_y c T / e B L_n$ ,  $L_n^{-1} = |d \ln N_0 / dx|$ ,  $x$  is the nonuniformity direction, and temperature gradients are ignored. Expanding  $\delta \bar{H}_0 = \delta H_0^{(1)} + \delta H_0^{(2)} + \delta H_0^{(3)} + \dots$ , we have, in the linear order,

$$\delta H_0^{(1)}(\underline{k}) = \frac{q}{T} \frac{(\omega + \omega_{\star})_{\underline{k}}}{\omega_{\underline{k}} - k_{\parallel} v_{\parallel}} F_0 \delta \bar{L}(\underline{k}) J_0(\gamma) . \quad (83)$$

To calculate  $\delta H_0^{(2),(3)}$ , we let  $(\omega_{\underline{k}}, \underline{k})$  and  $(\omega_{\underline{k}'}, \underline{k}')$  be the normal modes, while  $(\omega_{\underline{q}} = \omega_{\underline{k}} - \omega_{\underline{k}'}, \underline{q} = \underline{k} - \underline{k}')$  is the virtual or quasi mode. Also, we have in mind

$$|\omega_q| \sim |q_{\parallel} v_i| \sim |k_{\parallel}, k'_{\parallel}| v_i \ll |\omega_{\tilde{k}q}| \sim |\omega_{\tilde{k}, \tilde{k}'}| \ll |k_{\parallel}, k'_{\parallel}| v_e \quad (84)$$

Let us now calculate the  $q$ -mode response. For the ions we find, noting Eq. (84), that

$$\delta H_{oi}^{(2)}(q) = [\omega_{*qe} / (\omega_q - q_{\parallel} v_{\parallel})] \delta \tilde{P}_q \delta \tilde{\Phi}(k) \delta \tilde{\Phi}(k') J_o(\gamma) J_o(\gamma') F_{oi} \quad (85)$$

where

$$\delta \tilde{P}_q = i(\Omega_i / \omega_{k'}) \rho_g^2 [(k \times k') \cdot e_{\parallel}] \quad (86)$$

and  $\delta \tilde{\Phi} \equiv e \delta \tilde{\Phi} / T_e$ . In deriving Eq. (85), we have observed that  $|\delta \tilde{\Phi}| \sim |\omega \delta \tilde{A}_{\parallel} / ck_{\parallel}|$  for the  $\tilde{k}$  and  $\tilde{k}'$  modes. As to electron nonlinearity, it is  $O(|\omega / \omega_{*e}|_q)$  smaller and, hence, negligible. Furthermore, for  $\beta \ll 1$ , we have  $|\omega_q| \ll |q_{\parallel} v_A|$ , and the  $q$ -mode response is predominately electrostatic. Here  $v_A$  is the Alfvén speed. The quasi-neutrality condition then yields

$$\delta \tilde{\Phi}_q^{(1)} = -(F_1 / \Gamma_{oq}) \delta \tilde{P}_q \delta \tilde{\Phi}(k) \delta \tilde{\Phi}(k') \quad (87)$$

where

$$F_1(\tilde{k}, \tilde{k}') = 2\pi \int_0^{\infty} v_{\perp} dv_{\perp} J_o(\gamma) J_o(\gamma') J_o(\gamma_q) F_{Mz}(v_{\perp}) \equiv \langle J_o(\gamma) J_o(\gamma') J_o(\gamma_q) \rangle_{\perp} \quad (88)$$

$$\Gamma_{oq} = I_o(b_q) \exp(-b_q) \quad , \quad (89)$$

$\gamma_q = q_{\perp} v_{\perp} / \Omega_i$  and  $b_q = q_{\perp}^2 \rho_{i1}^2 / 2$ . Equations (83) and (87) then give  $\delta n_{oi}^{(1)}(q)$ .

For the  $\underline{k}$ -mode, the ion nonlinear response can be shown to be

$$\begin{aligned} \delta n_{oi}^{(3)}(\underline{k}) = & |\delta \tilde{\Phi}_q|^2 |\delta \tilde{\Phi}(\underline{k}')|^2 \delta \tilde{\Phi}(\underline{k}) F_{oi} \cdot \left( \frac{\omega_{*e}}{\omega - q_{\parallel} v_{\parallel}} \right)_q \\ & \cdot \left[ J_o^2(\gamma') J_o(\gamma) - F_1 J_o(\gamma_q) J_o(\gamma') / \Gamma_{oq} \right] \quad . \quad (90) \end{aligned}$$

From Eq. (90) we can readily calculate  $\delta n_i^{(3)}(\underline{k})$ . Meanwhile, the electron nonlinearity, again is found to be  $O(\omega / \omega_{*e})_q$  smaller. Substituting  $\delta n_i^{(3)}(\underline{k})$  and the linear response into the quasi-neutrality condition and parallel Ampere's law, it is straightforward to derive the following nonlinear dispersion relation

$$\begin{aligned} D_{\perp}(\underline{k}) = & - \frac{\omega_{*eq}}{|q_{\parallel}| v_{\parallel}} Z_{1q} \left[ 1 - \left( 1 - \frac{\omega_{*e}}{\omega} \right)_k \left( \frac{\omega}{k_{\parallel} v_A k_{\perp} \rho_B} \right)_k^2 \right] |\delta \tilde{\Phi}_q \delta \tilde{\Phi}(\underline{k}')|^2 \\ & \cdot G(\underline{k}, \underline{k}') \quad , \quad (91) \end{aligned}$$

where  $D_{\perp}(\underline{k})$  is the linear dispersion relation given by

$$\begin{aligned} D_{\perp}(\underline{k}) = & (\tau + \omega_{*e}/\omega)_k (1 - r_{ok}) + (1 - \omega_{*e}/\omega)_k \\ & \cdot \left[ 1 - \frac{\omega_{*e}^2}{\omega^2} (\tau + \omega_{*e}/\omega)_k (1 - r_{ok}) / (k_{\parallel} v_A k_{\perp} \rho_B)^2 \right] \quad , \quad (92) \end{aligned}$$

$$G(\underline{k}, \underline{k}') = \langle J_o^2(\gamma) J_o^2(\gamma') \rangle_{\perp} - F_1^2 / \Gamma_{oq} \quad , \quad (93)$$

$Z_{1q} \equiv Z(\omega / |q_{\parallel}| v_{\parallel})$  with  $Z$  being the standard plasma dispersion function.

In the low- $\beta$  limit of interest here,  $|\omega_{*e}/k_1 v_A|_{\underline{k}} < 1$ , and assuming  $\omega_{\underline{k}} = \omega_{r\underline{k}} + i\omega_{i\underline{k}}$  with  $|\omega_i/\omega_r| \ll 1$ , Eq. (92) gives the known result that there exist two branches of waves; the drift waves with

$$\omega_{r\underline{k},d} = \omega_{*e} \Gamma_{o\underline{k}} / [1 + \tau(1 - \Gamma_{o\underline{k}})] \quad (94)$$

and the kinetic shear Alfvén waves with

$$\omega_{r\underline{k},A}^2 = (k_1 v_A)^2 [1 + \tau(1 - \Gamma_{o\underline{k}})] k_1^2 c_s^2 / \tau(1 - \Gamma_{o\underline{k}}) \quad (95)$$

We note that nonlinear ion Landau damping of the drift waves, as described by Eq. (91), has previously been investigated.<sup>17</sup> Concentrating on the kinetic shear-Alfvén waves, we find

$$\left(\frac{\omega_i}{\omega_r}\right)_{\underline{k},A} = \frac{(\omega_{*k'} - \omega_{*k})_e}{2\tau(1 - \Gamma_{o\underline{k}})\{1 + \tau(1 - \Gamma_{o\underline{k}})\}} G(\underline{k}, \underline{k}') \frac{\text{Im } Z_{1q}}{|q_1 v_1|} |\delta\tilde{\omega}(\underline{k}') \delta\tilde{F}_q|^2 \quad (96)$$

Comparing the  $\omega_i$  given by Eq. (96) to that of Ref. 16 (which, we recall, does not include the  $\omega_*$  effects), we find the present parametric-decay growth rate is larger by a factor of  $O(|\omega_{*1q}|/|q_1 v_1|) > 1$ . Furthermore, noting that  $\text{Im } Z_{1q} = |q_1| v_1 \pi \delta(\omega_{k'} - \omega_k)$ ,  $G(\underline{k}, \underline{k}') \geq 0$ , and  $\omega_i = (\omega_{*k'} - \omega_{*k}) = (k'_y - k_y)$ ; hence, the daughter wave  $(\omega_k, \underline{k})$  has a smaller  $k_y$  but the same  $\omega$  as the pump wave  $(\omega_{k'}, \underline{k}')$ . This is qualitatively different from the results of Ref. 16, where  $\omega_i = (\omega_{k'} - \omega_k)$ . In fact, as might be expected, one can show that the results of Ref. 16 are valid for  $|\omega_{*1q}| < |q_1 v_1|$ , i.e., in the opposite limit.

Finally, we briefly discuss the implication of the present results on

the shear Alfvén wave heating scheme. Since the pump wave (i.e., the mode-converted kinetic shear Alfvén wave) has  $k'_x \rho_i \sim 0(1)$  and  $k'_y \rho_i \sim 0(\rho_i/a) \ll 1$  with  $a$  being the tokamak minor radius, we have  $\omega_i \approx |\underline{k} \times \underline{k}' \cdot \underline{e}_\parallel|^2 = (k'_x k_y)^2$ , and hence nonlinear coupling becomes appreciable for  $k_y \rho_i \sim 0(1) \gg k'_y \rho_i$ ; that is,  $\omega_i < \omega$ . Thus, our results suggest that parametric decay through nonlinear ion Landau damping will be, in general, unlikely to occur.

#### V. CONCLUSIONS AND DISCUSSIONS

In this work, a systematic formalism for the nonlinear interactions of microscopic low-frequency electromagnetic waves has been developed. This formalism extends the linear gyrokinetic formalism of Refs. 1 and 2 into the nonlinear regime. Corresponding nonlinear gyrokinetic equations, valid for general magnetic field configurations as well as the strong-turbulence regime, are derived. Effects due to fully electromagnetic perturbations, finite Larmor radii, plasma inhomogeneities, magnetic drifts, and magnetic trapping are retained. The results are thus rather general and should have wide application. Note that it is straightforward to extend the treatment to include collisional effects by retaining the Fokker-Planck collision operator in Eqs. (7) and (15). As a specific example of possible applications, we consider axisymmetric tokamaks and explore in more detail the properties of the nonlinear gyrokinetic equations via the ballooning-mode representation. Furthermore, a single nonlinear equation is derived for electrostatic drift waves in the limit of adiabatic electrons and cold fluid ions which retains the crucial features of toroidal geometry and nonlinear coupling and, therefore, may be exploited as a useful model equation. On the other hand, we have also applied the results to the



shear Alfvén wave heating scheme and considered the parametric decays of the mode-converted kinetic Alfvén waves via nonlinear ion Landau damping (ion-induced scattering). Here it is found that the diamagnetic drift effects not only enhance the parametric growth rate but also modify the decay process qualitatively. That is, the daughter waves tend to have smaller perpendicular (to  $\mathbf{B}$  and the density gradient) wavenumbers (i.e., smaller poloidal mode numbers for tokamak plasmas) instead of frequencies as suggested by uniform-plasma calculations. This property, therefore, suggests that the mode-converted kinetic Alfvén waves with small poloidal mode numbers will not, in general, parametrically decay via nonlinear ion Landau damping. Other possible channels, such as resonant decay to drift waves, of course, are not ruled out and need to be investigated.

Let us comment on some other possible applications of the general results obtained here. One possible application is the following: by taking the limit of adiabatic electrons and cold fluid ions, a sufficiently simple nonlinear equation may also be derived for the kinetic shear-Alfvén waves and hence could serve as a model equation for studying the nonlinear evolution and associated transport of kinetic ballooning-mode instabilities.<sup>18</sup> Another interesting application is to simulate plasmas employing the nonlinear gyrokinetic equation, Eq. (48) or, more transparently, Eq. (50). This application will further extend the gyrokinetic simulation scheme initially proposed by Lee<sup>19</sup> for electrostatic waves in simple (slab) geometries to fully electromagnetic perturbations in general plasma equilibria. The details of these applications, however, remain to be worked out.

ACKNOWLEDGMENTS

This work was supported by United States Department of Energy Contract  
No. DE-AC02-76-CR03073.

REFERENCES

1. P. H. Rutherford and E. A. Frieman, *Phys. Fluids* 11, 569 (1968).
2. J. B. Taylor and R. J. Hastie, *Plasma Phys.* 10, 479 (1968).
3. T. M. Antonsen and B. Lane, *Phys. Fluids* 23, 1205 (1980).
4. P. J. Catto, W. M. Tang, and D. E. Baldwin, Science Application, Inc. Report SAI-254-80-LJ PRI-14.
5. E. A. Frieman and L. Chen, *Bull. Am. Phys. Soc.* 24, 1009 (1979). A similar approach has been adopted by P. H. Diamond and M. N. Rosenbluth for a less general case in the University of Texas Report IFSR-24.
6. E. Mazzucato, *Phys. Rev. Lett.* 36, 702 (1976).
7. G. M. Surko and R. E. Slusher, *Phys. Rev. Lett.* 37, 1747 (1976).
8. J. W. Connor, R. J. Hastie, and J. B. Taylor, *Proc. R. Soc. Lond.* A365, 1 (1979).
9. Y. C. Lee and J. W. van Dam, in *Proceedings of the Finite-Beta Theory Workshop*, Varenna, Italy, edited by B. Coppi and W. Sadowski (National Technical Information Service, Springfield, VA., 1979), p. 93.
10. A. H. Glasser, in *Proceedings of the Finite-Beta Theory Workshop*, Varenna, Italy, edited by B. Coppi and W. Sadaowski (National Technical Information Service, Springfield, VA., 1979), p. 55. Also, R. L. Dewar, M. S. Chance, A. H. Glasser, J. M. Greene, and E. A. Frieman, Princeton Plasma Physics Laboratory Report PPPL-1587 (1979).
11. A. Hasegawa and L. Chen, *Phys. Rev. Lett.* 35, 370 (1975).
12. P. J. Catto, *Plasma Phys.* 20, 719 (1978).
13. F. L. Hinton and R. D. Hazeltine, *Rev. Mod. Phys.* 48, 239 (1976).
14. A. Hasegawa and K. Mima, *Phys. Rev. Lett.* 39, 205 (1977). Also, A. Hasegawa, C. G. MacLennan, and Y. Kodama, *Phys. Fluids* 22, 2122 (1979).
15. J. W. Connor, R. J. Hastie, and J. B. Taylor, *Plasma Phys.* 22, 757

- (1980). E. A. Frieman, G. W. Rewoldt, W. M. Tang, and A. H. Glasser, Phys. Fluids 23, 1750 (1980). F. Pegoraro and J. J. Schep, Phys. Fluids 24, 478 (1981).
16. A. Hasegawa and L. Chen, Phys. Rev. Lett. 36, 1362 (1976); also Phys. Fluids 19, 1924 (1976).
17. R. Z. Sagdeev and A. A. Galeev, Nonlinear Plasma Theory (Benjamin, New York, 1969), p. 103.
18. W. M. Tang, J. W. Connor, and R. J. Hastie, Nucl. Fusion 20, 1439 (1980).
19. W. W. Lee, Bull. Am. Phys. Soc. 25, 966 (1980).