

# On Implicit Taylor Series Methods for Stiff ODEs

G. Kirlinger  
 Institut für Angewandte  
 und Numerische Mathematik  
 Technische Universität Wien  
 Wiedner Hauptstrasse 8-10  
 A-1040 Wien, AUSTRIA

G. F. Corliss\*  
 Mathematics and Computer Science Division  
 Argonne National Laboratory  
 9700 S. Cass Avenue  
 Argonne, IL 60439-4801 USA

DEC 16 1991

## Abstract

Several versions of implicit Taylor series methods (ITSM) are presented and evaluated. Criteria for the approximate solution of ODEs via ITSM are given. Some ideas, motivations, and remarks on the inclusion of the solution of stiff ODEs are outlined.

## 1 Introduction

One approach for the validated solution of initial value problems for ODEs is built upon a Taylor series method. Moore [20, 21], Rall [22], Adams [1], Lohner [17], and Eijgenraam [14] were the first to use Taylor series methods to enclose the solutions of ODEs. One of the main advantages of using a Taylor series method is the simple representation of the local discretization error. For a Taylor series method, the discretization error is the remainder term of the series, which can easily be bounded by using automatic differentiation to generate as many derivatives as required. Further, the order of the Taylor series can also easily be adjusted to the needed accuracy.

This paper emphasizes approximation methods based on Taylor series, but we also give some initial ideas, motivations, and remarks on the inclusion of the solution of stiff ODEs (Section 5). We compare several numerical methods based on Taylor series to evaluate their suitability for use as methods for the verified inclusion of stiff differential equations.

This is a report on work in progress. Some numerical experiments have been done, but much more experimental and theoretical work remains to be done. Another manuscript [7] is in preparation describing in detail the relationship of the implicit Taylor series methods described here to implicit methods based on Padé approximations.

\*Supported in part by the National Science Foundation grant No. CCR-8802429, by the Applied Mathematical Sciences subprogram of the Office of Energy Research, U.S. Department of Energy, under Contract W-31-109-Eng-38, and through NSF Cooperative Agreement No. CCR-8809615. On leave from Marquette University, Milwaukee, WI 53233 USA.

**MASTER**

1

The submitted manuscript has been authored by a contractor of the U. S. Government under contract No. W-31-109-ENG-38. Accordingly, the U. S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or allow others to do so, for U. S. Government purposes.

REPRODUCTION OF THIS DOCUMENT IS UNLIMITED

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**DISCLAIMER**

## 2 Stiffness

In this paper, we present initial steps towards algorithms for solving stiff systems of ODEs based on Taylor series. We begin with a brief survey of theory and point algorithms for stiff problems. The modeling of evolution, or growth, processes in many applied sciences often leads to stiff ordinary differential equations. The initial value problem

$$y' = \lambda y, \quad y(0) = y_0 \quad \text{with } \lambda \ll 0 \quad (2.1)$$

is a simple example of a stiff problem. It shows the following characteristic behavior:

- (i) The Lipschitz constant  $L = |\lambda|$  of the right-hand side is very large.
- (ii) For  $y_0 = 0$ , the solution  $y(t) \equiv 0$  is smooth.
- (iii) For  $y_0 \neq 0$ , the solution  $y(t) = e^{\lambda t} y_0$  is rapidly decaying and very unsmooth in the sense that derivatives are large during the initial “transient phase.” Away from  $t = 0$ , the solution becomes smooth very quickly and tends towards the smooth solution (“smooth phase”).

In most practical situations, rapidly decaying components (corresponding to  $\lambda \ll 0$ ) occur together with smooth (nonstiff) components. Such a system quickly tends towards an “equilibrium,” that is, to a smooth solution. Rapid variations occur only while the state of equilibrium has not yet been reached or when the system is switched from one state to another (e.g., by nonlinear effects).

Although it is difficult to give a mathematically rigorous definition of stiffness, we call a system

$$\begin{aligned} y' &= f(t, y) \\ y(0) &= y_0 \end{aligned} \quad (2.2)$$

*stiff* if its Jacobian  $f_y$  (in a neighborhood of the solution) has eigenvalues  $\lambda_i$  with  $Re(\lambda_i) \ll 0$ , in addition to eigenvalues of moderate size.

Stiff systems are considered difficult because explicit numerical methods designed for nonstiff problems are forced to use very small steps. If “normal” steps are used, then perturbations in the computed solution are amplified by the influence of the Lipschitz constant  $L \gg 0$ . In order to retain the stability of the true solution in the computed solution, the step must be very small. It is not possible to use a step size that is adjusted to the smoothness of the solution sought. That is why authors as early as 1928 [8] or 1947 [9] were led to consider implicit methods in which the approximate solution  $y_{i+1}$  at  $t = t_{i+1}$  is given by the solution to some nonlinear system. Many implicit methods allow step sizes appropriate to the smoothness of the solution.

Historically, the first theoretical concept especially suited for the assessment of numerical methods for stiff problems was A-stability [11]. A-stability means that computed

numerical approximations to decreasing components are also decreasing. This analysis is based on the model linear constant coefficient problem  $y' = Ay$ . The more modern concept of *B-stability* [15] is the basis for a general convergence theory for nonlinear stiff problems. Nevertheless, we use the linear concept A-stability as a first criterion for the assessment of implicit procedures based on Taylor series methods.

The general algorithm for implicit methods for stiff systems is as follows:

```

Initialize
Loop for each integration step
    Guess step size
    Solve some nonlinear system for  $y_{i+1}$ 
    Estimate error
    Accept or reject step

```

Well-known examples of methods of this class that are suitable for the integration of stiff systems are backward differentiation formulas (BDFs) [10] and implicit Runge-Kutta methods [4] and [12]. Our contribution is to use Taylor series to formulate the nonlinear system for  $y_{i+1}$ .

Any numerical one-step method applied to (2.1) reduces to

$$y_{i+1} = R(z)y_i \quad z = h\lambda, \quad (2.3)$$

where  $R : \mathbb{C} \rightarrow \mathbb{C}$  is a polynomial or (in the case of implicit methods) a rational function with real coefficients.  $R(z)$  is called the *stability function* of the method.

**Definition:** A numerical method applied to (2.1) as well as its stability function  $R(z)$  is called *A-stable* if the left half-plane  $\{z : \operatorname{Re}(z) \leq 0\}$  is contained in the region  $\{z : |R(z)| \leq 1\}$ .

On the one hand, the definition of A-stability is too weak: only linear problems with constant coefficients are covered. On the other hand, the definition is too strong: many methods that perform well in practice are not A-stable. Thus we are led to the following generalization.

**Definition:** A numerical method is *A( $\alpha$ )-stable* if the sector  $\{z : |\arg(-z)| \leq \alpha, z \neq 0\}$  is contained in the stability region  $\{z : |R(z)| \leq 1\}$ .

In contrast to A-stable methods, there exists little *nonlinear* convergence theory in the literature for methods that are only A( $\alpha$ )-stable. A step in this direction for BDFs was given by Lubich [19].

We were led to consider implicit Taylor series methods for the validated solution of initial value problems for stiff ODEs because explicit Taylor series methods that have proven effective for nonstiff systems [6, 2] cannot be expected to handle highly stiff systems successfully and because Lohner's program AWA [18]) has proven so successful using Taylor series methods for computing inclusions.

### 3 Implicit Taylor Series Methods (ITSM)

The first formulations of ITSM are due to Chang and Corliss [5] and to Stetter [25]. Let us start by recalling explicit Taylor series methods. Let  $y_i$  be an approximation for the solution of (2.2) at  $t = t_i$ . The explicit Taylor series method uses recurrence relations derived from the ODE to generate the series for  $y(t)$  expanded at  $t_i$ . Then

$$y_{i+1} := y_i + \sum_{j=1}^k f_j(y_i) \frac{h^j}{j!}, \quad (3.4)$$

where  $f_j(y) := \frac{d^j y}{dt^j}$  for  $j = 0, \dots, k-1$ , and  $h := t_{i+1} - t_i$  is taken as large as possible consistent with error control constraints.

The simplest form of the ITSM is as follows:

```

Initialize
Loop for each integration step
  Guess step size
  Loop
    Guess  $y_{i+1}$ 
    Generate series for  $y$  expanded at  $t = t_{i+1}$ 
    Exit loop when  $y_i = y_{i+1} + \sum_{j=1}^k f_j(y_{i+1}) \frac{(t_i - t_{i+1})^j}{j!}$ 
      is satisfied according to tolerance parameters
  Estimate error
  Accept or reject step

```

The series at  $t = t_{i+1}$  is generated by using exactly the same recurrence relations as in the explicit method. That is, each  $f_j(y_{i+1})$  is some nonlinear function of  $y_{i+1}$ . The difference is that the “initial condition”  $y_{i+1}$  is determined by a Newton iteration, rather than being given by analytic continuation at the previous step. See Figure 1.

The Newton iteration for the equation

$$y_i = y_{i+1} + \sum_{j=1}^k f_j(y_{i+1}) \frac{(t_i - t_{i+1})^j}{j!} \quad (3.5)$$

requires

$$\frac{\partial}{\partial y_{i+1}} \left( y_{i+1} + \sum_{j=1}^k f_j(y_{i+1}) \frac{(t_i - t_{i+1})^j}{j!} \right) = \mathbf{1} + \sum_{j=1}^k \frac{\partial}{\partial y_{i+1}} (f_j(y_{i+1})) \frac{(t_i - t_{i+1})^j}{j!}, \quad (3.6)$$

which is computed from the solution of the variational equation  $U' = \frac{\partial f}{\partial y} U$  [17].

The stability function  $R(z)$  corresponding to this method is a  $(0, k)$ -Padé approximation to the exponential  $\exp(z)$  [12], where  $k$  is the degree of the Taylor series. It is well




Figure 1: Implicit Taylor series method

known that the  $(m, k)$ -Padé approximation is the unique rational function with numerator and denominator of degree  $m$  and  $k$ , respectively, which approximates  $\exp(z)$  to  $O(z^{k+m+1})$  as  $z \rightarrow 0$ .

Any  $(m, k)$ -Padé approximation to  $\exp(z)$  is A-stable if and only if  $k - 2 \leq m \leq k$ . That is, the diagonal and two subdiagonals in the Padé scheme are A-stable (see [12] and [13]).

In our case, only implicit Taylor series of degree 1 and 2 lead to A-stable ITSM. For degree  $k = 1$ , the resulting method is the implicit Euler method, whose stability and convergence properties are well known. For degree  $k = 2$ , the resulting method corresponds to a special Runge-Kutta scheme, Lobatto IIIC with the number of stages  $s = 2$ . The corresponding stability function is a  $(s-2, s)$ -Padé approximation. Schneid [24] showed that the Lobatto IIIC method for  $s = 2$  is not only A-stable, but also B-convergent of order 2 under some reasonable step-size restrictions.

The corresponding stability regions for the  $(0, k)$ -Padé approximation (see Figure 2) were drawn by using the software product S [3]. Sand and Østerby [23] give stability regions for certain Runge-Kutta methods that are qualitatively similar to Figure 2.

## 4 Implicit $\sigma$ Taylor Series Methods ( $I\sigma$ TSM)

The lack of A-stability of the simple ITSM for higher orders suggests that we try to increase the degree of the numerator in the corresponding rational stability function.

The ITSM matches the previously computed  $y_i$ , with the value obtained by expanding the solution at  $t_{i+1}$ . The  $I\sigma$ TSM generates the Taylor series at *both*  $t_i$  and  $t_{i+1}$ . The condition that the two series agree at  $t_i + \sigma(t_{i+1} - t_i)$ ,  $\sigma \in (0, 1)$ , provides the nonlinear equation for  $y_{i+1}$  (see Figure 3).




Figure 2: Stability regions of the  $(0, k)$ -Padé approximation for  $k = 1, 2, 3, 4, 5, 6$




Figure 3: Implicit  $\sigma$  Taylor series method

The test

$$y_i \approx y_{i+1} + \sum_{j=1}^k f_j(y_{i+1}) \frac{(t_i - t_{i+1})^j}{j!} \quad (4.7)$$

in the ITSM is now replaced by the test

$$y_i + \sum_{j=1}^m f_j(y_i) \frac{\sigma^j (t_{i+1} - t_i)^j}{j!} \approx y_{i+1} + \sum_{j=1}^k f_j(y_{i+1}) \frac{(\sigma - 1)^j (t_{i+1} - t_i)^j}{j!}. \quad (4.8)$$

In general,  $m$  need not be equal to  $k$ . The  $\text{I}\sigma\text{TSM}$  requires almost no work beyond that required by the ITSM. The series at  $t_{i+1}$  must be recomputed for each new iterate  $y_{i+1}$ , as in the ITSM described in Section 3. The series at  $t_i$ , however, is computed only once per step, and this computation has already been done at the end of the previous step. Here,  $\sigma$  is a tuning parameter of the method. The case  $\sigma = 1$  represents an explicit Taylor series, whereas  $\sigma = 0$  leads to the fully implicit form of Section 3. The case  $\sigma = \frac{1}{2}$  is the unique choice of  $\sigma$  for which the resulting method is A-stable, for any order  $m = k$ . Unfortunately, the resulting stability function is *not* a Padé approximation if the orders  $m, k$  of the Taylor series are higher than 1. Hence, the maximal achievable order of the local truncation error is reduced compared to the maximal possible order for the Padé approximation.

**Remark:** The  $\text{I}\sigma\text{TSM}$  with  $k = m = 1$  and  $\sigma = \frac{1}{2}$  is the well-known implicit trapezoidal rule, which is A-stable, but not B-stable.

In a forthcoming paper [7], a class of high-order stiff ordinary differential equations solvers based on Padé approximations is introduced and analyzed. In this approach, the advantages of Taylor series methods mentioned above are maintained and combined with highest possible order of the local truncation error. A general nonlinear stability analysis remains to be done.

## 5 Implications for Interval Methods

The first task when developing an algorithm for validated inclusions of solutions to stiff problems is to understand an interval analog of the implicit Euler scheme  $y_{i+1} = y_i + hf(t_{i+1}, y_{i+1})$ . Let  $\tilde{y}_i$  represent an approximate solution, and let  $[e_i]$  be an interval inclusion for the corresponding error. We represent an interval-valued solution at  $t_i$  as  $[y_i] = \tilde{y}_i + [e_i]$ . The implicit Euler method becomes

$$e_{n+1} - hf_y(\tilde{y}_{n+1} + [0, 1]e_{n+1}) \cdot e_{n+1} = e_n - \frac{h^2}{2} f_y(y(\tau)) \cdot f(y(\tau)), \quad (5.9)$$

for  $e_n \in [e_n]$ , and  $y(\tau)$  contained in an *a-priori* inclusion  $\tilde{Y}_{n+1}$ . This is a nonlinear system of interval equations. In the special case of a linear constant coefficient problem  $y' = Ay$ , the corresponding system reduces to a linear system with interval right-hand side. The methods presented in this paper can be viewed as higher-order generalizations of this simplest implicit scheme.

Another difficulty in the case of stiffness is the inevitably large Lipschitz constant  $L$  of the right-hand  $f(t, y)$  with respect to  $y$ . This would cause a severe and unrealistic step-size restriction whenever the Picard-Lindelöf existence theorem is applied to get an *a-priori* inclusion. Subtle algorithms have to be developed.

During the preparation of this manuscript, we received Kreuser's thesis [16], which presents an alternative approach to computing inclusions of solutions for stiff ODEs. Kreuser's approach transforms the original stiff system (2.2) into the new system

$$y' = Ay + (f(t, y) - Ay),$$

where  $A$  is a local approximation of the Jacobian  $f_y(t, y)$ . The *a-priori* inclusion is obtained by symbolic computation of the matrix exponential function, thus avoiding the severe step-size restriction inherent in Lohner's explicit approach [17]. To get an inclusion  $[y_i] = \tilde{y}_i + [e_i]$  for the solution at  $t_i$ , the approximate solution  $\tilde{y}_i$  is obtained by integrating the linear ODE and applying a Taylor series method to the nonlinear part. The inclusion  $[e_i]$  is obtained by iterating an interval integral operator. The method of Kreuser successfully handles stiff problems of the special form  $y' = Ay + g(t, y)$ , where  $g(t, y)$  is Lipschitz continuous in  $y$ . This excludes strongly nonlinear models.

## Acknowledgments

The authors thank Prof. H. J. Stetter and W. Auzinger for many valuable discussions. Special thanks are also due to an anonymous gardener from the Viennese Stadtpark who found and returned the first author's backpack after it had been stolen; it contained all written records and source material for the work reported here.

## References

- [1] E. Adams, Periodic solutions: Enclosure, verification, application, in *Computer Arithmetic and Self Validating Numerical Methods*, Proceedings of SCAN 1989 (Basel), Academic Press, Boston, 1990, 199–245.
- [2] D. Barton, I. M. Willers, R. V. M. Zahar, The automatic solution of ordinary differential equations by the method of Taylor series, *Comp. J.*, **14** (1971), 243–248.
- [3] R. A. Becker, J. M. Chambers, A. R. Wilks, *The New S Language: A Programming Environment for Data Analysis and Graphing*, Wadsworth and Brooks/Cole, Pacific Grove, Calif., 1988.
- [4] J. C. Butcher, Implicit Runge-Kutta processes, *Math. Comp.*, **18** (1964) 50–64.
- [5] Y. F. Chang, G. F. Corliss, Personal communication.



- [6] G. F. Corliss, Y. F. Chang, Solving ordinary differential equations using Taylor series, *ACM Trans. Math. Software*, **8** (1982), 114–144.
- [7] G. F. Corliss, A. Griewank, F. Potra, G. Kirlinger, H. J. Stetter, Personal communication.
- [8] R. Courant, K. Friedrichs, H. Lewy, Über die partiellen Differenzgleichungen der mathematischen Physik, *Math. Ann.*, **100** (1928), 32–74.
- [9] J. Crank, P. Nicholson, A practical method for numerical integration of solutions of partial differential equations of heat conduction type, *Proc. Cambridge Philos. Soc.*, **43** (1947), 50 ff.
- [10] C. F. Curtis, J. O. Hirschfelder, Integration of stiff equations, *Proc. Nat. Acad. Sci.*, **38** (1952), 235–243.
- [11] G. Dahlquist, A special stability problem for linear multistep methods, *BIT*, **3** (1963), 27–43.
- [12] K. Dekker, J. G. Verwer, *Stability of Runge-Kutta Methods for Stiff Nonlinear Differential Equations*, North-Holland, Amsterdam, 1984.
- [13] B. L. Ehle, A-stable methods and Padé approximations to the exponential, *SIAM J. Math. Anal.* **4** (1973), 671–680.
- [14] P. Eijgenraam, *The Solution of Initial Value Problems Using Interval Arithmetic*, Stichting mathematisch Centrum Amsterdam, Mathematical Center Tracts No. 144, 1981.
- [15] R. Frank, J. Schneid, Ch. W. Überhuber, The concept of B-convergence, *SIAM J. Numer. Anal.*, **18** (1981), 753–780.
- [16] A. Kreuser, *Zur Lösungseinschließung bei steifen Differentialgleichungen durch eine Verbindung von Computeralgebra und Intervallarithmetik*, Diplomarbeit (Universität Bonn), 1991.
- [17] R. Lohner, *Einschließung der Lösung gewöhnlicher Anfangs- und Randwertaufgaben und Anwendungen*, Dissertation (Universität Karlsruhe), 1988.
- [18] R. Lohner, *AWA, Software product in FORTRAN-SC for the inclusion of the solution of ODEs*, Karlsruhe, 1989.
- [19] Ch. Lubich, On the convergence of multistep methods for nonlinear stiff differential equations, *Numer. Math.* **58** (1991) 839–853.
- [20] R. E. Moore, *Interval Analysis*, Prentice-Hall, Englewood Cliffs, N.J., 1966.
- [21] R. E. Moore, *Methods and Applications of Interval Analysis*, SIAM, Philadelphia, 1979.

- [22] L. B. Rall, *Automatic Differentiation: Techniques and Applications*, Lecture Notes in Comp. Sci. No. 120, Springer-Verlag, Berlin, 1981.
- [23] J. Sand, O. Østerby, *Regions of absolute stability*, DAIMI PB-102, Computer Science Department, Aarhus University, September 1979.
- [24] J. Schneid, B-convergence of Lobatto IIC formulas, *Num. Math.*, **51** (1987), 229–235.
- [25] H. J. Stetter, Validated solution of initial value problems for ODE, in *Computer Arithmetic and Self Validating Numerical Methods*, Proceedings of SCAN 1989 (Basel), Academic Press, Boston, 1990, 171–187.

**END**

**DATE  
FILMED**

**01/27/92**

