



**Fermi National Accelerator Laboratory**

**FERMILAB-Conf-91/286-E**

## **Search for Top Quark at Fermilab Collider**

**The CDF Collaboration**

*Fermi National Accelerator Laboratory  
P.O. Box 500, Batavia, Illinois 60510*

presented by K. Sliwa

*Tufts University, Department of Physics  
Medford, Massachusetts 02155*

October 1991

\* Presented at the "4th International Symposium on Heavy Flavour Physics", Universite de Paris-Sud, Orsay, France, June 25-29, 1991.

TUFTS CONF 91/#13-1-E  
FERMILAB-CONF-91/286-E  
CDF/PUB/HEAVYFLAVOR/PUBLIC/1583

## SEARCH FOR TOP QUARK AT FERMILAB COLLIDER

The CDF Collaboration<sup>1)</sup>

presented by Krzysztof Sliwa  
Tufts University, Department of Physics  
Medford, Massachusetts 02155 USA

### ABSTRACT

The status of a search for the top quark with Collider Detector at Fermilab (CDF), based on a data sample recorded during the 1988-1989 run is presented. The plans for the next Fermilab Collider run in 1992-1993 and the prospects of discovering the top quark are discussed.

Presented at "4<sup>th</sup> International Symposium on Heavy Flavour Physics", Universite de Paris-Sud, Orsay, France, 25-29 June 1991.

## i. Introduction

The CDF limit on the mass of the top quark has not changed significantly from the one presented over a year ago<sup>2)</sup>. This is not because of lack of effort, simply, CDF has not accumulated any new data since the 1988-1989 run ended. Many systematic effects have been studied in detail, most importantly the total luminosity is now understood with uncertainty reduced by more than a factor of two (6.8 % now as opposed to 15% a year ago). A combined analysis of dilepton and e+multijet+ $\cancel{E}_t$  events, which led to the most stringent limit on the mass of the top quark, has been extended beyond the central region of the CDF detector to include the plug. However, the effect of increased acceptance has been almost exactly cancelled by the decrease of the total luminosity (from 4.4 pb<sup>-1</sup> to 4.1 pb<sup>-1</sup>) and the CDF limit on the mass of the top quark remains almost unchanged. Because the final details of this new analysis are still being discussed, the results presented at this conference correspond to the status of analysis as of Moriond 90', where the combined, from the di-leptons and low-P<sub>t</sub> searches, upper limit on top quark mass  $M_t > 89 \text{ GeV}/c^2$  @ 95% C.L. has been presented. (The final limit is only slightly better,  $M_t > 90\text{-}91 \text{ GeV}/c^2$  @ 95% C.L.).

## ii. Published CDF searches for $t\bar{t}$

At Fermilab Tevatron energy,  $\sqrt{s} = 1.8 \text{ TeV}$ , the dominant production mechanism of top quarks is the production of  $t\bar{t}$  pairs from a quark-antiquark or a gluon-gluon initial state through the strong interactions.

CDF has published<sup>3-4)</sup> results from two searches for the top quark, and preliminary results of analysis which combined several decay modes has been presented at various conferences.

An analysis<sup>3)</sup> of e+multijet+ $\cancel{E}_t$  events was based on the fact that, for top quark mass smaller than W mass, the transverse mass distribution of the e+ $\cancel{E}_t$  system,  $M_{t\bar{t}}^{e\nu}$ , is softer for  $t\bar{t}$  than for W+jets events. The observed distributions were found consistent with W boson production. The resulting upper limit on the  $t\bar{t}$  cross section excluded a Standard Model top quark mass in the range  $40 \text{ GeV}/c^2 < M_t < 77 \text{ GeV}/c^2$ . It is important to note this method loses its sensitivity with  $M_t$  approaching  $M_W$ , in which case the  $M_{t\bar{t}}^{e\nu}$  distributions become indistinguishable.

The other published analysis<sup>4)</sup> searched for high transverse energy e+ $\mu$  events, in which final state the background from other processes is very small. One candidate event was found. Translating a limit on  $t\bar{t}$  cross section found in this analysis yielded a limit on top quark mass of  $M_t > 72 \text{ GeV}/c^2$ .

## iii. Search for $t\bar{t}$ in dilepton and e+multijet+ $\cancel{E}_t$ events

In a natural extension of those two analyses additional decay modes were studied.

Dilepton analysis searched for events in which both top quarks decayed leptonically, for example  $t \rightarrow e^+ \nu_e b$  and  $\bar{t} \rightarrow e^- \bar{\nu}_e \bar{b}$ . Such events have a spectacular signature of two high transverse energy ( $E_t$ ) leptons accompanied with missing transverse energy ( $\cancel{E}_t$ ) and possibly soft b-quark jets.

If one of the top quarks decays leptonically,  $t \rightarrow be^+\nu_e$ , and the other hadronically,  $\bar{t} \rightarrow \bar{b}q\bar{q}$ , the final state will be that of an  $e+\text{multijet}+\cancel{E}_t$  event. The two quark jets from a  $W$  decay would be highly energetic, while the  $b$ -jets should be softer. In an attempt to tag one of the two  $b$ -jets a presence of an additional soft lepton was required.

The results of di-lepton and  $e+\text{multijet}+\cancel{E}_t+b\text{-tag}$  were combined. With increased acceptance, a more stringent limit on the mass of the top quark has been derived.

### *Dilepton search for $t\bar{t}$*

In the central region ( $|\eta| < 1.0$  for  $e^\pm$  and  $|\eta| < 1.2$  for  $\mu^\pm$ ;  $|\eta|$  - pseudorapidity), measuring the electron and muon parameters region is well understood and the knowledge of systematic errors is very good. CDF has analysed the high transverse energy  $e^+e^-$  and  $\mu^+\mu^-$  pairs in the central region, removing  $Z^0 \rightarrow e^+e^-$  and  $Z^0 \rightarrow \mu^+\mu^-$  candidates. The  $e+\mu$  channel has been reanalysed with the same cuts as for the other di-lepton channels. The details of event selection, in its lepton identification part, follow the  $e+\mu$  analysis<sup>4)</sup>. Events with opposite charge electrons or muons, and with  $E_t^e > 15$  GeV and  $P_t^\mu > 15$  GeV/ $c^2$ ) were selected. In the  $e^+e^-$  and  $\mu^+\mu^-$  events the  $Z^0$  events were removed with the help of:

- a) a cut on the di-lepton mass,  $75 \text{ GeV}/c^2 > M_{l+l-} > 105 \text{ GeV}/c^2$ ;
- b) a missing transverse energy cut,  $\cancel{E}_t > 20$  GeV;
- c) a cut on the azimuth angle between the leptons,  $20^\circ < \Delta\phi_{l+l-} < 160^\circ$ .

In Figure 1 we present the  $M_{e+e-}$  and  $\cancel{E}_t$  distributions for a large sample ( $730 \text{ pb}^{-1}$ ) of  $t\bar{t}$  Monte Carlo data with  $M_t=90 \text{ GeV}/c^2$ , and for CDF di-electron events. Figure 2 shows the  $\Delta\phi_{e+e-}$  distributions, and scatterplots of  $\Delta\phi_{e+e-}$  vs  $\cancel{E}_t$ , after  $M_{e+e-}$  cut, for Monte Carlo and data. No events pass the  $M_{e+e-}$ ,  $\Delta\phi$  and  $\cancel{E}_t$  cuts. Figure 3 shows the analogous  $\Delta\phi_{l+l-}$  vs  $\cancel{E}_t$  scatterplots for the  $\mu^+\mu^-$  and  $e^\pm\mu^\mp$  events. No  $\mu^+\mu^-$  events and one, already known<sup>4)</sup>,  $e\mu$  event is found after the  $M_{l+l-}$ ,  $\Delta\phi$  and  $\cancel{E}_t$  cuts are applied. The kinematics characteristics of the  $e\mu$  candidate event is presented in Table I.

Table II summarizes, as a function of top quark mass, the experimental acceptances (studied with ISAJET 6.21 and full CDF detector simulation Monte Carlo),  $t\bar{t}$  cross section<sup>5)</sup> and the expected number of events in  $4.4 \text{ pb}^{-1}$ . The total systematic error for the three di-lepton channels combined was 20%. The branching fraction of 4/81 has been assumed. The most important systematic error in the di-lepton analysis were uncertainty in luminosity (15%) and  $P_t$  spectrum uncertainties (10%).

After convoluting<sup>6,7,8)</sup> the 20% systematic error with a Poisson distribution a 95% C.L. upper limit of 5.16 events for a number of  $t\bar{t}$  events seen with CDF in the di-lepton channels has been found. An upper limit on the  $t\bar{t}$  cross section translates, when compared with the results of a theoretical calculation in the framework of the Standard Model<sup>5)</sup>, into a limit on the mass of the top quark,  $M_t > 84 \text{ GeV}/c^2$  @ 95% C.L.

### *Low- $P_t$ muon search*

A new technique was used to analyse the  $e$ +multijet and  $\mu$ +multijet events, which could occur when one of the top quarks quark decays leptonically and the other hadronically, leading to a lepton+ $\bar{\nu}_t$ +4jets final state. Selection cuts in electron+multijet case are virtually identical to those used in  $e$ +jets top search<sup>3</sup>). A sample of 105 events was selected by requiring  $\bar{E}_t > 20$  GeV,  $E_t^e > 20$  GeV, and at least two jets with  $E_t^j > 10$  GeV and within  $|\eta| < 2.0$ . The muon+multijet sample consisted of 87 events, with jet and  $\bar{E}_t$  cuts as in electron+multijet channel, and  $P_t^\mu > 20$  GeV/ $c^2$ . A new technique used here makes an attempt to tag one of the two b-quark jets by requiring a low transverse momentum muon to be well isolated from any of the two highest  $E_t$  jets,  $\Delta R > 0.5$ ,  $\Delta R \equiv [(\Delta\eta)^2 + (\Delta\phi)^2]^{1/2}$ . (According to a Monte Carlo study muons from b-jets have  $P_t^\mu < 15$  GeV/ $c^2$ .) Figure 4 shows the distributions of  $\Delta R$  between the low- $P_t$  muon and the nearest, in  $R$  space, of the two highest  $E_t$  jets for a)  $250 \text{ pb}^{-1}$  Monte Carlo  $t\bar{t}$  events generated with  $M_t=90$  GeV/ $c^2$ , and b)  $4.4 \text{ pb}^{-1}$  CDF data. No candidates were found.

In Table II we present, as a function of top quark mass, the experimental acceptance and the expected number of events in  $4.4 \text{ pb}^{-1}$  for the low- $P_t$  search. The combined branching fraction of 10.9% was assumed ( $2/3 \times 2/9 \times 2 \times (\text{probability of any of the two b-quarks to decay into a } \mu) = 8/27 \times 0.37$ ). The combined systematic error was 25%, with the largest contributions being the uncertainty in luminosity (15%), Monte Carlo statistics (12%) and B semileptonic branching fractions (9.4%)

### iii. Evaluation of a combined upper limit

Suppose one performs an analysis in  $m$  channels, with  $\epsilon_i$  being an efficiency for observing  $t\bar{t}$  events in each of the modes. The probability of observing  $n_i^o$  events in  $i$ -th channel is given by a likelihood function  $L \equiv \prod_{i=1}^m P(n_i^o; \epsilon_i N_{t\bar{t}})$ , where  $\epsilon_i N_{t\bar{t}}$  is the mean number of events expected in the  $i$ -th channel, and  $P(n; \mu)$  is Poisson probability distribution function with the mean  $\mu$ . Solving for  $\overline{N_{t\bar{t}}}$  that maximizes  $L$ , one finds:  $\overline{N_{t\bar{t}}} = \sum_{i=1}^m n_i^o / \sum_{i=1}^m \epsilon_i$ . To incorporate the systematic uncertainties we use a standard procedure<sup>6,7,8</sup>), where the correlations in uncertainties of acceptances were properly taken into account. Figure 5 shows the upper limits on the  $t\bar{t}$  cross section superimposed on the results of a theoretical calculation based on the Standard Model<sup>5</sup>). The combined upper limit on the  $t\bar{t}$  cross section, from the di-leptons and low- $P_t$  searches, leads to a lower limit on top quark mass  $M_t > 89$  GeV/ $c^2$  @ 95% C.L.

### iv. Physics reach of the future 1992-1993 Fermilab Collider run

The next physics run with Fermilab Collider is supposed to begin in March 1992. For those familiar with Fermilab schedule it means perhaps only that it will not begin before this date. With no more SppS runs, Fermilab's CDF and D0. are in a special position of being the only two experiments which may find the top quark in the near future.

D0 will have it's first physics run in 1992. Compared with CDF it has superior  $\mu$  coverage and good electromagnetic and hadronic calorimetry in a broader rapidity range. On the other hand, it does not have a magnetic field making particles' momentum and charge determination from tracking impossible. A crude momentum measurement is only available for  $\mu$ , in D0 magnetized

iron toroids outside the calorimeters. However, CDF has demonstrated that the signatures of possible  $t\bar{t}$  production are spectacular and clean, and it seems that D0 detector can do as well as CDF, if not better because of its better acceptance, in a search for top quark in the  $e\mu$  channel.

CDF is extending its  $\mu$  coverage in preparation for the next run. Also, its new silicon vertex detector may prove essential for attempts to tag the b-jets from  $t, \bar{t}$  decays.

Having two detectors, implementing different, and in the same time complementary, conceptual approach to detection of high energy physics processes may become very important if, say, more  $e\mu$  events are detected next year. It is relatively simple to set an upper limit, as shown in this paper. To *prove* that the observed events are *indeed* due to  $t\bar{t}$  production is more difficult. One would have to confirm the characteristics expected for  $t\bar{t}$  events in, most desirably, several independent channels.

Another reason why analysis becomes more difficult with increasing top mass is the phase space suppression. The higher the top quark mass the smaller the  $t\bar{t}$  production cross section. The backgrounds from other processes, even if found small so far in CDF searches, may overwhelm the possible signal. Let us recall, as an example, that for  $M_t=150$  GeV the cross section for  $t\bar{t}$  production is equal to that for  $W^+W^-$  production. A search for  $e\mu$  events from  $t\bar{t}$  will inevitably find an equal number of such events, with quite similar characteristics, from direct di-boson production.

The plan for the next Collider run is to accumulate events corresponding to about  $100 \text{ pb}^{-1}$ . The run may be divided into two parts, and in such a case  $25 \text{ pb}^{-1}$  are expected to be collected in the first running period.

Assuming no new candidates are found, the single most important factor in estimating how far CDF and D0 will be able to push the lower limit on the mass of the top quark is the integrated luminosity. The upper bound on the  $t\bar{t}$  cross section corresponding to one event (the  $e\mu$  candidate found in the last CDF run) would mean  $M_t > 120 \text{ GeV}/c^2$  for an integrated luminosity of  $25 \text{ pb}^{-1}$ ;  $M_t > 160 \text{ GeV}/c^2$  for a luminosity of  $75 \text{ pb}^{-1}$  and  $M_t > 170 \text{ GeV}/c^2$  for a luminosity of  $100 \text{ pb}^{-1}$ . (The above estimate may prove slightly conservative, since CDF is extending its  $\mu$  acceptance, and D0 detector has a good  $\mu$  coverage already.)

Alterelli<sup>9)</sup> has presented at this conference the most up-to-date results of a "consistency check" of the Standard Model. The mass of the top quark is now (compiling all available measurements and assuming the validity of the Standard Model) expected to be in the range  $M_t \in (95,190)$  GeV, with the central value of predictions varying with the Higgs mass, namely,  $M_t \approx 115$  GeV for a light Higgs and  $M_t \approx 150$  GeV for a heavy Higgs. This is *within* the reach of the  $100 \text{ pb}^{-1}$  run. Let the beams collide !

## REFERENCES

1. Argonne National Laboratory, Brandeis University, University of Chicago, Fermi National Accelerator Laboratory, Laboratori Nazionali di Frascati (INFN), Harvard University, University of Illinois, The Johns Hopkins University, National Laboratory for High Energy Physics (KEK), Lawrence Berkeley Laboratory, University of Padova, University of Pennsylvania, INFN University and Scuola Normale Superiore of Pisa, Purdue University, University of Rochester, Rockefeller University, Rutgers University, Texas A&M University, University of Tsukuba, Tufts University and University of Wisconsin.
2. "A new limit on the mass of the top quark", CDF Collaboration, K. Sliwa, Proceedings of the XXV-th Rencontres de Moriond, Leptonic Session, Les Arcs, Savoie, France, March 4-11, 1990; Editions Frontieres, M65, p459.
3. F. Abe *et al.*, Phys. Rev. Lett. **63** (1990) 142.
4. F. Abe *et al.*, Phys. Rev. Lett. **63** (1990) 147.
5. G. Altarelli, M. Diemoz, G. Martinelli, and P. Nason, Nucl. Phys. **B308** (1988) 724.
6. G. Zech, CERN-EP/88-164, November 1988.
7. Particle Data Group, Phys. Lett. **B204** (1988) 1.
8. CDF note CDF/ANAL/HEAVYFLAVOR/1109, January 1990.
9. G. Altarelli, these Proceedings.

**TABLE I: Characteristics of the CDF  $e\mu$  candidate event, Run 19250, event 20425. (Jet energies are the corrected energies, the uncorrected energies are listed in paranthesis for the sake of completeness.)**

	charge	E <sub>T</sub> (GeV)	η	φ radians
central electron	+	31.72	-0.81	2.30
central muon	-	42.54	-0.80	4.70
forward muon	+	7.58	-1.96	1.70
jet 1		19.73 (12.56)	1.07	5.96
jet 2		8.89 (5.0)	-2.76	1.49

**TABLE II:  $t\bar{t}$  cross section, acceptance×branching fractions, and predicted number of events in  $4.4 \text{ pb}^{-1}$  for the di-leptons and low- $P_t$  muon analyses.**

Top mass (GeV/c <sup>2</sup> )	σ( $t\bar{t}$ ) (pb)	acc×BF (di-leptons)	#events	acc×BF (low- $P_t$ μ)	#events
75	385	.00605	10.2	.00152	2.6
80	285	.00709	8.9	.00181	2.3
85	200	.00746	6.6	.00228	2.0
90	153	.00774	5.2	.00233	1.6







Figure 1. a)  $e^+e^-$  mass distribution for electrons with transverse energy  $\cancel{E}_t > 15$  GeV from a  $730 \text{ pb}^{-1}$  sample of Monte Carlo  $t\bar{t}$  events generated with top quark mass of  $90 \text{ GeV}/c^2$ ; b)  $e^+e^-$  mass distribution for electrons with  $\cancel{E}_t > 15$  GeV from a  $4.4 \text{ pb}^{-1}$  CDF data sample,  $Z^0$  peak is clearly visible; c) missing transverse energy ( $\cancel{E}_t$ ) distribution for  $e^+e^-$  events from a  $730 \text{ pb}^{-1}$  sample of Monte Carlo  $t\bar{t}$  events generated with  $M_t=90 \text{ GeV}/c^2$ ; d)  $\cancel{E}_t$  distribution for  $e^+e^-$  events from CDF data.









Figure 2. a) Distribution of the azimuth angle,  $\Delta\phi_{e^+e^-}$ , between the two electrons from a  $730 \text{ pb}^{-1}$  sample of Monte Carlo  $t\bar{t}$  events generated with top quark mass of  $90 \text{ GeV}/c^2$ ; b)  $\Delta\phi_{e^+e^-}$  distribution for CDF data sample; c)  $\Delta\phi_{e^+e^-}$  vs  $\cancel{E}_t$  scatterplot for  $90 \text{ GeV}/c^2$  top quark Monte Carlo events, after a  $M_{e^+e^-}$  cut to remove  $Z^0$  events; d)  $\Delta\phi_{e^+e^-}$  vs  $\cancel{E}_t$  scatterplot for CDF data,  $Z^0$  events removed.




Missing  $E_t$  for  $\mu^+\mu^-$  events (GeV)



Missing  $E_t$  for  $e^+\mu^-$  events (GeV)



Missing  $E_t$  for  $\mu^+\mu^-$  events (GeV)



Missing  $E_t$  for  $e^+\mu^-$  events (GeV)

Figure 3. a)  $\Delta\phi_{\mu+\mu^-}$  vs  $\cancel{E}_t$  scatterplot for 90 GeV/c<sup>2</sup> top quark Monte Carlo events, after a  $M_{\mu+\mu^-}$  cut to remove  $Z^0$  events; b)  $\Delta\phi_{\mu+\mu^-}$  vs  $\cancel{E}_t$  scatterplot for CDF data,  $Z^0$  events removed. c)  $\Delta\phi_{e+\mu^-}$  vs  $\cancel{E}_t$  scatterplot for 90 GeV/c<sup>2</sup> top quark Monte Carlo events, after a  $M_{e+\mu^-}$  cut; d)  $\Delta\phi_{e+\mu^-}$  vs  $\cancel{E}_t$  scatterplot for CDF data, after a  $M_{e+\mu^-}$  cut.




Figure 4. The distributions of  $\Delta R$ ,  $\Delta R \equiv [(\Delta\eta)^2 + (\Delta\phi)^2]^{1/2}$ , between the low- $P_t$  muon and the nearest, in  $R$  space, of the two highest  $E_t$  jets for a)  $250 \text{ pb}^{-1}$  Monte Carlo  $t\bar{t}$  events generated with  $M_t=90 \text{ GeV}/c^2$ , and b)  $4.4 \text{ pb}^{-1}$  CDF data.