

9-6-2012

# Synthesis and Characterization of Chain-End Functionalization of Glycosaminoglycans

Nikola Paulic

Cleveland State University, N.PAULIC90@csuohio.edu

Jacob J. Weingart

Cleveland State University, J.J.WEINGART@csuohio.edu

Follow this and additional works at: [https://engagedscholarship.csuohio.edu/u\\_poster\\_2012](https://engagedscholarship.csuohio.edu/u_poster_2012)

 Part of the [Organic Chemistry Commons](#)

**How does access to this work benefit you? Let us know!**

---

## Recommended Citation

Paulic, Nikola and Weingart, Jacob J., "Synthesis and Characterization of Chain-End Functionalization of Glycosaminoglycans" (2012). *Undergraduate Research Posters 2012*. 32.

[https://engagedscholarship.csuohio.edu/u\\_poster\\_2012/32](https://engagedscholarship.csuohio.edu/u_poster_2012/32)

This Book is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2012 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact [library.es@csuohio.edu](mailto:library.es@csuohio.edu).



This digital edition was prepared by MSL Academic Endeavors, the imprint of the Michael Schwartz Library at Cleveland State University.

# **Synthesis and Characterization of Chain-End Functionalization of Glycosaminoglycans**

College of Science and Health Professions

Department of Chemistry

**Student Researchers:** Nikola Paulic; Jacob Weingart

**Faculty Advisor:** Xue-Long Sun, Ph.D.

## **Abstract**

Chondroitin sulfate (CS) is a glycosaminoglycan (GAG) composed of two alternating sugars, glucuronic acid and *N*-acetylgalactosamine, with varying patterns of sulfation determining its type. Found in connective tissue extracellular matrices and on cell surfaces proteoglycans and is involved in a multitude of biological and chemical signaling events such as CNS development, coagulation, and cellular signal transduction. TM is a glycoprotein and its CS-containing domain contributes TM's thrombin binding and stability and plasminogen activation activities. Previous research has concluded that CS can inhibit the coagulation process by enhancing the binding affinity of the integral membrane protein thrombomodulin to thrombin 10- to 20-fold, subsequently activating Protein C and downregulating clotting Factors Va and VIIIa. We hypothesized that incorporation of CS into recombinant thrombomodulin (rTM), the active domains 4-6 of the native protein, should enhance the binding affinity to thrombin creating a more effective antithrombotic. This presentation reports the CSU Summer Undergraduate Research project, which is the synthesis and characterization of chain-end functionalized CS derivatives, which will be used to investigate various bio-orthogonal conjugation reactions in attempt to create a novel CS-rTM conjugate.