

9-5-2013

# Function of a Novel Checkpoint Protein in the Germ Line

Steven Drellishak  
*Cleveland State University*

Marina Bykova  
*Cleveland State University*

G. Valentin Börner  
*Cleveland State University*

Follow this and additional works at: [https://engagedscholarship.csuohio.edu/u\\_poster\\_2013](https://engagedscholarship.csuohio.edu/u_poster_2013)

 Part of the [Cell and Developmental Biology Commons](#), and the [Genetics Commons](#)

**How does access to this work benefit you? Let us know!**

---

## Recommended Citation

Drellishak, Steven; Bykova, Marina; and Börner, G. Valentin, "Function of a Novel Checkpoint Protein in the Germ Line" (2013).  
*Undergraduate Research Posters 2013*. 15.

[https://engagedscholarship.csuohio.edu/u\\_poster\\_2013/15](https://engagedscholarship.csuohio.edu/u_poster_2013/15)

This Article is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2013 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact [library.es@csuohio.edu](mailto:library.es@csuohio.edu).



This digital edition was prepared by MSL Academic Endeavors, the imprint of the Michael Schwartz Library at Cleveland State University.

## **Function of a novel checkpoint protein in the germ line**

**College of Sciences and Health Professions**

**Department of Biological, Geological & Environmental Sciences**

**Student Researchers: Steven Drellishak; Marina Bykova**

**Faculty Advisor: G. Valentin Börner, Ph.D.**

### **Abstract**

Successful reproduction of *Saccharomyces cerevisiae* relies on the organism's ability to complete the meiotic cell cycle and produce viable gametes. Zip1 is a protein that constitutes the central component of a protein structure that connects homologous chromosomes known as the synaptonemal complex. Zip1 is important for progression through the meiotic cell cycle. The C terminus of the coiled-coil Zip 1 protein is responsible for localization to the axes of the chromosomes. An internal deletion near the C terminus of Zip1, called zip1-c1, yields a stronger meiotic arrest than a mutation where Zip1 is completely deleted. The more efficient meiotic progression in a Zip1 deletion mutation versus the zip1-c1 mutant suggests that zip1-c1 prevents an alternative pathway of meiotic progression. A genomic screen of the Nasmyth genomic library revealed candidate plasmids N5 and N89 containing yeast genes which, when overexpressed, increase spore viability and bypass meiotic arrest in the zip1-c1 mutant. This has implications that the genes on the overexpression plasmids serve some function in correcting mistakes in meiosis when Zip1 is mutated.