

9-6-2012

Key Players in Reverse Cholesterol Transport: The Plasma Enzyme LCAT

Christopher J. Pechura

Cleveland State University, C.PECHURA@csuohio.edu

Follow this and additional works at: https://engagedscholarship.csuohio.edu/u_poster_2012

 Part of the [Enzymes and Coenzymes Commons](#)

How does access to this work benefit you? Let us know!

Recommended Citation

Pechura, Christopher J., "Key Players in Reverse Cholesterol Transport: The Plasma Enzyme LCAT" (2012). *Undergraduate Research Posters 2012*. 31.

https://engagedscholarship.csuohio.edu/u_poster_2012/31

This Book is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2012 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact library.es@csuohio.edu.



This digital edition was prepared by MSL Academic Endeavors, the imprint of the Michael Schwartz Library at Cleveland State University.

Key Players in Reverse Cholesterol Transport: The Plasma Enzyme LCAT

College of Sciences and Health Professions

Department of Chemistry

Student Researcher: Christopher Pechura

Faculty Advisor: Valentin Gogonea

Abstract

Lecithin cholesterol acyltransferase (LCAT) is a plasma enzyme that remodels nascent high density lipoprotein (HDL) into a mature form called spherical HDL. The impeding of this critical step in reverse cholesterol transport (RCT, the transport of cholesterol from periphery cells to liver), leads to atherosclerosis. Our goal is to use computational chemistry and molecular modeling tools to build molecular models for LCAT and its complex with nascent HDL. The molecular models are constructed from low resolution structures of LCAT and LCAT-HDL complex obtained by small angle neutron scattering (SANS) with contrast variation. Ultimately, we combine various experimental data (SANS data, hydrogen-deuterium exchange tandem mass spectrometry, other biochemical data) with computational techniques (bioinformatics, molecular modeling, SANS modeling, protein-protein docking, molecular dynamics simulation) to produce a molecular model for LCAT and its complex with nascent HDL. The model will be used in the future to map the amino acid residues from LCAT and the protein component of nascent HDL (apoA1) involved in mutual interaction, and to identify LCAT residues interacting with the lipid phase of nascent HDL.