

9-5-2013

# Analysis of Turning Behavior in the African Clawed Frog

Rachel Andrikanich  
*Cleveland State University*

William Hein  
*Cleveland State University*

Jeffrey Dean  
*Cleveland State University, J.DEAN@csuohio.edu*

Follow this and additional works at: [https://engagedscholarship.csuohio.edu/u\\_poster\\_2013](https://engagedscholarship.csuohio.edu/u_poster_2013)

 Part of the [Environmental Sciences Commons](#)

**How does access to this work benefit you? Let us know!**

---

## Recommended Citation

Andrikanich, Rachel; Hein, William; and Dean, Jeffrey, "Analysis of Turning Behavior in the African Clawed Frog" (2013).  
*Undergraduate Research Posters 2013*. 6.  
[https://engagedscholarship.csuohio.edu/u\\_poster\\_2013/6](https://engagedscholarship.csuohio.edu/u_poster_2013/6)

This Article is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2013 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact [library.es@csuohio.edu](mailto:library.es@csuohio.edu).



This digital edition was prepared by MSL Academic Endeavors, the imprint of the Michael Schwartz Library at Cleveland State University.

# **Analysis of turning behavior in the African Clawed Frog**

**College of Sciences and Health Professions**

**Department of Biological, Geological & Environmental Sciences**

**Student Researchers: Rachel Andrikanich; William Hein**

**Faculty Advisor: Jeffrey Dean**

## **Abstract**

**African Clawed Toads turn toward waves created by prey. We studied turning of four frogs. We concentrated on body movement, leg action and hip, knee and ankle movements, classified respectively as pushing/pulling or extension/flexion/return. As expected, we observed symmetrical patterns for left and right stimuli. Very rostral stimuli ( $\pm 30^\circ$ ) generally elicited a forward movement as the legs extended and pushed; hips extended while knee and ankle movement varied according to turn angle. More lateral and caudal stimuli commonly elicited an initial backward movement of the rear of the body as one or both legs pulled, reflecting hip flexion. Knee and ankle movement again varied by turn angle and together with the pulling initiated body rotation. Turning usually continued with a sweeping push by the contralateral leg, involving hip and knee extension and knee flexion. Overall leg movement patterns are similar when compared amongst frogs though the strength of joint flexion and extension vary. Variation in joint movement may be influenced by the overall size of the frog.**