

2011

Semen Quality and Chemical Oxidative Stress; Quantification and Remediation

Saleem Ali Bani-Hani
Cleveland State University

Follow this and additional works at: <https://engagedscholarship.csuohio.edu/etdarchive>

 Part of the [Chemistry Commons](#)

How does access to this work benefit you? Let us know!

Recommended Citation

Bani-Hani, Saleem Ali, "Semen Quality and Chemical Oxidative Stress; Quantification and Remediation" (2011). *ETD Archive*. 24.
<https://engagedscholarship.csuohio.edu/etdarchive/24>

This Dissertation is brought to you for free and open access by EngagedScholarship@CSU. It has been accepted for inclusion in ETD Archive by an authorized administrator of EngagedScholarship@CSU. For more information, please contact library.es@csuohio.edu.

SEMEN QUALITY AND CHEMICAL OXIDATIVE STRESS; QUANTIFICATION
AND REMEDIATION

SALEEM ALI BANI-HANI

Bachelor of Science in Analytical Chemistry

Jordan University of Science and Technology

February, 1999

Masters of Science in Medical Laboratory Sciences

Jordan University of Science and Technology

June, 2002

Submitted in Partial fulfillment of the requirements for the degree of
DOCTOR OF PHILOSOPHY IN CLINICAL BIOANALYTICAL CHEMISTRY

at the

CLEVELAND STATE UNIVERSITY

July 2011

This dissertation has been approved for the Department of Chemistry and
the College of Graduate Studies by:

Dissertation Committee Chairperson, Dr. Mekki Bayachou
Department of Chemistry, Cleveland State University

Dissertation Committee Member, Dr. Ashok Agarwal
Center for Reproductive Medicine, Cleveland Clinic Foundation

Dissertation Committee Member, Dr. David Anderson
Department of Chemistry, Cleveland State University

Dissertation Committee Member, Dr. Stan Duraj
Department of Chemistry, Cleveland State University

Dissertation Committee Member, Dr. John Turner
Department of Chemistry, Cleveland State University

DEDICATION

I dedicate this thesis to the memory of my father Ali Saleem Bani-Hani; my great mother Fatima Al-kofahi; my wonderful wife Hala Abdallah for her support; my daughters Rowena and Leya; and to my brothers Najeeb, Rezig and Marzoog; my sisters Zuhor, Fadwa, Fadya, Wafa' and Noor for their believing in me.

ACKNOWLEDGMENTS

I would like to acknowledge and extend my sincere and deepest gratitude to the people who made this work possible: my advisor, Dr. Mekki Bayachou, my advisor at Cleveland Clinic Dr. Ashok Agarwal, and my mentor Dr. Rakesh Sharma for all the help, guidance and motivation throughout those years; my PhD committee members, Dr. David Anderson, Dr. Stan Duraj and Dr. John Turner. I would like to thank my fellow Bayachou lab members at Cleveland State University and all my fellow Andrology lab members at Cleveland Clinic for their help and assistance. I would like to specifically thank Dr. Jean Boutros, Dr. Ling Li, Jinane Saliba, and Dr. Reda Mahfouz for their help throughout those years.

Last but not least, I would like to show my deep gratitude to Dr. Hussein Alzoubi, Mr. Abdullah Jordan, Dr. Abdullah Shatnawi, Dr. Huthaifa Al-Omari, Mr. Khaled Al-Shboul, Dr. Ziad Al-Quraan, Dr. Ahmed Tawfiq Al-Hammouri, Dr. Ahmed Abdel Nabi Al-Hammouri, and All of my colleagues at Cleveland Clinic and Cleveland State University, especially Amina Abbadi, Ghaith Al-Tawallbeh, and Bhagya Gunasekera. I wish everyone happiness in his life, and thanks again to all of you.

SEMEN QUALITY AND CHEMICAL OXIDATIVE STRESS; QUANTIFICATION
AND REMEDIATION

SALEEM A. BANI-HANI

ABSTRACT

The necessary multi-step process to prepare human semen for assisted reproduction (AR) such as *in vitro* fertilization (IVF) is known to induce oxidative stress in human spermatozoa and subsequent damage to the integrity of their cell components and, thus, to their function. Literature reports have viewed this as a potential link with the high frequency of birth defects among the IVF babies. In this work, we studied the effect of incubation and centrifugation on human sperm quality using established AR protocols. We found that *in vitro* incubation and centrifugation (20 min at 220g) of sperm generally degrade sperm quality in terms of motility and DNA oxidation. The total antioxidant capacity of the semen significantly decreased upon raising the centrifugation force from 220g to 400g. We then studied the efficacy of supplements such as L-carnitine (LC) on remediating the oxidative stress. We found that supplementation with LC counteracts the overall damage of quality brought by the multi-step method of sperm processing. We determined the LC concentration range (0.1-1.0 mg/mL) with optimum oxidative stress remediation and which enhances qualities such as motility. Among the various oxidative stress species, we particularly focused on nitric oxide as a species with known dual functional and cytotoxic properties. We have demonstrated a method to monitor its dynamic concentrations in spermatozoa. The method enabled us to establish a link between motility and nitric oxide levels in spermatozoa. The method also allowed us to

establish that the LC-induced enhancement of human sperm motility is also accompanied with a simultaneous increase in NO production.

TABLE OF CONTENTS

	Page
ABSTRACT.....	v
LIST OF TABLES.....	vi
LIST OF FIGURES.....	xi
CHAPTER	
I. INTRODUCTION AND OVERVIEW	
1.1 Assisted reproductive technologies.....	1
1.2 Human semen processing for ART.....	4
1.3 Reactive oxygen species and oxidative stress.....	4
1.4 Antioxidants role in human body.....	6
1.5 Oxidative stress, antioxidants and human sperm quality in assisted reproduction.....	7
1.6 Flow cytometry: an overview.....	9
1.7 References.....	18
II. SEMEN QUALITY DURING PREPARATION FOR ASSISTED REPRODUCTIVE TECHNOLOGIES	
2.1 Introduction.....	23
2.2 Research methods and experimental design.....	24
2.2.1 Sample collection.....	24
2.2.2 Effect of centrifugation on sperm motility and viability.....	25
2.2.3 Effect of centrifugation on total antioxidant capacity.....	28
2.2.4 Measurement of DNA oxidation by flow cytometry.....	30

2.2.5	Measurement of DNA oxidation by electrochemistry.....	32
2.2.5.1	Sperm DNA extraction.....	32
2.2.5.2	DNA film preparation.....	32
2.2.5.3	Square Wave Voltametry.....	35
2.2.6	Statistical analysis.....	35
2.3	Results.....	36
2.3.1	Effect of sperm centrifugation on sperm motility and viability.....	36
2.3.2	Effect of semen centrifugation on total antioxidant capacity of seminal plasma.....	38
2.3.3	Effect of centrifugation on sperm DNA oxidation as evaluated by flow cytometry.....	38
2.3.4	Effect of centrifugation on sperm DNA oxidation as evaluated by square wave voltammetry.....	41
2.4	Discussion.....	43
2.5	Conclusion.....	47
2.6	References.....	48

III. L-CARNITINE AS A PROTECTIVE MEDIA FOR SPERM IN ASSISTED REPRODUCTIVE TECHNOLOGIES

	Summary.....	52
3.1	Introduction.....	53
3.2	Materials and methods.....	56
3.2.1	Sample collection.....	56
3.2.2	Effect of various LC concentrations on sperm motility and viability.....	56

3.2.3	Measurement of sperm viability.....	57
3.2.4	Effect of LC on sperm motility and viability during centrifugation.....	57
3.2.5	Effect of LC on sperm motility and viability for Various sperm concentrations during sperm incubation.....	57
3.2.6	Effect of LC on sperm DNA oxidation.....	57
3.2.7	Statistical analysis.....	59
3.3	Results.....	60
3.3.1	Effect of LC on sperm motility and viability on incubation.....	60
3.3.2	Effect of LC on sperm motility and viability during Sperm centrifugation.....	60
3.3.3	Effect of LC on sperm motility and viability for various sperm concentrations.....	63
3.3.4	Effect of LC on sperm DNA oxidation.....	63
3.4	Discussion.....	66
3.5	References.....	72

IV. NITRIC OXIDE IN HUMAN MALE REPRODUCTION

	Summary.....	78
4.1	Nitric oxide, synthesis and function.....	79
4.2	NO synthases.....	81
4.3	NO and smooth muscle contraction.....	82
4.4	Nitric oxide mechanisms of action mediated by peroxinitrite.....	82
4.5	Nitric oxide and male infertility.....	90
4.6	Nitric oxide synthases in human spermatozoa.....	90

4.7	Nitric oxide and sperm motility.....	92
4.8	Nitric oxide mechanisms of action in human sperm....	94
4.9	Role of nitric oxide in human sperm capacitation and acrosomal reaction.....	95
4.10	Adverse effects of NO on sperm physiology and function.....	98
4.11	Measurement of NO in semen.....	99
4.12	Conclusion.....	101
4.13	References.....	102
V.	A NOVEL METHOD FOR MEASUREMENT OF NITRIC OXIDE PRODUCTION IN HUMAN SPERMATOZOA USING FLOW CYTOMETRY	
	Summary.....	114
5.1	Introduction.....	115
5.2	Materials and methods.....	116
5.3	Results and discussion.....	121
5.4	Conclusion.....	133
5.5	Effect of L-carnitine (LC) on NO production in human spermatozoa assessed by the DAF-2DA fluorometric method coupled with flow cytometry.....	133
5.6	References.....	136

LIST OF TABLES

Table	Page
4.1 Examples of cytosolic and membrane enzymes and proteins affected by Peroxynitrite.....	86

LIST OF FIGURES

Figure	Page
1.1 ICSI procedure.....	3
1.2 Lewis structure of oxygen and the main reactive oxygen species (ROS).....	5
1.3 Semen processing for ART such as IVF induces ROS against sperm.....	8
1.4 A representative scheme of flow cytometer system.....	11
1.5 Flow cytometry forward scatter-histogram.....	13
1.6 Flow cytometry side scatter-histogram.....	13
1.7 Flow cytometry fluorescece-histogram.....	13
1.8 Jablonski diagram (Energy statediagram).....	15
1.9 Flow cytometry Scatter plot.....	17
2.1 Sperm motility assessment.....	26
2.2 Sperm viability measurement using the Eosin-Nigrosin test.....	27
2.3 Cupric ion reducing antioxidant capacity (CUPRAC) principle.....	29
2.4 Experimental design of assessment of DNA oxidation using flow cytometry...	31
2.5 Electrochemical experiment design to explore the effect of sperm centrifugation on sperm DNA damage.....	34
2.6 Effect of semen centrifugation on sperm motility and viability.....	37
2.7 Effect of semen centrifugation on total antioxidant activity of seminal plasma as evaluated by Copper Reducing Antioxidant Capacity.....	39
2.8 Effect of centrifugation on sperm DNA oxidation as evaluated by flow cytometry.....	40
2.9 Square wave voltametry of (PDDA/DNA) ₂ films on PG electrode in pH 5.5 buffer containing 50 μM $\text{Ru}(\text{bpy})_3^{2+}$	42
3.1 L-carnitine function.....	55

3.2	Effect of various <i>in vitro</i> incubation with L-carnitine concentrations at 37°C on sperm motility and viability.....	61
3.3	Effect of centrifugation on A: sperm motility with and without L-carnitine; B: sperm viability with and without L-carnitine	62
3.4	Effect of L-carnitine (0.5 mg/mL) on A: sperm motility and B: viability following incubation with various sperm concentrations.....	64
3.5	Effect of LC on sperm DNA oxidation.....	65
4.1	Synthesis of nitric oxide.....	80
4.2	Nitric oxide actions mediated by peroxynitrite.....	87
4.3	Chemical structure of tetrahydrobiopterin (A); 7,8-dihydrobiopterin (B).....	89
4.4	Various compounds and factors increase sperm motility may be by increasing NO production.....	93
4.5	Synergistic effect of NO and progesterone.....	97
5.1	Schematic representation of fluorometric NO measurement procedure for human spermatozoa using the DAF-2DA precursor.....	122
5.2	Observed fluorescence as a function of sperm concentration with 20 µM dye Concentration.....	124
5.3	(A) Flow cytometry histograms of DAF-2T fluorescence as a function of concentrations of the DAF-2DA dye; (B) Mean areas of DAF-2T fluorescence at the various DAF-2DA doses.....	126
5.4	Effect of various DAF-2DA concentrations on sperm viability as evaluated by the Eosin-Nigrosin test.....	127
5.5	DAF-2T flow cytometry fluorescence in human sperm as a function of L-NAME concentration.....	129
5.6	DAF-2T flow cytometry fluorescence in human sperm as a function of NONOate concentration.....	130
5.7	Effect of varying concentration of NONOate on sperm viability as evaluated by the Eosin-Nigrosin test.....	132
5.8	Effect of LC on DAF-2T fluorescence in human spermatozoa.....	135

CHAPTER I

INTRODCUTION AND OVERVIEW

1.1 Assisted reproductive technologies

Assisted reproductive technologies (ART) are the artificial techniques used to facilitate the pregnancy for couples who are trying to conceive without success. The most common ones are *In vitro* fertilization (IVF), Intracytoplasmic sperm injection (ICSI) and intrauterine insemination (IUI) [1; 2].

In IVF, the fertilization process, i.e. sperm and egg fusion, occurs outside the female's body in the Petri dish. The ultimate goal of this procedure is to increase the chance of fertilization that is not achieved through the normal intercourse. In IVF, the oocyte production is first induced using hormonal therapy. After ovulation, the eggs are retrieved from the female's ovary; in the meantime, sperm, fresh or cryopreserved, from

the male are prepared and added to the female's oocytes. The fertilized egg (gamete) is then incubated at 37°C in the dark in highly sterilized conditions until the embryo is formed. After that, the embryo is transferred and placed in the center of female's uterus in order to achieve the pregnancy. In fact, several embryos are usually prepared and placed into the uterus so as to increase the success rate of the process.

The main primary steps of ovarian induction, oocyte retrieval technique, sperm preparations and embryo culture methods in IVF cycles are similar to those in ICSI cycles. Unlike IVF, in ICSI, the spermatozoa is selected based on certain criteria; spermatozoa of good motility and morphology is selected and injected artificially by the embryologist using a micro-needle inside the oocyte [3; 4] (figure 1-1). ICSI technique is mainly used to solve problems of severe male factor infertility as well as other problems that lead to failure of natural fertilization [3; 5].

In IUI, highly-motile washed sperm are injected into the female's uterus using a special plastic catheter [6]. The rationale of this method is to increase the number of highly-motile sperm at the site of fertilization, and hence increasing the probability of fertilization. IUI is used in cases where the sperm are unable, or barely able, to reach the oocyte in the female's body spontaneously. Such conditions are sexual dysfunction, asthenospermia (low sperm motility), and antagonistic cervical conditions such as increased cervical mucus thickness [6; 7; 8; 9].





Figure 1-1: ICSI procedure. (A) the selected spermatozoa is immobilized and aspirated into a microinjection pipette;(B) the oocyte is secured using a holding pipette and a microinjection pipette; (C) the trapped spermatozoa is injected gently into the oocyte. (Adapted from M.C.Schiewe I, 1996).

1.2 Human semen processing for ART


Whether the sperm is fresh or cryopreserved, semen processing is routinely performed in all ART. Contemporary processing techniques of sperm preparation for ART require prolonged incubation as well as repeated centrifugation of human semen to separate sperm from the seminal plasma at speeds of 200-400g for 10 to 20 minutes. The processing of sperm should be gentle to allow the recovery of highly motile spermatozoa that are morphologically and functionally normal. Nevertheless, reports show that sperm centrifugation may generate low sperm quality, likely, by affecting the DNA integrity of sperm [10; 11]. New findings suggest that some birth defects occur more often among infants conceived with ART [12].

1.3 Reactive oxygen species and oxidative stress



Free radicals are highly reactive molecules or atoms that contain an unpaired electron. Among free radicals in biological systems, reactive oxygen species (ROS) are the most important due to their high reactivity in the body (figure 1-2). They are associated with the onset of a variety of diseases including cancer, arteriosclerosis, cardiovascular disease, infertility, and a host of inflammatory diseases [13]. The most important ROS in the human body are superoxide radical ($O_2^{\cdot-}$), hydroxyl radical ($\cdot OH$), Nitric oxide (NO), and the non-radical derivative hydrogen peroxide (H_2O_2) [14]. Endogenous entities such as mitochondria [15], peroxisomes, lipoxygenases, NADPH oxidase and cytochrome P450 as well as exogenous factors such as ultraviolet light, ionizing radiation, chemotherapeutics inflammatory cytokines, and environmental toxins are considered as to be among major factors that generate ROS in living cells [16].



Oxygen



Hydrogen peroxide




Superoxide ion



Nitric oxide



Hydroxyl radical



Hypochlorite ion



Figure 1-2: Lewis structure of dioxygen and some reactive oxygen species (ROS).

In live cells, the disturbance in the balance between oxidants and antioxidants in the favor of the former is termed as oxidative stress. Increased levels of free radicals such as O_2^- , NO and $\cdot OH$ are capable of damaging DNA as well as the proteins and lipids, and ultimately leads to cell injury or cell death (Figure 1-3) [17; 18]. The mechanisms of the products formed by oxidative stress vary depending on the nature of free radical and its molecular target [18]. Some of oxidative products could be a rigid sign of oxidative stress and can be used as markers to assess the extent of the oxidative damage. For example, 8-oxo-guanine (8-oxodeoxyguanosine) serves as a sensitive marker to estimate the level of DNA damage caused by hydroxyl radicals [19; 20].

1.4 Antioxidants role in human body

Antioxidants are substances capable of counteracting the free radical oxidative damage in body tissues [17]. They could be dietary, such as α -tocopherol (vitamin E) and ascorbic acid (vitamin C), or synthesized in the body like glutathione [21]. Antioxidants have attracted a great deal of attention for the defense they provide against oxidative stress [17]. Several studies showed that supplementation of dietary antioxidants such as vitamin E and β -carotene is associated with decreasing the risk of many diseases, such as heart diseases [22], cancer [23], Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis [24; 25; 26].

1.5 Oxidative stress, antioxidants and human sperm quality in assisted reproduction

Oxidative stress in human spermatozoa may be due to the lack of molecular defense mechanisms and increased sources of ROS [27]. Various studies have reported the role of oxidative stress in male infertility pathogenesis [28; 29]. Male infertility problems such as asthenozoospermia and necrozoospermia may due to oxidative stress [27]. Similar to the effect of oxidative stress in other types of cell types, both sperm DNA and sperm membrane are affected negatively by oxidative stress (Figure 1-3), which consequently decreases the change of reproductive outcomes [27].

Sperm preparation for assisted reproductive technologies includes various routine steps such as incubation, cryopreservation and centrifugation. This induces oxidative stress and subsequently affects negatively the quality of sperm. The use of antioxidants to improve human sperm quality during processing for ART have been used extensively in the last 2 decades. Although there is a great evidence supporting the use of antioxidants to improve sperm quality in assisted reproduction [30; 31; 32; 33], optimal concentrations and the correct combination of the antioxidants should be cautiously standardized. In chapter III we will report on the efficacy of L-carnitine, a potent antioxidant, in improving sperm quality during centrifugation and prolonged incubation *in vitro* for use in ART protocols.




Figure 1-3: semen processing for ART such as IVF induces ROS against sperm.

1.6 Flow cytometry: an overview

In our experimental work, we intensively used flow cytometry to explore the effect of various factors affecting human spermatozoa in assisted reproduction. In addition, we standardized a method to detect the intracellular nitric oxide in human spermatozoa using flow cytometry. Therefore, there is a need to provide an overview of the main principles and fundamentals of this technique.

Flow cytometry is a technique for the analysis of various parameters of individual cells. It is used in a wide range of applications to measure different cell characteristics such as size, fluorescence, shape, viability, intracellular pH, intracellular redox state, and others. Besides, a variety of cell components can be determined using flow cytometry such as DNA, RNA, protein, Lipid, antigens, surface receptors, calcium and others. The flow cytometer achieves this analysis via passing a representative population of cells from the sample through a laser beam and capturing the emerged light from each individual cell as it passes through [34].

The main components of the flow cytometer are: (a) the fluidics (hydrodynamic focusing system). This system allows the chosen cell population to pass the laser beam one at a time (Figure 1-4). The intersection point between the laser beam and the cell is termed as interrogation point. (b) The laser, which represent the light source in the flow cytometer. (c) The optics, their main function is to gather the light from the cells after crossing the laser beam. (d) The detectors; they receive the accumulated light from the optics. (e) The computer system and other electronics, which convert the detector's signals into digital data and analyzes the data [34; 35; 36].

At interrogation point, when the cell crosses the laser beam, two types of light scatter are formed: (a) forward scatter (FSC), which is the light, scattered in the forward direction, and (b) side scatter (SSC), which is the light scattered on the sides (Figure 1-4). In general, the forward scatter provides information about the size of the cell, whereas the side scatter provides information about characteristics within the cell such as granularity inside the cytoplasm, shape of nucleus, and membrane irregularity [37; 38; 39].




Figure 1-4: a representative scheme of flow cytometer system (see the text for details).

FSC is proportional to the size of cells that are passing the laser [38]. The bigger the cell size, the higher the FSC intensity. The detector converts the forward scattered light into a voltage pulse. Hence, the size of cell is finally proportional to the magnitude of the voltage pulse. The plot of cell count versus FSC is called forward scatter histogram (FSCH) (Figure 1-5). In this histogram, the larger cells appear on the right.

SSC is the light scattered at higher angles compared to the FSC. The main determinants of the angles of the scattered light are the structural complexity of the cell, as we have mentioned above. A separate detector in the flow cytometer, other than one used for FSC, is used to gather the side scattered light; it is usually placed 90° from the laser beam path. The plot of cell count versus SCC is also a one-dimensional histogram named side scatter histogram (SSCH) (Figure 1-6). It is similar to the FSCH, but it provides information about the cells characteristics under investigation.




Figure 1-5: Flow cytometry forward scatter-histogram. Cells are sorted according to their sizes




Figure 1-6: Flow cytometry side scatter-histogram. Cells are sorted according to their complexity fluorescence.




Figure 1-7: Flow cytometry fluorescence-histogram.

Fluorescence is another phenomenon used as handle to quantify a given property within cells. At the fundamental levels, it is a term used to describe light emission, mainly visible, from an excited fluorophore. The emission occurs when the fluorophore returns to its ground state from an excited state. Figure 1-8 is the Jablonski diagram (energy state diagram) of a fluorophore excitation and emission. The emitted light from a fluorophore occurs at a higher wavelength. For example, the fluorophore fluorescein isothiocyanate (FITC) that we used in this work to detect DNA oxidation can be excited with the blue light at 494 nm; while it emits the green light at 518 nm.

Fluorescence emitted from cells labeled with a fluorophore travels in the same path in the flow cytometer as the side scatter light. This path contains a number of light filters so as to deliver each range of wavelengths to the proper detector. Same as data collection in the cases of FSC and SSC, the light directed to the detector is translated into a voltage pulse. The more the light emitted from the cell, the higher the voltage pulse translated. The plot of intensity of fluorescence versus cell count is termed as flow cytometry fluorescence-histogram, which provides information about cell fluorescence that is increasing in the right direction (Figure 1-7).




Figure 1-8: Jablonski diagram (Energy state diagram) - fluorophore excitation and emission. The fluorophore emission usually occurs at a lower wavelength; for example, the fluorophore Fluorescein isothiocyanate (FITC) can be excited at 494 nm, while it emits the light at 518 nm.

Forward scatter histogram is not necessarily showing the complexity of the cells. In other words, the peak population that appears in the FSCH could be a representative of multiple population. This problem can be resolved by looking at the data in the second dimension using a two-dimensional scatter plot or dot plot (Figure 1-9). In this plot (SSC versus FSC), we can look at both cellular parameters together: cell size and cell complexity.

Another important problem in flow cytometry is to analyze only the cells of interest. In many cases, cell fragments such as, small immature cells and debris, and sometimes bigger cells that are not of interest interfere with the measurement. This problem can be solved by establishing a threshold, or discriminator before start analyzing the cells. This will exclude interferences for compounds that are not of interest.




Figure 1-9: flow cytometry Scatter or dot plot. Cells are analyzed for both complexity and size.

1.7 References

- [1]M.C. Schiewe, An effective, simplified, and practical approach to intracytoplasmic sperm injection at multiple IVF centers. *J Assist Reprod Genet* 13 (1996) 238-45.
- [2]D. Royere, [Intrauterine insemination: state-of-the-art in humans]. *Gynecol Obstet Fertil* 32 (2004) 873-9.
- [3]R. Orvieto, Z. Ben-Rafael, J. Ashkenazi, R. Yoeli, B. Messing, T. Perri, Y. Shalev, and I. Bar-Hava, Outcome of pregnancies derived from assisted reproductive technologies: IVF versus ICSI. *J Assist Reprod Genet* 17 (2000) 385-7.
- [4]C.B. Coulam, M.S. Opsahl, R.J. Sherins, L.P. Thorsell, A. Dorfmann, L. Krysa, E. Fugger, and J.D. Schulman, Comparisons of pregnancy loss patterns after intracytoplasmic sperm injection and other assisted reproductive technologies. *Fertil Steril* 65 (1996) 1157-62.
- [5]O. Harari, H. Bourne, M. McDonald, N. Richings, A.L. Speirs, W.I. Johnston, and H.W. Baker, Intracytoplasmic sperm injection: a major advance in the management of severe male subfertility. *Fertil Steril* 64 (1995) 360-8.
- [6]G.I. Meniru, *Cambridge guide to infertility management and assisted reproduction*, Cambridge University Press, Cambridge ; New York, 2001.
- [7]E. Simsek, B. Haydardedeoglu, S.O. Hacivelioglu, T. Cok, A. Parlakgumus, and T. Bagis, Effect of cervical mucus aspiration before intrauterine insemination. *Int J Gynaecol Obstet* 103 (2008) 136-9.

- [8]V.I. Grishchenko, S.M. Nabil, and F.V. Dakhno, [Intrauterine insemination as a method to surmount the immunological barrier of the cervical mucus]. *Akush Ginekol (Mosk)* (1982) 55-7.
- [9]X.W. Nie, Y. Qian, C.Y. Liu, and Y. Tan, [Semen cryopreservation applied to intrauterine insemination cycles for oligospermia and asthenospermia in infertile men]. *Zhonghua Nan Ke Xue* 16 232-5.
- [10]A. Agarwal, and S.S. Allamaneni, The effect of sperm DNA damage on assisted reproduction outcomes. A review. *Minerva Ginecol* 56 (2004) 235-45.
- [11]E.S. Sills, K.M. Wittkowski, M.J. Tucker, M. Perloe, C.R. Kaplan, and G.D. Palermo, Comparison of centrifugation- and noncentrifugation-based techniques for recovery of motile human sperm in assisted reproduction. *Arch Androl* 48 (2002) 141-5.
- [12]J. Reefhuis, M.A. Honein, L.A. Schieve, A. Correa, C.A. Hobbs, and S.A. Rasmussen, Assisted reproductive technology and major structural birth defects in the United States. *Hum Reprod* 24 (2009) 360-6.
- [13]B. Halliwell, Oxidative stress, nutrition and health. Experimental strategies for optimization of nutritional antioxidant intake in humans. *Free Radic Res* 25 (1996) 57-74.
- [14]B. Halliwell, P.J. Evans, H. Kaur, and O. Aruoma, Free radicals, tissue injury, and human disease - a potential for therapeutic use of antioxidants, Raven Press, New York, NY., 1994.
- [15]S. Raha, and B.H. Robinson, Mitochondria, oxygen free radicals, and apoptosis. *Am J Med Genet* 106 (2001) 62-70.

- [16]T. Finkel, and N.J. Holbrook, Oxidants, oxidative stress and the biology of ageing. Nature 408 (2000) 239-47.
- [17]H. Sies, Oxidative stress: oxidants and antioxidants. Exp Physiol 82 (1997) 291-5.
- [18]P. Therond, [Oxidative stress and damages to biomolecules (lipids, proteins, DNA)]. Ann Pharm Fr 64 (2006) 383-9.
- [19]M.K. Shigenaga, C.J. Gimeno, and B.N. Ames, Urinary 8-hydroxy-2'-deoxyguanosine as a biological marker of in vivo oxidative DNA damage. Proc Natl Acad Sci U S A 86 (1989) 9697-701.
- [20]K.C. Cundy, R. Kohen, and B.N. Ames, Determination of 8-hydroxydeoxyguanosine in human urine: a possible assay for in vivo oxidative DNA damage. Basic Life Sci 49 (1988) 479-82.
- [21]VertuaniS, AngustiA, and ManferdiniS, The antioxidants and pro-antioxidants network: an overview. Curr Pharm Des 10 (2004) 1677-94.
- [22]E.B. Rimm, M.J. Stampfer, A. Ascherio, E. Giovannucci, G.A. Colditz, and W.C. Willett, Vitamin E consumption and the risk of coronary heart disease in men. N Engl J Med 328 (1993) 1450-6.
- [23]A.J. McMichael, Food, nutrition, physical activity and cancer prevention. Authoritative report from World Cancer Research Fund provides global update. Public Health Nutr 11 (2008) 762-3.
- [24]V. Di Matteo, and E. Esposito, Biochemical and therapeutic effects of antioxidants in the treatment of Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis. Curr Drug Targets CNS Neurol Disord 2 (2003) 95-107.

- [25]A.V. Rao, and B. Balachandran, Role of oxidative stress and antioxidants in neurodegenerative diseases. *Nutr Neurosci* 5 (2002) 291-309.
- [26]S.A. Stanner, J. Hughes, C.N. Kelly, and J. Buttriss, A review of the epidemiological evidence for the 'antioxidant hypothesis'. *Public Health Nutr* 7 (2004) 407-22.
- [27]A. Agarwal, T.M. Said, M.A. Bedaiwy, J. Banerjee, and J.G. Alvarez, Oxidative stress in an assisted reproductive techniques setting. *Fertil Steril* 86 (2006) 503-12.
- [28]S.C. Sikka, Role of oxidative stress and antioxidants in andrology and assisted reproductive technology. *J Androl* 25 (2004) 5-18.
- [29]A. Agarwal, and R.A. Saleh, Role of oxidants in male infertility: rationale, significance, and treatment. *Urol Clin North Am* 29 (2002) 817-27.
- [30]H.A. Askari, J.H. Check, N. Peymer, and A. Bollendorf, Effect of natural antioxidants tocopherol and ascorbic acids in maintenance of sperm activity during freeze-thaw process. *Arch Androl* 33 (1994) 11-5.
- [31]A.P. Brennan, and C.A. Holden, Pentoxifylline-supplemented cryoprotectant improves human sperm motility after cryopreservation. *Hum Reprod* 10 (1995) 2308-12.
- [32]P. Stanic, Z. Sonicki, and E. Suchanek, Effect of pentoxifylline on motility and membrane integrity of cryopreserved human spermatozoa. *Int J Androl* 25 (2002) 186-90.
- [33]F.J. Pena, A. Johannisson, M. Wallgren, and H. Rodriguez Martinez, Antioxidant supplementation in vitro improves boar sperm motility and mitochondrial

membrane potential after cryopreservation of different fractions of the ejaculate.
Anim Reprod Sci 78 (2003) 85-98.

[34]M. Rieseberg, C. Kasper, K.F. Reardon, and T. Scheper, Flow cytometry in biotechnology. Appl Microbiol Biotechnol 56 (2001) 350-60.

[35]R. Nunez, Flow cytometry: principles and instrumentation. Curr Issues Mol Biol 3 (2001) 39-45.

[36]G.V. Chapman, Instrumentation for flow cytometry. J Immunol Methods 243 (2000) 3-12.

[37]J.P. McCoy, Jr., Basic principles of flow cytometry. Hematol Oncol Clin North Am 16 (2002) 229-43.

[38]A.L. Givan, Principles of flow cytometry: an overview. Methods Cell Biol 63 (2001) 19-50.

[39]M.J. Jaroszeski, and G. Radcliff, Fundamentals of flow cytometry. Mol Biotechnol 11 (1999) 37-53.

CHAPTER II

SEMEN QUALITY DURING PREPARATION FOR ASSISTED REPRODUCTIVE TECHNOLOGIES

2.1 Introduction

Sperm centrifugation, required for sperm washing, is a routine technique during sperm processing for assisted reproduction technologies (ART). Reports from our laboratory showed a significant increase in reactive oxygen species (ROS) during sperm centrifugation. Studies have shown a decrease in fertilizing capacity of human spermatozoa, and an increase in the birth defects among the ART babies compared to the normal babies. Here, we examine the effect of centrifugation on total antioxidant capacity (TAC) of human semen on one hand, and on DNA oxidation, motility, and viability of sperm, on the other. We measured sperm motility, viability TAC and sperm DNA oxidative damage after centrifugation of semen samples for 20 min at 220g. DNA oxidative damage in sperm was measured using both electrochemical and flow cytometry methods. Centrifugation significantly decreased sperm motility. The total antioxidant

capacity of the semen significantly decreased upon raising the centrifugation force from 220g to 400g. Concomitantly, the baseline of DNA oxidative damage in sperm increased after centrifugation. Together, sperm centrifugation during processing for ART increases oxidative stress in sperm, which negatively affects their quality. This could be one factor behind the negative reproductive outcomes.

2.2 Research methods and experimental design

In this chapter we aimed to examine the effect of centrifugation on sperm quality according to the assisted reproduction protocols. We used DNA oxidation, motility and viability of sperm, as well as total antioxidant capacity (TAC) of semen as measurement criteria to study this effect. The oxidative DNA damage in sperm was evaluated using both electrochemical and flow cytometry techniques. We also adapted an assay based on cupric reducing antioxidant capacity (CUPRAC) to measure the TAC in human semen [1].

2.2.1 Sample collection

The present study was approved by the institutional review board of Cleveland Clinic Hospital. Forty normospermic men were chosen and analyzed on the basis of normal semen analysis according to the World Health Organization (WHO, 1999) guidelines [2]. The effect of centrifugation on TAC of semen as well as DNA oxidation, motility and viability of sperm were examined. Samples were centrifuged at 1600 rpm (220g) for 20 minutes similar to sperm preparation protocols used in ART. Samples were collected by masturbation after a minimum of 48 to 72 hours of sexual abstinence. Following

liquefaction at 37 °C for 20 minutes, semen specimens were evaluated for volume, sperm concentration, total cell count, motility, and morphology.

2.2.2 Effect of centrifugation on sperm motility and viability

Two equal aliquots (5×10^6 sperm) were prepared from each of the sixteen normospermic semen samples, and mixed with human tubal fluid (HTF; 1:1 vol. / vol.). The first aliquot was centrifuged at 1600 rpm at 25 °C; while the second aliquot (control) was just incubated at the same temperature. Sperm motility and viability was measured after 20 min. A total amount of 5 μ L of the sample was used for evaluation of concentration and motility using a micro-cell slide chamber (Figure 2-1) (Conception Technologies, San Diego, CA).

Sperm viability was measured by the Eosin-Nigrosin staining method (Figure 2-2). To one drop of liquefied semen placed on a Boerner slide, two drops of 1% aqueous Eosin was added and mixed with a wooden stirrer for 15 seconds. Then, three drops of 10% aqueous Nigrosin were added and mixed well. Thin smear was made by pipetting 10 μ L onto a slide and allowed to air dry. Slides were mounted with a coverslip using Accu-mount media. Measurements were performed on two hundred sperm counted on each slide in duplicate sets using an X100 objective. Percentage of viable (unstained) sperm and non-viable (stained) were calculated.

Motility: (normal: >50%)

$$\text{Motility (\%)} = \frac{\text{Motile sperm} \times 100\%}{\text{Total sperm}}$$

If we have one motile spermatozoa
In the first two rows, then the motility:
 $(1/7) (100\%) = 14.3\%$





Figure 2-1: Sperm motility assessment. 5µL of semen sample is used to assess motility using a disposable micro-cell slide chamber (an example is shown in the figure).




Figure 2-2: Sperm viability measurement using the Eosin-Nigrosin test. Stained sperm is considered viable, while unstained one (white) is considered non-viable.

2.2.3 Effect of centrifugation on total antioxidant capacity

Semen samples (n=10) from healthy donors, were divided into two aliquots; each contained 0.5 mL of semen. We centrifuged the first aliquot at 220g for 20 min at 25 °C, while the second aliquot (control) was incubated for 20 min at the same temperature. Semen samples (n=10), in another experimental set, were subjected to the same treatment as described above, but centrifuged at a higher centrifugal force (400g).

We adapted the cupric reducing antioxidant capacity (CUPRAC) method to measure the total antioxidant capacity in semen [1]. Briefly, 1 mL of working solution (CuCl₂ solution; 0.02 M), neocuproine alcoholic solution (0.0075 M) and 1-mL ammonium acetate (NH₄Ac) buffer solution (1:1:1 vol./vol./vol.) was mixed in a centrifugation tube with 12.5 µL of seminal plasma. All reaction tubes were centrifuged 3 min at 4000 rpm, and the entire supernatant was collected for use in absorbance measurements. The absorbance of the colored complex formed was measured against the reagent blank at 450 nm after 30 min incubation at room temperature (Figure 2-3) [3].




Figure 2-3: Cupric ion reducing antioxidant capacity (CUPRAC) principle. The colored complex (Neocuproine-Cu⁺¹) is formed after reducing cupric ion by semen antioxidants. The intensity of the color is proportional to the total antioxidant activity in semen.

2.2.4 Measurement of DNA oxidation by Flow cytometry

Semen samples (n=5) from healthy donors were divided into 2 aliquots (5×10^6 sperm) and centrifuged at 1600 rpm for 20 min. at 25°C. Controls were incubated for 20 min. at 25°C. Oxidative damage to DNA was evaluated using the flow cytometric OxiDNA assay kit (Calbiochem, San Diego). The assay is based on fluorescent protein that binds to oxidized moieties on DNA (Figure 2-4). Briefly, both treated sperm samples and controls were washed in phosphate saline buffer, resuspended in 1% paraformaldehyde at a concentration of $2-4 \times 10^6$ cell/ mL, and placed on ice for 15 to 30 minutes. Sperm were again washed and resuspended in 70% ice-cold ethanol by centrifugation at 300g for 5 minutes as per kit instructions. The ethanol supernatant was removed, and the sperm pellets were washed twice in the wash buffer and resuspended in 100 μ L of the staining solution for 1 hour at room temperature in dark. The staining solution contained FITC-labeled protein conjugate in distilled water. All cells were further washed using rinse buffer, resuspended in 250 μ L and incubated for 30 minutes in the dark on ice for flow cytometry measurements.

Data acquisition was performed within 30 min on a flow cytometer equipped with a 515-nm argon laser as a light source (FACScan; Becton Dickinson, San Jose, CA). 10,000 cells were examined for each assay at a flow rate of 100 cells/ second. The FITC fluorescence was measured and data analysis was performed using FlowJo v4.4.4 software (Tree Star Inc., Ashland, OR).




Figure 2-4: Experimental design of assessment of DNA oxidation using flow cytometry.

2.2.5 Measurement of DNA oxidation by Electrochemistry

Semen samples (n=5) from healthy donors were divided into 2 aliquots (sample and control). Each aliquot (5×10^6 Sperm) was exposed to 20 min centrifugation at 1600 rpm under 25 °C. Controls of the same sperm concentration were incubated for 20 min at 25°C without centrifugation.

2.2.5.1 Sperm DNA Extraction

Cell lysis buffer (0.5 mL) and 60 μ L of proteinase K were added to both samples and controls and incubated overnight at 55 °C. A 0.5 mL aliquot of chloroform: phenol mixture (1:1 vol./ vol.) was added and centrifuged for 5 min at 6000 rpm. The supernatant (aqueous layer) was collected, and 0.5 mL of chloroform was added, vortexed and centrifuged. The top layer was collected and mixed with 0.5 mL isopropyl alcohol to salt-out the white DNA pellet. The precipitated DNA was collected by centrifugation and the supernatant was gently decanted. Finally, the DNA was dissolved in 200 μ L Tris-EDTA buffer and kept at 4°C for future use.

2.2.5.2 DNA film preparation

DNA films were prepared by layer-by-layer electrostatic assembly method [4]. The method is simply based on adsorbing thin layers from two polyelectrolytes with opposite charge (in this case: poly (diallyldimethylammonium chloride) (PDDA) and the DNA sample under investigation). Disks of ordinary basal plane pyrolytic graphite (PG) were abraded with 400 grit silicon carbide (SiC) paper and sonicated in water for 1 min, then rinsed with water and dried. The graphitic surfaces of these electrodes are then

oxidized in 2.5% $\text{K}_2\text{Cr}_2\text{O}_7$ /10% HNO_3 by scanning the potential once from 1.5V to 1.75V versus Ag/AgCl to form negative carboxylate groups on the surface [5]. Layers were formed by placing a drop of each adsorbate solution on the PG electrode for 15-20 min. The electrode was then rinsed with water to remove weakly adsorbed DNA and dried between adsorption steps. The times for steady-state adsorption, volume and concentration of each adsorbate solution were: (1) 15 min, 25 μL of 2 mg mL^{-1} poly(diallyldimethylammonium chloride) (PDDA) in 50 mM NaCl; (2) 20 min, 25 μL of 0.1-0.2 mg mL^{-1} DNA in 5 mM pH 7.1 Tris buffer and 0.50 M NaCl. Final DNA film architecture was denoted as the order of layer adsorption, (PDDA/DNA)₂. Alternate adsorption cycles were repeated until the desired number of layers was made. The number of layers of DNA/PDDA was optimized for the best time-sensitivity ratio. In this work films with 2 layers of DNA/PDDA ([DNA/PDDA]₂ films) were used (Figure 2-5).




Figure 2-5: Electrochemical experiment design to explore the effect of sperm centrifugation on sperm DNA damage. Final DNA film architecture is denoted as the order of layer adsorption; (PDDA/DNA)₂.

2.2.5.3 Square Wave Voltametry

Square Wave Voltametry (SWV) was performed using CH electrochemical work station (CH Instruments 440A). The electrochemical cell employed an Ag/AgCl reference electrode, a platinum wire counter, and a film-coated PG disk as the working electrode. Unless otherwise noted, the SWV conditions were 4 mV step height, 25 mV pulse height, and 5 Hz. The electrolyte solution was 10 mM acetate buffer, pH 5.5, containing 50 mM NaCl and 50 μ M Tris(2,2'-bipyridine)ruthenium(II), (Ru (bpy)₃²⁺) as the reporting redox probe.

2.2.6 Statistical analysis

All values were reported as mean \pm standard deviation. Difference was considered significant at $P < 0.05$ against the control group. Statistical analyses were operated using one-way analysis of variance (ANOVA) followed by an unpaired Student's *t*-test using the SPSS/PC computer software (SPSS 10.0.7, SPSS Inc.).

2.3 Results

2.3.1 Effect of sperm centrifugation on sperm motility and viability

Figure 2-6 shows the effect of semen centrifugation (at 220g, 20 min) on sperm motility and viability. Figure 1A shows the mean values \pm standard deviations of sperm motility for both samples and controls (i.e. without centrifugation). Sperm motility was significantly different between centrifuged and uncentrifuged samples (24.0 ± 12.8 vs. $33.3 \pm 14.2\%$) ($P < 0.05$; t -test). In contrast, no significant change was seen in sperm viability between centrifuged and uncentrifuged samples (34.2 ± 12.6 vs. $39.1 \pm 9.7\%$, respectively; Fig.1B).




Figure 2-6: Effect of semen centrifugation on sperm motility (A) and viability (B). Sperm samples centrifuged 20 min at 1600 rpm; control assayed without centrifugation. Data are mean values \pm standard deviations obtained from analysis of 16 semen samples.

2.3.2 Effect of semen centrifugation on total antioxidant capacity of seminal plasma

Figure 2-7 shows the effect of semen centrifugation on the total antioxidant activity (TAC) in seminal plasma as measured by the CUPRAC assay. The absorbance measured at 450 nm is directly proportional with the TAC of semen sample. We saw an insignificant difference ($P > 0.05$, t -test) in average absorbance (0.67 ± 0.08 vs. 0.71 ± 0.09) at 450nm, and thus in total antioxidant capacity of seminal plasma in semen samples (exposed to 20 min centrifugation at 220g) and the control, Figure 2A. On the other hand, we observed a significant decrease in average absorbance (0.78 ± 0.1 vs 0.62 ± 0.03) ($P < 0.05$; t -test) and thus in antioxidant activity in semen samples centrifuged at forces between 220g and 400g under the same conditions, Figure 2B.

2.3.3 Effect of centrifugation on sperm DNA oxidation as evaluated by flow cytometry

The effect of centrifugation on sperm DNA oxidation as evaluated by flow cytometry is shown in Figure 2-8. The flow cytometry analysis of FITC-labeled sperm showed that the intensity of the FITC fluorescence increased significantly ($P < 0.5$, t -test) upon centrifugation of sperm at 220g for 20 min. The intensity of fluorescence, represented as mean area of FITC-fluorescence \pm standard deviation, for uncentrifuged and centrifuged sperm were 58.5 ± 3.4 vs 210.5 ± 14.8 , respectively. This result shows that the onset of DNA oxidative stress is significant at forces as low as 220g.




Figure 2-7: Effect of semen centrifugation on total antioxidant activity of seminal plasma as evaluated by Copper Reducing Antioxidant Capacity (CUPRAC). (A) Sperm centrifuged 20 min at 220g; (B) sperm centrifuged 20 min at 400g. Controls assayed without centrifugation. The absorbance at 450 nm is proportional to the total antioxidant activity in semen. P -value < 0.05 is considered statistically significant




Figure 2-8: Effect of centrifugation on sperm DNA oxidation as evaluated by flow cytometry. Sperm centrifuged 20 minutes at 1600 rpm; control assays without centrifugation. The X-axis represents the intensity of FITC fluorescence. Data are representative of 4 independent experiments.

2.3.4 Effect of centrifugation on sperm DNA oxidation as evaluated by square wave voltammetry

DNA damage can also be detected through the -now standard- electrocatalytic oxidation of guanine bases by ruthenium tris-bipyridine as introduced by Thorp et al.[6; 7; 8]. Damaged DNA, with open double helices or broken strands, is expected to provide the reporter ruthenium redox probe higher access to guanine sites, and thus would generate oxidation currents commensurate with the extent of DNA damage. This criterion (i.e. large electrocatalytic currents equivalent to large DNA damage) is used here as such without explanation of the fundamental concepts, which can be found in numerous original papers using this method [6; 7; 8]. All DNA damage measurements are carried out in 10 mM acetate buffer, pH 5.5 solutions containing 50 mM NaCl and 50 μ M Ru(bpy)₃²⁺ as the reporter probe. All experiments are triplicates and data points reported are means of 4 independent experiments.

Figure 2-9 shows the voltametric response of DNA modified electrode for both centrifuged and uncentrifuged sperm for the desired period of time. As seen in the figure, an increase in the oxidation current by 17.8% ($I_{\text{centrifuged sperm}}/I_{\text{uncentrifuged sperm}} = 1.178$), and thus in sperm DNA oxidative damage, was seen in sperm samples subjected to centrifugation (220g) compared with the controls (without centrifugation).




Figure 2-9: Square wave voltammetry of (PDDA/DNA)₂ films on PG electrode in pH 5.5 buffer containing 50 μM Ru(bpy)₃²⁺. DNA samples were extracted from sperm exposed to 20 min centrifugation at 220g; control samples were DNA extracted from uncentrifuged sperm. Data are representative of 4 independent experiments.

2.4 Discussion

It has been demonstrated that centrifugation has an adverse effect on sperm recovery rates [9]. Centrifugation < 330 g for about 15 min had insignificant effect on sperm motility. However a significant decrease in sperm motility was observed with centrifugation above 580g at the same length of time [10]. Our study shows that centrifugation forces as low as 220g for 20 min result in a significant decrease in sperm motility. We applied a longer centrifugation period (20 min) since this is an IVF protocol used in many andrology labs. A recent study reported by Lampiao *et al.* (2010) found that sperm centrifugation from 10 to 30 min at 400g also decreased sperm motility [11]. Our results are in agreement with these findings.

In an earlier study, we reported that centrifugation of motile sperm induces the generation of reactive oxygen species (ROS) [12]. In sperm washing procedures, depriving the sperm of seminal plasma –their antioxidant reservoir-, is a main contributing factor behind increased levels of ROS [13]. Induced ROS, mainly hydroxyl radicals, were shown to affect sperm function and motility [14]. The mechanism by which the elevated levels of ROS decreases sperm function and motility could be via increasing sperm's lipid peroxidation as well as direct oxidation of sperm DNA bases [14].

The effect of centrifugation on sperm viability was assessed using the one-step Eosin-Nigrosin test [15]. This test is principally based the membrane permeability for the supravital Eosin stain. Membrane injury allows the Eosin to enter inside the spermatozoa and stain it. The intensity of the color is proportional with the magnitude of the damage

[15]. The unstained spermatozoa (white-colored under the microscope) were considered viable, while the stained spermatozoa, regardless the intensity of the stain color, considered non-viable. Our results from this experiment show that viability is not affected by centrifugation (at least for a 20 min centrifugation period at 220g).

Namik *et al.* (2000), in their study measured sperm viability after cryopreservation-thawing cycles, using annexin V-Cy3 and eosin-Y staining. They noticed a significant variance between the percentages of sperm accepted as viable between both tests, in which, Eosin-Y staining test showed lower percent of live sperm compared to the annexin V-Cy3 test [16]. This variation is attributed to the different mechanisms through which these tests function. In the present study, we assessed the viability of sperm using the Eosin stain test that is based on the membrane permeability to Eosin. The mechanism by which this test categorizes the viable versus non-viable sperm could be a contributing factor behind the insignificant effect of centrifugation on sperm viability. Using other reliable viability tests may be needed for conclusive results on the effect of centrifugation on viability.

As a criterion of oxidative damage exposure, we have adapted the CUPRAC method, developed by Apak *et al.* (2008), to measure the total antioxidant capacity (TAC) in human semen. We used this assay because the reaction is carried out at nearly physiological conditions. In addition, the CUPRAC method is straightforward and reliable [1].

At 220g centrifugation force, our results did not show a significant effect of centrifugation on TAC of semen samples compared to controls. However, we observed a significant decrease in semen TAC at 400g. It was reported that repeated sperm


centrifugation, the key step in sperm washing during processing for ART, leads to an excessive formation of ROS [12]. Along the same lines, a higher centrifugation force may induce more ROS formation [11]. Accordingly, the increased levels of ROS explain the decreased TAC that we observed in seminal plasma of sperm subjected to centrifugation at 400g.

Our flow cytometry experiment was designed to study the effect of centrifugation (20 min at 220g) on sperm DNA oxidation. Sperm samples were fixed, permabilized and the DNA oxidation was determined via specific detection of 8-oxoguanines, biomarkers of DNA oxidative damage, using FITC-labeled conjugate.

The results from this experiment showed an increase in the baseline of sperm DNA oxidation in the centrifuged sperm compared to the control sample. It is instructive to compare our findings to results obtained by other methods. In this regard, G.D. Smith *et al.* reported that sperm centrifuged during washing steps with HTF show an increase in the DNA fragmentation using sperm chromatin dispersion assay (SCD) [17]. Our findings are in line with these findings, and may indicate that the increase in DNA damage that we observe may be the result of an increase in DNA breaks.

In live cells, including sperm, the presence of metal ions, such as ferrous and cupric ions, and hydrogen peroxide (H_2O_2) is known to produce hydroxyl radicals through what is termed as the Fenton reaction [18; 19]. Hydroxyl radicals are highly reactive oxygen species that induce lesions in DNA bases and deoxyribose sugars in the double helix backbone, which leads to unwinding and DNA fragmentation and breaks [20].

In the electrochemical experiment, the $\text{Ru}(\text{bpy})_3^{2+}$ redox probe reports on exposed guanine bases in DNA [21]. The $\text{Ru}(\text{bpy})_3^{2+}$ is oxidized at the electrode surface to $\text{Ru}(\text{bpy})_3^{3+}$. In the presence of exposed guanine bases, the oxidized ruthenium probe is reduced back to Ru (II) state (eq.1), which is again reoxidized at the electrode. This process produce a catalytic current proportional to the number of exposed guanine bases [22]. The double-stranded DNA shields guanine moieties from effective contact with $\text{Ru}(\text{bpy})_3^{3+}$. Guanine becomes more available after chemical damage occurs in the DNA this provides more exposure, and thus, access to the resulting catalyst [23].



These results show an increase in the baseline of sperm DNA damage after centrifugation. This is confirms our results obtained with the flow cytometry method.

2.5 Conclusion

In summary, semen centrifugation during sperm preparation for ART may negatively affect sperm quality, via increasing baseline of sperm DNA oxidation and decreasing seminal TAC and sperm motility. This might be one factor behind the negative reproductive outcomes such as the increased birth defects among the IVF babies compared to the normal babies.

2.6 References

- [1]R. Apak, K. Guclu, M. Ozyurek, B. Bektas Oglu, and M. Bener, Cupric ion reducing antioxidant capacity assay for food antioxidants: vitamins, polyphenolics, and flavonoids in food extracts. *Methods Mol Biol* 477 (2008) 163-93.
- [2]W.H. Organization, WHO Laboratory Manual for the Examination of Human Semen and Sperm–Cervical Mucus Interaction. Cambridge University Press, Cambridge. (1999).
- [3]R. Apak, K. Guclu, M. Ozyurek, and S.E. Karademir, Novel total antioxidant capacity index for dietary polyphenols and vitamins C and E, using their cupric ion reducing capability in the presence of neocuproine: CUPRAC method. *J Agric Food Chem* 52 (2004) 7970-81.
- [4]P. He, and M. Bayachou, Layer-by-layer fabrication and characterization of DNA-wrapped single-walled carbon nanotube particles. *Langmuir* 21 (2005) 6086-92.
- [5]W.F. Blakely, A.F. Fuciarelli, B.J. Wegher, and M. Dizdaroglu, Hydrogen peroxide-induced base damage in deoxyribonucleic acid. *Radiation Research* 121 (1990) 338-343.
- [6]H.H. Thorp, Cutting out the middleman: DNA biosensors based on electrochemical oxidation. *Trends in Biotechnology* 16 (1998) 117-121.
- [7]B.T. Farrer, and H.H. Thorp, Redox Pathways in DNA Oxidation: Kinetic Studies of Guanine and Sugar Oxidation by Para-Substituted Derivatives of Oxoruthenium(IV). *Inorganic Chemistry* 39 (2000) 44-49.

- [8]M.F. Sistare, S.J. Codden, G. Heimlich, and H.H. Thorp, Effects of Base Stacking on Guanine Electron Transfer: Rate Constants for G and GG Sequences of Oligonucleotides from Catalytic Electrochemistry. *Journal of the American Chemical Society* 122 (2000) 4742-4749.
- [9]C. Matas, G. Decuadro, S. Martinez-Miro, and J. Gadea, Evaluation of a cushioned method for centrifugation and processing for freezing boar semen. *Theriogenology* 67 (2007) 1087-91.
- [10]A. Makler, and P. Jakobi, Effects of shaking and centrifugation on human sperm motility. *Arch Androl* 7 (1981) 21-6.
- [11]F. Lampiao, H.S. 1, and S.d. Plessis², Effects of Sperm Processing Techniques Involving Centrifugation on Nitric Oxide, Reactive Oxygen Species Generation and Sperm Function. *The Open Andrology Journal*, 2 (2010) 1-5.
- [12]A. Agarwal, I. Ikemoto, and K.R. Loughlin, Effect of sperm washing on levels of reactive oxygen species in semen. *Arch Androl* 33 (1994) 157-62.
- [13]R.J. Aitken, D.W. Buckingham, and K.M. West, Reactive oxygen species and human spermatozoa: analysis of the cellular mechanisms involved in luminol- and lucigenin-dependent chemiluminescence. *J Cell Physiol* 151 (1992) 466-77.
- [14]C.S. Chen, H.T. Chao, R.L. Pan, and Y.H. Wei, Hydroxyl radical-induced decline in motility and increase in lipid peroxidation and DNA modification in human sperm. *Biochem Mol Biol Int* 43 (1997) 291-303.

- [15]L. Bjorndahl, I. Soderlund, and U. Kvist, Evaluation of the one-step eosin-nigrosin staining technique for human sperm vitality assessment. *Hum Reprod* 18 (2003) 813-6.
- [16]N.K. Duru, M. Morshedi, A. Schuffner, and S. Oehninger, Semen treatment with progesterone and/or acetyl-L-carnitine does not improve sperm motility or membrane damage after cryopreservation-thawing. *Fertil Steril* 74 (2000) 715-20.
- [17]R.E. Jackson, C.L. Bormann, P.A. Hassun, A.M. Rocha, E.L. Motta, P.C. Serafini, and G.D. Smith, Effects of semen storage and separation techniques on sperm DNA fragmentation. *Fertil Steril* 94 2626-30.
- [18]M.J. Burkitt, ESR spin trapping studies into the nature of the oxidizing species formed in the Fenton reaction: pitfalls associated with the use of 5,5-dimethyl-1-pyrroline-N-oxide in the detection of the hydroxyl radical. *Free Radic Res Commun* 18 (1993) 43-57.
- [19]R.V. Lloyd, P.M. Hanna, and R.P. Mason, The origin of the hydroxyl radical oxygen in the Fenton reaction. *Free Radic Biol Med* 22 (1997) 885-8.
- [20]M. Fojta, L. Havran, T. Kubiarova, and E. Palecek, Electrode potential-controlled DNA damage in the presence of copper ions and their complexes. *Bioelectrochemistry* 55 (2002) 25-7.
- [21]D.H. Johnston, K.C. Glasgow, and H.H. Thorp, Electrochemical Measurement of the Solvent Accessibility of Nucleobases Using Electron Transfer between DNA and Metal Complexes.

- [22]L. Zhou, and J.F. Rusling, Detection of chemically induced DNA damage in layered films by catalytic square wave voltammetry using Ru(bpy)₃(2+). Anal Chem 73 (2001) 4780-6.
- [23]H.H. Thorp, Cutting out the middleman: DNA biosensors based on electrochemical oxidation. Trends Biotechnol 16 (1998) 117-121.

CHAPTER III

L-CARNITINE AS PROTECTIVE MEDIA FOR SPERM IN ASSISTED REPRODUCTIVE TECHNOLOGIES

Summary

In vitro incubation and centrifugation are known to decline human sperm quality. In our bodies, besides its antioxidant effects, L-carnitine facilitates the transport of the activated fatty acids from cytosol to the mitochondrial matrix and has antioxidant effects. Here, we investigated the effect of L-carnitine (LC) on human sperm motility, viability and DNA oxidation after incubation and centrifugation, following the sperm preparation protocols of assisted reproduction. Normospermic semen samples (n=55) were analyzed according to the World Health Organization (WHO) guidelines. LC concentrations that are not toxic to sperm using sperm motility and viability were standardized after 2- and 4-hour incubation at 37°C. Semen samples were also centrifuged for 20 min at 1600 rpm with the optimized LC concentration and analyzed for sperm motility, viability and DNA oxidation. Optimal improvement in sperm motility was seen at 0.5 mg/mL LC after

incubation and centrifugation with 5×10^6 sperm/mL. Higher concentration of LC (50 mg/mL) significantly decreased sperm motility and viability. LC did not alter the baseline of sperm DNA oxidation during both incubation and centrifugation. In conclusion, LC may enhance sperm motility following incubation and centrifugation, while it might not affect sperm viability and DNA oxidation.

3.1 Introduction

Infertility affects one in seven couples who are trying to conceive. Male factor infertility is involving in up to half of all infertile couples [1; 2]. Decreased semen quality, a measure of the ability of semen to achieve fertilization, is a primary cause of male infertility. Usually, poor semen quality is characterized by low sperm motility and viability. Assisted reproductive techniques (ART) such as intrauterine insemination (IUI), *in vitro* fertilization (IVF) with or without intracytoplasmic sperm injection (ICSI) are the most successful therapeutic means for male factor infertility.

Sperm preparation protocols used in ART involve sperm incubation and centrifugation. It was reported that sperm motility and viability decrease over time after ejaculation [3]. Further, a rapid decline in motility was noted after incubating periods lasting more than 4-hours at 37°C [4]. A recent study has found that different centrifugation adversely affects sperm recovery [5]. Increased centrifugation speeds (> 500g) significantly increases the number of dead sperm [6]. Centrifugation has also been shown to increase the reactive oxygen species (ROS) formation in semen, which may affect sperm survival [7].

L-Carnitine (LC) is a naturally occurring molecule, derived from the amino acid lysine. In cellular systems, LC serves as a facilitator for the transport of activated fatty acids into the mitochondrial matrix, so that they can be broken down through β -oxidation to produce ATP [8]. A number of studies have examined the role of LC in reducing the cellular oxidative stress [9; 10; 11] (figure 3-1). Studies reported an association between lower levels of LC in semen and male infertility [12; 13]. Dietary supplementation with LC has been reported to improve sperm quality [14; 15]. Further, a positive correlation between free LC in human semen and sperm count and motility has been reported [16]. In addition, it was suggested that LC might be regarded as an index of androgenization [17]. *In vitro* as well as *in vivo* studies show that LC decreases the oxidative stress-induced DNA damage [18; 19; 20]. Furthermore, LC was reported to decrease actinomycin-D, hydrogen peroxide and TNF-alpha-induced DNA damage and improve the *in vitro* blastocyst development rate in mouse embryos [21].

The main goal of this study was first to standardize the LC doses that are not toxic to sperm, and examine LC effectiveness in improving sperm motility and viability, and decreasing sperm DNA oxidation induced by *in vitro* incubation and centrifugation.




Figure 3-1: LC function. LC facilitates the fatty acid transport into mitochondrial matrix and has an antioxidant effect.

3.2 Materials and methods

3.2.1 Sample collection

This study was approved by the institutional review board of Cleveland Clinic Hospital. Fifty five normozoospermic men were selected and analyzed on the basis of normal semen analysis according to the World Health Organization (WHO, 1999) guidelines (section 2.2.1), and used to study the effectiveness of LC on sperm motility, viability and sperm DNA oxidation during incubation and centrifugation. Samples were collected by masturbation following a minimum of 48 to 72 hours of sexual abstinence. Following liquefaction at 37 °C for 20 minutes, semen specimens were evaluated for volume, sperm concentration, total cell count, motility and morphology. A 5µL aliquot of the sample was used for evaluation of concentration and motility using a microcell slide chamber (Conception Technologies, San Diego, CA).

3.2.2 Effect of various LC concentrations on sperm motility and viability

Sixteen semen samples were examined. Each sample was divided into 6 equal aliquots; each contained 5×10^6 sperm/ mL. Samples were supplemented with various concentrations of LC dissolved in phosphate buffered saline, pH 7.4 (PBS). LC concentrations tested were 0.1, 0.5, 1.0, 10 and 50 mg/mL. The control contained PBS in place of LC. Sperm motility and viability were assessed after 2 and 4-hours incubation at 37°C.

3.2.3 Measurement of sperm viability

Sperm viability was measured by Eosin-Nigrosin staining (see page 25, chapter II)

3.2.4 Effect of LC on sperm motility and viability during centrifugation

Semen samples (n=19) were divided into two equal aliquots; each aliquot (5×10^6 sperm) was supplemented with 0.5mg/mL of LC dissolved in human tubal fluid (HTF); control was supplemented with HTF in the place of LC. Sperm motility and viability were assessed after centrifugation at 1600 rpm for 20 min.

3.2.5 Effect of LC on sperm motility and viability for various sperm concentrations during sperm incubation

Semen samples from 12 donors were used to examine the optimal sperm concentration for studying the effects of LC. Sperm concentrations used were 5, 10, 20 and 40×10^6 /mL. Each aliquot was supplemented with 0.5 mg/mL LC dissolved in PBS. Control contained PBS in place of LC. Sperm motility and viability were measured immediately (time 0) and after 2h incubation at 37°C.

3.2.6 Effect of LC on sperm DNA oxidation

Oxidative damage to DNA was evaluated using the flow cytometric OxiDNA assay kit (Calbiochem, San Diego). The assay is based on utilizing a direct fluorescent

protein binding method targeting oxidized sites in DNA. Briefly, sperm samples were treated with LC concentration at 0.5 and 50 mg/mL, incubated for 2h at 37°C, washed twice in PBS, resuspended in 1% paraformaldehyde at a concentration of $2-4 \times 10^6$ cell/mL, and placed on ice for 15 to 30 minutes. Sperm were again washed and resuspended in 70% ice-cold ethanol by centrifugation at X300g for 5 minutes as per the kit instructions. The ethanol supernatant was removed, and the sperm pellets were washed twice in wash buffer and resuspended in 100 μ L of the staining solution for 1 hour at room temperature in dark. The staining solution contained FITC-labeled protein conjugate and distilled water. All cells were further washed using rinse buffer, resuspended in 250 μ L and incubated for 30 minutes in the dark on ice for flow cytometry measurements. Control samples without LC supplementation were subjected to the same assay. Negative controls were prepared without FITC staining, while the positive controls were prepared in the presence of 0.5 mM H₂O₂ and 0.25 mM FeSO₄. To study the effect of LC on sperm DNA oxidation during centrifugation, sperm samples (5×10^6 cell/mL) were treated with one volume HTF supplemented with LC concentration at 0.5 mg/mL, centrifuged for 20 min at 1600 rpm, then submitted to the same treatment as described above.

Data acquisition was performed within 30 min on a flow cytometer equipped with a 515-nm argon laser as a light source (FACScan; Becton Dickinson, San Jose, CA). 10,000 cells were examined for each assay at a flow rate of 100 cells/second. The FITC (log green fluorescence) was measured on FL1 channel. Data analysis was performed using FlowJo v4.4.4 software (Tree Star Inc., Ashland, OR).

3.2.7 Statistical analysis

All values were reported as mean \pm S.D. Differences were considered significant at $P < 0.05$ against control group. Statistical analyses were operated using one-way analysis of variance (ANOVA) followed by an unpaired Student's t -test using the SPSS/PC computer software (SPSS 10.0.7, SPSS Inc.).

3.3 Results

3.3.1 Effect of LC on Sperm Motility and Viability on Incubation

The effect of various LC concentrations (0.1, 0.5, 1.0, 10, and 50 mg/mL) on sperm motility and viability was examined after 2 and 4-hour incubation periods at 37°C (Figure 3-2). Sperm motility was significantly increased after LC supplementation at 0.1, 0.5 and 1.0 mg/mL after 2 h incubation ($55.0\% \pm 16.9\%$, $58.4\% \pm 13.9\%$ and $52.4\% \pm 19\%$ respectively) compared with the control ($39.4\% \pm 15\%$; $P < 0.05$) (Figure 3-2A). Semen samples supplemented with LC and evaluated after the 4-hours incubation period did not significantly increase in sperm motility compared with the control (Figure 3-2A). Semen supplementation with LC did not show a statistically significant increase in sperm viability at any concentration compared with the control at both incubation periods (Figure 3-2B). LC concentration of 50 mg/mL significantly reduced sperm motility and viability compared with the control at the two incubation periods ($P < 0.05$; Figure 3-2A-B).

3.3.2 Effect of LC on Sperm Motility and Viability during Sperm Centrifugation

Figure 3-3 shows the effect of LC on sperm motility and viability after sperm centrifugation for 20 min at 1600 rpm. Figure 3-3A shows the mean values \pm standard deviations of sperm motility for semen samples centrifuged in presence of LC (0.5 mg/mL) and samples centrifuged in absence of LC (control). A significant difference ($P < 0.05$) in average sperm motility (37.5 ± 16.6 vs. 24.0 ± 12.8 %) was observed between sperm centrifuged with LC compared to the control. The viability was comparable

between sperm samples treated with LC compared to those without LC treatment (38.0 ± 12.0 vs. 34.2 ± 12.6 %, respectively) (Figure 3-3B).




Figure 3-2: Effect of various *in vitro* incubation with L-carnitine concentrations at 37°C on sperm motility for A: 2h and B: 4 h; viability C: 2 h and D: 4h. Data are mean values \pm standard deviations of sperm motility and viability obtained from analysis of sixteen different semen samples. *P*-values for samples and their controls were measured for all LC concentrations (**P*-value < 0.05; ***P*-value <0.01); *P*<0.05 was considered statistically significant.




Figure 3-3: Effect of centrifugation on A: sperm motility with and without L-carnitine; B: sperm viability with and without L-carnitine. Sperm aliquots were centrifuged for 20 min at 1600 rpm. All data are mean values \pm standard deviations. $P < 0.05$ are considered statistically significant.

3.3.3 Effect of LC on Sperm Motility and Viability for various Sperm Concentrations

The effect of LC at 0.5 mg/mL concentration on sperm motility and viability was evaluated for various sperm concentrations (5, 10, 20, and 40 million/mL) (Figure 3-4). Control samples without LC supplementation were subjected to the same assay conditions. Motility and viability of sperm were assessed immediately after 2h incubation at 37°C. Statistically significant increase in sperm motility was observed in semen samples supplemented with LC compared with the control only for a sperm concentration of 5×10^6 /mL ($P < 0.05$; Figure 3-4A). No increase in sperm viability was seen after addition of LC at all sperm concentrations tested (Figure 3-4B).

3.3.4 Effect of LC on Sperm DNA Oxidation

The effect of LC on sperm DNA oxidation during incubation and centrifugation is shown in Figure 3-5. Flow cytometry analysis of FITC-labeled sperm showed that sperm treated with the LC at 0.5 and 50 mg/mL for 2h at 37 °C did not increase the intensity of the FITC fluorescence, indicating an insignificant effect on the intrinsic baseline of DNA oxidation (Figure 3-5 A1-A2).

Figure 3-5B shows the effect of LC on the baseline of sperm DNA oxidation during centrifugation as evaluated by flow cytometry. Sperm samples centrifuged with LC at 0.5 mg/mL concentration did not show a significant difference in the intensity of FITC fluorescence, and thus in sperm DNA oxidation, compared to those without LC supplementation.




Figure 3-4: Effect of L-carnitine (0.5 mg/mL) on A: sperm motility and B: viability following incubation with various sperm concentration (5, 10, 20 and 40 X 10⁶/mL. Motility and viability of sperm were assessed immediately after 2h *in vitro* incubation at 37°C. The histograms are mean values ± standard deviations obtained from analysis of twelve different semen samples (yellow). Controls (blue) were assayed without addition of L-carnitine. *P*-values for samples and their controls were measured for all sperm concentrations (* *P*-value < 0.05); *P*<0.05 was considered statistically significant.




Figure 3-5: Effect of LC on sperm DNA oxidation during sperm incubation and centrifugation and evaluated by flow cytometry. A: Sperm incubated for 2h with 0.5 mg/mL (A1) and 50 mg/mL L-carnitine concentrations (A2) at 37°C. B: Sperm centrifuged in the presence of 0.5 mg/mL LC for 20 min at 1600 rpm. Sample controls assayed in the absence of LC in both incubation and centrifugation. Negative controls were analyzed without FITC staining; positive controls contained H₂O₂-induced sperm DNA oxidative damage. The X-axis represents the intensity of sperm-labeled FITC fluorescence. Data are representative of 4 independent experiments.

3.4 Discussion

Studies examining effects of LC on sperm quality in general have focused on the effects of LC *in vivo*. Recent findings in this context show that administration of LC might antagonize the oxidative, as well as the pro-inflammatory, pathways that attenuate sperm motility [15; 23; 24]. In the present study, we have focused on the role of LC in improving sperm quality *in vitro*. In the first series of experiments, we standardized the LC concentrations, utilizing semen from normozoospermic donors and established that 0.1, 0.5 and 1.0 mg/mL concentrations were not toxic. Next, we examined sperm motility and viability after 2- and 4-hours of incubation.

To study the effect of LC on sperm motility and viability during sperm centrifugation, we used 0.5 mg/mL LC concentration. Sperm aliquots with and without LC (control) were centrifuged for 20 min at 1600 rpm. Sperm motility and viability were then assessed. Our results suggest that LC at the standardized concentrations was beneficial and improved sperm motility, in particular, after the 2h incubation period. A significant increase in sperm motility was observed in the centrifuged sperm supplemented with LC compared to the control. Similar enhancement in sperm motility *in vitro* has been reported after adding acetylcarnitine, an acetylated form of carnitine, to semen at 37°C [25]. This positive effect of LC on sperm motility is due to its role in sperm metabolism as well as its antioxidant properties [26; 27]. LC facilitates the transport of long chain fatty acids across the inner membrane of mitochondria, so that they can be broken down through β -oxidation to produce ATP [8]; therefore, providing readily available energy for use by spermatozoa. Accumulation of the reactive oxygen species (ROS) in sperm leads to ATP depletion, lipid peroxidation and insufficient

axonemal phosphorylation [28]. The property of LC as a scavenger for ROS could thus be responsible for its positive effect on sperm motility.

The effect of LC on sperm viability was measured using the one-step Eosin-Nigrosin Test [29]. This test is primarily based on the structural integrity of the spermatozoa plasma membrane; damage in the membrane provides an access for Eosin to stain the entire spermatozoa or part of it based on the size of the damage [29]. A stained spermatozoa is considered non-viable, while the unstained one is counted as viable.

LC contributes in shielding sperm membranes against harmful ROS because of its antioxidant activity [27]. It also decreases the availability of lipids for peroxidation *via* easing the transport of fatty acids to the mitochondrial matrix for β -oxidation [30]. Neuman *et al.* (2002) reported that dietary supplementation with LC reduces sperm lipid peroxidation in roosters; this may preserve sperm membranes, thereby increasing their viability [31]. We therefore anticipated that semen supplementation with LC would improve sperm viability. Contrary to our expectation, we observed that semen supplementation with LC showed only a slight but statistically insignificant effect in improving sperm viability during incubation. Similar insignificant effects were reported earlier by Namik *et al.* (2000) who studied the influence of acetyl-L-carnitine on sperm membrane damage after cryopreservation-thawing cycles [32]. Similarly, using *in vivo* systems, it was found that dietary supplementation with LC did not improve the membrane integrity of pony stallion sperm [33]. A more recent study by Yeste *et al.* (2010) showed that the addition of LC to the diet of boar males had no effect on their sperm viability [34]. In contrast, De Rosa *et al.* (2005) noted that LC administration to infertile human males with poor sperm motility (<50%) was effective in improving

membrane integrity of the ejaculated sperm. In our *in vitro* study, we only added LC to semen samples with normal motility (>50%). A similar *in vitro* study, utilizing semen samples from infertile subjects is needed to assess LC effect on sperm viability during incubation and substantiate our findings.

Namik et al. (2000) used two assays eosin-Y staining and annexin V-Cy3 in their study to measure sperm viability after cryopreservation-thawing. These investigators observed significant discrepancy between the percentages of sperm marked as “live” between eosin-Y staining and annexin V-Cy3 assay. The eosin-Y staining showed lower percent of viable sperm compared to the annexin V-Cy3 assay [32]. These investigators attributed this discrepancy to the different mechanisms through which these assays operate. In our study, we measured sperm viability using the eosin-staining assay, which is based on the membrane permeability for the eosin stain. Therefore, the mechanism by which this assay recognizes live sperm could be a contributing factor in this insignificant effect of LC on sperm viability.

The influence of LC on sperm motility and viability of various sperm concentrations was also investigated (Figure 3). We supplemented different concentrations of sperm with LC at 0.5 mg/mL. Our results from this experiment suggest that LC was effective in increasing sperm motility only when utilizing sperm concentration of 5×10^6 cell/mL. At higher sperm concentrations, the decrease in the ratio of the sperm number against available LC molecules may explain the lack of effect in improving the sperm motility. On the other hand, sperm viability did not increase utilizing any of the sperm concentrations. These results are in agreement with our experiment above. This information may be exploited in exploring and designing

protocols that aim to improve sperm motility in patients with low sperm count and especially oligozoospermic patients.

LC has been shown to improve resistance to oxidative stress by decreasing DNA damage in *Ataxia telangiectasia* cells [35]. Moreover, Thangasamy *et al.* (2009) reported that LC significantly reduces DNA damage in lymphocytes of aged rats [18]. We also recently demonstrated the effectiveness of LC in reducing H₂O₂-induced DNA damage and improvement of *in vitro* blastocyst development rate in mouse embryos [21]. In the present study, we examined if LC had similar effects on human sperm DNA *in vitro*. We incubated the sperm samples with the standardized LC concentration 0.5 mg/mL, at 37°C for 2h. We did not see the decrease in the extent of the sperm DNA oxidation (Figure 3-5-A1). Similarly, we did not see a significant effect of LC on the baseline of sperm DNA oxidation after centrifugation compared with control samples (i.e. centrifugation without LC) (Figure 3-5-B).

Studies *in vitro* show that semen supplementation with an antioxidant *does not* affect sperm DNA integrity. In fact, Donnelly *et al.* (1999) showed that neither addition of ascorbate nor addition of α -tocopherol to sperm preparation medium affected DNA integrity of sperm. However, both antioxidants provided protection against the H₂O₂-induced DNA damage in sperm [36]. Similar reports showed that addition of vitamin E to cryopreservation medium did not alter the post-thaw DNA fragmentation of sperm [37]. These results are in line with our observations, which don't show a significant effect of LC as an antioxidant LC in reducing DNA oxidative damage in sperm.

Very low concentrations of LC have been reported in azoospermic men compared with normospermic control [38]. Further, lower concentrations of LC were noted in oligoasthenozoospermic males, and LC supplementation was shown to improve semen quality in these people [39]. One explanation for the undetectable effect of LC on sperm DNA oxidation may be the fact that we used only semen samples from normozoospermic subjects in our study. Furthermore, contrary to majority of reported studies [21; 23; 35; 36], we measured the effect of LC on the baseline levels of oxidative DNA damage in sperm without any induction of oxidative damage as in other studies this might be another likely reason for the lack of significant effect of LC on sperm DNA oxidation. Another reason for lack of improvement in DNA oxidation may be the fact that mammalian sperm DNA is highly compacted (six-fold more highly condensed) compared to DNA in somatic cells [40]. Also, the majority of antioxidants in somatic cells are present within their cytoplasm, while in sperm, a significant amount of the antioxidants are present in seminal plasma outside the sperm because spermatozoa lose most of its cytoplasm during maturation [41]. Therefore, the differences in the DNA packaging and the distribution of antioxidants between spermatozoa and somatic cells might introduce variations in outcomes when measuring the effect of *in vitro* supplementation of antioxidants on DNA oxidation.

Given that LC is a substantial antioxidant [27; 42], the role of antioxidant therapy in improving male fertility could be a double-edged sword with unfavorable effects if the safety threshold dose is exceeded [42]. For example, extreme levels of the known antioxidant L-ascorbic acid are thought to be associated with a decline in fertility [43], even though it is the principal antioxidant in seminal plasma of fertile males [44]. Further

research is needed before the role for antioxidants in the treatment of male infertility is established [45]. The *in vitro* studies on human sperm by Donnelly et al. (1999) showed that higher concentrations of L-ascorbic acid (> 20 µM) negatively affect sperm motility in both normozoospermic and asthenozoospermic samples. Our findings are in agreement with this concept, and we have shown that sperm quality *in vitro* is lower at high concentrations of LC.

It has been shown that cell treatment with high concentrations of antioxidants *in vitro* increases the levels of 8-oxoguanine moieties on DNA, a biomarker for DNA oxidative damage [46]. Increasing DNA oxidative damage may lead to apoptosis and cell death [47; 48]. We therefore examined if high LC concentration decreased *in vitro* sperm quality by inducing sperm DNA oxidation. We measured the 8-oxoguanine moieties in the sperm DNA after incubating the sperm with 50 mg/mL LC using flow cytometry. Our results suggest that this might not be the likely mechanism by which the high LC level affects negatively sperm quality *in vitro* (Figure 3-5-A2). Again, the fact that we used samples from normozoospermic subjects may contribute to the absence of significant effects in our work.

In conclusion, LC enhances sperm motility following *in vitro* incubation and centrifugation, while it might not significantly affect sperm viability and DNA oxidation baseline of sperm. Our results provide a solid ground to explore and design protocols to test potentially beneficial effect of LC supplementation for sperm prepared for ART.

3.5 References

- [1]D.J. Lamb, and L.I. Lipshultz, Male infertility: recent advances and a look towards the future. *Curr Opin Urol* 10 (2000) 359-62.
- [2]V.M. Brugh, 3rd, and L.I. Lipshultz, Male factor infertility: evaluation and management. *Med Clin North Am* 88 (2004) 367-85.
- [3]R. Singer, M. Sagiv, B. Landau, D. Allalouf, E. Segenreich, and C. Servadio, Motility and vitality of human spermatozoa at various time intervals after ejaculation. *Experientia* 36 (1980) 578-9.
- [4]C. Jeulin, C. Serres, and P. Jouannet, The effects of centrifugation, various synthetic media and temperature on the motility and vitality of human spermatozoa. *Reprod Nutr Dev* 22 (1982) 81-91.
- [5]C. Matas, G. Decuadro, S. Martinez-Miro, and J. Gadea, Evaluation of a cushioned method for centrifugation and processing for freezing boar semen. *Theriogenology* 67 (2007) 1087-91.
- [6]A. Makler, and P. Jakobi, Effects of shaking and centrifugation on human sperm motility. *Arch Androl* 7 (1981) 21-6.
- [7]M. Shekarriz, D.M. DeWire, A.J. Thomas, Jr., and A. Agarwal, A method of human semen centrifugation to minimize the iatrogenic sperm injuries caused by reactive oxygen species. *Eur Urol* 28 (1995) 31-5.
- [8]A. Steiber, J. Kerner, and C.L. Hoppel, Carnitine: a nutritional, biosynthetic, and functional perspective. *Mol Aspects Med* 25 (2004) 455-73.
- [9]D. Silva-Adaya, V. Perez-De La Cruz, M.N. Herrera-Mundo, K. Mendoza-Macedo, J. Villeda-Hernandez, Z. Binienda, S.F. Ali, and A. Santamaria, Excitotoxic

- damage, disrupted energy metabolism, and oxidative stress in the rat brain: antioxidant and neuroprotective effects of L-carnitine. *J Neurochem* 105 (2008) 677-89.
- [10]R. Apak, K. Guclu, M. Ozyurek, and S.E. Karademir, Novel total antioxidant capacity index for dietary polyphenols and vitamins C and E, using their cupric ion reducing capability in the presence of neocuproine: CUPRAC method. *J Agric Food Chem* 52 (2004) 7970-81.
- [11]A. Vanella, A. Russo, R. Acquaviva, A. Campisi, C. Di Giacomo, V. Sorrenti, and M.L. Barcellona, L -propionyl-carnitine as superoxide scavenger, antioxidant, and DNA cleavage protector. *Cell Biol Toxicol* 16 (2000) 99-104.
- [12]K. Li, W. Li, Y.F. Huang, and X.J. Shang, [Level of free L-carnitine in human seminal plasma and its correlation with semen quality]. *Zhonghua Nan Ke Xue* 13 (2007) 143-6.
- [13]I. Matalliotakis, Y. Koumantaki, A. Evageliou, G. Matalliotakis, A. Goumenou, and E. Koumantakis, L-carnitine levels in the seminal plasma of fertile and infertile men: correlation with sperm quality. *Int J Fertil Womens Med* 45 (2000) 236-40.
- [14]D.M. Kozink, M.J. Estienne, A.F. Harper, and J.W. Knight, Effects of dietary L-carnitine supplementation on semen characteristics in boars. *Theriogenology* 61 (2004) 1247-58.
- [15]M. Yeste, S. Sancho, M. Briz, E. Pinart, E. Bussalleu, and S. Bonet, A diet supplemented with l-carnitine improves the sperm quality of Pietrain but not of Duroc and Large White boars when photoperiod and temperature increase. *Theriogenology* (2009).

- [16]G.F. Menchini-Fabris, D. Canale, P.L. Izzo, L. Olivieri, and M. Bartelloni, Free L-carnitine in human semen: its variability in different andrologic pathologies. *Fertil Steril* 42 (1984) 263-7.
- [17]G. Abbaticchio, V.A. Giagulli, M. Defini, F.M. Micale, and R. Giorgino, Free l-carnitine in human semen. *Arch Androl* 15 (1985) 137-42.
- [18]T. Thangasamy, P. Jeyakumar, S. Sittadjody, A.G. Joyee, and P. Chinnakannu, L-carnitine mediates protection against DNA damage in lymphocytes of aged rats. *Biogerontology* 10 (2009) 163-72.
- [19]D. Haripriya, P. Sangeetha, A. Kanchana, M. Balu, and C. Panneerselvam, Modulation of age-associated oxidative DNA damage in rat brain cerebral cortex, striatum and hippocampus by L-carnitine. *Exp Gerontol* 40 (2005) 129-35.
- [20]S.N. Qi, Z.F. Zhang, Z.Y. Wang, A. Yoshida, and T. Ueda, L-carnitine inhibits apoptotic DNA fragmentation induced by a new spin-labeled derivative of podophyllotoxin via caspase-3 in Raji cells. *Oncol Rep* 15 (2006) 119-22.
- [21]H. Abdelrazik, R. Sharma, R. Mahfouz, and A. Agarwal, L-carnitine decreases DNA damage and improves the in vitro blastocyst development rate in mouse embryos. *Fertil Steril* 91 (2009) 589-96.
- [22]M.J. Burkitt, ESR spin trapping studies into the nature of the oxidizing species formed in the Fenton reaction: pitfalls associated with the use of 5,5-dimethyl-1-pyrroline-N-oxide in the detection of the hydroxyl radical. *Free Radic Res Commun* 18 (1993) 43-57.
- [23]A.R. Abd-Allah, G.K. Helal, A.A. Al-Yahya, A.M. Aleisa, S.S. Al-Rejaie, and S.A. Al-Bakheet, Pro-inflammatory and oxidative stress pathways which compromise

- sperm motility and survival may be altered by L-carnitine. *Oxid Med Cell Longev* 2 (2009) 73-81.
- [24]A. Garolla, M. Maiorino, A. Roverato, A. Roveri, F. Ursini, and C. Foresta, Oral carnitine supplementation increases sperm motility in asthenozoospermic men with normal sperm phospholipid hydroperoxide glutathione peroxidase levels. *Fertil Steril* 83 (2005) 355-61.
- [25]N. Tanphaichitr, In vitro stimulation of human sperm motility by acetylcarnitine. *Int J Fertil* 22 (1977) 85-91.
- [26]A. Agarwal, and T.M. Said, Carnitines and male infertility. *Reprod Biomed Online* 8 (2004) 376-84.
- [27]I. Gulcin, Antioxidant and antiradical activities of L-carnitine. *Life Sci* 78 (2006) 803-11.
- [28]D. Dokmeci, Oxidative stress, male infertility and the role of carnitines. *Folia Med (Plovdiv)* 47 (2005) 26-30.
- [29]L. Bjorndahl, I. Soderlund, and U. Kvist, Evaluation of the one-step eosin-nigrosin staining technique for human sperm vitality assessment. *Hum Reprod* 18 (2003) 813-6.
- [30]Thangavel Kalaiselvia, and C. Panneerselvama, Effect of l-carnitine on the status of lipid peroxidation and antioxidants in aging rats. 9 (1998) 575-581.
- [31]S.L. Neuman, T.L. Lin, and P.Y. Heste, The effect of dietary carnitine on semen traits of white Leghorn roosters. *Poult Sci* 81 (2002) 495-503.

- [32]N.K. Duru, M. Morshedi, A. Schuffner, and S. Oehninger, Semen treatment with progesterone and/or acetyl-L-carnitine does not improve sperm motility or membrane damage after cryopreservation-thawing. *Fertil Steril* 74 (2000) 715-20.
- [33]K. Deichsel, F. Palm, P. Koblichke, S. Budik, and C. Aurich, Effect of a dietary antioxidant supplementation on semen quality in pony stallions. *Theriogenology* 69 (2008) 940-5.
- [34]M. Yeste, S. Sancho, M. Briz, E. Pinart, E. Bussalleu, and S. Bonet, A diet supplemented with L-carnitine improves the sperm quality of Pietrain but not of Duroc and Large White boars when photoperiod and temperature increase. *Theriogenology* 73 577-86.
- [35]A. Berni, R. Meschini, S. Filippi, F. Palitti, A. De Amicis, and L. Chessa, L-carnitine enhances resistance to oxidative stress by reducing DNA damage in Ataxia telangiectasia cells. *Mutat Res* 650 (2008) 165-74.
- [36]E.T. Donnelly, N. McClure, and S.E. Lewis, The effect of ascorbate and alpha-tocopherol supplementation in vitro on DNA integrity and hydrogen peroxide-induced DNA damage in human spermatozoa. *Mutagenesis* 14 (1999) 505-12.
- [37]K. Taylor, P. Roberts, K. Sanders, and P. Burton, Effect of antioxidant supplementation of cryopreservation medium on post-thaw integrity of human spermatozoa. *Reprod Biomed Online* 18 (2009) 184-9.
- [38]L.M. Lewin, D.P. Shalev, R. Weissenberg, and Y. Soffer, Carnitine and acylcarnitines in semen from azoospermic patients. *Fertil Steril* 36 (1981) 214-8.
- [39]C.M. Ng, M.R. Blackman, C. Wang, and R.S. Swerdloff, The role of carnitine in the male reproductive system. *Ann N Y Acad Sci* 1033 (2004) 177-88.

- [40]W.S. Ward, and D.S. Coffey, DNA packaging and organization in mammalian spermatozoa: comparison with somatic cells. *Biol Reprod* 44 (1991) 569-74.
- [41]E.T. Donnelly, N. McClure, and S.E. Lewis, Antioxidant supplementation in vitro does not improve human sperm motility. *Fertil Steril* 72 (1999) 484-95.
- [42]Tarin JJ, Brines J, and Cano A, Antioxidants may Protect against infertility. *Hum Reprod* 13 (1998) 1419-21.
- [43]M. Igarashi, Augmentative effect of ascorbic acid upon induction of human ovulation in clomiphene-ineffective anovulatory women. *Int J Fertil* 22 (1977) 168-73.
- [44]S.E. Lewis, E.S. Sterling, I.S. Young, and W. Thompson, Comparison of individual antioxidants of sperm and seminal plasma in fertile and infertile men. *Fertil Steril* 67 (1997) 142-7.
- [45]R.C. Martin-Du Pan, and D. Sakkas, Is antioxidant therapy a promising strategy to improve human reproduction? Are anti-oxidants useful in the treatment of male infertility? *Hum Reprod* 13 (1998) 2984-5.
- [46]J.M. Gutteridge, Antioxidants, nutritional supplements and life-threatening diseases. *Br J Biomed Sci* 51 (1994) 288-95.
- [47]Y.J. Hurh, Z.H. Chen, H.K. Na, S.Y. Han, and Y.J. Surh, 2-Hydroxyestradiol induces oxidative DNA damage and apoptosis in human mammary epithelial cells. *J Toxicol Environ Health A* 67 (2004) 1939-53.
- [48]X. Wang, R.K. Sharma, S.C. Sikka, A.J. Thomas, Jr., T. Falcone, and A. Agarwal, Oxidative stress is associated with increased apoptosis leading to spermatozoa DNA damage in patients with male factor infertility. *Fertil Steril* 80 (2003) 531-5.

CHAPTER IV

NITRIC OXIDE IN HUMAN MALE REPRODUCTION

Summary

In this chapter, we provide a literature review of the role of nitric oxide (NO), one of the main reactive oxygen species, in human male reproduction. The goal of this review is to enrich our understanding of the molecular basis of the function of NO in human spermatozoa. In the beginning of this chapter, we discuss nitric oxide synthesis and mechanisms of action and function in cellular systems. Then, we provide a comprehensive literature review of the role of NO in human spermatozoa physiology and function, as well as its contribution as a marker in male infertility diagnosis. In addition, we provide an in-depth review of NO antagonistic effects on spermatozoa function and physiology. Moreover, we review the methods that are commonly used to measure NO in semen. The collective information from this literature survey points to the fact that some level of NO is vital for human sperm function and physiology and mainly for the motility

of spermatozoa. However, increased levels of NO are considered detrimental to spermatozoa and contribute to male infertility.

4.1 Nitric oxide, function and synthesis

Nitric oxide (NO) is a diatomic free radical. It is similar to oxygen in its maximum solubility in water. NO is also a non-polar molecule and able to diffuse easily through cellular membranes [1]. One of the most distinctive chemical features of NO is that it is paramagnetic, does not have a tendency to make dimers at the standard conditions [2].

Currently, it is one of the extensively studied species in biomedicine because of its various functions in biological systems. It has been referred to as the endothelium-derived relaxing factor (EDRF) because it causes vasodilation via a cascade that leads to vascular smooth muscle relaxation [3; 4; 5]. In addition, nitric oxide prevents platelets aggregation [6], acts as a neurotransmitter [7; 8], and mediates macrophage bactericidal activity.

Reports have shown that NO has a major contribution in human reproduction. It is recognized as a key molecule in controlling hypothalamic-pituitary-gonadal axis [1]; it activates the release of luteinizing hormone-releasing hormone (LHRH), which triggers the pituitary gland to release the gonadotropins via activating nNOS in the pituitary [9]. In the gonad, it has a central role in inducing ovulation in human's female and in causing luteolysis [9]. NO has also a significant contribution in human's sexual behavior; in males, it induces the erection via generating cyclic guanosine monophosphate (cGMP) in

corpora cavernosa [10]. In females, it relaxes vaginal smooth muscle and induces vaginal secretions [9].




Figure 4-1: Synthesis of nitric oxide.

NO is synthesized by a family of enzymes called nitric oxide synthases (NOSs) from the oxidation of arginine. NO is released after a two-step catalytic cycle along with citrulline as a byproduct. O_2 and NADPH, the reduced form of nicotinamide adenine dinucleotide phosphate, are required in this reaction (figure 4-1). Flavin adenine dinucleotide (FAD), flavin mononucleotide (FMN), and tetrahydrobiopterin, are also needed for the NOS reaction [11]. In tissues, NO has a very short half-life (3 to 10 seconds) because it reacts with oxygen (O_2) and converts into nitrite and nitrate. It also reacts with superoxide ion ($O_2^{\cdot-}$) and thiol sites to yield peroxynitrite and nitrosothiols respectively.

4.2 NO synthases

Three isoenzymes of NO synthase have been identified. Endothelium NOS (eNOS), primarily present in endothelium, neuronal NOS (nNOS), primarily present in neural tissue, and inducible NOS (iNOS), can be expressed in various cells such as macrophages, monocytes, neutrophils and hepatocytes [11; 12].

The isoforms eNOS and nNOS are constitutive, synthesized usually at a steady rate in spite of physiologic need, and Ca^{2+} -calmodulin-dependent enzymes [11; 12]. iNOS is a Ca^{2+} -independent enzyme; it has specific inducers (e.g. inflammatory cytokines, tumor necrosis factor, bacterial endotoxins) which vary by cell type. These inducers increase iNOS synthesis, which can result in a large amount of NO being formed [13; 14; 15].

4.3 NO and smooth muscle contraction:


In endothelial cells, NO is synthesized by eNOS and diffuses into vascular smooth muscle (VSM), and activates the cytosolic *–soluble–* guanylate cyclase (sGC), which increases the production of cGMP, and consequently, protein kinase G (PKG). PKG then phosphorylates myosin light-chain kinase and leaves it inactive, thereby lowering smooth muscle contraction.

4.4 Nitric oxide mechanisms of action mediated by peroxynitrite

In biological systems, peroxynitrite, which results from the combination of NO with superoxide ion ($O_2^{\cdot-}$), is a *double-edged sword* anion. It has both beneficial and detrimental actions. These actions, endogenous and exogenous, are mostly due to oxidation and nitration reactions with other biomolecules, mainly proteins, lipids and nucleic acids (Figure 4-2) [16]. The effects of peroxynitrite could be both direct and indirect. For example, peroxynitrite produces small concentrations of secondary NO when reacting with other biomolecules and promotes vasodilation, inhibition of platelet aggregation and hypotension [17].

1. Role in bactericidal activity

Normally, iNOS activity in macrophages is not high. In response to infection, by the effect of interferon-alpha and bacterial lipopolysaccharides (LPS), its synthesis increases resulting in higher NO production. NO combines with $O_2^{\cdot-}$ and form peroxynitrite ($ONOO^-$), which, among other reaction pathways, breaks down in the presence of hydrogen ion into HO^{\cdot} , a highly bactericidal radical (scheme 1).



Scheme 1: Formation of hydroxyl radical from nitric oxide.

2. Effects on proteins and enzymes

Various proteins in humans and other biological systems are targets for peroxynitrite attacks. Oxidation and nitration of proteins may down-regulate the positive feedback cycles of injury and decreases the ability of the cell to fight against free radical damage. This will lead to antioxidant depletion, oxidative stress, and consequently, cell necrosis or apoptosis. Such cases are more likely when peroxynitrite attacks antioxidant enzymes such as the superoxide dismutase and glutathione reductase, as well as, other species of antioxidant defense mechanisms like glutathione and cysteine [16; 18-22]. Inhibition of cytosolic as well as membrane proteins and enzymes because of peroxynitrite attack leads to impairment of cell function, and subsequently disease onset (Table 4-1).

3. Lipid oxidation and nitration by peroxynitrite

The decomposition of peroxynitrite, in the presence of hydrogen ion, to form $\bullet\text{OH}$ and $\bullet\text{NO}_2$ may initiate lipid peroxidation when it occurs in hydrophobic phases [16; 23]. The reaction results in Lipid peroxidation products such as malondialdehyde, hydroperoxide and conjugated diene [23], as well as in the nitro, nitrito-, nitrosoperoxo- and nitrated lipid oxidation products. Low-density lipoprotein (LDL) is also a target for peroxynitrite insult. LDL oxidation by peroxynitrite leads to the formation of F₂-isoprostanes [16; 24], a family of prostaglandin-like compounds formed via non-enzymatic lipid peroxidation. Fatty acids nitration may secondarily inhibit protein function through thiol modifications [25]. Conversely, nitrated lipids can have anti-

inflammatory functions via *peroxisome proliferator-activated receptor gamma* (PPAR- γ)-dependent and independent pathways [16; 26].

4. *Effect on DNA*

DNA is also another target for peroxynitrite and its derived radicals. They can cause DNA base modification; for example, 8-oxoguanine and 8-nitroguanine are main products of purine oxidation [27]. Furthermore, it can activate Ca^{2+} - Mg^{2+} -endonucleases [16]. In addition, peroxynitrite can oxidize deoxyribose sugar and lead to double and single strand breaks [16; 28]. Such DNA injury may lead to genotoxic damage, which may develop into pathological disorders. Peroxynitrite may also induce secondary DNA repair and activate the suicidal pathway including poly ADP-ribose polymerase activation (PARP-activation), and consequently cell necrosis (Figure 4-2) [29].

Table 4-1: Examples of cytosolic and membrane enzymes and proteins affected by peroxynitrite (adapted from Casaba Szabo *et al.*, 2007).

Enzyme or protein	Mechanism of action	Consequences and related health disorders	Ref.
Glyceraldehyde-3-phosphate dehydrogenase	Oxidation of cysteine residues	Impairment of cellular function	[30]
Tyrosine hydroxylase	Nitration of tyrosine residues	Impaired the production of dopamine in the dopaminergic neurons (e.g. in Parkinson disease)	[31]
Creatine kinase	Oxidation of cysteine residues	Impairment of myocytes function, which may leads to myocardial infarction or chronic heart failure.	[16; 32]
Calcium pumps: including sarco/endoplasmic reticulaum Ca^{2+} -ATPase (SERCA)	Tyrosine nitration and cysteine oxidation	Impairment of cellular calcium handling and cellular ionic balance, which is linked with heart failure.	[33; 34]
Epidermal growth factor receptor (EGF)	Dimerization, nitration and oxidation	Cellular signaling impairment including a possible pathway leading to insulin resistance	[16; 35]
Prostacyclin synthase	Oxidation and nitration	Prostaglandin formation inhibition.	[36]
Alpha-synuclein and microtubule related tau protein modification in CNS	Nitration and oxidation	Initiation pathogenesis of alpha-synuclein aggregates	[16; 37]




Figure 4-2: nitric oxide actions mediated by peroxynitrite.

5. *Effect on NOS cofactors*

Peroxynitrite can impair enzymatic cofactors directly or indirectly via free radical-dependent mechanisms. It can oxidize tetrahydrobiopterin (BH4) (Figure 4-3), a cofactor of NOS enzyme that serves as an electron relay for timed redox reactions used in oxygen activation. The oxidation of BH4 leads to the inactive form quinonoid 5,6-dihydrobiopterin [38] which is subsequently converted to 7, 8-dihydrobiopterin. This oxidation affects negatively the function of NOS, and thus NO production [16]. Milstien and Katusic (1999) have suggested that abnormally low concentrations of BH4 may promote a cycle of its own destruction arbitrated by NOS-dependent formation of peroxynitrite; such mechanism might have a contribution in vascular endothelial dysfunction stimulated by oxidative stress [38].




Figure 4-3: Chemical structure of tetrahydrobiopterin (A), and 7, 8-dihydrobiopterin (B).

4.5 Nitric oxide and male infertility

Studies have shown that NO has a major contribution in male reproduction [39-44]. Higher levels of nitric oxide were recognized in the semen of infertile men compared to fertile [42]. It was found that NO generation is very important in gamete interaction and fertilization route [45]. Xu *et al.* (2008) studied the relationship between nitric oxide synthase and nitric oxide with varicocele-based infertility; they found that NO mediates the decrease in semen quality induced by varicocele [46]. A recent study reported that the missense Glu298Asp polymorphism of eNOS gene is present at a higher frequency in infertile asthenozoospermic subjects [47].

Supplementation of asthenozoospermic infertile men ejaculates with Ginsenoside R (e) improves human's male fertility by increasing sperm motility [48]. The mechanism by which this occurs may involve NOS stimulation and enhanced NO synthesis [48]. Huang *et al.* (2006) studied the correlation between NO levels and sperm morphology; they found that low concentrations of NO are associated with defects in sperm morphology [49]. A very recent study showed that increased levels of NO is linked with a decrease in sperm function in asthenospermia, normospermia and leukocytospermia, whereas, lower levels of NO may contribute in the etiology of oligospermia [50].

4.6 Nitric oxide synthases in human spermatozoa

NOSs are ubiquitous, present in almost all tissues of the human body [51]. NOS activity has been verified in human testes, seminal vesicles, and prostates [52]. These findings imply a possible function of NO in spermatogenesis and the maturation of

sperm. It has been recognized that nitric oxide synthase (NOS) is present in human spermatozoa and NO production at endogenous levels is very important for adequate sperm function [48; 51; 53-55]. Lewis *et al.* (1996) reported that human spermatozoa express eNOS as well as nNOS [53]. At a later date (1998), Moria *et al.* confirmed that human spermatozoa express eNOS. The same group, using immunohistochemistry, detected the presence of eNOS in sperm and showed that the majority of sperm display postacrosomal and equatorial eNOS staining (head and midpiece). In 2000, Francavilla *et al.* concluded that constitutive NOS (eNOS and nNOS) plays a role in the capability of human sperm to fuse with oocyte in the *zona pellucida* (ZP), a glycoprotein membrane that surrounds the plasma membrane of an oocyte. They showed that inhibited constitutive NOS activity using N^G-nitro-L-arginine methyl ester (L-NAME) significantly decreased the oocyte fusion without affecting the ZP binding [56]. Newly published study showed that increased NOS activity is present in mature sperm compared to immature [57].

NOS activity was also studied in seminal plasma. Both calcium-dependent NOS activity as well as calcium-independent NOS activity were detected in human seminal plasma [58]. There was no association between NOS activity in seminal plasma and sperm motility and concentration [58]. Later, another study supported these findings by showing that seminal plasma nitrite does not correlate with sperm count and rapid-forward motility [59]. In addition, positive sperm culture did not increase the concentration of nitrite in seminal plasma [59].

On a related front, mRNA transcripts of eNOS and nNOS have been detected in semen samples, and they have a clinical significance in the diagnosis of fertilization [60].

For example, eNOS and nNOS transcripts were present in low motile sperm samples, while they are undetectable in the high motile ones [60].

4.7 Nitric oxide and sperm motility

Flagellum beating is very important to sperm motion; defects in sperm motility leads to infertility and ectopic pregnancy [61]. Several studies showed that NO production at endogenous levels is very important for adequate sperm motion [48; 51; 53-55]. Insulin, a central hormone in regulating carbohydrate metabolism in the human body, and leptin, an adipose-derived hormone, significantly increased *in vitro* motility and nitric oxide production in sperm [15] (Figure 4-4). Ankri *et al.* (2009) proposed that exposure of human spermatozoa to visible light may increase NO formation, which enhances the motility of sperm, and thus the rate of fertilization. According to the same group, this effect may be attributable to increase of light absorbance by the NOS coenzymes flavin and heme. Lately, an *in vitro* study observed that adding L-arginine to asthenospermic patient's semen samples increases NO production and sperm motility. This effect happens only in the presence of an active anion transport system [62].

It was also reported that sperm recovery is also altered upon using dissimilar methods to prepare sperm *in vitro*. Ding *et al.* (2002) used two different methods, Percoll and IxaPrep, to prepare human sperm *in vitro* and they found that the IxaPrep method provides a better recovery rate of sperm. They concluded that the better sperm quality by the IxaPrep method may be the result of decreased levels of NO generated, because increased levels of nitric oxide affects negatively the quality of the sperm [63].




Figure 4-4: various compounds and factors increase sperm motility may be by increasing NO production.

4.8 Nitric oxide mechanisms of action in human sperm

Miraglia *et al.* (2010) showed that NO stimulates human spermatozoa motility by activation of soluble guanylate cyclase (sGC), subsequent production of cyclic guanosine monophosphate (cGMP) and activation of cGMP-dependent protein kinases (cGMP/protein kinase G signaling pathway) [64]. An earlier study suggested that NO induces acrosomal reaction in human spermatozoa via stimulating sGC, cGMP, and PKG activation, and extracellular Ca^{2+} is required in this activation [65].

Alternatively, by a mechanism that does not require sGC activation; L. Lefievre *et al.* (2007) demonstrated that many proteins (signaling, regulatory, kinases, receptors, transport, structural and others) in human spermatozoa are possible targets for S-nitrosylation, a dynamic mechanism analogous to phosphorylation in which a nitrosyl group is added to a protein (25). In general, S-nitrosylation by nitric oxide is very important reaction in biological systems; it refers usually to the conversion of thiol groups, involving cysteine residues in proteins, to S-nitrosothiols, such examples are shown above (Table 4-1) (26, 27).

A subsequent study reported that NO has a key function in the gamete interaction and fertilization within human female's tract. It regulates the mobilization of stored Ca^{2+} ions in human spermatozoa via protein S-nitrosylation, which ultimately lead to the regulation of flagellar beating. This action is synergistic with the action of progesterone (Figure 4-5) [66]. The pre-treatment of sperm with NO increased significantly the effect of progesterone on intracellular free calcium ions, possibly by binding on a membrane surface receptor, which activates Ca^{2+} influx using a plasma membrane channels [45].

Increasing intracellular Ca^{2+} induces mobilization of stored calcium via a Ca^{2+} -induced Ca^{2+} release process (CICR).

On another front, reports have shown an association between apoptosis mechanisms in sperm and NO synthesis. For example, activation of apoptosis signaling in human spermatozoa is negatively correlated with NO production [57].

4.9 Role of nitric oxide in human sperm capacitation and acrosomal reaction

Sperm capacitation is an ultimate biochemical event that occurs typically in the female's reproductive tract after ejaculation. It is required for the spermatozoa to be competent for oocyte penetration. A number of studies reported that NO is important for human sperm capacitation [67; 68; 69]. It was indicated that NO is formed by capacitating human spermatozoa [70]. NO may act as a messenger via modulating the cAMP/protein tyrosine phosphorylation pathway involved in spermatozoa capacitation [70]. Herrero *et.al.* (2001) emphasized that capacitation is part of an oxidation process that involves NO. They demonstrated that the nitration of tyrosine in sperm proteins by NO takes place in capacitated human spermatozoa [71]. Later study found that NO regulates the phosphorylation of threonine-Glutamine-Tyrosine motif in human spermatozoa proteins of 80 and 105 kDa through capacitation [72]. Others have shown the effect of Ginsenoside Re, a bioactive component from ginseng, on sperm capacitation and acrosomal reaction occurs through NO-dependent cGMP/ protein kinase G signaling pathway [73]. De Lamirande and Lamothe (2009) showed a close interaction between NO and superoxide ion ($\text{O}_2^{\cdot-}$) during human sperm capacitation is needed for acquisition of fertility [69]. Inhibition of NO generation as well as other ROS by semenogelin, the

predominant protein in human semen, blocks sperm capacitation [74]. On the other hand, nitric oxide at low concentrations increased human sperm capacitation through enhancement of sperm-zona pillucida (ZP) binding capacity [75].




Figure 4-5: Synergistic effect of NO and progesterone (modified from Linda *et al.*, 2009).

4.10 Adverse effects of NO on sperm physiology and function

It has been shown that NO is vital for appropriate sperm function, while overproduction of this reactive free radical has also pathologic implications, particularly on the motility of sperm [55]. Induced oxidative stress to human sperm *in vitro* using high doses of H₂O₂ showed a significant increase in NO levels and a significant decrease in sperm motility [55; 76]. Rosselli *et al.* (1995) found that higher levels of nitric oxide decrease the quality of sperm *in vitro* [40]. Later study on varicocele suggested that nitric oxide synthase released within the dilated spermatic vein establishes a high oxidative stress state as a result of increased production of NO. Formation of peroxynitrite from the reaction of NO with superoxide ion also affects sperm function in varicocele patients [77]. On the other hand, no significant relationship was found between NO concentrations and the severity of athenozoospermia, oligospermia, and abnormal sperm morphology in these people [78].


Treatment of purified spermatozoa with NO-doses from NO-donors such as S-nitroso-N-acetylpenicillamine and sodium nitroprusside decreased both sperm motility and viability [40; 42; 79]. The nitric oxide generator, SIN-1 (3-morpholinopyridone hydrochloride), was used as a spermicidal agent [80]. Adding the calcium ionophore A23187, an activator of both eNOS and nNOS, to human sperm medium increased significantly the production of NO [54] and, as a result, decreased sperm motility. Others showed that the overproduction of NO in the genital tract of infertile human subjects decreases the integrity of sperm DNA [81]. A later study demonstrated that TNF- α and IL-6, two cytokines involved in inflammation, decrease sperm motility possibly via inducing NO formation [82].

In contrast, melatonin (*N*-acetyl-5-methoxytryptamine), a hormone secreted by pineal gland and responsible for circadian rhythm, is able to scavenge directly and indirectly NO in human sperm. Accordingly, it was concluded that supplementation of sperm medium with melatonin might be able to counteract the adverse effects of NO on sperm [83].

4.11 Measurement of NO in semen

The most common method used to measure NO in human semen is *nitrite colorimetric assay*, also known as the Griess assay. In this assay, NO is assessed indirectly via measuring its breakdown product nitrite (NO_2^-), formed by spontaneous oxidation of NO [84]. The detection limit for this assay is 1 μM nitrite [84]. Sulfanilic acid first reacts with NO_2^- , in the semen sample to produce an intermediate diazonium, which then reacts with N-(1-naphthyl) ethylenediamine and form a stable azo dye, which absorbs light at 540nm [85]. To accurately measure all nitric oxide produced in the sample, some reports first convert all nitrate to nitrite using nitrate reductase enzyme, before using the Griess assay [85].

Chemiluminescence is another method commonly used to determine NO in semen. The detection is based on NO reaction with ozone (O_3), which leads to red light emission (Scheme 4-2). Photons detected from this reaction, using a photomultiplier tube are transformed to an electrical signal that is proportional to the amount of measured NO [63].



Scheme 4-2: reaction of NO with ozone.

Others have used other techniques to measure NO production at cellular level such as *electron paramagnetic resonance (EPR) spectroscopy and spin trapping* [70]. Such techniques are not commonly used in the andrology laboratories because they are more complicated and not readily available to common andrology laboratories [86; 87]. In the following chapter, we have standardized a direct method to measure nitric oxide in human spermatozoa using flow cytometry.

4.12 Conclusion

Nitric oxide synthases, mainly the constitutive isoforms, are present in human spermatozoa. Physiologic levels of NO are vital for human sperm function and physiology, mainly for adequate flagellar beating and progressive motion. The effects of NO on human spermatozoa could be direct or indirect (mediated by peroxynitrite and/or its derivatives). Decreased levels of NO impair the function of spermatozoa. Likewise, excessive levels of NO contribute to oxidative stress and are detrimental to sperm quality. Although the evidence cited in this chapter shows that NO is very important for human sperm quality such as adequate sperm motility, many of its roles and mechanisms of action in sperm are yet to be fully unveiled.

4.13 References

- [1]V.D. Dixit, and N. Parvizi, Nitric oxide and the control of reproduction. *Anim Reprod Sci* 65 (2001) 1-16.
- [2]J.S. Stamler, D.J. Singel, and J. Loscalzo, Biochemistry of nitric oxide and its redox-activated forms. *Science* 258 (1992) 1898-902.
- [3]P.A. Bush, W.J. Aronson, G.M. Buga, J. Rajfer, and L.J. Ignarro, Nitric oxide is a potent relaxant of human and rabbit corpus cavernosum. *J Urol* 147 (1992) 1650-5.
- [4]J. Rajfer, W.J. Aronson, P.A. Bush, F.J. Dorey, and L.J. Ignarro, Nitric oxide as a mediator of relaxation of the corpus cavernosum in response to nonadrenergic, noncholinergic neurotransmission. *N Engl J Med* 326 (1992) 90-4.
- [5]L.J. Ignarro, Mechanism of action of nitric oxide: heme-dependent activation of guanylate cyclase represents a unifying signal transduction mechanism. *Jpn J Pharmacol* 58 Suppl 2 (1992) 207P-212P.
- [6]A.F. Varela, A. Runge, L.J. Ignarro, and G. Chaudhuri, Nitric oxide and prostacyclin inhibit fetal platelet aggregation: a response similar to that observed in adults. *Am J Obstet Gynecol* 167 (1992) 1599-604.
- [7]U. Simonsen, D. Prieto, J.A. Delgado, M. Hernandez, L. Resel, I. Saenz de Tejada, and A. Garcia-Sacristan, Nitric oxide is involved in the inhibitory neurotransmission and endothelium-dependent relaxations of human small penile arteries. *Clin Sci (Lond)* 92 (1997) 269-75.

- [8]M.G. Belvisi, J.K. Ward, J.A. Mitchell, and P.J. Barnes, Nitric oxide as a neurotransmitter in human airways. *Arch Int Pharmacodyn Ther* 329 (1995) 97-110.
- [9]S.M. McCann, C. Mastronardi, A. Walczewska, S. Karanth, V. Rettori, and W.H. Yu, The role of nitric oxide in reproduction. *Braz J Med Biol Res* 32 (1999) 1367-79.
- [10]S.M. McCann, and V. Rettori, The role of nitric oxide in reproduction. *Proc Soc Exp Biol Med* 211 (1996) 7-15.
- [11]R.G. Knowles, and S. Moncada, Nitric oxide synthases in mammals. *Biochem J* 298 (Pt 2) (1994) 249-58.
- [12]W.K. Alderton, C.E. Cooper, and R.G. Knowles, Nitric oxide synthases: structure, function and inhibition. *Biochem J* 357 (2001) 593-615.
- [13]S.R. Paludan, L. Malmgaard, S. Ellermann-Eriksen, L. Bosca, and S.C. Mogensén, Interferon (IFN)-gamma and Herpes simplex virus/tumor necrosis factor-alpha synergistically induce nitric oxide synthase 2 in macrophages through cooperative action of nuclear factor-kappa B and IFN regulatory factor-1. *Eur Cytokine Netw* 12 (2001) 297-308.
- [14]R.A. Shafer, and S. Murphy, Activated astrocytes induce nitric oxide synthase-2 in cerebral endothelium via tumor necrosis factor alpha. *Glia* 21 (1997) 370-9.
- [15]F. Lampiao, and S.S. du Plessis, Insulin and leptin enhance human sperm motility, acrosome reaction and nitric oxide production. *Asian J Androl* 10 (2008) 799-807.
- [16]C. Szabo, H. Ischiropoulos, and R. Radi, Peroxynitrite: biochemistry, pathophysiology and development of therapeutics. *Nat Rev Drug Discov* 6 (2007) 662-80.

- [17]L.M. Villa, E. Salas, V.M. Darley-Usmar, M.W. Radomski, and S. Moncada, Peroxynitrite induces both vasodilatation and impaired vascular relaxation in the isolated perfused rat heart. *Proc Natl Acad Sci U S A* 91 (1994) 12383-7.
- [18]H. Ischiropoulos, L. Zhu, J. Chen, M. Tsai, J.C. Martin, C.D. Smith, and J.S. Beckman, Peroxynitrite-mediated tyrosine nitration catalyzed by superoxide dismutase. *Arch Biochem Biophys* 298 (1992) 431-7.
- [19]L.A. MacMillan-Crow, J.P. Crow, and J.A. Thompson, Peroxynitrite-mediated inactivation of manganese superoxide dismutase involves nitration and oxidation of critical tyrosine residues. *Biochemistry* 37 (1998) 1613-22.
- [20]B. Alvarez, V. Demicheli, R. Duran, M. Trujillo, C. Cervenansky, B.A. Freeman, and R. Radi, Inactivation of human Cu,Zn superoxide dismutase by peroxynitrite and formation of histidinyl radical. *Free Radic Biol Med* 37 (2004) 813-22.
- [21]S.N. Savvides, M. Scheiwein, C.C. Bohme, G.E. Arteel, P.A. Karplus, K. Becker, and R.H. Schirmer, Crystal structure of the antioxidant enzyme glutathione reductase inactivated by peroxynitrite. *J Biol Chem* 277 (2002) 2779-84.
- [22]A. Van der Vliet, D. Smith, C.A. O'Neill, H. Kaur, V. Darley-Usmar, C.E. Cross, and B. Halliwell, Interactions of peroxynitrite with human plasma and its constituents: oxidative damage and antioxidant depletion. *Biochem J* 303 (Pt 1) (1994) 295-301.
- [23]S. Bartsaghi, V. Valez, M. Trujillo, G. Peluffo, N. Romero, H. Zhang, B. Kalyanaraman, and R. Radi, Mechanistic studies of peroxynitrite-mediated tyrosine nitration in membranes using the hydrophobic probe N-t-BOC-L-tyrosine tert-butyl ester. *Biochemistry* 45 (2006) 6813-25.

- [24]K.P. Moore, V. Darley-Usmar, J. Morrow, and L.J. Roberts, 2nd, Formation of F2-isoprostanes during oxidation of human low-density lipoprotein and plasma by peroxynitrite. *Circ Res* 77 (1995) 335-41.
- [25]C. Batthyany, F.J. Schopfer, P.R. Baker, R. Duran, L.M. Baker, Y. Huang, C. Cervenansky, B.P. Branchaud, and B.A. Freeman, Reversible post-translational modification of proteins by nitrated fatty acids in vivo. *J Biol Chem* 281 (2006) 20450-63.
- [26]M.M. Wright, F.J. Schopfer, P.R. Baker, V. Vidyasagar, P. Powell, P. Chumley, K.E. Iles, B.A. Freeman, and A. Agarwal, Fatty acid transduction of nitric oxide signaling: nitrolinoleic acid potently activates endothelial heme oxygenase 1 expression. *Proc Natl Acad Sci U S A* 103 (2006) 4299-304.
- [27]J.C. Niles, J.S. Wishnok, and S.R. Tannenbaum, Peroxynitrite-induced oxidation and nitration products of guanine and 8-oxoguanine: structures and mechanisms of product formation. *Nitric Oxide* 14 (2006) 109-21.
- [28]L.J. Kennedy, K. Moore, Jr., J.L. Caulfield, S.R. Tannenbaum, and P.C. Dedon, Quantitation of 8-oxoguanine and strand breaks produced by four oxidizing agents. *Chem Res Toxicol* 10 (1997) 386-92.
- [29]P. Jagtap, and C. Szabo, Poly(ADP-ribose) polymerase and the therapeutic effects of its inhibitors. *Nat Rev Drug Discov* 4 (2005) 421-40.
- [30]S. Mohr, J.S. Stamler, and B. Brune, Mechanism of covalent modification of glyceraldehyde-3-phosphate dehydrogenase at its active site thiol by nitric oxide, peroxynitrite and related nitrosating agents. *FEBS Lett* 348 (1994) 223-7.

- [31]B. Blanchard-Fillion, J.M. Souza, T. Friel, G.C. Jiang, K. Vrana, V. Sharov, L. Barron, C. Schoneich, C. Quijano, B. Alvarez, R. Radi, S. Przedborski, G.S. Fernando, J. Horwitz, and H. Ischiropoulos, Nitration and inactivation of tyrosine hydroxylase by peroxynitrite. *J Biol Chem* 276 (2001) 46017-23.
- [32]E.A. Konorev, N. Hogg, and B. Kalyanaraman, Rapid and irreversible inhibition of creatine kinase by peroxynitrite. *FEBS Lett* 427 (1998) 171-4.
- [33]A.K. Grover, C.Y. Kwan, and S.E. Samson, Effects of peroxynitrite on sarco/endoplasmic reticulum Ca²⁺ pump isoforms SERCA2b and SERCA3a. *Am J Physiol Cell Physiol* 285 (2003) C1537-43.
- [34]P. Pacher, J.S. Beckman, and L. Liaudet, Nitric oxide and peroxynitrite in health and disease. *Physiol Rev* 87 (2007) 315-424.
- [35]A. van der Vliet, M. Hristova, C.E. Cross, J.P. Eiserich, and T. Goldkorn, Peroxynitrite induces covalent dimerization of epidermal growth factor receptors in A431 epidermoid carcinoma cells. *J Biol Chem* 273 (1998) 31860-6.
- [36]M. Zou, A. Yesilkaya, and V. Ullrich, Peroxynitrite inactivates prostacyclin synthase by heme-thiolate-catalyzed tyrosine nitration. *Drug Metab Rev* 31 (1999) 343-9.
- [37]E. Paxinou, Q. Chen, M. Weisse, B.I. Giasson, E.H. Norris, S.M. Rueter, J.Q. Trojanowski, V.M. Lee, and H. Ischiropoulos, Induction of alpha-synuclein aggregation by intracellular nitrative insult. *J Neurosci* 21 (2001) 8053-61.
- [38]S. Milstien, and Z. Katusic, Oxidation of tetrahydrobiopterin by peroxynitrite: implications for vascular endothelial function. *Biochem Biophys Res Commun* 263 (1999) 681-4.

- [39]R. Middendorff, D. Muller, S. Wichers, A.F. Holstein, and M.S. Davidoff, Evidence for production and functional activity of nitric oxide in seminiferous tubules and blood vessels of the human testis. *J Clin Endocrinol Metab* 82 (1997) 4154-61.
- [40]M. Rosselli, R.K. Dubey, B. Imthurn, E. Macas, and P.J. Keller, Effects of nitric oxide on human spermatozoa: evidence that nitric oxide decreases sperm motility and induces sperm toxicity. *Hum Reprod* 10 (1995) 1786-90.
- [41]M.G. Yang, Y. Yang, P. Huang, X.K. Hao, S.L. Zheng, Z.Y. Zhang, and G.Z. Rao, [Relationship between the amount of nitric oxide and apoptosis of germ cells in the semen of infertile men]. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi* 21 (2005) 510-2.
- [42]T. Nobunaga, Y. Tokugawa, K. Hashimoto, Y. Kubota, K. Sawai, T. Kimura, K. Shimoya, M. Takemura, N. Matsuzaki, C. Azuma, and F. Saji, Elevated nitric oxide concentration in the seminal plasma of infertile males: nitric oxide inhibits sperm motility. *Am J Reprod Immunol* 36 (1996) 193-7.
- [43]J.B. Weinberg, E. Doty, J. Bonaventura, and A.F. Haney, Nitric oxide inhibition of human sperm motility. *Fertil Steril* 64 (1995) 408-13.
- [44]J. Ji, Y. Zhao, and G. Chen, [Nitric oxide and nitric oxide synthase related to male reproduction]. *Wei Sheng Yan Jiu* 36 (2007) 636-9.
- [45]G. Machado-Oliveira, L. Lefievre, C. Ford, M.B. Herrero, C. Barratt, T.J. Connolly, K. Nash, A. Morales-Garcia, J. Kirkman-Brown, and S. Publicover, Mobilisation of Ca²⁺ stores and flagellar regulation in human sperm by S-nitrosylation: a role for NO synthesised in the female reproductive tract. *Development* 135 (2008) 3677-86.

- [46]Y. Xu, Q.Y. Xu, B.H. Yang, X.M. Zhu, and Y.F. Peng, [Relationship of nitric oxide and nitric oxide synthase with varicocele infertility]. *Zhonghua Nan Ke Xue* 14 (2008) 414-7.
- [47]E. Buldreghini, R.Z. Mahfouz, A. Vignini, L. Mazzanti, G. Ricciardo-Lamonica, A. Lenzi, A. Agarwal, and G. Balercia, Single nucleotide polymorphism (SNP) of the endothelial nitric oxide synthase (eNOS) gene (Glu298Asp variant) in infertile men with asthenozoospermia. *J Androl* 31 482-8.
- [48]H. Zhang, Q.M. Zhou, X.D. Li, Y. Xie, X. Duan, F.L. Min, B. Liu, and Z.G. Yuan, Ginsenoside R(e) increases fertile and asthenozoospermic infertile human sperm motility by induction of nitric oxide synthase. *Arch Pharm Res* 29 (2006) 145-51.
- [49]I. Huang, J. Jones, and O. Khorram, Human seminal plasma nitric oxide: correlation with sperm morphology and testosterone. *Med Sci Monit* 12 (2006) CR103-6.
- [50]T. Ramya, M.M. Misro, D. Sinha, D. Nandan, and S. Mithal, Altered levels of seminal nitric oxide, nitric oxide synthase, and enzymatic antioxidants and their association with sperm function in infertile subjects. *Fertil Steril* 95 135-40.
- [51]M.K. O'Bryan, A. Zini, C.Y. Cheng, and P.N. Schlegel, Human sperm endothelial nitric oxide synthase expression: correlation with sperm motility. *Fertil Steril* 70 (1998) 1143-7.
- [52]A. Zini, M.K. O'Bryan, M.S. Magid, and P.N. Schlegel, Immunohistochemical localization of endothelial nitric oxide synthase in human testis, epididymis, and vas deferens suggests a possible role for nitric oxide in spermatogenesis, sperm maturation, and programmed cell death. *Biol Reprod* 55 (1996) 935-41.

- [53]S.E. Lewis, E.T. Donnelly, E.S. Sterling, M.S. Kennedy, W. Thompson, and U. Chakravarthy, Nitric oxide synthase and nitrite production in human spermatozoa: evidence that endogenous nitric oxide is beneficial to sperm motility. *Mol Hum Reprod* 2 (1996) 873-8.
- [54]E.T. Donnelly, S.E. Lewis, W. Thompson, and U. Chakravarthy, Sperm nitric oxide and motility: the effects of nitric oxide synthase stimulation and inhibition. *Mol Hum Reprod* 3 (1997) 755-62.
- [55]G. Balercia, S. Moretti, A. Vignini, M. Magagnini, F. Mantero, M. Boscaro, G. Ricciardo-Lamonica, and L. Mazzanti, Role of nitric oxide concentrations on human sperm motility. *J Androl* 25 (2004) 245-9.
- [56]F. Francavilla, R. Santucci, B. Macerola, G. Ruvolo, and R. Romano, Nitric oxide synthase inhibition in human sperm affects sperm-oocyte fusion but not zona pellucida binding. *Biol Reprod* 63 (2000) 425-9.
- [57]C. Roessner, U. Paasch, H.J. Glander, and S. Grunewald, Activity of nitric oxide synthase in mature and immature human spermatozoa. *Andrologia* 42 132-7.
- [58]A. Zini, M.K. O'Bryan, and P.N. Schlegel, Nitric oxide synthase activity in human seminal plasma. *Urology* 58 (2001) 85-9.
- [59]A. Revelli, L. Bergandi, M. Massobrio, B. Lindblom, A. Bosia, and D. Ghigo, The concentration of nitrite in seminal plasma does not correlate with sperm concentration, sperm motility, leukocytospermia, or sperm culture. *Fertil Steril* 76 (2001) 496-500.
- [60]S. Lambard, I. Galeraud-Denis, G. Martin, R. Levy, A. Chocat, and S. Carreau, Analysis and significance of mRNA in human ejaculated sperm from

normozoospermic donors: relationship to sperm motility and capacitation. *Mol Hum Reprod* 10 (2004) 535-41.

- [61]P.C. Chiu, S. Liao, K.K. Lam, F. Tang, J.C. Ho, P.C. Ho, W.S. O, Y.Q. Yao, and W.S. Yeung, Adrenomedullin regulates sperm motility and oviductal ciliary beat via cyclic adenosine 5'-monophosphate/protein kinase A and nitric oxide. *Endocrinology* 151 3336-47.
- [62]S. Srivastava, and A. Agarwal, Effect of anion channel blockers on L-arginine action in spermatozoa from asthenospermic men. *Andrologia* 42 76-82.
- [63]D.C. Ding, Y.C. Huang, J.Y. Liu, and G.J. Wu, Comparison of nitric oxide production and motion characteristics after 3-layer percoll and IxaPrep preparation methods of human sperm. *Arch Gynecol Obstet* 266 (2002) 210-3.
- [64]E. Miraglia, F. De Angelis, E. Gazzano, H. Hassanpour, A. Bertagna, E. Aldieri, A. Revelli, and D. Ghigo, Nitric oxide stimulates human sperm motility via activation of the cyclic GMP/protein kinase G signaling pathway. *Reproduction* 141 (2011) 47-54.
- [65]A. Revelli, C. Costamagna, F. Moffa, E. Aldieri, S. Ochetti, A. Bosia, M. Massobrio, B. Lindblom, and D. Ghigo, Signaling pathway of nitric oxide-induced acrosome reaction in human spermatozoa. *Biol Reprod* 64 (2001) 1708-12.
- [66]K. Bedu-Addo, S. Costello, C. Harper, G. Machado-Oliveira, L. Lefievre, C. Ford, C. Barratt, and S. Publicover, Mobilisation of stored calcium in the neck region of human sperm--a mechanism for regulation of flagellar activity. *Int J Dev Biol* 52 (2008) 615-26.

- [67]M.B. Herrero, E. de Lamirande, and C. Gagnon, Nitric oxide regulates human sperm capacitation and protein-tyrosine phosphorylation in vitro. *Biol Reprod* 61 (1999) 575-81.
- [68]A. Zini, E. De Lamirande, and C. Gagnon, Low levels of nitric oxide promote human sperm capacitation in vitro. *J Androl* 16 (1995) 424-31.
- [69]E. de Lamirande, and G. Lamothe, Reactive oxygen-induced reactive oxygen formation during human sperm capacitation. *Free Radic Biol Med* 46 (2009) 502-10.
- [70]M. Belen Herrero, S. Chatterjee, L. Lefievre, E. de Lamirande, and C. Gagnon, Nitric oxide interacts with the cAMP pathway to modulate capacitation of human spermatozoa. *Free Radic Biol Med* 29 (2000) 522-36.
- [71]M.B. Herrero, E. de Lamirande, and C. Gagnon, Tyrosine nitration in human spermatozoa: a physiological function of peroxynitrite, the reaction product of nitric oxide and superoxide. *Mol Hum Reprod* 7 (2001) 913-21.
- [72]J. Thundathil, E. de Lamirande, and C. Gagnon, Nitric oxide regulates the phosphorylation of the threonine-glutamine-tyrosine motif in proteins of human spermatozoa during capacitation. *Biol Reprod* 68 (2003) 1291-8.
- [73]H. Zhang, Q. Zhou, X. Li, W. Zhao, Y. Wang, H. Liu, and N. Li, Ginsenoside Re promotes human sperm capacitation through nitric oxide-dependent pathway. *Mol Reprod Dev* 74 (2007) 497-501.
- [74]E. de Lamirande, and G. Lamothe, Levels of semenogelin in human spermatozoa decrease during capacitation: involvement of reactive oxygen species and zinc. *Hum Reprod* 25 1619-30.

- [75]K. Sengoku, K. Tamate, T. Yoshida, Y. Takaoka, T. Miyamoto, and M. Ishikawa, Effects of low concentrations of nitric oxide on the zona pellucida binding ability of human spermatozoa. *Fertil Steril* 69 (1998) 522-7.
- [76]S.S. du Plessis, D.A. McAllister, A. Luu, J. Savia, A. Agarwal, and F. Lampiao, Effects of H₂O₂ exposure on human sperm motility parameters, reactive oxygen species levels and nitric oxide levels. *Andrologia* 42 206-10.
- [77]D. Mitropoulos, G. Deliconstantinos, A. Zervas, V. Villiotou, C. Dimopoulos, and J. Stavrides, Nitric oxide synthase and xanthine oxidase activities in the spermatic vein of patients with varicocele: a potential role for nitric oxide and peroxynitrite in sperm dysfunction. *J Urol* 156 (1996) 1952-8.
- [78]M. Rodriguez Pena, L. Alescio, A. Russell, J. Lourenco da Cunha, G. Alzu, and E. Bardoneschi, Predictors of improved seminal parameters and fertility after varicocele repair in young adults. *Andrologia* 41 (2009) 277-81.
- [79]D.M. Perera, M. Katz, S.R. Heenbanda, and S. Marchant, Nitric oxide synthase inhibitor NG-monomethyl-L-arginine preserves sperm motility after swim-up. *Fertil Steril* 66 (1996) 830-3.
- [80]T.R. Green, J.H. Fellman, and D.P. Wolf, Human spermicidal activity of inorganic and organic oxidants. *Fertil Steril* 76 (2001) 157-62.
- [81]I. Amiri, N. Sheikh, and R. Najafi, Nitric oxide level in seminal plasma and its relation with sperm DNA damages. *Iran Biomed J* 11 (2007) 259-64.
- [82]F. Lampiao, and S.S. du Plessis, TNF-alpha and IL-6 affect human sperm function by elevating nitric oxide production. *Reprod Biomed Online* 17 (2008) 628-31.

- [83]S.S. du Plessis, K. Hagenaar, and F. Lampiao, The in vitro effects of melatonin on human sperm function and its scavenging activities on NO and ROS. *Andrologia* 42 112-6.
- [84]W.R. Tracey, J. Linden, M.J. Peach, and R.A. Johns, Comparison of spectrophotometric and biological assays for nitric oxide (NO) and endothelium-derived relaxing factor (EDRF): nonspecificity of the diazotization reaction for NO and failure to detect EDRF. *J Pharmacol Exp Ther* 252 (1990) 922-8.
- [85]R.W. Nims, J.C. Cook, M.C. Krishna, D. Christodoulou, C.M. Poore, A.M. Miles, M.B. Grisham, and D.A. Wink, Colorimetric assays for nitric oxide and nitrogen oxide species formed from nitric oxide stock solutions and donor compounds. *Methods Enzymol* 268 (1996) 93-105.
- [86]H. Strijdom, C. Muller, and A. Lochner, Direct intracellular nitric oxide detection in isolated adult cardiomyocytes: flow cytometric analysis using the fluorescent probe, diaminofluorescein. *J Mol Cell Cardiol* 37 (2004) 897-902.
- [87]J.F. Leikert, T.R. Rathel, C. Muller, A.M. Vollmar, and V.M. Dirsch, Reliable in vitro measurement of nitric oxide released from endothelial cells using low concentrations of the fluorescent probe 4,5-diaminofluorescein. *FEBS Lett* 506 (2001) 131-4.

CHAPTER V

A NEW METHOD FOR MEASUREMENT OF NITRIC OXIDE PRODUCTION IN HUMAN SPERMATOOZOA USING FLOW CYTOMETRY

Summary

Studies showed that nitric oxide (NO) has a major contribution in male reproduction. However, many of NO's mechanisms of action in sperm remain unclear; one important reason is that methods for direct NO measurement at the level single cells and/or collections of cells are not straightforward. Here we demonstrated a method to measure intracellular NO in human sperm by means of flow cytometry using NO-specific probe 4,5-diaminofluorescein-2-diacetate (DAF-2DA). Motile sperm are freshly prepared using a modified direct swim-up procedure. Both sperm and DAF-2DA concentrations are standardized. Sperm samples incubated with L-nitro-arginine methyl ester (L-NAME), a Nitric Oxide Synthase (NOS) inhibitor, showed lower fluorescence in a dose-dependent manner. Sperm samples supplemented with Diethylamine NONOate (DEA NONOate), a nitric oxide donor, showed higher fluorescence compared to the control. At last, we used this method to measure the effect of L-carnitine, a motility stimulant as we

have showed in the third chapter, on NO production in human spermatozoa. We found that LC may increases sperm motility via increasing the production of NO.

5.1 Introduction

Nitric oxide (NO) is one of the most studied molecules in biomedicine because of its various physiological functions. As we have reported in the previous chapter, it has been referred to as an endothelium-derived relaxing factor because it causes vasodilation via relaxing vascular smooth muscle [1; 2; 3]. In addition, it prevents platelets aggregation [4], acts as a neurotransmitter [5; 6], and mediates macrophage bactericidal activity [7].

Reports have shown that NO has a significant involvement in male reproduction [8; 9]. Various studies showed that endogenous NO is very important for sperm motion [10- 14]. NO generation was found to be crucial in gamete interaction and fertilization route [15]. A number of studies reported that NO is involved in human sperm capacitation [16; 17; 18]. In contrast, others have shown that higher levels of nitric oxide were observed in infertile men compared to control groups [19].

Although the evidence above points to the fact that NO is very important for human sperm physiology and function, its exact mechanisms of action in sperm are yet to be fully understood. The majority of measurement of NO in sperm relied on nitrite determinations using colorimetric assays. This may not relay an accurate picture on NO concentration and its fluctuations in spermatozoa. Methods that have the potential to follow NO concentration under varying conditions within cells are of particular importance to investigate the mechanisms of actions of NO in sperm function.

Analysis by flow cytometry has entered many andrology laboratories and has been used to evaluate sperm count and viability [20], to detect antisperm antibody [21; 22] and to study the chromatin structure of sperm [23]. The considerable advantage of flow cytometry is that millions of cells can be analyzed in seconds, providing a statistically more precise assessment of the level of analyte under consideration. Here, we demonstrated a method to detect the intracellular NO in human sperm using the NO-specific probe 4,5-diaminofluorescein-2-diacetate (DAF-2DA). The specific detection of NO and the sensitivity of fluorescence are coupled to the high throughput capability of flow cytometry; this provides a specific, sensitive, and fast method of monitor NO concentration in spermatozoa under varying experimental conditions.

5.2 Materials and methods

Sample collection

Our study was approved by the institutional review board of Cleveland Clinic Hospital. Normospermic men were selected and analyzed based on normal semen analysis according to the World Health Organization (WHO) (1999) guidelines (section 2.2.1). Fresh semen samples were used to standardize the measurement of intracellular NO in sperm using DAF-2DA by means of flow cytometry.

Samples were collected by masturbation following a minimum of 48 to 72 hours of sexual abstinence. Following liquefaction at 37 °C for 20 minutes, semen specimens were evaluated for volume, sperm concentration, total cell count, motility and morphology. A total amount of 5µL of the sample was used for evaluation of

concentration and motility using a microcell slide chamber (Conception Technologies, San Diego, CA).

Preparation of motile sperm samples

Motile sperm were retrieved from semen samples using direct swim-up technique. This procedure was adapted from the one described by World Health Organization (1999) (section 2.2.1). Briefly, spermatozoa are selected based on the ability to swim out of seminal plasma into the human tubal fluid (HTF). We use only untreated, fresh semen in this process. 0.8 ml of homogenized semen was pipetted carefully under 1 ml of HTF placed in a 15 ml conical centrifuge tube while avoiding the mixing between the semen and the supplemented medium. Afterward, the tube was left at 90° angle for 1 hr at 37°C. The uppermost (~ 0.6 – 0.8 µL) medium containing only the highly motile sperm is transferred to another test tube, and centrifuged for 4 min at 1600 rpm. Then, the top layer was discarded, and the sperm in the bottom resuspended in 0.5 mL HTF. The resulting motile sperm are washed and their concentration adjusted as needed.

DAF-2DA preparation

DAF-2DA was dissolved in dimethyl sulfoxide (DMSO) because it has a very low solubility in water. Although DMSO is a very suitable solvent for DAF-2DA, it is considered toxic to human sperm when added at higher concentrations [24]. Therefore, in our experimental design, we added DAF-2DA/DMSO solution to sperm medium at a concentration of $\leq 5\%$ (v/v). We did not see a statistical difference in sperm viability after adding DMSO of $\leq 5\%$ (v/v) compared to sperm without DMSO (Figure 6-5). We

run our experiments in the dark conditions in order to avoid the exposure of DAF-2DA to the light, because fluorescein can be easily photobleached, a photochemical destruction of the fluorophore [25] .

Measurement of DAF-2T fluorescence at various sperm concentrations

The concentration of washed motile sperm is adjusted with HTF to within 0.5- 8.0 x 10⁶/ mL range. The samples (n=6) are then supplemented with 20 µM of DAF-2DA, incubated for 90 min at 37 °C in dark, and analyzed with flow cytometry. Identical samples without addition of DAF-2DA are used as controls.

Measurement of DAF-2T fluorescence at various DAF-2DA concentrations

Samples (n=6) of 2 x 10⁶/mL sperm concentration are incubated with various concentrations (0.5 nM, 5 µM, 50 µM, 0.5 mM and 1 mM) of DAF-2DA for 90 min at 37 °C in dark. After incubation, the fluorescence of DAF-2T was measured using flow cytometry along with control samples.

Measurement of sperm viability

Sperm viability was measured by the Eosin-Nigrosin staining procedure (see page 25, chapter II)

NO measurement in sperm after NOS inhibition

Sperm samples (n=8) at 2 x 10⁶/mL concentration are incubated at 37°C with various concentrations (100, 400 and 800 µM) of the NOS inhibitor N^w-nitro -L-arginine

methyl ester (L-NAME) for 120 min in the dark. 15 min after the addition of L-NAME, DAF-2DA is added to the samples to a final concentration of 20 μ M. The fluorescence of DAF-2T is measured immediately after incubation using flow cytometry. Samples without incubation with L-NAME are taken as controls for this series of experiments.

NO measurement in the presence of an exogenous NO donor.

Sperm samples (n=6) at 2×10^6 /mL concentration were incubated at 37°C with various concentrations (10, 100, 300, 500 and 800 μ M) of the NO donor 2-(N,N-diethylamino)-diazene-2-oxide, sodium salt (DEA-NONOate) for 45 min in dark. DAF-2DA at 20 μ M is added to the samples 45 min prior to the addition of the NO donor. The fluorescence of DAF-2T was measured immediately after incubation using flow cytometry. Samples in absence of NONOate are taken as controls.

Flow cytometry

Flow cytometry data acquisition is completed within 10-15 min from sample preparation. The flow cytometer (FACScan; Becton Dickinson, San Jose, CA) is equipped with a 515-nm argon laser as a light source. Samples are adjusted to have 20,000 cells examined for each assay at a flow rate of 100 cells/second. Data analysis is performed using FlowJo v4.4.4 software (Tree Star Inc., Ashland, OR).

In our flow cytometry data, we used 20,000 count from the 2×10^6 /mL total sperm in each sample to obtain consistent measurements. The threshold value of the forward scatter (FSC) was set < 40 FSC to exclude small cells, cell fragments, and debris from the analyzed sperm. The sperm window was established after running multiple

unstained sperm samples (instrumental control) submitted to swim up and followed by washing with HTF.

Statistical analysis

All values were recorded as mean \pm S.D. Differences between samples and their control were considered significant at $P < 0.05$. Statistical analyses were performed by one-way analysis of variance (ANOVA) and an unpaired Student's *t*-test using the SPSS/PC computer software (SPSS 10.0.7, SPSS Inc.)

5.3 Results and discussion

The work we presented here is a modification of what Strijdom *et al.* (2004) and S. du Plessis *et al.* (2006) have described [26; 27]. The fluorescent NO-indicators diaminofluoresceins (DAFs) were first designed and synthesized by Kojima *et al.* (1998) [25]. DAF-2DA, a non-fluorescent dye, is able to cross the membrane of live cells and is hydrolyzed to 4,5-diaminofluorescein (DAF-2) by intracellular esterases [28]. DAF-2 reacts rapidly with NO (in the presence of O₂; the reaction is actually with an NO⁺ equivalent likely from a nitrogen oxide derivative of NO) to form the fluorescent dye triazolofluorescein (DAF-2T) (Figure 5-1). DAF-2T fluoresces at 515 nm upon excitation at 495 nm. Because of the specificity of the reaction of DAF-2 with NO, the intensity of the fluorescence is proportional to the amount of intracellular NO generated.




Figure 5-1: Schematic representation of fluorometric NO measurement procedure for human spermatozoa using the DAF-2DA precursor.

Effect of sperm concentrations on DAF-2T fluorescence at 20 μ M DAF-2DA

To determine the optimum sperm concentration of samples used in our flow cytometry measurements, we monitored DAF-2T fluorescence as a function sperm concentration for a set concentration of the dye DAF-2DA at 20 μ M (Figure 5-2). No significant difference ($P > 0.05$; t -test) in average fluorescence is seen between sample with sperm concentrations 0.5, 1.0 and 2.0 $\times 10^6$ /mL. On the other hand, a significant difference ($P < 0.05$; t -test) in average fluorescence is observed at higher concentrations (4 and 8 $\times 10^6$ /mL). Unstained control samples (i.e. 0.0 μ M DAF-2DA) exhibit very negligible fluorescence.

Our experiments indicate that concentrations in the range of 0.5-2.0 $\times 10^6$ /mL could all be used for a set DAF-2DA concentration of 20 μ M. Sample with higher sperm concentrations such as 4x or 8x10⁶/mL showed lower fluorescence, likely a direct result of a lower ratio of DAF-2DA molecules per sperm cell.




Figure 5-2: Observed fluorescence as a function of sperm concentration with 20 μM dye concentration. The values reported are averages \pm standard deviations; $n=6$. The control is the unstained sample.

Effect of DAF-2DA concentrations on observed fluorescence

Figure 5-3 shows flow cytometry histograms of DAF-2DT fluorescence (A) as well as the mean areas for these histograms (B) at various concentrations of DAF-2DA (0.5 nM -1 mM). All DAF-2DA concentrations used show a statistical increase ($P < 0.05$; t -test) in fluorescence compared to unstained control samples (without DAF-2DA). Sperm samples incubated with 50 μ M of DAF-2DA show the highest intensity of DAF-2T fluorescence, while the lowest fluorescence is seen for 0.5 nM dye concentration. The intensity of the fluorescence decreased at higher doses (above 0.5 mM) of DAF-2DA used.

Our results showed a statistically significant increase in fluorescence even at a low concentration of DAF-2DA of ~ 0.5 nM compared to the control sample (without DAF-2DA). Because concentrations of 0.5 mM and 1 mM of DAF-2DA exhibited lower fluorescence compared to 50 μ M, we examined if sperm viability is compromised at these higher concentrations. Indeed, we find that the millimolar range of dye concentration significantly affects sperm viability, which explains the observed decrease in fluorescence (Figure 5-4).




Figure 5-3: (A) Flow cytometry histograms of DAF-2T fluorescence as a function of concentrations of the DAF-2DA dye (0.5nM, 5µM, 50µM, 0.5mM, 1mM); (B) Mean areas of DAF-2T fluorescence at the various DAF-2DA doses; data are means (for n=6) ± standard deviations; the control is the unstained sample (i.e. no DAF-2DA). Sperm concentration used in the sample was 2×10^6 /mL with 20,000 cells acquired in flow cytometry.




Figure 5-4: Effect of various DAF-2DA concentrations on sperm viability as evaluated by the Eosin-Nigrosin test. The evaluation of sperm viability performed immediately after the flow cytometry measurements. Data are means (n=6) \pm standard deviations.

Dynamic NO sensitivity of DAF-2DA in the presence of NOS inhibitor and an NO donor.

To determine the dynamic sensitivity of this fluorometric method to sudden variations of intracellular NO concentration in spermatozoa, we monitored fluorescence changes in response to an NO donor as well as in the presence of a NOS-inhibitor.

a. Effect of the NOS inhibitor L-NAME on observed fluorescence

Figure 5-5 shows the effect of various concentrations (100, 400, and 800 μM) of L-NAME on the observed DAF-2T fluorescence. Data represented are mean areas of fluorescence \pm standard deviations. The figure shows a significant decrease ($P < 0.05$) in the intensity of fluorescence of DAF-2T in the presence of L-NAME in a dose dependent manner. The fluorescence decrease of DAF-2T did not significantly change for L-NAME concentrations between 100 and 400 μM ($P > 0.05$; *t*-test).

b. DAF-2T fluorescence in response to varying concentration of the NO donor

The specificity of DAF-2DA to measure NO and to respond to increases of intracellular NO concentration was validated using incubation with DEA-NONOate, a known NO donor.

Figure 5-6 shows the effect of various concentrations of DEA-NONOate, the NO releasing compound, on observed fluorescence in human sperm. We observed significant increase in fluorescence in samples with 100 μM NO donor concentration while the change was not significant with 10 μM . Also, the mean area of DAF-2T fluorescence did not increase significantly above the level measured with 100 μM NONOate for concentrations above 300 μM .




Figure 5-5: DAF-2T flow cytometry fluorescence in human sperm as a function of L-NAME concentration. Data are means (n=10) \pm standard deviation.




Figure 5-6: DAF-2T flow cytometry fluorescence in human sperm as a function of NONOate concentration. Data are means (n=6) \pm standard deviations.

Our results show that the fluorescence of DAF-2T increases after adding the NO donor, indicating that the method follows dynamic increases of intracellular NO. However, the method shows that the fluorescence did not increase significantly beyond the level seen for 100 μM of the NO donor. It is known that concentrations of NONOates (in the range of 0.3- 1.0 mM) affect sperm motility [17]. In addition to sperm motility, we suspected that sperm viability might be behind the sluggish increase (for 300 μM) and then decrease of DAF-2 fluorescence (for 500-800 μM). To confirm this, we measured the viability of sperm immediately after the flow cytometry analysis (Figure 5-7). As expected, the figure shows a significant decrease in sperm viability after 100 μM in NO donor. This is not a surprise since literature reports established that NO donors can have both a protective role (at low concentration) but can also induce cell death in a variety of cell types at high concentrations [29; 30]. The cytotoxic effects of NO are mainly due to higher oxides that derive from NO and that are more reactive. Peroxynitrite is one of them and has many reactivity facets under physiologic conditions which contribute to cell dysfunction, including the disruption of the cell's ability to sustain healthy levels of endogenous NO, as well as other vital functions.




Figure 5-7: Effect of varying concentration of NONOate on sperm viability as evaluated by the Eosin-Nigrosin test. The viability test is conducted immediately after the flow cytometry analysis. Sperm concentration was $2 \times 10^6/\text{mL}$. Data are means ($n=6$) \pm standard deviations.

5.4 Conclusion

Collectively, these findings show that measurement of intracellular nitric oxide in human sperm can be performed using the DAF-2DA fluorometric method coupled to flow cytometry. The method has now been standardized for human spermatozoa and will be of assistance to elucidate the NO mechanisms of action in human sperm as well as its role in maintaining sperm quality or in inducing cytotoxicity.

5.5 Effect of L-Carnitine (LC) on NO production in human spermatozoa assessed by the DAF-2DA fluorometric method coupled with flow cytometry

In this section, we measured the effect of L-carnitine (LC) on NO production in human sperm using the method that we have standardized as described in the previous section of this chapter. As we have seen in chapter III, supplementation of sperm with LC at 0.5 mg/mL increased sperm motility. On the other hand, as we described previously, nitric oxide has been implicated for adequate sperm function including motility. Our goal here is to test whether LC supplementation causes the levels of endogenous NO to increase in sperm. This may be a mechanism by which LC stimulates sperm motility.

Methodology and experimental design

Highly motile sperm were prepared using direct swim up and adjusted to 2×10^6 /mL concentration with HTF. After that, Semen Samples (n=4) were supplemented with 0.1 mg/mL LC, followed by addition of DAF-2DA at 20 μ M concentration. Samples were then incubated for 90 min at 37°C in the dark. After incubation, the fluorescence of DAF-2T formed was measured immediately using flow cytometry as described in above.

Results and discussion

In chapter 2, we have shown that 0.5 mg/mL LC concentration is beneficial to sperm at 5 million/mL. Here we supplemented the sperm with LC at 0.1 mg/mL because we treated lower sperm number (2 million/mL). Figure 6-7 shows the effect of 0.1 mg/mL LC on DAF-2T fluorescence in human spermatozoa. The figure shows an increase in DAF-2T fluorescence, and thus an increase in the intracellular NO produced in the sperm supplemented with LC. Control samples assayed without LC supplementation do not show any significant increase in fluorescence. Data in the figure are representative of four independent experiments. As reported in chapter 4, NOS in all of its isoforms is present in human spermatozoa. NO has a major role in the physiology and function of the spermatozoa [10; 31]. A number of studies showed that NO is very important for adequate motility of human spermatozoa [12; 17; 32]. Our results from the fluorometric determination of intracellular NO production in spermatozoa upon incubation with LC show a causal relationship between LC supplementation and NO production. Whether the LC-induced NO production is direct or via an indirect pathway cannot be concluded and awaits further investigation. For now, the evidence shows that LC induces intracellular NO production, and this may explain the enhanced motility of spermatozoa incubated with L-Carnitine.




Figure 5-8: Effect of LC on DAF-2T fluorescence in human spermatozoa. Sperm ($2 \times 10^6/\text{mL}$) incubated 90 min with $20 \mu\text{M}$ DAF-2DA in the presence of 0.1 mg/mL LC (a), and without LC (b). The control histogram is for sperm population without DAF-2DA, and without LC. Data are representative of four independent experiments.

5.6 References

- [1]P.A. Bush, W.J. Aronson, G.M. Buga, J. Rajfer, and L.J. Ignarro, Nitric oxide is a potent relaxant of human and rabbit corpus cavernosum. *J Urol* 147 (1992) 1650-5.
- [2]J. Rajfer, W.J. Aronson, P.A. Bush, F.J. Dorey, and L.J. Ignarro, Nitric oxide as a mediator of relaxation of the corpus cavernosum in response to nonadrenergic, noncholinergic neurotransmission. *N Engl J Med* 326 (1992) 90-4.
- [3]L.J. Ignarro, Mechanism of action of nitric oxide: heme-dependent activation of guanylate cyclase represents a unifying signal transduction mechanism. *Jpn J Pharmacol* 58 Suppl 2 (1992) 207P-212P.
- [4]A.F. Varela, A. Runge, L.J. Ignarro, and G. Chaudhuri, Nitric oxide and prostacyclin inhibit fetal platelet aggregation: a response similar to that observed in adults. *Am J Obstet Gynecol* 167 (1992) 1599-604.
- [5]U. Simonsen, D. Prieto, J.A. Delgado, M. Hernandez, L. Resel, I. Saenz de Tejada, and A. Garcia-Sacristan, Nitric oxide is involved in the inhibitory neurotransmission and endothelium-dependent relaxations of human small penile arteries. *Clin Sci (Lond)* 92 (1997) 269-75.
- [6]M.G. Belvisi, J.K. Ward, J.A. Mitchell, and P.J. Barnes, Nitric oxide as a neurotransmitter in human airways. *Arch Int Pharmacodyn Ther* 329 (1995) 97-110.

- [7]A. Gross, S. Spiesser, A. Terraza, B. Rouot, E. Caron, and J. Dornand, Expression and bactericidal activity of nitric oxide synthase in *Brucella suis*-infected murine macrophages. *Infect Immun* 66 (1998) 1309-16.
- [8]R. Middendorff, D. Muller, S. Wichers, A.F. Holstein, and M.S. Davidoff, Evidence for production and functional activity of nitric oxide in seminiferous tubules and blood vessels of the human testis. *J Clin Endocrinol Metab* 82 (1997) 4154-61.
- [9]M. Rosselli, R.K. Dubey, B. Imthurn, E. Macas, and P.J. Keller, Effects of nitric oxide on human spermatozoa: evidence that nitric oxide decreases sperm motility and induces sperm toxicity. *Hum Reprod* 10 (1995) 1786-90.
- [10]S.E. Lewis, E.T. Donnelly, E.S. Sterling, M.S. Kennedy, W. Thompson, and U. Chakravarthy, Nitric oxide synthase and nitrite production in human spermatozoa: evidence that endogenous nitric oxide is beneficial to sperm motility. *Mol Hum Reprod* 2 (1996) 873-8.
- [11]E.T. Donnelly, S.E. Lewis, W. Thompson, and U. Chakravarthy, Sperm nitric oxide and motility: the effects of nitric oxide synthase stimulation and inhibition. *Mol Hum Reprod* 3 (1997) 755-62.
- [12]M.K. O'Bryan, A. Zini, C.Y. Cheng, and P.N. Schlegel, Human sperm endothelial nitric oxide synthase expression: correlation with sperm motility. *Fertil Steril* 70 (1998) 1143-7.
- [13]G. Balercia, S. Moretti, A. Vignini, M. Magagnini, F. Mantero, M. Boscaro, G. Ricciardo-Lamonica, and L. Mazzanti, Role of nitric oxide concentrations on human sperm motility. *J Androl* 25 (2004) 245-9.

- [14]H. Zhang, Q.M. Zhou, X.D. Li, Y. Xie, X. Duan, F.L. Min, B. Liu, and Z.G. Yuan, Ginsenoside R(e) increases fertile and asthenozoospermic infertile human sperm motility by induction of nitric oxide synthase. *Arch Pharm Res* 29 (2006) 145-51.
- [15]G. Machado-Oliveira, L. Lefievre, C. Ford, M.B. Herrero, C. Barratt, T.J. Connolly, K. Nash, A. Morales-Garcia, J. Kirkman-Brown, and S. Publicover, Mobilisation of Ca²⁺ stores and flagellar regulation in human sperm by S-nitrosylation: a role for NO synthesised in the female reproductive tract. *Development* 135 (2008) 3677-86.
- [16]M.B. Herrero, E. de Lamirande, and C. Gagnon, Nitric oxide regulates human sperm capacitation and protein-tyrosine phosphorylation in vitro. *Biol Reprod* 61 (1999) 575-81.
- [17]A. Zini, E. De Lamirande, and C. Gagnon, Low levels of nitric oxide promote human sperm capacitation in vitro. *J Androl* 16 (1995) 424-31.
- [18]E. de Lamirande, and G. Lamothe, Reactive oxygen-induced reactive oxygen formation during human sperm capacitation. *Free Radic Biol Med* 46 (2009) 502-10.
- [19]T. Nobunaga, Y. Tokugawa, K. Hashimoto, Y. Kubota, K. Sawai, T. Kimura, K. Shimoya, M. Takemura, N. Matsuzaki, C. Azuma, and F. Saji, Elevated nitric oxide concentration in the seminal plasma of infertile males: nitric oxide inhibits sperm motility. *Am J Reprod Immunol* 36 (1996) 193-7.
- [20]F. Ferrara, R. Daverio, G. Mazzini, P. Bonini, and G. Banfi, Automation of human sperm cell analysis by flow cytometry. *Clin Chem* 43 (1997) 801-7.

- [21]M.L. Rasanen, O.L. Hovatta, I.M. Penttila, and Y.P. Agrawal, Detection and quantitation of sperm-bound antibodies by flow cytometry of human semen. *J Androl* 13 (1992) 55-64.
- [22]G.G. Haas, Jr., and M.E. Cunningham, Identification of antibody-laden sperm by cytofluorometry. *Fertil Steril* 42 (1984) 606-13.
- [23]D.P. Evenson, R.J. Emerick, L.K. Jost, H. Kayongo-Male, and S.R. Stewart, Zinc-silicon interactions influencing sperm chromatin integrity and testicular cell development in the rat as measured by flow cytometry. *J Anim Sci* 71 (1993) 955-62.
- [24]R.S. Jeyendran, V.C. Acosta, S. Land, and C.B. Coulam, Cryopreservation of human sperm in a lecithin-supplemented freezing medium. *Fertil Steril* 90 (2008) 1263-5.
- [25]H. Kojima, N. Nakatsubo, K. Kikuchi, S. Kawahara, Y. Kirino, H. Nagoshi, Y. Hirata, and T. Nagano, Detection and imaging of nitric oxide with novel fluorescent indicators: diaminofluoresceins. *Anal Chem* 70 (1998) 2446-53.
- [26]H. Strijdom, C. Muller, and A. Lochner, Direct intracellular nitric oxide detection in isolated adult cardiomyocytes: flow cytometric analysis using the fluorescent probe, diaminofluorescein. *J Mol Cell Cardiol* 37 (2004) 897-902.
- [27]F. Lampiao, H. Strijdom, and S.S. du Plessis, Direct nitric oxide measurement in human spermatozoa: flow cytometric analysis using the fluorescent probe, diaminofluorescein. *Int J Androl* 29 (2006) 564-7.
- [28]K. Taniguchi, K. Mamba, and M. Kitahama, [On the localization of enzymes in human spermatozoa. III. Esterase and acrosomal proteinase]. *Nihon Hoigaku Zasshi* 40 (1986) 24-9.

- [29]P. Pacher, J.S. Beckman, and L. Liaudet, Nitric oxide and peroxynitrite in health and disease. *Physiol Rev* 87 (2007) 315-424.
- [30]S. Figueroa, E. Lopez, C. Arce, M.J. Oset-Gasque, and M.P. Gonzalez, SNAP, a NO donor, induces cellular protection only when cortical neurons are submitted to some aggression process. *Brain Res* 1034 (2005) 25-33.
- [31]A. Zini, M.K. O'Bryan, M.S. Magid, and P.N. Schlegel, Immunohistochemical localization of endothelial nitric oxide synthase in human testis, epididymis, and vas deferens suggests a possible role for nitric oxide in spermatogenesis, sperm maturation, and programmed cell death. *Biol Reprod* 55 (1996) 935-41.
- [32]E. Miraglia, F. De Angelis, E. Gazzano, H. Hassanpour, A. Bertagna, E. Aldieri, A. Revelli, and D. Ghigo, Nitric oxide stimulates human sperm motility via activation of the cyclic GMP/protein kinase G signaling pathway. *Reproduction* 141 47-54.