

Cleveland State University
EngagedScholarship@CSU



Chemistry Faculty Publications

Chemistry Department

12-1-2009

Synthesis and Cytotoxic Analysis of Some Disodium $3\beta,6\beta$ -Dihydroxysterol Disulfates

Jianguo Cui

Guangxi Teachers Education University

Hui Wang

Guangxi Teachers Education University

Yi Xin

Cleveland State University

Aimin Zhou

Cleveland State University, A.ZHOU@csuohio.edu

Follow this and additional works at: https://engagedscholarship.csuohio.edu/scichem_facpub

 Part of the [Chemistry Commons](#)

How does access to this work benefit you? Let us know!

Recommended Citation

Cui, Jianguo; Wang, Hui; Xin, Yi; and Zhou, Aimin, "Synthesis and Cytotoxic Analysis of Some Disodium $3\beta,6\beta$ -Dihydroxysterol Disulfates" (2009). *Chemistry Faculty Publications*. 414.

https://engagedscholarship.csuohio.edu/scichem_facpub/414

This Article is brought to you for free and open access by the Chemistry Department at EngagedScholarship@CSU. It has been accepted for inclusion in Chemistry Faculty Publications by an authorized administrator of EngagedScholarship@CSU. For more information, please contact library.es@csuohio.edu.

Synthesis and cytotoxic analysis of some disodium 3 β ,6 β -dihydroxysterol disulfates

Jianguo Cui , Hui Wang , Yanmin Huang , Yi Xin , Aimin Zhou

Introduction

In recent years, a variety of steroids with unusual and interesting structures have been isolated from a wide range of marine organisms, particularly from sponges and echinoderms [1–3]. Among these steroidal compounds, many are sulfated polyhydroxysterols in sodium salt forms [4,5]. These compounds have exhibited a broad spectrum of biologic activities, such as suppression of HIV replication [6–9], inhibition of protein tyrosine kinases [10] and antitumor activities [11–13]. Therefore they greatly attract the attention of organic chemists and biomedical scientists [14–16]. Most of the sulfated polyhydroxysteroids isolated from these echinoderms or ophiuroids are found that the disulfation occurs on two hydroxyls located at C-2 and C-3 or C-3 and C-7, or C-3 and C-21 [17]. In this report, we synthesized three disulfated steroids with sulfate groups located at C-3 and C-6 on the ring A and B as shown in Fig. 1 and determined their cytotoxicity against cancer cells. Interestingly we found that the cytotoxicity of these compounds was differentially changed along with the structure of the R group. Our finding provides new evidence for the relationship between chemical structure and biofunctions.


Results and discussion

Chemistry

The synthesis of sulfated 3,6-dihydroxysterol was based on the methodology developed by Arnostova et al. [18]. Here we present a more efficient method for the synthesis of disodium 3 β ,6 β -dihydroxysterol disulfates with a cholesterol-like or stigmasterol-like or sitosterol-like side chain at position 17 [19,20]. As shown in Scheme 1, cholesterol, stigmasterol, and β -sitosterol were used as raw materials.

Compounds (**4a–4c**) were converted to the corresponding 4-en-3, 6-dioxysteroids (**5a–5c**) via oxidation with PCC in CH₂Cl₂. The reduction of **5a–5c** by NaBH₄ in the presence of NiCl₂ gave 3 β ,6 β -dihydroxysteroids (**6a–6c**) according to the synthetic method we previously developed [21]. The compounds **6a–6c** presented a typical difference in their spectrum from their parent compounds **5a–5c**. In the IR spectra the absorption at 1714 and 1686 cm⁻¹ for the original diketone carbonyl groups (CO) of **5b**, was replaced by a new broad absorption at 3427 cm⁻¹ for **6b**, indicating that the carbonyl groups in **5b** had been changed to the hydroxy groups in **6b**. In addition, the double bond of **5b** at 1609 cm⁻¹ had been eliminated in **6b**. In the ¹H NMR spectrum the chemical shifts at 3.650 ppm (1H, tt, *J* = 11.0, 5.0, C₃- α H) and 3.803 ppm (1H, dd, *J* = 5.5, 2.5, C₆- α H) for **6b** showed a α -configuration of 3-H and 6-H.

Based on the synthetic method developed by Comin et al. [14], the treatment of **6a–6c** with triethylammonium-sulfur tri-



Scheme 1. Reagents and conditions: (a) PCC/CH₂Cl₂, (b) NaBH₄/NiCl₂·6H₂O, (c) Et₃N-SO₃/DMF, and (d) cation exchange resin 732 (sodium form)(Na⁺)/MeOH.

oxide complex gave the ammonium sulfate of compounds 1–3, which was converted to their disodium salt via ion exchange (Scheme 1). The presence of the sulfate group in 1–3 was confirmed by the downfield shift of 3-H and 6-H to 3.961–3.966 ppm and 4.140–4.152 ppm, and by comparing to the ¹H NMR spectra of 6a and 6c (3.65–3.70 ppm and 3.80–3.82 ppm, respectively).

Antitumor activities

The antitumor activities of all these sulfated 3β,6β-dihydroxysterols were determined *in vitro* on Sk-Hep-1 (human liver carcinoma), H-292 (human lung carcinoma), PC-3 (human prostate carcinoma) and Hey-1B (human ovarian carcinoma) tumor cells. The results were summarized as IC₅₀ values in nmol/mL in Table 1.

Compound 1 displayed significantly higher cytotoxicity against these cancer cells when compared to compounds 2 and 3. Interestingly, the cytotoxicity of the compounds against these cancer cells was increased along with the order of the side chain at 17-C: cholesterol-like side chain (1), stigmasterol-like side chain

Table 1
In vitro antitumor activities (IC₅₀ in nmol/mL) of compounds 1, 2 and 3.^a

Compound	Sk-Hep-1	H-292	PC-3	Hey-1B
1	21	63	35	33
2	160	120	121	170
3	>200	>200	>200	>200

^a The MTS method was used to analyze the antiproliferative activity.

(2), and sitosterol-like side chain (3). Obviously the presence of a cholesterol-type side chain at 17-C is necessary for the best biological activity. The morphological change of the cancer cells induced by compound 1 was shown in Fig. 1A and B.

Experiments

The sterol and NaBH₄ were purchased from Merck Co. All chemicals and solvents were analytical grade and solvents were purified by general methods before being used. Melting points were determined on an X₄ apparatus and were uncorrected. Infrared spectra were measured with a Nicolet FT-360 Spectrophotometer. The ¹H and ¹³C NMR spectra were recorded in CDCl₃ or DMSO on a Bruker AV-500 spectrometer at working frequencies 500 and 125 MHz, respectively. Chemical shifts are expressed in ppm (δ) values and coupling constants (*J*) in Hz. Mass spectra (ESI) were recorded on an LCMS-2010A instrument. The cell viability was determined by the MTS method using 96-well plates in a LD400 AD/LD analysis spectrometer (Beckman Coulter).

The synthesis of compounds 5a–5c

Cholest-4-en-3,6-dione (5a): Pyridinium chlorochromate (PCC) (2.564 g, 11.87 mmol) was added to a solution of cholesterol (4a) (0.924 g, 2.2 mmol) in dried CH₂Cl₂ (40 mL) in one portion at room temperature. The reaction was completed in 28 h. To the mixture was then added 30 mL of CH₂Cl₂, and the suspension was poured over a silica gel column and eluted with CH₂Cl₂. The eluate was washed with cold water and saturated brines. After drying over anhydrous sodium sulfate, the solvent was removed under reduced pressure, and the crude product was purified by chromatography on silica gel using petroleum ether (60–90 °C)/EtOAc (5: 1) as the eluent to give 0.795 g (84%) of 5a as pale yellow crystals, m.p. 90–91 °C; IR (KBr) ν 2953, 2865, 1693, 1600, 1486, 1249, 1221, 1117, 942 cm⁻¹; ¹H NMR(CDCl₃, 500 MHz): 0.746(3H, s, 18-CH₃), 0.886(3H, d, *J* = 6.4, 26 or 27-CH₃), 0.899(3H, d, *J* = 6.4, 26 or 27-CH₃), 0.952(3H, d, *J* = 6.5, 21-CH₃), 1.172(3H, s, 19-CH₃), 2.546(1H, dd, *J* = 14.6, 5.2, C₂-βH), 2.706(1H, dd, *J* = 16.0, 4.0, C₇-αH), 6.196(1H, s, C₄-H).




Fig. 1. The morphology of cancer cells. Hey 1B (A) and Sk-Hep-1 (B) cells were treated with compound 1 at various doses for 24 h and 48 h, respectively. The photos of cell morphology were taken under an Olympus CKX31 microscope at 100× magnification.

24-Ethylcholest-4,22-dien-3,6-dione (5b): Yield: 83%, m.p. 134–135 °C; IR (KBr) ν : 2959, 1714, 1686, 1609, 969, 864 cm^{-1} ; ^1H NMR(CDCl_3 , 500 MHz): 0.743(3H, s, 18- CH_3), 0.805(3H, t, $J=7.0$, 29- CH_3), 0.798(3H, d, $J=6.5$, 26- or 27- CH_3), 0.849(3H, d, $J=6.5$, 26- or 27- CH_3), 1.036(3H, d, $J=7.0$, 21- CH_3), 1.169(3H, s, 19- CH_3), 5.040(1H, dd, $J=15.2$, 9.0, $\text{C}_{22}\text{-H}$), 5.150(1H, dd, $J=15.2$, 8.5, $\text{C}_{23}\text{-H}$), 6.171(1H, s, $\text{C}_4\text{-H}$).

24-Ethylcholest-4-en-3,6-dione (5c): Yield: 86%, m.p. 172–174 °C. IR (KBr) ν : 2959, 1683, 1601, 1581, 1461, 1377, 1246, 1124, 948, 871 cm^{-1} ; ^1H NMR(CDCl_3 , 500 MHz): 0.724(3H, s, 18- CH_3), 0.816(3H, d, $J=7.0$, 26- or 27- CH_3), 0.841(3H, d, $J=7.0$, 26- or 27- CH_3), 0.848(3H, t, $J=8.0$, 29- CH_3), 0.935(3H, d, $J=6.5$, 21- CH_3), 1.167(3H, s, 19- CH_3), 2.13–2.17(1H, m, $\text{C}_2\text{-}\alpha\text{H}$), 2.44–2.58 (2H, m, $\text{C}_7\text{-}\beta\text{H}$ and $\text{C}_2\text{-}\beta\text{H}$), 2.682(1H, dd, $J=15.5$, 4.5, $\text{C}_7\text{-}\alpha\text{H}$), 6.170(1H, s, $\text{C}_4\text{-H}$).

The synthesis of compounds 6a–6c

Cholest-3 β ,6 β -diol (6a): NaBH_4 (90 mg, 2.38 mmol) was added to a solution of **5a** (200 mg, 0.50 mmol) and $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (120 mg, 0.50 mmol) in CH_3OH (20 mL) in 1 min at room temperature. After 20 min, the reaction was stopped. The solution was neutralized with 1 M HCl. After evaporating the majority of the MeOH under reduced pressure, ethyl acetate (40 mL) was added to the residue. The resulting solution was washed with cold water and saturated brines. After drying over anhydrous sodium sulfate, the solvent was removed under reduced pressure, and the crude product was then purified by flash chromatography on silica gel using petroleum ether/ethyl acetate (1:1) as the eluent, to give **6a** as a white solid (164 mg, 81%), m.p. 161–162 °C; IR (KBr) ν : 3399, 2925, 2868, 1470, 1360, 1319, 1172, 1078, 1041, 1021, 955, 767 cm^{-1} ; ^1H NMR(CDCl_3 , 500 MHz): 0.709(3H, s, 18- CH_3), 0.878(3H, d, $J=2.2$, 26 or 27- CH_3), 0.891(3H, d, $J=2.2$, 26 or 27- CH_3), 0.927(3H, d, $J=6.4$, 21- CH_3), 1.052(3H, s, 19- CH_3), 3.64–3.82(1H, m, $\text{C}_3\text{-}\alpha\text{H}$), 3.825(1H, d, $J=2.4$, $\text{C}_6\text{-}\alpha\text{H}$).

24-Ethylcholest-22-en-3 β ,6 β -diol (6b): Yield: 72%; m.p. 207–209 °C; IR (KBr) ν : 3427, 2929, 2864, 1581, 1462, 1380, 1037, 967 cm^{-1} ; ^1H NMR(CDCl_3 , 500 MHz): 0.708(3H, s, 18- CH_3), 0.795(3H, d, $J=6.5$, 26 or 27- CH_3), 0.845(3H, d, $J=6.5$, 26 or 27- CH_3), 0.802(3H, t, $J=7.5$, 29- CH_3), 1.013(3H, d, $J=7.0$, 21- CH_3), 1.033(3H, s, 19- CH_3), 3.650(1H, tt, $J=11.0$, 5.0, $\text{C}_3\text{-}\alpha\text{H}$), 3.803(1H, dd, $J=5.5$, 2.5, $\text{C}_6\text{-}\alpha\text{H}$), 5.022(1H, dd, $J=15.0$, 8.5, $\text{C}_{22}\text{-H}$), 5.140(1H, dd, $J=15.0$, 8.5, $\text{C}_{23}\text{-H}$).

24-Ethylcholest-3 β ,6 β -diol (6c): Yield: 72%; m.p. 172–173 °C; IR (KBr) ν : 3419, 3125, 2941, 2872, 1397, 1172, 1033 cm^{-1} ; ^1H NMR(CDCl_3 , 500 MHz): 0.728(3H, s, 18- CH_3), 0.831(6H, d, $J=7.6$ Hz, 26 and 27- CH_3), 0.866(3H, d, $J=6.2$ Hz, 29- CH_3), 0.896(3H, s, 21- CH_3), 1.027(3H, s, 19- CH_3), 3.667(1H, t, $J=5.5$, $\text{C}_3\text{-H}$), 3.821(1H, s, $\text{C}_6\text{-H}$).

General procedure for the synthesis of compounds 1–3

Disodium 3 β ,6 β -dihydroxy-5 α -cholestane disulfate (1): The triethylamine-sulfur trioxide complex (173 mg, 0.96 mmol) was added to a solution of cholest-3 β ,6 β -diol (**6a**) (50 mg, 0.12 mmol) in DMF (0.6 mL) under an argon atmosphere, and the mixture was stirred at 95 °C for 3 h. Then the reaction mixture was quenched with water (0.2 mL). The solution was poured over a silica gel column to remove excess $\text{SO}_3 \cdot \text{NEt}_3$. The product was eluted by using petroleum ether/ CH_2Cl_2 (1:1) as the eluent, and then continued to be eluted with MeOH/ CHCl_3 (1:12), and followed by evaporation of the solvent to yield a white solid (diammonium 3 β ,6 β -dihydroxy-5 α -cholestane disulfate). To the solution of the solid in methanol (15 mL) was added Cation exchange resin 732 (sodium form) (Na^+) (10 g), and stirred for 5 h at room temperature. The resin was removed by filtration, and the filtrate was concen-

trated. This process was repeated one more time with 15 g of the resin. Finally compound **1** was obtained as a colorless solid (45 mg), yield: 60%, m.p. 173–174 °C; IR (KBr) ν : 3464, 2937, 2868, 1470, 1383, 1226, 1066, 980, 947, 853, 820, 624 cm^{-1} ; ^1H NMR(DMSO, 500 MHz): 0.657(3H, s, 18- CH_3), 0.851(3H, d, $J=6.5$, 26- or 27- CH_3), 0.855(3H, d, $J=6.5$, 26- or 27- CH_3), 0.896(3H, d, $J=6.5$, 21- CH_3), 0.932(3H, s, 19- CH_3), 1.935(1H, brd, $J=12.5$, $\text{C}_2\text{-}\beta\text{H}$), 3.966(1H, m, $\text{C}_3\text{-}\alpha\text{H}$), 4.140(1H, s, $\text{C}_6\text{-}\alpha\text{H}$); ^{13}C NMR(DMSO, 125 MHz): 36.1(1-C), 28.2(2-C), 70.1(3-C), 33.4(4-C), 35.4(5-C), 76.1(6-C), 42.8(7-C), 30.5(8-C), 47.7(9-C), 38.6(10-C), 21.1(11-C), 39.4(12-C), 40.6(13-C), 56.2(14-C), 24.4(15-C), 27.8(16-C), 54.2(17-C), 12.4(18-C), 15.8(19-C), 35.6(20-C), 19.0(21-C), 36.1(22-C), 23.7(23-C), 40.5(24-C), 29.2(25-C), 22.8(26-C), 23.1(27-C); (-)LRESIMS m/z : 585[M–Na] $^{-1}$, 563[M–2Na+H] $^{-1}$.

Disodium 24-ethyl-3 β ,6 β -dihydroxycholest-22-ene disulfate (2): Yield: 66%; m.p. 174–175 °C; IR (KBr) ν : 2937, 2864, 1646, 1466, 1368, 1209, 1066, 967, 853, 812 cm^{-1} ; ^1H NMR(DMSO, 500 MHz): 0.679(3H, s, 18- CH_3), 0.778 (3H, t, $J=4.4$, 29- CH_3), 0.795(3H, d, $J=6.1$, 26- or 27- CH_3), 0.835(3H, d, $J=6.1$, 26- or 27- CH_3), 0.935(3H, s, 19- CH_3), 0.998(3H, d, $J=6.5$, 21- CH_3), 1.917(1H, d, $J=12.5$, $\text{C}_2\text{-}\beta\text{H}$), 2.031(1H, m, $\text{C}_{20}\text{-H}$), 3.961(1H, m, $\text{C}_3\text{-}\alpha\text{H}$), 4.152(1H, d, $J=3.4$, $\text{C}_6\text{-}\alpha\text{H}$), 5.031(1H, dd, $J=15.1$, 8.6, $\text{C}_{22}\text{-H}$), 5.161(1H, dd, $J=15.1$, 8.8, $\text{C}_{23}\text{-H}$); ^{13}C NMR(DMSO, 125 MHz): 35.5(1-C), 27.4(2-C), 70.1(3-C), 33.4(4-C), 31.8(5-C), 76.1(6-C), 45.7(7-C), 29.2(8-C), 54.2(9-C), 38.6(10-C), 21.4(11-C), 40.6(12-C), 44.2(13-C), 56.4(14-C), 25.3(15-C), 29.0(16-C), 56.0(17-C), 12.7(18-C), 15.9(19-C), 42.6(20-C), 21.6(21-C), 138.6(22-C), 129.3(23-C), 47.8(24-C), 30.5(25-C), 21.1(26-C), 19.4(27-C), 24.4(28-C), 12.6(29-C); (-)LRESIMS m/z : 611[M–Na] $^{-1}$, 588[M–2Na] $^{-2}$.

Disodium 24-ethyl-3 β ,6 β -dihydroxycholestane disulfate (3): Yield: 65%; m.p. 242–249 °C; IR (KBr) ν : 3427, 2925, 1634, 1401, 1209, 815 cm^{-1} ; ^1H NMR(DMSO, 500 MHz): 0.650(3H, s, 18- CH_3), 0.798(3H, d, $J=7.0$, 26- or 27- CH_3), 0.818(3H, d, $J=7.0$, 26- or 27- CH_3), 0.825(3H, t, $J=6.5$, 29- CH_3), 0.896(3H, d, $J=6.3$, 21- CH_3), 0.923(3H, s, 19- CH_3), 1.924(1H, brad, $J=12.5$, $\text{C}_2\text{-}\beta\text{H}$), 3.961(1H, m, $\text{C}_3\text{-H}$), 4.181(1H, d, $J=3.5$, $\text{C}_6\text{-H}$); (-)LRESIMS m/z : 613[M–Na] $^{-1}$, 590[M–2Na] $^{-2}$.

Assay for cell viability

Materials and methods

Stock solutions of compounds **1**, **2** and **3**, were prepared in sterile dimethyl sulfoxide (DMSO) (Sigma) at a concentration of 10 mg/mL, and diluted to proper concentrations according to the experimental design with the cell culture medium.

Cell culture

Sk-Hep-1, H-292, PC-3 (ATCC) and Hey-1B (a gift from Dr. Yan Xu, University of Indiana) cells were cultured in a proper medium supplemented with 10% fetal bovine serum and 0.1 g/L penicillin G + 0.1 g/L streptomycin sulfate in a humidified atmosphere of 5% CO_2 at 37 °C.

Treatment of cancer cells

Cancer cells (4×10^4 cells/mL, 200 μL) were seeded into each well of a 96-well microtiter plate. After incubation for 24 h, the compounds with a series of concentrations (range 5–100 $\mu\text{g}/\text{mL}$) were added to the cells. An equal amount of DMSO was added to the cells as negative controls. All were treated in triplicate.

Determination of cell viability

MT Stetrazolium salt ((3-4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulphophenyl)-2H-tetrazolium) (CellTiter 96 AQueous Non-Radioactive Cell Proliferation Assay, Cat #G5421, Promega Corporation) dye reduction assay was used to determine the cell viability. The assay is dependent on the

reduction of MTS by mitochondrial dehydrogenases of viable cells to a blue water soluble formazan product, which can be measured spectrophotometrically. The absorbance of the formazan product at 490 nm is in linear proportion to cell numbers. Briefly, after treatment (see 3.4.3) with the compounds for 48 h, the medium was removed and the cells were incubated with 100 μ L of fresh medium plus 20 μ L of MTS solution for additional 4 h according to the instruction provided by the manufacturer. The absorbance (A) at 490 nm was measured using a LD400 AD/LD analysis spectrometer (Beckman coulter). The IC_{50} value was calculated as the concentration of drug yielding 50% cell survival.

Acknowledgments

The authors acknowledge the financial support of the National Natural Science Foundation of China (No.: 20562001), the Natural Science Foundation of Guangxi Province (No.: 0575054) of China.

References

- [1] Blunt JW, Copp BR, Hu WP, Munro MHG, Northcote PT, Prinsep MP. Marine natural products. *Nat Prod Rep* 2007;24:31–86.
- [2] Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MP. Marine natural products. *Nat Prod Rep* 2005;22:15–61.
- [3] Faulkner DJ. Marine natural product. *Nat Prod Rep* 2001;18:1–49.
- [4] Huang SF, Cui JG, Liu ZP, Gan CF. Recent Progresses in natural sulfated polyhydroxysteroids from marine organism and their bioactivity. *Tianran Chanwu Yanjiu Yu Kaifa* (Chinese) 2006;18:681–5.
- [5] Han LD, Cui JG, Huang CS. Bioactive polyhydroxy sterols and their sapogenins for marine organisms. *Youji Huaxue* (Chinese) 2003;23:305–11.
- [6] McKee TC, Cardellina JH, Riccio R, D'Auria MV, Iorizzi M, Minale L, et al. HIV-inhibitory natural products, 11. Comparative studies of sulfated sterols from marine invertebrates. *J Med Chem* 1994;37:793–7.
- [7] Lerch ML, Faulkner DJ. Unusual polyoxygenated sterols from a Philippines sponge *Xestospongia* sp. *Tetrahedron* 2001;57:4091–4.
- [8] Rudi A, Yosief TM, Loya S, Hizi A, Schleyer M, Kashman Y, et al. A novel anti-HIV-1 RT sulfated sterol from the sponge *Clathria Species*. *J Nat Prod* 2001;64:1451–3.
- [9] Qureshi A, Faulkner DJ. Haploamates A and B: new steroidal sulfamate esters from two sponges. *Tetrahedron* 1999;55:8323.
- [10] Fu X, Schmitz FJ, Lee RH, Papkoff JS, Slate DL. Inhibition of protein tyrosine kinase pp60v-src: sterol sulfates from the brittle star *Ophiarachna incrassata*. *J Nat Prod* 1994;57:1591–4.
- [11] Roccatagliata AJ, Maier MS, Seldes AM. New sulfated polyhydroxysteroids from the Antarctic ophiuroid *Astrotoma agassizii*. *J Nat Prod* 1998;61:370–4.
- [12] Ivanchina NV, Kicha AA, Kalinovsky AI, Dmitrenok PS, Stonik VA, Riguera R, et al. Hemolytic polar steroidal constituents of the starfish *Astropecten latespinosus*. *J Nat Prod* 2000;63:1178.
- [13] Yang SW, Alexei B, Chan TM, Michelle S, Jean L, Shirley AP, et al. A new sterol sulfate, Sch 572423, from a marine sponge, *Topsentia* sp. *Bioorg Med Chem Lett* 2003;13:1791–4.
- [14] Comin MJ, Maier MS, Roccatagliata AJ, Pujol CA, Damonte EB. Evaluation of the antiviral activity of natural sulfated polyhydroxysteroids and their synthetic derivatives and analogs. *Steroids* 1999;64:335–40.
- [15] Santos GA, Garrido M, Ana P, Pujol CA, Damonte EB, Maier MS. Synthesis and antiviral activity of sulfated and acetylated derivatives of 2,3-dihydroxy-5-cholestane. *Steroids* 2003;68:125–32.
- [16] Oishi T, Tsuchikawa H, Murata M, Yoshida M, Morisawa M. Synthesis and identification of an endogenous sperm activating and attracting factor isolated from eggs of the ascidian *Ciona intestinalis*; an example of nanomolar-level structure elucidation of novel natural compound. *Tetrahedron* 2004;60:6971–80.
- [17] Levina EV, Andriyaschenko PV, Kalinovsky AI, Stonik VA. New ophiuroid-type sterols from the starfish *Pteraster tesselatu*. *J Nat Prod* 1998;61:1423.
- [18] Arnostova LM, Pouzar V, Drasar P. Synthesis of the sulfates derived from 5-cholestane-3,6-diol. *Steroids* 1992;57:233–5.
- [19] Wang H, Cui JG, Huang LL. Synthesis of disodium 3 β ,6 β -dihydroxystigmast-22-ene disulfate. *Huaxue Shiji* 2007;29:566–8.
- [20] Wang H, Cui JG, Huang LL. Synthesis of disodium 3 β ,6 β -dihydroxystigmast-22-ene disulfate. *Huagong jishu Yu Kaifa* (Chinese) 2007;36(9):1–3.
- [21] Cui JG, Zeng LM, Su JY. Synthesis of some polyhydroxysterols and investigation of relationship between their structure and cytotoxicity. *Chem J Chin Univ (Chin)* 2000;21:1399–404.