

Cleveland State University  
**EngagedScholarship@CSU**

Undergraduate Research Posters 2015

Undergraduate Research Posters

2015

# Dual Stimuli Response Frequency and Stimulus Choice of the African Clawed Frog, *Xenopus laevis*, when presented with two stimuli

Mingo Rolince  
*Cleveland State University*

Heidi Pignolet  
*Cleveland State University*

Alexa Hoy  
*Cleveland State University*

Follow this and additional works at: [https://engagedscholarship.csuohio.edu/u\\_poster\\_2015](https://engagedscholarship.csuohio.edu/u_poster_2015)

**How does access to this work benefit you? Let us know!**

## Recommended Citation

Rolince, Mingo; Pignolet, Heidi; and Hoy, Alexa, "Dual Stimuli Response Frequency and Stimulus Choice of the African Clawed Frog, *Xenopus laevis*, when presented with two stimuli" (2015). *Undergraduate Research Posters 2015*. 10.  
[https://engagedscholarship.csuohio.edu/u\\_poster\\_2015/10](https://engagedscholarship.csuohio.edu/u_poster_2015/10)

This Book is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2015 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact [library.es@csuohio.edu](mailto:library.es@csuohio.edu).



This digital edition was prepared by MSL Academic Endeavors, the imprint of the Michael Schwartz Library at Cleveland State University.

*\*Supported by the McNair Scholars Program*

***Dual Stimuli Response Frequency and Stimulus Choice of the African Clawed Frog, *Xenopus laevis*, when presented with two stimuli***

College of Sciences and Health Professions

**Student Researchers:** Mingo Rolince, Heidi Pignolet, and Alexa Hoy

**Faculty Advisor:** Jeffrey Dean

**Abstract**

This preliminary study examines responses of African Clawed Frogs to simultaneous presentation of two stimuli. Frogs were tested in a round arena with water 4 cm deep. Four stimulus rods driven via computer-controlled stepper motors were concealed in a screen suspended above the water. These rods could present a lateral line stimulus, a visual stimulus, or a combination of both. Overall, reactions and no reactions were evenly distributed--51.1% and 48.9%, respectively. Frogs responded more frequently to rostral than caudal stimuli (chi-squared 20.8, df=11,  $p < 0.04$ ). Frogs reacted more to stimuli between  $-90^\circ$  and  $90^\circ$ . Turn angle depended linearly on stimulus angle (e.g., Turn angle =  $0.44 + 0.64 \times$  Stimulus angle;  $p_{\text{slope}} < 0.0001$ ;  $R^2_{\text{adj}} = 69.5\%$ ). (The distributions of stimulus angles appear slightly skewed as often the same stimulus rod was retested after the frog responded and partially oriented towards it, prompting a second test with smaller angles.) Two stimuli did not elicit more responses than one stimuli ( $P=0.25$ ). The frogs' choice of stimulus depended primarily on stimulus proximity and angle, not stimulus type. When presented with two stimuli, the frog chose the nearer stimulus and the more rostral stimulus ( $p < 0.0001$ ). The largest factor in predicting the frog's choice of stimuli seems to be the linear distance from the stimuli to the frog.