

The University of San Francisco  
USF Scholarship: a digital repository @ Gleeson Library |  
Geschke Center

---

Mathematics

College of Arts and Sciences

---

2009

# Generalized Helmholtz-Kirchhoff Model for Two-Dimensional Distributed Vortex Motion

Raymond J. Nagem

Guido Sandri

David Uminsky

*University of San Francisco*, [duminsky@usfca.edu](mailto:duminsky@usfca.edu)

C. Eugene Wayne

Follow this and additional works at: <http://repository.usfca.edu/math>

 Part of the [Fluid Dynamics Commons](#), and the [Mathematics Commons](#)

---

## Recommended Citation

Nagem, Raymond, Guido Sandri, David Uminsky, and C. Eugene Wayne. 2009. "Generalized Helmholtz-Kirchhoff Model for Two-Dimensional Distributed Vortex Motion." *SIAM Journal On Applied Dynamical Systems* 8, no. 1: 160-179.

This Article is brought to you for free and open access by the College of Arts and Sciences at USF Scholarship: a digital repository @ Gleeson Library | Geschke Center. It has been accepted for inclusion in Mathematics by an authorized administrator of USF Scholarship: a digital repository @ Gleeson Library | Geschke Center. For more information, please contact [repository@usfca.edu](mailto:repository@usfca.edu).

## Generalized Helmholtz–Kirchhoff Model for Two-Dimensional Distributed Vortex Motion\*

Raymond Nagem<sup>†</sup>, Guido Sandri<sup>†</sup>, David Uminsky<sup>‡</sup>, and C. Eugene Wayne<sup>‡</sup>

**Abstract.** The two-dimensional Navier–Stokes equations are rewritten as a system of coupled nonlinear ordinary differential equations. These equations describe the evolution of the moments of an expansion of the vorticity with respect to Hermite functions and of the centers of vorticity concentrations. We prove the convergence of this expansion and show that in the zero viscosity and zero core size limit we formally recover the Helmholtz–Kirchhoff model for the evolution of point vortices. The present expansion systematically incorporates the effects of both viscosity and finite vortex core size. We also show that a low-order truncation of our expansion leads to the representation of the flow as a system of interacting Gaussian (i.e., Oseen) vortices, which previous experimental work has shown to be an accurate approximation to many important physical flows [P. Meunier, S. Le Dizès, and T. Leweke, *C. R. Phys.*, 6 (2005), pp. 431–450].

**Key words.** vortex dynamics, Navier–Stokes equations, fluid dynamics

**AMS subject classifications.** 76D05, 76D17, 35Q30

**DOI.** 10.1137/080715056

**1. Introduction.** In this paper we represent solutions of the two-dimensional Navier–Stokes equations as a system of interacting vortices. This expansion, which generalizes the Helmholtz–Kirchhoff model of interacting point vortices in an inviscid fluid, systematically incorporates the effects of both vorticity and finite vortex core size. Furthermore, we give conditions which guarantee the convergence of our expansion. Incompressible viscous flow has two standard analytic representations: a formulation in terms of the primitive velocity and pressure variables, and a formulation in terms of the velocity and vorticity variables [8]. The velocity-vorticity representation has particular advantages when boundaries are unimportant, since vorticity cannot be created or destroyed in the interior of a fluid. The vorticity field can also be directly related to physically observed flow structures such as line and ring vortices.

In two space dimensions, the vorticity field has the additional advantage of reducing to a scalar. An early representation of two-dimensional flow in terms of moving point vortices was developed by Kirchhoff [5] and by Von Helmholtz [17]. The point vortex model has been studied extensively—a thorough review of the model and recent developments is provided in the monograph by Newton [13]. While the Helmholtz–Kirchhoff point vortex model captures many of the basic physical phenomena observed in two-dimensional rotational flows, experi-

---

\*Received by the editors February 1, 2008; accepted for publication (in revised form) by T. Kaper October 1, 2008; published electronically January 23, 2009.

<http://www.siam.org/journals/siads/8-1/71505.html>

<sup>†</sup>Department of Aerospace and Mechanical Engineering, Boston University, Boston, MA 02215 ([nagem@bu.edu](mailto:nagem@bu.edu), [sandri@bu.edu](mailto:sandri@bu.edu)).

<sup>‡</sup>Department of Mathematics and Statistics and Center for Biodynamics, Boston University, Boston, MA 02215 ([duminsky@math.bu.edu](mailto:duminsky@math.bu.edu), [cew@math.bu.edu](mailto:cew@math.bu.edu)). The work of these authors was supported in part by DMS-0405724.

ments with even relatively simple vortex configurations exhibit complications far beyond the point vortex predictions [11]. Additionally, the classical point vortex model neglects the effects of *viscosity*. However, these experiments also reinforce the idea that in many circumstances the fluid flow may be well approximated by a collection of interacting vortices, albeit vortices with finite core size, subject to the effects of viscosity. Earlier work to introduce a finite core size model for Euler’s equations was done in [9]. In this paper the vortex patches that are modeled are described as disjoint piecewise-constant elliptical finite-area vortex regions (FAVORs). In this model, higher spatial moments of the ellipse are allowed to evolve under the flow. The moment model is truncated to second order, and the corotating vortex example is analyzed. Later in [10] the authors generalize the spatial moment model found in [9] by incorporating weak Newtonian viscosity to derive equations of motions for two like signed vortices.

Other recent studies of the interaction of viscous vortices can be found in [1, 2, 6, 7, 11, 16]. The main focus of most of these papers is the merger of two like signed vortices. A metastable state is found before merger which consists of two rotating near-circular vortices. More recently, it has been conjectured that in a two-vortex system the profiles relax to a pair of Gaussian vortices before merging [7]. An alternative method for incorporating viscous effects was proposed in [1] by adding a random forcing term to the ODEs which describe the motion of the vortex centers. Thus, it is of interest to extend and generalize the Helmholtz–Kirchhoff point vortex model to a model that incorporates nonzero vortex core size and viscous effects while retaining its basic form. Such an extension is the goal of this paper.

The governing equations for the velocity ( $\mathbf{u}$ ) and pressure ( $p$ ) variables are

$$(1) \quad \frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\nabla \frac{p}{\rho} + \nu \Delta \mathbf{u},$$

$$(2) \quad \nabla \cdot \mathbf{u} = 0,$$

where  $\rho$  is the fluid density and  $\nu$  is the kinematic viscosity. Taking the curl of (1) and (2) gives

$$(3) \quad \frac{\partial \boldsymbol{\omega}}{\partial t} + (\mathbf{u} \cdot \nabla) \boldsymbol{\omega} - (\boldsymbol{\omega} \cdot \nabla) \mathbf{u} = \nu \Delta \boldsymbol{\omega},$$

$$(4) \quad \boldsymbol{\omega} = \nabla \times \mathbf{u}, \quad \nabla \cdot \boldsymbol{\omega} = 0,$$

which are the governing equations for the velocity-vorticity variables. For two-dimensional flows, the vorticity vector  $\boldsymbol{\omega}$  is perpendicular to the plane of the flow, and the third term on the left-hand side of (3) vanishes. The condition  $\nabla \cdot \boldsymbol{\omega} = 0$  is identically satisfied, and (3) then reduces to the single scalar equation:

$$(5) \quad \frac{\partial \omega}{\partial t} + (\mathbf{u} \cdot \nabla) \omega = \nu \Delta \omega,$$

where  $\omega$  is the single nonzero component of the vorticity. A drawback of the formulation in (5) is that the velocity of the fluid is still present in the equation. However, assuming that

the vorticity field is sufficiently localized, the velocity vector can be computed in terms of the vorticity  $\omega$  by the Biot–Savart law

$$(6) \quad \mathbf{u}(\mathbf{x}) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{(\mathbf{x} - \mathbf{y})^\perp}{|\mathbf{x} - \mathbf{y}|^2} \omega(\mathbf{y}) d\mathbf{y},$$

where for a two-vector  $\mathbf{z} = (z_1, z_2)$ ,  $\mathbf{z}^\perp = (-z_2, z_1)$ .

In this paper we use (5) and (6) to develop a vorticity representation of two-dimensional viscous flow. Our representation is based on a decomposition of the vorticity field into a set of moving distributed vortices. Differential equations are derived for the motion of the vortex centers and for the time evolution of the vortex distributions. The evolution of each individual vortex is represented as an expansion with respect to a sequence of Hermite functions. Such expansions have proven useful in theoretical studies of two-dimensional fluid flows [3, 4] and the leading-order term in this expansion is precisely the Gaussian vortex (i.e., Oseen vortex [15, 14]) whose utility as an approximation for vortex interaction was shown in [11]. We show that the coefficients in this expansion satisfy a system of ODEs whose coefficients can be explicitly represented in terms of a fixed, computable kernel function. We also prove the convergence of this expansion. It is shown that our representation reduces in the appropriate limit to the Helmholtz–Kirchhoff model and allows at the same time arbitrarily complex evolution and interaction of the moving vortices.

In the present paper we concentrate on the mathematical formulation of the generalized Helmholtz–Kirchhoff model. In future work we will explore the predictions of this model both numerically and analytically in a number of different physical settings.

**2. The “multivortex” expansion.** In this section we separate the solution of the vorticity equation into  $N$  components and derive separate evolution equations for each component. From a physical point of view this decomposition will be most useful when each of the components corresponds to a localized region of vorticity (e.g., a vortex) well separated from the other lumps, but the mathematical development described below is well defined without regard to these physical considerations. However, with this application in mind, we will often refer to each of the components as a “vortex.”

Consider the initial value problem for the two-dimensional vorticity equation

$$(7) \quad \begin{aligned} \frac{\partial \omega}{\partial t} &= \nu \Delta \omega - \mathbf{u} \cdot \nabla \omega, \\ \omega &= \omega(\mathbf{x}, t), \quad \mathbf{x} \in \mathbb{R}^2, \quad t > 0, \\ \omega(\mathbf{x}, 0) &= \omega_0(\mathbf{x}), \end{aligned}$$

where  $\mathbf{u}$  is the velocity field associated with the vorticity field  $\omega$ . We begin by decomposing the initial vorticity distribution by writing

$$(8) \quad \omega_0(\mathbf{x}) = \sum_{j=1}^N \omega_0^j(\mathbf{x}).$$

Of course, this decomposition is not unique—even the number of pieces,  $N$ , into which we decompose the vorticity is up to us to choose. In general the choice we make will be motivated by physical considerations; however, for the development below, all we require of the

decomposition is that the total vorticity of each vortex be nonzero, i.e.,

$$(9) \quad m^j = \int_{\mathbb{R}^2} \omega_0^j(\mathbf{x}) d\mathbf{x} \neq 0, \quad j = 1, \dots, N.$$

If (9) is satisfied we define  $\mathbf{x}_0^j$  by

$$(10) \quad \int_{\mathbb{R}^2} (\mathbf{x} - \mathbf{x}_0^j) \omega_0^j(\mathbf{x}) d\mathbf{x} = 0,$$

or equivalently,

$$(11) \quad \mathbf{x}_0^j = \frac{1}{m^j} \int_{\mathbb{R}^2} \mathbf{x} \omega_0^j(\mathbf{x}) d\mathbf{x}.$$

We now write the vorticity for  $t > 0$  as

$$(12) \quad \omega(\mathbf{x}, t) = \sum_{j=1}^N \omega^j(\mathbf{x} - \mathbf{x}^j(t); t)$$

and the velocity field as

$$(13) \quad \mathbf{u}(\mathbf{x}, t) = \sum_{j=1}^N \mathbf{u}^j(\mathbf{x} - \mathbf{x}^j(t); t),$$

where  $\mathbf{u}^j(\mathbf{y}, t)$  is the velocity field associated with  $\omega^j(\mathbf{y}, t)$  by the Biot–Savart law. Of course we still have to define the equations of motion for  $\omega^j(\mathbf{y}, t)$  and  $\mathbf{x}^j(t)$ .

The centers of the vorticity regions,  $\mathbf{x}^j(t)$ , and the vorticity regions themselves evolve via a coupled system of ordinary-partial differential equations constructed so that in the limit of zero viscosity and when the different components of the vorticity happen to be point vortices (i.e., Dirac-delta functions) we recover the Helmholtz–Kirchhoff point vortex equations. If we take the partial derivative of (12) and use the equation satisfied by the vorticity, we find

$$(14) \quad \begin{aligned} \partial_t \omega(\mathbf{x}, t) &= \sum_{j=1}^N \partial_t \omega^j(\mathbf{x} - \mathbf{x}^j(t), t) - \sum_{j=1}^N \dot{\mathbf{x}}^j(t) \cdot \nabla \omega^j(\mathbf{x} - \mathbf{x}^j(t), t) \\ &= \sum_{j=1}^N \nu \Delta \omega^j(\mathbf{x} - \mathbf{x}^j(t), t) - \sum_{j=1}^N \left( \sum_{\ell=1}^N \mathbf{u}^\ell(\mathbf{x} - \mathbf{x}^\ell(t), t) \right) \cdot \nabla \omega^j(\mathbf{x} - \mathbf{x}^j(t), t). \end{aligned}$$

Given this equation, it is natural to define  $\omega^j$  as the solution of

$$(15) \quad \begin{aligned} \frac{\partial \omega^j}{\partial t}(\mathbf{x} - \mathbf{x}^j(t), t) &= \nu \Delta \omega^j(\mathbf{x} - \mathbf{x}^j(t), t) - \left( \sum_{\ell=1}^N \mathbf{u}^\ell(\mathbf{x} - \mathbf{x}^\ell(t), t) \right) \cdot \nabla \omega^j(\mathbf{x} - \mathbf{x}^j(t), t) \\ &+ \dot{\mathbf{x}}^j(t) \cdot \nabla \omega^j(\mathbf{x} - \mathbf{x}^j(t), t), \quad j = 1, \dots, N. \end{aligned}$$

To close this system of equations we must specify how the centers of vorticity  $\mathbf{x}^j(t)$  evolve. We impose the condition that the first moment of each vorticity region must vanish at every time  $t > 0$ ; i.e., we require that

$$(16) \quad \int_{\mathbb{R}^2} (\mathbf{x} - \mathbf{x}^j(t)) \omega^j(\mathbf{x} - \mathbf{x}^j(t), t) d\mathbf{x} = \mathbf{0} \quad \text{for all } t > 0, j = 1, \dots, N.$$

(Note that this equation really contains two conditions, one for each component of  $\mathbf{x} - \mathbf{x}^j(t)$ .) We impose this condition to fix the evolution of  $\mathbf{x}^j(t)$  because Galloway and Wayne have recently shown [3] that if one considers the evolution of general solutions of (7), the solution will approach an Oseen vortex, and the rate of the approach will be faster if the vorticity distribution has first moment equal to zero. Solutions of (7) preserve the first moment, and hence if the initial conditions have first moment equal to zero, the solution will have first moment zero for all time. The equations (15) no longer preserve the first moment, and thus we impose this condition for all time, which then defines the motion of the center of vorticity.

Note that if we change variables in (16) to  $\mathbf{z} = \mathbf{x} - \mathbf{x}^j(t)$ , we find

$$(17) \quad \int_{\mathbb{R}^2} \mathbf{z} \omega^j(\mathbf{z}, t) d\mathbf{z} = \mathbf{0}.$$

Since this equation holds for all  $t > 0$ , we can differentiate both sides with respect to  $t$  to obtain

$$(18) \quad \int_{\mathbb{R}^2} \mathbf{z} \partial_t \omega^j(\mathbf{z}, t) d\mathbf{z} = \mathbf{0}.$$

Using (15), we can insert the formula for  $\partial_t \omega^j$  into this integral, and we obtain

$$(19) \quad \nu \int_{\mathbb{R}^2} \mathbf{z} \Delta \omega^j(\mathbf{z}, t) d\mathbf{z} - \int_{\mathbb{R}^2} \mathbf{z} \left( \sum_{\ell=1}^N \mathbf{u}^\ell(\mathbf{z} + \mathbf{x}^j(t) - \mathbf{x}^\ell(t), t) \right) \cdot \nabla \omega^j(\mathbf{z}, t) d\mathbf{z} + \int_{\mathbb{R}^2} \mathbf{z} (\dot{\mathbf{x}}^j(t) \cdot \nabla \omega^j(\mathbf{z}, t)) d\mathbf{z} = \mathbf{0}.$$

We first note that if we integrate twice by parts, we have

$$(20) \quad \int_{\mathbb{R}^2} \mathbf{z} \Delta \omega^j(\mathbf{z}, t) d\mathbf{z} = \mathbf{0}.$$

Next if we take the  $n$ th component and integrate by parts, we find

$$(21) \quad \int_{\mathbb{R}^2} z_n (\dot{\mathbf{x}}^j(t) \cdot \nabla \omega^j(\mathbf{z}, t)) d\mathbf{z} = \sum_{m=1}^2 \dot{\mathbf{x}}_m^j(t) \int_{\mathbb{R}^2} z_n \partial_{z_m} \omega^j(\mathbf{z}, t) d\mathbf{z} = -m^j \dot{\mathbf{x}}_n^j(t),$$

where  $m^j = \int_{\mathbb{R}^2} \omega^j(\mathbf{z}, t) d\mathbf{z}$  and  $n = 1, 2$ .

**Remark 2.1.** Note that (15) does preserve the total integral (“mass”) of the solution, so this definition of  $m^j$  is consistent with (9).

Finally, recalling that the velocity field is incompressible, we can rewrite

$$\left( \sum_{\ell=1}^N \mathbf{u}^\ell(\mathbf{z} + \mathbf{x}^j(t) - \mathbf{x}^\ell(t), t) \right) \cdot \nabla \omega^j(\mathbf{z}, t) = \nabla \cdot \left( \sum_{\ell=1}^N \mathbf{u}^\ell(\mathbf{z} + \mathbf{x}^j(t) - \mathbf{x}^\ell(t), t) \omega^j(\mathbf{z}, t) \right),$$

and so, again considering the  $n$ th component and integrating by parts, we have

$$(22) \quad \int_{\mathbb{R}^2} z_n \left( \sum_{\ell=1}^N \mathbf{u}^\ell(\mathbf{z} + \mathbf{x}^j(t) - \mathbf{x}^\ell(t), t) \right) \cdot \nabla \omega^j(\mathbf{z}, t) d\mathbf{z} \\ = - \int_{\mathbb{R}^2} \left( \sum_{\ell=1}^N \mathbf{u}_n^\ell(\mathbf{z} + \mathbf{x}^j(t) - \mathbf{x}^\ell(t), t) \right) \omega^j(\mathbf{z}, t) d\mathbf{z}.$$

Thus, if we combine (20), (21), and (22), we see that (19) reduces to the system of ODEs for the centers of the vorticity distributions,

$$(23) \quad \frac{d\mathbf{x}_n^j}{dt}(t) = \frac{1}{m^j} \sum_{\ell=1}^N \int_{\mathbb{R}^2} \left( \mathbf{u}_n^\ell(\mathbf{z} + \mathbf{x}^j(t) - \mathbf{x}^\ell(t), t) \omega^j(\mathbf{z}, t) \right) d\mathbf{z},$$

supplemented by the initial conditions (10), while the  $N$  components of the vorticity evolve according to the PDEs (15) with initial conditions

$$(24) \quad \omega^j(\mathbf{z}, 0) = \omega_0^j(\mathbf{z} + \mathbf{x}_0^j),$$

obtained by combining (8) and (12).

**Remark 2.2.** Consider (23) in the limit in which the components  $\omega^j$  are all point vortices, i.e.,  $\omega^j(\mathbf{z}, t) = m^j \delta(\mathbf{z})$ , with  $\delta(\mathbf{z})$  the Dirac-delta function. Recall that the velocity field associated with such a point vortex is

$$U^j(\mathbf{z}, t) = m^j \frac{\mathbf{n} \times \mathbf{z}}{\|\mathbf{z}\|^2},$$

where  $\mathbf{n}$  is the unit normal vector to the plane. Then if we ignore the (singular) term with  $\ell = j$  in the sum on the right-hand side of (23), we find

$$(25) \quad \frac{d\mathbf{x}^j}{dt}(t) = \sum_{\ell=1; \ell \neq j}^N m^\ell \frac{\mathbf{n} \times (\mathbf{x}^j(t) - \mathbf{x}^\ell(t))}{\|\mathbf{x}^j(t) - \mathbf{x}^\ell(t)\|^2}.$$

These, of course, are just Helmholtz–Kirchhoff equations for the inviscid motion of a system of point vortices. Thus, our expansion can be regarded as a generalization of this approximation, which allows for both nonzero viscosity and vortices of finite size. To justify omitting the term with  $\ell = j$  on the right-hand side of (25) we note that if we approximate the delta function with a narrow Gaussian vorticity distribution, and  $\mathbf{u}^j$  by the corresponding velocity field, this term will vanish by symmetry.

**3. The moment expansion: Case of a single center.** In this section we introduce another idea—an expansion of the vorticity in terms of Hermite functions. Then, in the next section we will combine the Hermite expansion with the multivortex expansion of the previous section.

The moment expansion is an expansion of the solution of the vorticity equation in terms of Hermite functions. Define

$$(26) \quad \phi_{00}(\mathbf{x}, t; \lambda) = \frac{1}{\pi\lambda^2} e^{-|\mathbf{x}|^2/\lambda^2},$$

where  $\lambda^2 = \lambda_0^2 + 4\nu t$ . Three simple facts that we will use repeatedly are the following:

- (i)  $\partial_t \phi_{00} = \nu \Delta \phi_{00}$ .
- (ii)  $\int_{\mathbb{R}^2} \phi_{00}(\mathbf{x}, t; \lambda) d\mathbf{x} = 1$  for all  $t \geq 0$ .
- (iii) Finally, and crucially for what follows, the vorticity function  $\omega(\mathbf{x}, t) = \alpha \phi_{00}(\mathbf{x}, t)$  is an exact solution (called the Oseen, or Lamb, vortex) of the two-dimensional vorticity equation for all values of  $\alpha$ .

Note that we will often suppress the dependence of  $\phi_{00}$  on  $\lambda$  when there is no fear of confusion.

We now define the Hermite functions of order  $(k_1, k_2)$  by

$$(27) \quad \phi_{k_1, k_2}(\mathbf{x}, t; \lambda) = D_{x_1}^{k_1} D_{x_2}^{k_2} \phi_{00}(\mathbf{x}, t; \lambda)$$

and the corresponding moment expansion of a function by

$$(28) \quad \omega(\mathbf{x}, t) = \sum_{k_1, k_2=0}^{\infty} M[k_1, k_2; t] \phi_{k_1, k_2}(\mathbf{x}, t; \lambda).$$

We define the Hermite polynomials via their generating function:

$$(29) \quad H_{n_1, n_2}(\mathbf{z}; \lambda) = \left( D_{t_1}^{n_1} D_{t_2}^{n_2} e^{\left( \frac{2\mathbf{t} \cdot \mathbf{z} - |\mathbf{t}|^2}{\lambda^2} \right)} \right) \Big|_{t=0}.$$

Note that the “standard” Hermite polynomials correspond to taking  $\lambda = 1$ .

Then using the standard orthogonality relationship for the Hermite polynomials,

$$(30) \quad \int_{\mathbb{R}^2} H_{n_1, n_2}(\mathbf{z}; 1) H_{m_1, m_2}(\mathbf{z}; 1) e^{-z^2} d\mathbf{z} = \pi 2^{n_1+n_2} (n_1!) (n_2!) \delta_{n_1, m_1} \delta_{n_2, m_2},$$

we see that the coefficients in the expansion (28) are defined by the projection operators:

$$(31) \quad M[k_1, k_2; t] = (P_{k_1, k_2} \omega)(t) = \frac{(-1)^{k_1+k_2} \lambda^{2(k_1+k_2)}}{2^{k_1+k_2} (k_1!) (k_2!)} \int_{\mathbb{R}^2} H_{k_1, k_2}(\mathbf{z}; \lambda) \omega(\mathbf{z}, t) d\mathbf{z}.$$

Note that if the function  $\omega(\mathbf{x}, t)$  in (28) is the vorticity field of some fluid, the linearity of the Biot–Savart law implies that we can expand the associated velocity field as

$$(32) \quad \mathbf{V}(\mathbf{x}, t) = \sum_{k_1, k_2=0}^{\infty} M[k_1, k_2; t] \mathbf{V}_{k_1, k_2}(\mathbf{x}, t; \lambda),$$



where

$$(33) \quad \mathbf{V}_{k_1, k_2}(\mathbf{x}, t; \lambda) = D_{x_1}^{k_1} D_{x_2}^{k_2} \mathbf{V}_{00}(\mathbf{x}, t; \lambda)$$

and  $\mathbf{V}_{00}(\mathbf{x}, t; \lambda)$  is the velocity field associated with the Gaussian vorticity distribution  $\phi_{00}$ —explicitly we have

$$(34) \quad \mathbf{V}_{00}(\mathbf{x}, t; \lambda) = \frac{1}{2\pi} \frac{(-x_2, x_1)}{|\mathbf{x}|^2} \left(1 - e^{-|\mathbf{x}|^2/\lambda^2}\right).$$

**3.1. Convergence of the moment expansion.** In this subsection we derive a criterion for the convergence of the moment expansion derived above, and we show that if this criterion is satisfied for  $t = 0$ , then it is satisfied for all subsequent times  $t > 0$ .

Our convergence criterion is based on the observation that the Hermite functions  $\phi_{k_1, k_2}(\mathbf{x}, t; \lambda)$  are, for any value of  $t$ , the eigenfunctions of the linear operator

$$(35) \quad \mathcal{L}^\lambda \psi = \frac{1}{4} \lambda^2 \Delta \psi + \frac{1}{2} \nabla \cdot (\mathbf{x} \psi).$$

This fact can be verified by direct computation and is related to the fact that  $\mathcal{L}^\lambda$  can be transformed into the Hamiltonian quantum mechanical harmonic oscillator.

The Gaussian function  $\phi_{00}$  plays a crucial role in the convergence proof, and its dependence on the parameter  $\lambda$  is particularly important in this discussion, so for this subsection only we will define

$$\phi_{00}(\mathbf{x}, t; \lambda) = \Phi_\lambda(\mathbf{x}, t)$$

to emphasize this dependence.

If one now proceeds as in Lemma 4.7 of [4], one can prove the following.

**Proposition 3.1.** *The operator  $\mathcal{L}^\lambda$  is self-adjoint in the Hilbert space*

$$X^\lambda = \{f \in L^2(\mathbb{R}^2) \mid \Phi_\lambda^{-1/2} f \in L^2(\mathbb{R}^2)\}$$

with inner product  $(f, g)_\lambda = \int_{\mathbb{R}^2} \Phi_\lambda^{-1} \bar{f} g d\mathbf{x}$ .

An immediate corollary of this proposition and the general theory of self-adjoint operators is the following.

**Corollary 3.2.** *The eigenfunctions of  $\mathcal{L}^\lambda$  form a complete orthogonal set in the Hilbert space  $X^\lambda$ .*

As a corollary of this result and the observation that the eigenfunctions of  $\mathcal{L}^\lambda$  are precisely our Hermite functions  $\{\phi_{k_1, k_2}\}$ , we have finally the next claim.

**Proposition 3.3.** *Suppose that*

$$\|f\|_\lambda^2 = \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) |f(\mathbf{x})|^2 d\mathbf{x} < \infty;$$

then the expansion

$$f(\mathbf{x}) = \sum_{k_1, k_2} M[k_1, k_2] \phi_{k_1, k_2}(\mathbf{x})$$

converges with respect to the norm on the Hilbert space  $X^\lambda$ .

Thus, the following criterion guarantees that the expansion (28) for the vorticity converges:

$$(36) \quad \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x})(\omega(\mathbf{x}, t))^2 d\mathbf{x} < \infty.$$

The main result of this subsection is the following theorem, which proves that if our initial vorticity distribution satisfies (36) for some  $\lambda = \lambda_0$ , then the solution of the vorticity equation with that initial condition will satisfy (36) for all time  $t$  with  $\lambda = \sqrt{4\nu t + \lambda_0^2}$ , and hence as a corollary if the initial vorticity distribution satisfies (36), then our moment expansion converges for all times  $t$ .

**Theorem 3.4.** *Define the weighted enstrophy function*

$$\mathcal{E}(t) = \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x})(\omega(\mathbf{x}, t))^2 d\mathbf{x}.$$

*If the initial vorticity distribution  $\omega_0$  is such that  $\mathcal{E}(0) < \infty$  for some  $\lambda_0$ , and  $\omega_0$  is bounded (in the  $L^\infty$  norm), then  $\mathcal{E}(t)$  is finite for all times  $t > 0$ .*

*Proof.* The idea of the proof is to derive a differential inequality for  $\mathcal{E}(t)$  which guarantees that if  $\mathcal{E}(0)$  is finite, then  $\mathcal{E}(t)$  will be finite for all  $t$ . Differentiating  $\mathcal{E}(t)$ , we obtain

$$(37) \quad \begin{aligned} \frac{d\mathcal{E}}{dt}(t) &= \frac{4\nu}{\lambda^2} \mathcal{E}(t) - \frac{4\nu}{\lambda^4} \int_{\mathbb{R}^2} |\mathbf{x}|^2 \Phi_\lambda^{-1}(\mathbf{x})(\omega(\mathbf{x}, t))^2 d\mathbf{x} \\ &\quad + 2 \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) \omega(\mathbf{x}, t) \partial_t \omega(\mathbf{x}, t) d\mathbf{x} \\ &= \frac{4\nu}{\lambda^2} \mathcal{E}(t) - \frac{4\nu}{\lambda^4} \int_{\mathbb{R}^2} |\mathbf{x}|^2 \Phi_\lambda^{-1}(\mathbf{x})(\omega(\mathbf{x}, t))^2 d\mathbf{x} \\ &\quad + 2 \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) \omega(\mathbf{x}, t) (\nu \Delta \omega - \mathbf{u} \cdot \nabla \omega) d\mathbf{x}. \end{aligned}$$

We now consider the last term in (37). First note that upon integration by parts we have

$$(38) \quad 2 \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) \omega(\mathbf{x}, t) (\nu \Delta \omega(\mathbf{x}, t)) d\mathbf{x} = -2\nu \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) \left( |\nabla \omega|^2 + \frac{2}{\lambda^2} \omega \mathbf{x} \cdot \nabla \omega \right) d\mathbf{x}.$$

The right-hand side of the expression in (38) can again be broken up into two pieces, and the second can be bounded by

$$(39) \quad 2\nu \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) \left( \frac{2}{\lambda^2} \omega \mathbf{x} \cdot \nabla \omega \right) d\mathbf{x} \leq \nu \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) |\nabla \omega|^2 d\mathbf{x} + \frac{4\nu}{\lambda^4} \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) (\mathbf{x}^2 \omega^2) d\mathbf{x}.$$

Finally, we bound the last term in (37), which comes from the nonlinear term in the vorticity equation. In this estimate we use the fact (see [3, Lemma 2.1]) that the  $L^\infty$  norm of the velocity field  $\mathbf{u}$  can be bounded by a constant times the sum of the  $L^1$  and  $L^\infty$  norms of the vorticity field—i.e., by  $C(\|\omega\|_{L^1(\mathbb{R}^2)} + \|\omega\|_{L^\infty(\mathbb{R}^2)})$ . This observation, combined with the fact that  $\|\omega(\cdot, t)\|_{L^p(\mathbb{R}^2)} \leq \|\omega_0\|_{L^p(\mathbb{R}^2)}$ , which is a consequence of the maximum principle, implies that

$$\|\mathbf{u}(\cdot, t)\|_{L^\infty(\mathbb{R}^2)} \leq C(\|\omega_0\|_{L^1(\mathbb{R}^2)} + \|\omega_0\|_{L^\infty(\mathbb{R}^2)})$$

and hence that

$$(40) \quad \begin{aligned} 2 \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) \omega(\mathbf{x}, t) (\mathbf{u} \cdot \nabla \omega) \, d\mathbf{x} &\leq 2C(\omega_0) \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) |\omega(\mathbf{x}, t)| |\nabla \omega| \, d\mathbf{x} \\ &\leq \frac{4C(\omega_0)}{\nu} \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) (\omega(\mathbf{x}, t))^2 \, d\mathbf{x} + \nu \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) |\nabla \omega|^2 \, d\mathbf{x}. \end{aligned}$$

If we now combine the inequalities in (38), (39), and (40) with the expression for  $\frac{d\mathcal{E}}{dt}$  in (37), we obtain

$$(41) \quad \frac{d\mathcal{E}}{dt}(t) \leq \left( \frac{4C(\omega_0)}{\nu} + \frac{4\nu}{\lambda^2} \right) \mathcal{E}(t),$$

from which we see immediately that if  $\mathcal{E}(0)$  is bounded,  $\mathcal{E}(t)$  remains bounded for all time. ■

**3.2. Differential equations for the moments.** Assuming that the function  $\omega(\mathbf{z}, t)$  is a solution of (7), we can derive differential equations satisfied by the moments  $M[k_1, k_2, t]$  in (28). Surprisingly the expressions for the coefficients in these expansions are quite simple and explicit.

If we differentiate (28) and assume that  $\omega$  is a solution of the two-dimensional vorticity equation, we obtain

$$(42) \quad \begin{aligned} \partial_t \omega &= \sum_{k_1, k_2=0}^{\infty} \frac{dM[k_1, k_2; t]}{dt} \phi_{k_1, k_2}(\mathbf{x}, t; \lambda) + \sum_{k_1, k_2=0}^{\infty} M[k_1, k_2; t] \partial_t \phi_{k_1, k_2}(\mathbf{x}, t; \lambda) \\ &= \sum_{k_1, k_2=0}^{\infty} M[k_1, k_2; t] (\nu \Delta \phi_{k_1, k_2}(\mathbf{x}, t; \lambda)) \\ &\quad - \left( \sum_{\ell_1, \ell_2=0}^{\infty} M[\ell_1, \ell_2; t] \mathbf{V}_{\ell_1, \ell_2}(\mathbf{x}, t; \lambda) \right) \cdot \nabla \left( \sum_{k_1, k_2=0}^{\infty} M[k_1, k_2; t] \phi_{k_1, k_2}(\mathbf{x}, t; \lambda) \right). \end{aligned}$$

From the first of the “simple facts” we stated about  $\phi_{00}$ , we see that the last term on the first line cancels the middle line, and hence if we apply the projection operators defined in (31), we are left with the system of ODEs for the moments,

$$(43) \quad \frac{dM[k_1, k_2; t]}{dt} = -P_{k_1, k_2} \left[ \left( \sum_{\ell_1, \ell_2=0}^{\infty} M[\ell_1, \ell_2; t] \mathbf{V}_{\ell_1, \ell_2}(\mathbf{x}, t; \lambda) \right) \cdot \nabla \left( \sum_{m_1, m_2=0}^{\infty} M[m_1, m_2; t] \phi_{m_1, m_2}(\mathbf{x}, t; \lambda) \right) \right].$$

The somewhat surprising fact which, in our opinion, makes the preceding straightforward calculations interesting is that the projection on the right-hand side of (43) can be computed explicitly in terms of the derivatives of a relatively simple function.

We now explain how this is done. First recall that

$$(44) \quad \phi_{m_1, m_2}(\mathbf{x}, t; \lambda) = D_{x_1}^{m_1} D_{x_2}^{m_2} \phi_{00}(\mathbf{x}, \lambda), \quad \mathbf{V}_{\ell_1, \ell_2}(\mathbf{x}, t; \lambda) = D_{x_1}^{\ell_1} D_{x_2}^{\ell_2} \mathbf{V}_{00}(\mathbf{x}, \lambda).$$

In order to avoid confusing the two sets of derivatives we will rewrite these formulas as

$$(45) \quad \begin{aligned} \phi_{m_1, m_2}(\mathbf{x}, t; \lambda) &= (D_{a_1}^{m_1} D_{a_2}^{m_2} \phi_{00}(\mathbf{x} + \mathbf{a}, \lambda))|_{\mathbf{a}=0}, \\ \mathbf{V}_{\ell_1, \ell_2}(\mathbf{x}, t; \lambda) &= (D_{b_1}^{\ell_1} D_{b_2}^{\ell_2} \mathbf{V}_{00}(\mathbf{x} + \mathbf{b}, \lambda))|_{\mathbf{b}=0}. \end{aligned}$$

Inserting these formulas into the right-hand side of (43) and using the formula for the projection operator  $P_{k_1, k_2}$  in terms of the integration against a Hermite polynomial, we obtain

$$(46) \quad \begin{aligned} \frac{dM}{dt}[k_1, k_2, t] &= -\frac{(-1)^{(k_1+k_2)} \lambda^{2(k_1+k_2)}}{2^{k_1+k_2} (k_1!) (k_2!)} \sum_{\ell_1, \ell_2} \sum_{m_1, m_2} M[\ell_1, \ell_2, t] M[m_1, m_2, t] \\ &\quad \times \int_{\mathbb{R}^2} H_{k_1, k_2}(\mathbf{x}) (D_{x_1}^{m_1} D_{x_2}^{m_2} \mathbf{V}_{00}(\mathbf{x}; \lambda)) \cdot \nabla_{\mathbf{x}} (D_{x_1}^{\ell_1} D_{x_2}^{\ell_2} \phi_{00}(\mathbf{x}; \lambda)) d\mathbf{x} \\ &= -\frac{(-1)^{(k_1+k_2)} \lambda^{2(k_1+k_2)}}{2^{k_1+k_2} (k_1!) (k_2!)} \sum_{\ell_1, \ell_2} \sum_{m_1, m_2} M[\ell_1, \ell_2, t] M[m_1, m_2, t] \\ &\quad \times D_{t_1}^{k_1} D_{t_2}^{k_2} D_{b_1}^{m_1} D_{b_2}^{m_2} D_{a_1}^{\ell_1} D_{a_2}^{\ell_2} \nabla_{\mathbf{a}} \\ &\quad \cdot \left( \int_{\mathbb{R}^2} e^{\frac{(-t_1^2 - t_2^2 + 2t_1 x_1 + 2t_2 x_2)}{\lambda^2}} \mathbf{V}_{00}(\mathbf{x} + \mathbf{b}; \lambda) \phi_{00}(\mathbf{x} + \mathbf{a}; \lambda) d\mathbf{x} \right) \Big|_{\mathbf{t}=0, \mathbf{a}=0, \mathbf{b}=0}. \end{aligned}$$

The last equality in this expression results from rewriting the Hermite polynomial  $H_{k_1, k_2}$  in terms of its generating function.

The last step in deriving the equations for the moments is to evaluate the integral in the last line of (46). The key step in this evaluation is to recall that for these incompressible flows the velocity field can be written in terms of the derivatives of the stream function and that the Laplacian of the stream function is minus the vorticity. Thus, we can write

$$(47) \quad \mathbf{V}_{00}(\mathbf{x} + \mathbf{b}; \lambda) = -\nabla_{\mathbf{b}}^* (\Delta_{\mathbf{b}})^{-1} \phi_{00}(\mathbf{x} + \mathbf{b}),$$

where  $\nabla_{\mathbf{b}}^* f = (\partial_{x_2} f, -\partial_{x_1} f)$ .

Inserting this into the integral in (46), we find

$$(48) \quad \begin{aligned} \int_{\mathbb{R}^2} e^{\frac{(-t_1^2 - t_2^2 + 2t_1 x_1 + 2t_2 x_2)}{\lambda^2}} \mathbf{V}_{00}(\mathbf{x} + \mathbf{b}; \lambda) \phi_{00}(\mathbf{x} + \mathbf{a}; \lambda) d\mathbf{x} \\ = -\nabla_{\mathbf{b}}^* (\Delta_{\mathbf{b}})^{-1} \int_{\mathbb{R}^2} e^{\frac{(-t_1^2 - t_2^2 + 2t_1 x_1 + 2t_2 x_2)}{\lambda^2}} \phi_{00}(\mathbf{x} + \mathbf{b}; \lambda) \phi_{00}(\mathbf{x} + \mathbf{a}; \lambda) d\mathbf{x}. \end{aligned}$$

Now note that all three factors in the integrand are Gaussians, and thus the integral can be evaluated explicitly, and we find

$$(49) \quad \begin{aligned} \int_{\mathbb{R}^2} e^{\frac{(-t_1^2 - t_2^2 + 2t_1 x_1 + 2t_2 x_2)}{\lambda^2}} \phi_{00}(\mathbf{x} + \mathbf{b}; \lambda) \phi_{00}(\mathbf{x} + \mathbf{a}; \lambda) d\mathbf{x} \\ = \frac{1}{2\pi\lambda^2} e^{-\frac{1}{2\lambda^2} (a_1^2 + a_2^2 - 2a_1 b_1 + b_1^2 - 2a_2 b_2 + b_2^2 + 2a_1 t_1 + 2b_1 t_1 + t_1^2 + 2a_2 t_2 + 2b_2 t_2 + t_2^2)}. \end{aligned}$$

We next compute the expression

$$(50) \quad -\nabla_{\mathbf{b}}^*(\Delta_{\mathbf{b}})^{-1} \frac{1}{2\pi\lambda^2} e^{-\frac{1}{2\lambda^2}(a_1^2+a_2^2-2a_1b_1+b_1^2-2a_2b_2+b_2^2+2a_1t_1+2b_1t_1+t_1^2+2a_2t_2+2b_2t_2+t_2^2)}.$$

Recall that, given a vorticity field  $\omega$ ,  $-(\Delta)^{-1}\omega$  is the associated stream function and  $-\nabla^*(\Delta)^{-1}\omega$  the velocity field associated with  $\omega$ . Since the inverse Laplacian and derivatives in (50) act only on the  $\mathbf{b}$ -dependent parts of the expression, we need to evaluate

$$(51) \quad -\nabla_{\mathbf{b}}^*(\Delta_{\mathbf{b}})^{-1} \frac{1}{2\pi\lambda^2} e^{-\frac{1}{2\lambda^2}(-2a_1b_1+b_1^2-2a_2b_2+b_2^2+2b_1t_1+2b_2t_2)} \\ = -e^{\frac{1}{2\lambda^2}((t_1-a_1)^2+(t_2-a_2)^2)} \nabla_{\mathbf{b}}^*(\Delta_{\mathbf{b}})^{-1} \frac{1}{2\pi\lambda^2} e^{-\frac{1}{2\lambda^2}((b_1+(t_1-a_1))^2+(b_2+(t_2-a_2))^2)}.$$

However,

$$-\nabla_{\mathbf{b}}^*(\Delta_{\mathbf{b}})^{-1} \frac{1}{\pi\lambda^2} e^{-\frac{1}{2\lambda^2}((b_1+(t_1-a_1))^2+(b_2+(t_2-a_2))^2)}$$

is just the velocity field associated with a Gaussian vorticity distribution (i.e., an Oseen vortex) centered at the point  $-((t_1 - a_1), (t_2 - a_2))$  which we know explicitly. Hence, the expression on the right-hand side of (48) has the explicit representation

$$(52) \quad \frac{1}{2\pi} e^{-\frac{2}{\lambda^2}(t_1a_1+t_2a_2)} \frac{(-(b_2 + (t_2 - a_2)), (b_1 + (t_1 - a_1)))}{(b_1 + (t_1 - a_1))^2 + (b_2 + (t_2 - a_2))^2} \\ \times \left(1 - e^{-\frac{1}{2\lambda^2}((b_1+(t_1-a_1))^2+(b_2+(t_2-a_2))^2)}\right).$$

If we now return to (46), we see that in order to compute the coefficients in the moment equations we need to evaluate the divergence of this last expression with respect to  $\mathbf{a}$ , which gives

$$K(a_1, a_2, b_1, b_2, t_1, t_2; \lambda) = \frac{1}{\pi\lambda^2} e^{-\frac{2}{\lambda^2}(t_1a_1+t_2a_2)} \\ \times \left( \frac{-a_2 t_1 + b_2 t_1 + a_1 t_2 - b_1 t_2}{((b_1 + (t_1 - a_1))^2 + (b_2 + (t_2 - a_2))^2)} \right) \left(1 - e^{-\frac{1}{2\lambda^2}((b_1+(t_1-a_1))^2+(b_2+(t_2-a_2))^2)}\right).$$

Returning to (46), we finally conclude that

$$(53) \quad \frac{dM}{dt}[k_1, k_2, t] = -\frac{(-1)^{(k_1+k_2)}\lambda^{2(k_1+k_2)}}{2^{k_1+k_2}(k_1!)(k_2!)} \sum_{\ell_1, \ell_2} \sum_{m_1, m_2} \Gamma[k_1, k_2; \ell_1, \ell_2, m_1, m_2; \lambda] M[\ell_1, \ell_2, t] M[m_1, m_2, t],$$

where

$$(54) \quad \Gamma[k_1, k_2; \ell_1, \ell_2, m_1, m_2; \lambda] \\ = D_{t_1}^{k_1} D_{t_2}^{k_2} D_{b_1}^{m_1} D_{b_2}^{m_2} D_{a_1}^{\ell_1} D_{a_2}^{\ell_2} K(a_1, a_2, b_1, b_2, t_1, t_2; \lambda)|_{t=0, a=0, b=0}.$$

Thus, we have succeeded in rewriting the two-dimensional vorticity equation as a system of ODEs with simple, quadratic nonlinear terms whose coefficients can be evaluated in terms of derivatives of a single explicit function. Furthermore, we have given a sufficient condition on the initial vorticity distribution to guarantee that the expansion of the vorticity generated by the solution of these ODEs converges for all time.

**4. The moment expansion for several vortex centers.** In this section we extend the Hermite moment expansion of the previous section to the case in which there are two or more centers of vorticity by combining this expansion with the multivortex representation of section 2. For simplicity of exposition we limit the discussion here to the case of two vortices, but the expansion can be extended to any finite number of vortices.

The basic idea is just to consider (15) for the evolution of each vortex and then expand each of the functions  $\omega^j$  in Hermite moments as in the previous section. Thus, we define

$$(55) \quad \omega^j(\mathbf{z}, t) = \sum_{k_1, k_2=0}^{\infty} M^j[k_1, k_2; t] \phi_{k_1, k_2}(\mathbf{z}, t; \lambda)$$

for  $j = 1, 2$ . We make a similar expansion for the velocity field in terms of the functions  $\mathbf{V}_{\ell_1, \ell_2}$  and insert the expansions into (15). Letting  $\mathbf{z} = \mathbf{x} - \mathbf{x}^j(t)$  and recalling that  $\partial_t \phi_{k_1, k_2} = \nu \Delta \phi_{k_1, k_2}$ , we obtain

$$(56) \quad \frac{dM^j[k_1, k_2; t]}{dt} = -P_{k_1, k_2} \left[ \left( \sum_{j'=1}^2 \sum_{\ell_1, \ell_2=0}^{\infty} M^{j'}[\ell_1, \ell_2; t] \mathbf{V}_{\ell_1, \ell_2}(\mathbf{z} + \mathbf{s}_{j, j'}, t; \lambda) \right) \cdot \nabla \left( \sum_{m_1, m_2=0}^{\infty} M^j[m_1, m_2; t] \phi_{m_1, m_2}(\mathbf{z}, t; \lambda) \right) \right] + P_{k_1, k_2} \left[ \dot{\mathbf{x}}^j(t) \cdot \nabla \left( \sum_{m_1, m_2=0}^{\infty} M^j[m_1, m_2; t] \phi_{m_1, m_2}(\mathbf{z}, t; \lambda) \right) \right],$$

where  $\mathbf{s}_{j, j'} = \mathbf{x}^{j'}(t) - \mathbf{x}^j(t)$ . For the first projection term we proceed as in the previous section, and using (46), we finally conclude that the first term is equal to

$$-\frac{(-1)^{(k_1+k_2)} \lambda^{2(k_1+k_2)}}{2^{k_1+k_2} (k_1!) (k_2!)} \sum_{j'=1}^2 \sum_{\ell_1, \ell_2} \sum_{m_1, m_2} \Gamma^{j, j'}[k_1, k_2; \ell_1, \ell_2, m_1, m_2; \mathbf{s}_{j, j'}, \lambda] \times M^j[\ell_1, \ell_2, t] M^{j'}[m_1, m_2, t],$$

where

$$(57) \quad \Gamma^{j, j'}[k_1, k_2; \ell_1, \ell_2, m_1, m_2; \mathbf{s}_{j, j'}, \lambda] = D_{t_1}^{k_1} D_{t_2}^{k_2} D_{b_1}^{m_1} D_{b_2}^{m_2} D_{a_1}^{\ell_1} D_{a_2}^{\ell_2} K^{multi}(a_1, a_2, b_1, b_2, t_1, t_2; \mathbf{s}_{j, j'}, \lambda)|_{\mathbf{t}=0, \mathbf{a}=0, \mathbf{b}=0}$$

and

$$(58) \quad K^{multi}(a_1, a_2, b_1, b_2, t_1, t_2; \mathbf{s}, \lambda) = \nabla_{\mathbf{a}} \cdot \left( \int_{\mathbb{R}^2} e^{\frac{(-t_1^2 - t_2^2 + 2t_1 x_1 + 2t_2 x_2)}{\lambda^2}} \mathbf{V}_{00}(\mathbf{x} - \mathbf{s} + \mathbf{b}; \lambda) \phi_{00}(\mathbf{x} + \mathbf{a}; \lambda) d\mathbf{x} \right).$$

**Remark 4.1.** Note that, comparing (57) with the calculation leading up to (53), we see that  $K^{multi}$  can be written in terms of the expression for  $K$  via the simple formula

$$K^{multi}(a_1, a_2, b_1, b_2, t_1, t_2; s_1, s_2, \lambda) = K(a_1, a_2, b_1 - s_1, b_2 - s_2, t_1, t_2; \lambda).$$

The second projection term in (56) can be reduced to

$$(59) \quad \dot{x}_1^j(t)M^j[k_1 - 1, k_2, t] + \dot{x}_2^j(t)M^j[k_1, k_2 - 1, t];$$

thus we can combine the above equations to write

$$(60) \quad \begin{aligned} \frac{dM^j}{dt}[k_1, k_2, t] &= -\frac{(-1)^{(k_1+k_2)}\lambda^{2(k_1+k_2)}}{2^{k_1+k_2}(k_1!)(k_2!)} \sum_{j'=1}^2 \sum_{\ell_1, \ell_2} \sum_{m_1, m_2} \Gamma^{j,j'}[k_1, k_2; \ell_1, \ell_2, m_1, m_2; \mathbf{s}_{j,j'}, \lambda] \\ &\quad \times M^j[\ell_1, \ell_2, t]M^{j'}[m_1, m_2, t] \\ &\quad + \dot{x}_1^j(t)M^j[k_1 - 1, k_2, t] + \dot{x}_2^j(t)M^j[k_1, k_2 - 1, t]. \end{aligned}$$

We now derive a similar expansion for the evolution of the centers of each vortex. We begin with (23),

$$\frac{d\mathbf{x}^j}{dt}(t) = \frac{1}{m^j} \sum_{j'=1}^N \int_{\mathbb{R}^2} \left( \mathbf{u}^\ell(\mathbf{z} + \mathbf{x}^j(t) - \mathbf{x}^{j'}(t), t)\omega^j(\mathbf{z}, t) \right) d\mathbf{z}.$$

(Recall that this is really a pair of equations, one for each component of  $\mathbf{x}^j$ .) Now insert the moment expansion of  $\mathbf{u}^{j'}$  and  $\omega^j$  into this expression, to obtain

$$(61) \quad \frac{d\mathbf{x}^j}{dt}(t) = \sum_{j'=1}^N \sum_{\ell_1, \ell_2} \sum_{m_1, m_2} M^j[\ell_1, \ell_2, t]M^{j'}[m_1, m_2, t] \int_{\mathbb{R}^2} \mathbf{V}_{\ell_1, \ell_2}(\mathbf{z} - \mathbf{s}_{j,j'}, t)\phi_{\ell_1, \ell_2}(\mathbf{z}, t)d\mathbf{z},$$

where, as before,  $\mathbf{s}_{j,j'} = \mathbf{x}^{j'}(t) - \mathbf{x}^j(t)$ . The integral of the velocity and vorticity can be evaluated just as in the preceding section, and we find

$$(62) \quad \frac{d\mathbf{x}^j}{dt}(t) = \sum_{j'=1}^N \sum_{\ell_1, \ell_2} \sum_{m_1, m_2} \Xi^{j,j'}[\ell_1, \ell_2, m_1, m_2; \mathbf{s}_{j,j'}, \lambda]M^j[\ell_1, \ell_2, t]M^{j'}[m_1, m_2, t],$$

where in this case the coefficients  $\Xi^{j,j'}[\ell_1, \ell_2, m_1, m_2; \mathbf{s}_{j,j'}, \lambda]$  are given by the expression

$$(63) \quad \begin{aligned} &\Xi^{j,j'}[\ell_1, \ell_2, m_1, m_2; s, \lambda] \\ &= \frac{1}{2\pi} D_{b_1}^{m_1} D_{b_2}^{m_2} D_{a_1}^{\ell_1} D_{a_2}^{\ell_2} \left( \frac{(-(b_2 - s_2 - a_2), (b_1 - s_1 - a_1))}{(b_2 - s_2 - a_2)^2 + (b_1 - s_1 - a_1)^2} \right. \\ &\quad \left. \times \left( 1 - e^{-\frac{1}{2\lambda^2}((b_2 - s_2 - a_2)^2 + (b_1 - s_1 - a_1)^2)} \right) \right) \Big|_{\mathbf{a}=\mathbf{0}, \mathbf{b}=\mathbf{0}}. \end{aligned}$$

**Remark 4.2.** Equations (62) and (63) could be expressed in terms of Cartesian tensors.

**4.1. Convergence of multivortex expansion.** We note that it is easy to extend our previous result on the convergence of the moment expansion to the multivortex expansion. To do so, we change variables to recenter each vortex at the origin. Thus if we let

$$\omega^j(\mathbf{x} - \mathbf{x}^j(t), t) = w^j(\mathbf{x}, t), \quad \mathbf{u}^j(\mathbf{x} - \mathbf{x}^j(t), t) = \mathbf{v}^j(\mathbf{x}, t),$$

then (15) becomes

$$\frac{\partial w^j}{\partial t}(\mathbf{x}, t) = \nu \Delta w^j(\mathbf{x}, t) - \left( \sum_{\ell=1}^N \mathbf{v}^\ell(\mathbf{x} - \mathbf{x}^\ell(t) + \mathbf{x}^j(t), t) \right) \cdot \nabla w^j(\mathbf{x}, t),$$

for  $j = 1, \dots, N$ .

We are now ready to state our result.

**Theorem 4.3.** *Define*

$$\mathcal{E}^j(t) = \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) w^j(\mathbf{x}, t)^2 d\mathbf{x}.$$

If the initial vorticity distribution  $\omega_0 \equiv \sum_{\ell=1}^N \omega^\ell(\mathbf{x} - \mathbf{x}^\ell(0), 0)$  is such that  $\mathcal{E}^j(0) < \infty$  for some  $\lambda_0$  and for all  $j = 1, 2, \dots, N$ , and if  $\omega_0$  is bounded (in the  $L^\infty$  norm), then each  $\mathcal{E}^j(t)$  is finite for all times  $t > 0$ .

*Proof.* We use the same idea as in the single vortex case and differentiate  $\mathcal{E}^j(t)$ :

$$\begin{aligned} \frac{d\mathcal{E}^i}{dt}(t) &= \frac{4\nu}{\lambda^2} \mathcal{E}^i(t) - \frac{4\nu}{\lambda^4} \int_{\mathbb{R}^2} |\mathbf{x}|^2 \Phi_\lambda^{-1}(\mathbf{x}) w^i(\mathbf{x})^2 d\mathbf{x} \\ &\quad + 2 \int_{\mathbb{R}^2} \Phi_\lambda^{-1}(\mathbf{x}) w^i(\mathbf{x}, t) \partial_t w^i(\mathbf{x}, t) d\mathbf{x} \\ &= \frac{4\nu}{\lambda^2} \mathcal{E}^i(t) - \frac{4\nu}{\lambda^4} \int_{\mathbb{R}^2} |\mathbf{x}|^2 \Phi_\lambda^{-1}(\mathbf{x}) w^i(\mathbf{x})^2 d\mathbf{x} \\ &\quad + 2 \int_{\mathbb{R}^2} \Phi_\lambda^{-1} w^i \left( \nu \Delta w^i - \left( \sum_{\ell=1}^N \mathbf{v}^\ell(\mathbf{x} - \mathbf{x}^\ell(t) + \mathbf{x}^j(t), t) \right) \cdot \nabla w^i \right) d\mathbf{x}. \end{aligned}$$

From here the proof proceeds identically to the proof of Theorem 3.4 except we must examine the nonlinear term that comes from the vorticity equation a bit more closely. In general, it is unknown whether  $w^j(\cdot, t)$  satisfies a maximum principle, but since  $(\sum_{\ell=1}^N \mathbf{v}^\ell(\mathbf{x} - \mathbf{x}^\ell(t) + \mathbf{x}^j(t), t)) = \mathbf{u}$ , the solution to (1), it does satisfy a maximum principle. Hence, as in subsection 3.1, we can bound the  $L^\infty$  norm of  $\sum_{\ell=1}^N \mathbf{v}^\ell(\mathbf{x} - \mathbf{x}^\ell(t) + \mathbf{x}^j(t), t)$  by a constant depending only on the initial vorticity distribution. Hence we proceed to bound the integral

$$\int_{\mathbb{R}^2} \Phi_\lambda^{-1} w^i \left( \left( \sum_{\ell=1}^N \mathbf{v}^\ell(\mathbf{x} - \mathbf{x}^\ell(t) + \mathbf{x}^j(t), t) \right) \cdot \nabla w^i \right) d\mathbf{x} \leq 2C(\omega_0) \int_{\mathbb{R}^2} \Phi_\lambda^{-1} |w^i| |\nabla w^i| d\mathbf{x},$$

and thus the rest of the bounds are the same. Again, putting everything together, we arrive at

$$(64) \quad \frac{d\mathcal{E}^i}{dt}(t) \leq \left( \frac{4\nu}{\lambda^2} + \frac{4C(\omega_0)}{\nu} \right) \mathcal{E}^i(t). \quad \blacksquare$$



**4.2. Interaction of Gaussian vortices.** The experimental and numerical work of [11] has shown that widely separated regions of vorticity can be well approximated by Gaussians for long periods of time. This corresponds to truncating our expansions for the vorticity so that they contain only a single term. In this subsection we analyze the equations that result from this truncation. This approximation can be viewed as a generalization of the Helmholtz–Kirchhoff approximation in which we include the effects of vorticity and finite core size to lowest order. We show that the total vorticity of each of the vortices is constant, while the centers of vorticity evolve along either straight lines or circles.

*Remark 4.4.* It should be noted here that the effect of truncating our expansion after one term allows for only viscosity as the driving force in vortex merger. Allowing more terms in the expansion introduces convective forces.

If we start with two Gaussian distributions for our initial vorticity with the same value of  $\lambda_0$ , and truncate the equations of motion for the moments so that all terms containing higher-order moments are omitted, then we can conclude that the two vortices travel along circular or straight line orbits around the “center of vorticity.” Moreover, the leading coefficients  $M^j[0, 0 : t]$  of the expansions are constant. To be precise, we let

$$(65) \quad \omega^1(\mathbf{x}, t) = M^1[0, 0; t]\phi_{0,0}(x - x^1(t), t; \lambda),$$

$$(66) \quad \omega^2(\mathbf{x}, t) = M^2[0, 0; t]\phi_{0,0}(x - x^2(t), t; \lambda).$$

Let us also write  $\mathbf{s}_{i,j} = \mathbf{x}^j(t) - \mathbf{x}^i(t)$ ; then using (56)–(58), we first calculate the evolution of  $M^1(0, 0 : t)$ :

$$\begin{aligned} \frac{dM^1}{dt}[0, 0; t] &= -M^1[0, 0; t]^2 K(0, 0, 0, 0, 0, 0, \lambda) \\ &\quad - M^1[0, 0; t]M^2[0, 0; t]K^{multi}(0, 0, 0, 0, 0, 0, \mathbf{s}_{1,2}, \lambda) \\ &= 0, \end{aligned}$$

since  $K(0, 0, 0, 0, 0, 0, \lambda) = 0$  and  $K^{multi}(0, 0, 0, 0, 0, 0, \mathbf{s}_{1,2}, \lambda) = 0$ , respectively. The calculation for  $M^2[0, 0; t]$  is the same. Thus the evolutions of the coefficients of leading order are constant.

*Remark 4.5.* The fact that  $M^1[0, 0; t]$  is constant in time is not a consequence of the truncation of the moment equations to first order. One can show that the equations of (14) conserve the zeroth moment of  $\omega^j$ , independent of any truncation.

More interesting, though, is the calculation for the evolution of  $\mathbf{x}^j(t)$ , the centers of these vortices. Again if we denote  $\mathbf{x}^j = (x_1^j, x_2^j)$ , then the evolution for each component as defined by (23) can be written as

$$(67) \quad \frac{dx_i^j}{dt} = \frac{1}{M^j[0, 0, t]} \int V_i^k(\mathbf{y} - \mathbf{s}_{1,2}, t)\omega^j(\mathbf{y}, t)d\mathbf{y}.$$

Now using (47) to evaluate  $V_i^k(\mathbf{y} - \mathbf{s}_{1,2}, t)$ , equation (67) yields the following equations for

each  $x_i^j$ ,  $j = 1, 2$ , and  $i = 1, 2$ :

$$\begin{aligned}
 \dot{x}_1^1 &= -\frac{M^2}{2\pi} \frac{\left( e^{-\frac{(x_1^1 - x_1^2)^2 + (x_2^1 - x_2^2)^2}{2\lambda(t)^2}} - 1 \right) (x_2^1 - x_2^2)}{(x_1^1 - x_1^2)^2 + (x_2^1 - x_2^2)^2}, \\
 \dot{x}_2^1 &= -\frac{M^2}{2\pi} \frac{\left( e^{-\frac{(x_1^1 - x_1^2)^2 + (x_2^1 - x_2^2)^2}{2\lambda(t)^2}} - 1 \right) (-x_1^1 + x_1^2)}{(x_1^1 - x_1^2)^2 + (x_2^1 - x_2^2)^2}, \\
 \dot{x}_1^2 &= \frac{M^1}{2\pi} \frac{\left( e^{-\frac{(x_1^1 - x_1^2)^2 + (x_2^1 - x_2^2)^2}{2\lambda(t)^2}} - 1 \right) (x_2^1 - x_2^2)}{(x_1^1 - x_1^2)^2 + (x_2^1 - x_2^2)^2}, \\
 \dot{x}_2^2 &= \frac{M^1}{2\pi} \frac{\left( e^{-\frac{(x_1^1 - x_1^2)^2 + (x_2^1 - x_2^2)^2}{2\lambda(t)^2}} - 1 \right) (-x_1^1 + x_1^2)}{(x_1^1 - x_1^2)^2 + (x_2^1 - x_2^2)^2},
 \end{aligned}
 \tag{68}$$

where  $M^j \equiv M^j[0, 0, t]$  is constant and represents the total vorticity of the  $j$ th vortex.

**Remark 4.6.** *If the vortices have different  $\lambda(t)$  values, say  $\lambda_1(t)$  and  $\lambda_2(t)$ , then one just needs to replace the  $2\lambda(t)^2$  with  $\lambda_1(t)^2 + \lambda_2(t)^2$  in the exponential to arrive at the correct system.*

We now state the main result of this section.

**Theorem 4.7.** *System (68) admits only circular or straight-line trajectories.*

*Proof.* Away from rest points our system (68) can be transformed into

$$\begin{aligned}
 \frac{\partial x_2^1}{\partial x_1^1} &= \frac{-x_1^1 + x_1^2}{x_2^1 - x_2^2}, \\
 \frac{\partial x_2^2}{\partial x_1^2} &= \frac{-x_1^1 + x_1^2}{x_2^1 - x_2^2}, \\
 \frac{\partial x_2^1}{\partial x_2^2} &= -\frac{M^2}{M^1}, \\
 \frac{\partial x_1^1}{\partial x_1^2} &= -\frac{M^2}{M^1}.
 \end{aligned}
 \tag{69}$$

Integrating out the bottom two equations, we get

$$\begin{aligned}
 x_2^1 &= -\frac{M^2}{M^1}(x_2^2 + k_2), \\
 x_1^1 &= -\frac{M^2}{M^1}(x_1^2 + k_1).
 \end{aligned}
 \tag{70}$$

First let us assume that  $M^2 \neq -M^1$ . Then plugging back into the first two equations

of (69), we arrive at

$$(71) \quad \begin{aligned} \frac{\partial x_2^1}{\partial x_1^1} &= \frac{\left(-1 - \frac{M^2}{M^1}\right) x_1^1 - k_1}{\left(1 + \frac{M^2}{M^1}\right) x_2^1 - k_2} = -\frac{x_1^1 - \tilde{k}_1}{x_2^1 - \tilde{k}_2}, \\ \frac{\partial x_2^2}{\partial x_1^2} &= \frac{\left(1 + \frac{M^2}{M^1}\right) x_1^2 - \frac{M^2}{M^1} k_1}{\left(-1 - \frac{M^2}{M^1}\right) x_2^2 - \frac{M^2}{M^1} k_2} = -\frac{x_1^2 - \hat{k}_1}{x_2^2 - \hat{k}_2} \end{aligned}$$

for appropriate constants  $\tilde{k}_1, \tilde{k}_2, \hat{k}_1, \hat{k}_2$ . Thus we integrate again and get

$$(72) \quad \begin{aligned} (x_2^1(t) - \tilde{k}_2)^2 + (x_1^1(t) - \tilde{k}_1)^2 &= C_1, \\ (x_2^2(t) - \hat{k}_2)^2 + (x_1^2(t) - \hat{k}_1)^2 &= C_2. \end{aligned}$$

If  $M^2 = -M^1$ , then we have equal but opposite size vortices, and (68) becomes

$$(73) \quad \begin{aligned} \frac{\partial x_2^1}{\partial x_1^1} &= -\frac{k_2}{k_1}, \\ \frac{\partial x_2^2}{\partial x_1^2} &= -\frac{k_2}{k_1}, \end{aligned}$$

which gives us straight-line solutions with slope  $-\frac{k_2}{k_1}$ , as desired. ■

If we now consider the case of equal total vorticity for the two vortices ( $M^2 = M^1 \equiv M$ ), the classical point vortex result is that the vortices will rotate around the center of vorticity at a constant frequency,  $\frac{2M}{2\pi D^2}$ , where  $D$  is the distance between the vortex centers. We will now compute the viscous and finite core size effects on the frequency of rotation predicted by our model. For simplicity we will center the vortices at the origin and place them on the circle of radius  $r$ . We apply the polar change of variables,

$$(74) \quad x_1^i = r \cos(\theta_i), \quad x_2^i = r \sin(\theta_i), \quad i = 1, 2;$$

then, using the fact that our vortices are out of phase by  $\pi$ , we can compute that

$$(75) \quad (x_1^1 - x_1^2) = r \cos(\theta_1) - r \cos(\theta_2) = 2r \cos(\theta_1),$$


$$(76) \quad (x_2^1 - x_2^2) = r \sin(\theta_1) - r \sin(\theta_2) = 2r \sin(\theta_1),$$

and thus we arrive at the expression for the frequency of rotation

$$(77) \quad \Omega = \frac{M}{4\pi r^2} \left(1 - e^{\frac{-2r^2}{\lambda(t)^2}}\right).$$

Since  $\lambda(t)^2 = \lambda_0^2 + 4\nu t$  we notice three things: the first is that viscosity,  $\nu$ , slows the frequency of rotation down and, second, in the formal limit as  $\nu \rightarrow 0$  and  $\lambda_0^2 \rightarrow 0$  we recover the constant frequency,  $\frac{2M}{2\pi D^2}$ , which is what the Helmholtz–Kirchhoff model for the rotation of two point vortices predicts. The third property we notice is that the frequency predicted

“Typical” frequency of rotation for two localized vortices



**Figure 1.** Here we plot both the frequency of rotation predicted by single vortex expansion up to quadrupole order (dashed) and the frequency predicted by two vortex expansions truncated to leading order (solid). The parameter values used are  $\nu = .01$ ,  $M = 1$ ,  $r = 1$ , and  $\lambda_0 = .01$ .

by (77) is different than  $\Omega = \frac{M}{4\pi r^2}(1 - e^{-4r^2/\lambda(t)^2})$ , which is obtained by naively computing the velocity field of each Gaussian vortex in the center of the other vortex.

With this calculation we may now compare the results of expanding a two-Gaussian initial distribution as a single vortex or as two independent vortices. In [12] a two-Gaussian initial distribution at a distance of  $2r$  apart with core size  $\lambda_0$ , each with mass  $M$ , is approximated by a single vortex expansion using (28). The authors truncate the expansion to a quadrupole moment ( $n = 2$ ) and calculate the frequency of rotation to be

$$(78) \quad \Omega = \frac{M}{8\pi} \left[ \frac{1}{2\nu t} \ln \left( 1 + \frac{4\nu t}{r^2} \right) - \frac{\lambda_0^2}{r^4} \frac{1}{1 + 4\nu t/r^2} \right].$$

This equation is directly comparable to (77). Notice that both equations for frequency of rotation indicate slowing of rotation over time, albeit at different rates. In addition, both equations recover the Helmholtz–Kirchhoff approximation of  $\frac{2M}{2\pi D^2}$  in the limit as  $\nu \rightarrow 0$  and  $\lambda_0 \rightarrow 0$ . In fact, in the sufficiently localized regime,  $r \ll \lambda_0$ , both equations have similar initial frequencies and remain close asymptotically. A typical example is shown in Figure 1.

**5. Conclusions.** In this paper we have derived a system of ODEs whose solutions give a representation of solutions of the two-dimensional vorticity equation in terms of a system of interacting vortices. We have also derived a sufficient condition on the initial vorticity distribution which guarantees that this representation in terms of interacting vortices is equivalent to the original solution of the two-dimensional vorticity equation. This model generalizes the classical Helmholtz–Kirchhoff model of interacting, inviscid, point vortices to include the effects of both finite core size and viscosity. A given initial vorticity distribution may be represented in various ways by our model. As an example, we have considered the predicted frequency of rotation by representing a spinning vortex as a pair of interacting Gaussian vortices versus a single vortex but including higher-order moments. While the predictions of the

two approaches were similar in this example, we expect that in general the preferred approach will be governed by the details of the initial conditions.

**Acknowledgments.** CEW wishes to acknowledge many useful discussions about two-dimensional fluid flows with Th. Gallay. DU and CEW wish also to acknowledge a very helpful discussion of Hermite expansions with G. Van Baalen. Any findings, conclusions, opinions, or recommendations are those of the authors and do not necessarily reflect the views of the NSF.

## REFERENCES

- [1] O. AGULLO AND A. VERGA, *Effect of viscosity in the dynamics of two point vortices: Exact results*, Phys. Rev. E, 63 (2001), paper 056304.
- [2] C. CERRETELLI AND C. H. K. WILLIAMSON, *The physical mechanism for vortex merging*, J. Fluid Mech., 475 (2003), pp. 41–77.
- [3] TH. GALLAY AND C. E. WAYNE, *Invariant manifolds and the long-time asymptotics of the Navier-Stokes and vorticity equations on  $\mathbf{R}^2$* , Arch. Ration. Mech. Anal., 163 (2002), pp. 209–258.
- [4] TH. GALLAY AND C. E. WAYNE, *Global stability of vortex solutions of the two-dimensional Navier-Stokes equation*, Comm. Math. Phys., 255 (2005), pp. 97–129.
- [5] G. R. KIRCHHOFF, *Vorlesungen über Mathematische Physik. Mechanik*, Teubner, Leipzig, 1876.
- [6] S. LE DIZÈS, *Non-axisymmetric vortices in two-dimensional flows*, J. Fluid Mech., 406 (2000), pp. 175–198.
- [7] S. LE DIZÈS AND A. VERGA, *Viscous interactions of two co-rotating vortices before merging*, J. Fluid Mech., 467 (2002), pp. 389–410.
- [8] A. J. MAJDA AND A. L. BERTOZZI, *Vorticity and Incompressible Flow*, Cambridge Texts Appl. Math. 27, Cambridge University Press, Cambridge, UK, 2002.
- [9] M. V. MELANDER, A. S. STYCZEK, AND N. J. ZABUSKY, *Elliptically desingularized vortex model for the two-dimensional Euler equations*, Phys. Rev. Lett., 53 (1984), pp. 1222–1225.
- [10] M. V. MELANDER, N. J. ZABUSKY, AND J. C. MCWILLIAMS, *Symmetric vortex merger in two dimensions: Causes and conditions*, J. Fluid Mech., 195 (1988), pp. 303–340.
- [11] P. MEUNIER, S. LE DIZÈS, AND T. LEWEKE, *Physics of vortex merging*, C. R. Phys., 6 (2005), pp. 431–450.
- [12] R. J. NAGEM, G. SANDRI, AND D. UMINSKY, *Vorticity dynamics and sound generation in two-dimensional fluid flow*, J. Acoust. Soc. Amer., 122 (2007), pp. 128–134.
- [13] P. K. NEWTON, *The N-Vortex Problem*, Appl. Math. Sci. 145, Springer-Verlag, New York, 2001.
- [14] C. W. OSEEN, *Über Wirbelbewegung in einer reibenden Flüssigkeit*, Ark. f. Mat. Astron. och Fys., 7 (1911), pp. 1–11.
- [15] C. W. OSEEN, *Neuere Methoden und Ergebnisse in der Hydrodynamik*, Akademische Verlagsgesellschaft, Leipzig, 1927.
- [16] L. TING AND R. KLEIN, *Viscous Vortical Flows*, Lecture Notes in Phys. 374, Springer-Verlag, Berlin, 1991.
- [17] H. VON HELMHOLTZ, *Über Integrale der hydrodynamischen Gleichungen, welche den Wirbelbewegungen entsprechen*, J. Reine Angew. Math., 55 (1858), pp. 25–55.

Copyright of *SIAM Journal on Applied Dynamical Systems* is the property of Society for Industrial & Applied Mathematics and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.