

The University of Maine  
DigitalCommons@UMaine

---

Earth Science Faculty Scholarship

Earth Sciences

---

7-15-2004

# A Record of Atmospheric Co<sub>2</sub> During the Last 40,000 Years from the Siple Dome, Antarctica Ice Core

Jinho Ahn

Martin Wahlen

Bruce L. Deck

Ed J. Brook

Paul Andrew Mayewski

University of Maine - Main, [paul.mayewski@maine.edu](mailto:paul.mayewski@maine.edu)

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.library.umaine.edu/ers\\_facpub](https://digitalcommons.library.umaine.edu/ers_facpub)

 Part of the [Earth Sciences Commons](#)

---

## Repository Citation

Ahn, Jinho; Wahlen, Martin; Deck, Bruce L.; Brook, Ed J.; Mayewski, Paul Andrew; Taylor, Kendrick C.; and White, James W.C., "A Record of Atmospheric Co<sub>2</sub> During the Last 40,000 Years from the Siple Dome, Antarctica Ice Core" (2004). *Earth Science Faculty Scholarship*. 6.

[https://digitalcommons.library.umaine.edu/ers\\_facpub/6](https://digitalcommons.library.umaine.edu/ers_facpub/6)

This Article is brought to you for free and open access by DigitalCommons@UMaine. It has been accepted for inclusion in Earth Science Faculty Scholarship by an authorized administrator of DigitalCommons@UMaine. For more information, please contact [um.library.technical.services@maine.edu](mailto:um.library.technical.services@maine.edu).

---

**Authors**

Jinho Ahn, Martin Wahlen, Bruce L. Deck, Ed J. Brook, Paul Andrew Mayewski, Kendrick C. Taylor, and James W.C. White

## A record of atmospheric CO<sub>2</sub> during the last 40,000 years from the Siple Dome, Antarctica ice core

Jinho Ahn,<sup>1</sup> Martin Wahlen,<sup>1</sup> Bruce L. Deck,<sup>1</sup> Ed J. Brook,<sup>2</sup> Paul A. Mayewski,<sup>3</sup> Kendrick C. Taylor,<sup>4</sup> and James W. C. White<sup>5</sup>

Received 3 December 2003; revised 21 February 2004; accepted 13 April 2004; published 15 July 2004.

[1] We have measured the CO<sub>2</sub> concentration of air occluded during the last 40,000 years in the deep Siple Dome A (hereafter Siple Dome) ice core, Antarctica. The general trend of CO<sub>2</sub> concentration from Siple Dome ice follows the temperature inferred from the isotopic composition of the ice and is mostly in agreement with other Antarctic ice core CO<sub>2</sub> records. CO<sub>2</sub> rose initially at ~17.5 kyr B.P. (thousand years before 1950), decreased slowly during the Antarctic Cold Reversal, rose during the Younger Dryas, fell to a local minimum at around 8 kyr B.P., and rose continuously since then. The CO<sub>2</sub> concentration never reached steady state during the Holocene, as also found in the Taylor Dome and EPICA Dome C (hereafter Dome C) records. During the last glacial termination, a lag of CO<sub>2</sub> versus Siple Dome isotopic temperature is probable. The Siple Dome CO<sub>2</sub> concentrations during the last glacial termination and in the Holocene are at certain times greater than in other Antarctic ice cores by up to 20 ppm ( $\mu\text{mol CO}_2/\text{mol air}$ ). While in situ production of CO<sub>2</sub> is one possible cause of the sporadic elevated levels, the mechanism leading to the enrichment is not yet clear. *INDEX TERMS:* 0325

Atmospheric Composition and Structure: Evolution of the atmosphere; 0365 Atmospheric Composition and Structure: Troposphere—composition and chemistry; 1610 Global Change: Atmosphere (0315, 0325); 1615 Global Change: Biogeochemical processes (4805); *KEYWORDS:* CO<sub>2</sub>, paleoclimate, Siple Dome

**Citation:** Ahn, J., M. Wahlen, B. L. Deck, E. J. Brook, P. A. Mayewski, K. C. Taylor, and J. W. C. White (2004), A record of atmospheric CO<sub>2</sub> during the last 40,000 years from the Siple Dome, Antarctica ice core, *J. Geophys. Res.*, 109, D13305, doi:10.1029/2003JD004415.

### 1. Introduction

[2] Understanding how the atmospheric concentration of CO<sub>2</sub> changed in the past in response to other changes in the climate system provides us with a better understanding of how current and future changes in the carbon cycle will influence our future climate. The concentration of CO<sub>2</sub> in the atmosphere during previous times can be determined by measurements made on the air trapped in ice cores [e.g., Delmas *et al.*, 1980; Neftel *et al.*, 1982; Fischer *et al.*, 1999; Petit *et al.*, 1999]. CO<sub>2</sub> records from Antarctic ice cores are considered to be representative of paleoatmospheric concentrations. Greenland ice cores are not useful for determining the paleoatmospheric concentration because of in situ production of CO<sub>2</sub> due to high dust content [Barnola *et al.*, 1995; Anklin *et al.*, 1995, 1997; Smith *et al.*, 1997a, 1997b; Tschumi and Stauffer, 2000].

[3] The Siple Dome ice core from West Antarctica was drilled from 1997 to 1999. The site is at 81.66°S, 148.82°W, at an altitude of 621 meters asl, with an annual mean temperature of -25.4°C and an accumulation rate of 12.4 g cm<sup>-2</sup> yr<sup>-1</sup> as water equivalent (Figure 1 and Table 1). The total depth of the core is 1003.8 m. The high accumulation rate at Siple Dome allows rapid climate changes to be more accurately preserved and permits more accurate assessment of the relative timing between ice-isotopic variations and of variations in occluded gases compared to the low accumulation rate sites like Vostok, Dome C, or Dome Fuji (Table 1). In this study, we compare the CO<sub>2</sub> record from the Siple Dome ice core for the last 40,000 years with CO<sub>2</sub> records from other Antarctic ice cores using common time scales based on the GISP2 gas age and correlation to Antarctica using CH<sub>4</sub> concentrations variations. We also examine the relative timing of the CO<sub>2</sub> increase and temperature change. The highest resolution CO<sub>2</sub> record from Law Dome covering only the last thousand years [Etheridge *et al.*, 1996], and the recent Dome Fuji CO<sub>2</sub> record over the last 320 kyr [Kawamura *et al.*, 2003] measured by the wet extraction method are not discussed.

### 2. Method and Timescales

[4] Measurements were made on 462 samples from 86 depths in the top 832 meters in the Siple Dome ice core.

<sup>1</sup>Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California, USA.

<sup>2</sup>Department of Geology and Program in Environmental Science, Washington State University, Vancouver, Washington, USA.

<sup>3</sup>Climate Change Institute, University of Maine, Orono, Maine, USA.

<sup>4</sup>Desert Research Institute, University of Nevada, Reno, Nevada, USA.

<sup>5</sup>Institute of Arctic and Alpine Research (INSTAAR), University of Colorado, Boulder, Colorado, USA.



**Figure 1.** Location map of Antarctic ice cores referred to in the text.

These measurements span over the last 40,000 years. The ice core has a diameter of 13 cm. The outer 0.5 to 2 cm of the samples was removed with a band saw to reduce the possibility of contamination from present atmospheric CO<sub>2</sub> or postdrilling chemical reactions. The Siple Dome core samples contain bubbles and did not contain clathrates. Post collection temperature monitoring confirmed that the ice was kept below  $-15^{\circ}\text{C}$  from the time it was removed from the drill site to arrival at Scripps Institution of Oceanography (SIO). The storing, cutting, and loading of samples into a mechanical crusher at SIO was done at  $-25^{\circ}\text{C}$ . At most depths four to six ice samples, taken from a depth interval spanning 1 to 3 cm (mostly less than several years in ice age), were analyzed for each reported data point. At some depths samples of 6 to 12 cm depth intervals were used because of poor quality (extensive fracturing in the ice). Many fractures were observed below 500 m and the samples were more extensively trimmed in this region to assure sampling of unfractured ice. Contamination through undetected fractures results in higher CO<sub>2</sub> values due to high CO<sub>2</sub> concentration in the ambient air.

[5] For outliers in a group of samples cut from the same depth interval, we applied Grubb's test at the 95% confidence level to the suspected data. In this test a ratio of the difference between the outlier and the mean and the standard deviation of the data for the group was compared to a critical value [Iglewicz and Hoaglin, 1993]. We rejected results from 12 individual ice samples (usually the highest values) out of the total of 462 analyzed.

[6] The gas extraction and IR spectroscopic methods used are well established [Wahlen *et al.*, 1991; Smith *et al.*, 1997a, 1997b; Fischer *et al.*, 1999]. Occluded air was dry extracted by mechanical crushing of samples in a double walled crusher cooled using an ethanol-liquid nitrogen mixture at about  $-50^{\circ}\text{C}$ . The liberated air was collected in cold traps chilled by closed cycle He refrigerators to a temperature of about 32 K. The frozen air samples held in the cold traps were liberated by heating and transferred to an IR absorption cell held at a constant pressure and temperature. IR absorption measurements were made several times on each gas sample with a tunable diode laser by scanning a single vibrational-rotational CO<sub>2</sub> absorption line at Doppler resolution in the R branch of the 4.3  $\mu\text{m}$  CO<sub>2</sub> absorption band. To calibrate the instrument, measurements were made

with three air standards of precisely known CO<sub>2</sub> concentrations of 163, 240, and 330 ppm ( $\pm 0.01$  ppm, C. D. Keeling, personal communication) that were introduced over three of the crushed ice samples, thus duplicating the crushing procedure. This calibration procedure was performed each day. A group of samples from the same depth interval were measured on the same day. The average standard deviation of the data of the samples from the same depth interval measured on the same day (5.9 ppm) is greater than the daily internal precision (1.7 ppm) measured on crushed ice samples with reference gas added. Considering the daily uncertainty in calibration and the reproducibility, we expect the internal precision to be better than 2 ppm when measured on different days for a group of samples from the same depth interval with constant CO<sub>2</sub> concentration among the samples. Directly introduced standards (not over ice) were run to check laser linearity and performance.

[7] The gas age dating of the Siple Dome core over the time period of 40 to 8.2 kyr B.P. is established by correlation of CH<sub>4</sub> data with those of the GISP2 ice core [Taylor *et al.*, 2004b; E. Brook *et al.*, manuscript in preparation, 2004]. The methods for CH<sub>4</sub> measurements are described by Brook *et al.* [2000, manuscript in preparation, 2004]. For ice younger than 8.2 kyr B.P. the age of the ice was determined by counting annual layers and the corresponding gas ages were assigned by subtracting the ice age – gas age difference ( $\Delta\text{age}$ ) from the ice ages [Taylor *et al.*, 2004a]. The  $\Delta\text{age}$  was calculated with a firn densification model (Brook *et al.*, manuscript in preparation, 2004). The  $\Delta\text{age}$  is about  $280 \pm 80$  years in the early Holocene, increasing to about  $750 \pm 220$  years during the Last Glacial Maximum (LGM). The gas age of bubble ice varies according to ice bubble close-off depth. We estimate that the width of the age distribution is on the order of 10% of the  $\Delta\text{age}$  value [Goujon *et al.*, 2003].

### 3. Results

#### 3.1. Data Quality

[8] Measurements were made over a period of 2.5 years at random depth intervals to eliminate the possibility that

**Table 1.** Characteristics of the Antarctic Ice Cores From Which CO<sub>2</sub> Gases Have Been Measured

Core Name	Mean Annual Temperature, $^{\circ}\text{C}$	Mean Accumulation Rate as Water Equivalent, $\text{g cm}^{-2} \text{yr}^{-1}$	Elevation, m
Law Dome	$-22 \sim -19^{\text{a,b}}$	$60 \sim 110^{\text{b}}$	1390 <sup>b</sup>
Siple Dome	$-25.4^{\text{c}}$	$12.4^{\text{d}}$	621 <sup>c</sup>
Byrd <sup>f</sup>	$-28$	16	1530
Taylor Dome	$-42^{\text{e}}$	$7^{\text{h}}$	2374 <sup>h</sup>
EPICA Dome C	$-54^{\text{i}}$	$3^{\text{j}}$	3233 <sup>k</sup>
Vostok <sup>f</sup>	$-55.5$	2.3	3490
Dome Fuji <sup>l</sup>	$-58$	3.2	3810

<sup>a</sup>Etheridge and Wooley [1989].

<sup>b</sup>Etheridge *et al.* [1996].

<sup>c</sup>Severinghaus *et al.* [2001].

<sup>d</sup>Hamilton [2002].

<sup>e</sup>Taylor *et al.* [2004b].

<sup>f</sup>Raynaud *et al.* [1993].

<sup>g</sup>Waddington and Morse [1994].


<sup>h</sup>As unit of cm ice equivalent per year [Steig *et al.*, 1998].

<sup>i</sup>Schwander *et al.* [2001].

<sup>j</sup>EPICA Dome C 2001-02 Science and Drilling Teams [2002].

<sup>k</sup>Tabacco *et al.* [1988].

<sup>l</sup>Dome-F Deep Coring Group [1998].




**Figure 2.** Antarctic ice core CO<sub>2</sub> plotted on GISP2 gas age scale. (a) Siple Dome  $\delta D_{ice}$  on ice age scale. (b) Ice core CO<sub>2</sub> records on synchronized gas age scale with GISP2 via CH<sub>4</sub>. Shown are Vostok data [Petit *et al.*, 1999], EPICA Dome C [Monnin *et al.*, 2001; Flückiger *et al.*, 2002], Taylor Dome [Indermühle *et al.*, 1999, 2000; Smith *et al.*, 1999], and Byrd [Neftel *et al.*, 1988; Staffelbach *et al.*, 1991] and Siple Dome (this study). Error bars represent one standard error deviation of the replicate measurements at the same depth interval. The brackets indicate averages of two anomalously high values of the six samples from the same depth intervals. Taylor Dome gas ages of 40 to 20 kyr B.P. are from Brook *et al.* [2000]; Byrd gas ages are from Blunier and Brook [2001]; Vostok gas ages for 40 to 10 kyr B.P. are from Blunier *et al.* [1998]. The Vostok GT4 timescales [Petit *et al.*, 1999] are younger than those of GISP2 by about 1 kyr for the latter of the last termination and by up to 5 kyr during 40 to 20 kyr B.P. The Holocene depth-gas age scales of Vostok and Dome C are linearly interpolated between the youngest age points and synchronized gas age points in the early Holocene. Dome C gas ages for 17.2 to 11 kyr B.P. are synchronized with GISP2 in this study and for 22.5 to 17.2 kyr B.P. are extended from the synchronized gas age at 17.2 kyr B.P. by adding the depth-gas age intervals of the original paper [Monnin *et al.*, 2001]. YD, Younger Dryas; ACR, Antarctic Cold Reversal.

unnoticed analytical changes would create trends in the data. Measurements of Siple Dome CO<sub>2</sub> made in 2001 (black open squares) at SIO agree well with those made in 2002 and early 2003 (solid circles) as shown in Figure 2b.

[9] The results can be compared to other Antarctic ice cores as shown in Figure 2b. The CO<sub>2</sub> concentration of Dome C, Taylor Dome and Byrd cores were measured in Bern [Monnin *et al.*, 2001; Flückiger *et al.*, 2002; Indermühle *et al.*, 1999, 2000; Neftel *et al.*, 1988; Staffelbach *et al.*, 1991] and compare well with the SIO data. This is expected because Bern's gas extraction method (dry crushing) is similar to that at SIO.

[10] High accumulation rate provides a small width of gas age distribution and therefore a record of high temporal resolution. Similarly, t of small pieces of ice poten-

tially also improves temporal resolution. On the other hand, this potentially increases the scatter in the data since inhomogeneities are not smoothed out as they are in larger samples covering longer intervals of time [Smith *et al.*, 1997a]. The accumulation rate at Siple Dome is two and five times larger than that at Taylor Dome and Dome C, respectively. Thus the high accumulation rate at Siple Dome (Table 1) and the small sample size (4 to 6 cm<sup>3</sup>) used in the measurement at SIO may increase the scattering of the Siple Dome data. Sample sizes used in the Siple Dome CO<sub>2</sub> study at SIO are half of those used for Taylor Dome and Dome C at Bern. Some Taylor Dome samples were also measured at SIO (Figure 2b) and there is good agreement between the SIO and the Bern results, even though the volume of samples for a depth interval used at SIO was about one



**Figure 3.** CO<sub>2</sub> change during the last Termination. (a) Siple Dome  $\delta D_{ice}$  on ice age scale. (b) CO<sub>2</sub> records from Siple Dome (solid circles) and Dome C (open squares) [Monnin *et al.*, 2001]. The line between Siple Dome CO<sub>2</sub> is the running average of three adjacent points. (c) Melt layers (vertical arrows) [Das, 2003]. The horizontal bars at top are gas age ranges around age tie points based on CH<sub>4</sub> records between Siple Dome and GISP2. The thick vertical dashed bar indicates where the argon and nitrogen isotopic anomaly occurs [Severinghaus *et al.*, 2003]. YD, Younger Dryas; ACR, Antarctic Cold Reversal.

fourth of what was used at Bern, where calibration was routinely done using reference gases from SIO at 251.7 ppm and 321.06 ppm [Indermöhle *et al.*, 1999].

### 3.2. Trend of the Siple Dome CO<sub>2</sub> Record

[11] In general, the trends in the Siple Dome CO<sub>2</sub> record (solid circles and open squares in Figure 2b) follow temperature trends, indicated by  $\delta D_{ice}$  (Figure 2a), which is taken as a proxy for local temperature. The general time series of Siple Dome CO<sub>2</sub> concentration confirms previous studies in Antarctic ice cores (Figures 2 and 3). The Siple Dome CO<sub>2</sub> record shows a continuous decrease from 214 ppm at 38 kyr B.P. to 186 ppm at LGM (~18 kyr B.P.) and then a rapid increase up to 247 ppm during the early part of the last termination. It is important to note that the CO<sub>2</sub> increase is reversed during the Antarctic Cold Reversal (15 to 13 kyr B.P.) reaching a local minimum of 239 ppm, which has been observed in the Dome C, Taylor Dome and Byrd records. After the Antarctic Cold Reversal, Siple Dome CO<sub>2</sub> increases again reaching up to 284 ppm at the beginning of the Holocene (11.5 kyr B.P.). This increase occurs during the Younger Dryas interval as shown in the Dome C record [Monnin *et al.*, 2001]. During the Holocene, the Siple Dome CO<sub>2</sub> concentrations decrease to a local minimum of 261 ppm at about 8 kyr B.P., and then increase to 285 ppm in the late Holocene (mean of the scattered values). There is good agreement of this local minimum at about 8 kyr B.P. between the different Antarctic cores from Siple Dome, Taylor Dome, Dome C and Vostok. The CO<sub>2</sub> concentration never reaches a steady state during the Holo-

cene after the end of the last termination as was also found in the Taylor Dome and Dome C records [Indermöhle *et al.*, 1999; Flückiger *et al.*, 2002].


[12] Figure 3 shows an enlarged portion of Figure 2 covering the last termination. Again, Siple Dome CO<sub>2</sub> follows the temperature proxy ( $\delta D_{ice}$ ) as seen in other Antarctic ice cores. For most of the record, the Siple Dome CO<sub>2</sub> time trend is almost the same as that of the high resolution Dome C record.

[13] However, at the depth (677.61 m) corresponding to around 15 kyr B.P., CO<sub>2</sub> in Siple Dome is higher than in Dome C by about 21 ppm. We believe that this anomalous CO<sub>2</sub> value is suspicious as an atmospheric CO<sub>2</sub> record because of the following two reasons. First, near the depth of the anomalously high CO<sub>2</sub> value (~20 cm apart in depth), the  $\delta^{40}Ar$  and  $\delta^{15}N$  of N<sub>2</sub> values reach atmospheric values, indicating there was no diffusive column and a possible hiatus in deposition [Severinghaus *et al.*, 2003]. Second, the difference of 21 ppm is too large to be explained by age uncertainties. The anomaly occurs at a greater depth than that corresponding to the beginning of Antarctic Cold Reversal (defined by the rapid increase in CH<sub>4</sub> concentration from Siple Dome ice, as defined in the Dome C ice core, where abrupt CO<sub>2</sub> and CH<sub>4</sub> increases happened [Monnin *et al.*, 2001]). Thus the age uncertainty at this depth does not affect the CO<sub>2</sub> difference between Siple Dome and Dome C significantly (<5 ppm).

[14] Except for this anomalous period, we calculate the rates of CO<sub>2</sub> change and time lag of CO<sub>2</sub> compared to temperature. During the last termination, CO<sub>2</sub> changed at the rate of 19.1 ppm/kyr from 17.7 to 15 kyr B.P., -5.2 ppm/kyr during the Antarctic Cold Reversal (15 to 13 kyr B.P.) and 26.8 ppm/kyr from 13.0 to 11.3 kyr B.P., respectively. These values are similar to CO<sub>2</sub> changes in the Dome C record [Monnin *et al.*, 2001].

### 3.3. Phase Relationship Between Changes in CO<sub>2</sub> and Siple Dome Temperature

[15] To compare the phasing of changes in CO<sub>2</sub> and surface temperature (using  $\delta D_{ice}$  as a temperature proxy), we interpolated the CO<sub>2</sub> and  $\delta D_{ice}$  data on a 10-year spacing. We calculated the change in CO<sub>2</sub> and  $\delta D_{ice}$  ( $dCO_2/dt$  and  $d(\delta D_{ice})/dt$ ) for 10-year intervals and smoothed them by averaging over time windows ranging from 0 to 500 years. The correlation coefficient between the  $dCO_2/dt$  and  $d(\delta D_{ice})/dt$  was then calculated for different lag times and for different degrees of smoothing. The maximum correlation coefficients were obtained with a 210 ~ 330 year lag of  $dCO_2/dt$  behind  $d(\delta D_{ice})/dt$  with a smoothing of 200 ~ 500 years as shown in Figure 4a. Several maxima in correlation coefficients for different time lags are observed over less than 200-year windows, possibly due to the average spacing of 230 years in the CO<sub>2</sub> data. To check the sensitivity of the time lag by the uncertainties in CO<sub>2</sub> concentration and  $\Delta age$ , we carried out Monte Carlo simulations for CO<sub>2</sub> concentrations in two conditions, namely maximum  $\Delta age$  (estimated  $\Delta age +$  uncertainty) and minimum  $\Delta age$  (estimated  $\Delta age -$  uncertainty). For each situation, we produced 3000 different sets of CO<sub>2</sub> concentrations, which vary randomly with Gaussian propagation in their uncertainties. After interpolation and smoothing both  $dCO_2/dt$  and  $d(\delta D_{ice})/dt$  with



**Figure 4.** Time lag analysis of (dCO<sub>2</sub>/dt) after (dδD<sub>ice</sub>/dt) in the Siple Dome ice core during the last termination. (a) Correlation coefficients for different time lags after averaging data with windows of 200 ~ 500 years. (b) Histograms of the time lags after Monte Carlo simulation for CO<sub>2</sub> concentrations in two situations, maximum Δ<sub>age</sub> and minimum Δ<sub>age</sub>. For each situation, 3000 different sets of CO<sub>2</sub> concentrations were generated and used. The derived dCO<sub>2</sub>/dt and d(δD<sub>ice</sub>)/dt were smoothed with 500-year window for each data set.

500-year window, maximum correlation coefficients were obtained with a  $369 \pm 23$  years lag (mean  $\pm 1\sigma$ ) for maximum Δ<sub>age</sub> and  $196 \pm 25$  (mean  $\pm 1\sigma$ ) for minimum Δ<sub>age</sub> as seen in Figure 4b. The time lags for CO<sub>2</sub> behind temperature are greater than the uncertainty of the Δ<sub>age</sub> of 80 years at the end of the last termination but close to 220 years at LGM. The time lags roughly agree with those of Fischer *et al.* [1999] ( $600 \pm 400$  years during the last three terminations in Byrd, Taylor Dome and Vostok records) and are slightly less than the value proposed by Monnin *et al.* [2001] ( $410$  years during the last termination in Dome C record). Indermühle *et al.* [2000] found higher values ( $1200 \pm 700$  years) for the period 60 to 20 kyr B.P. in the Taylor Dome record, as did Caillon *et al.* [2003] ( $800 \pm 200$  years) across termination III in the Vostok record. Although the correlation method discussed above suggests that changes in surface temperature at Siple Dome could lead changes in the concentration of atmospheric CO<sub>2</sub> by  $\sim 300$  years, this lead is small relative to both our sampling interval and possible systematic error in Δ<sub>age</sub>, for example due to inaccuracies in estimating past accumulation rates, which are derived from a thermodynamic model employing temperatures calculated from the δD<sub>ice</sub> record (Brook *et al.*, manuscript in preparation, 2004). Our main conclusion is therefore that a lead of CO<sub>2</sub> versus Siple Dome temperature is unlikely, a lag of CO<sub>2</sub> versus Siple Dome temperature is likely, and that our results provide strong support for previous suggestions of a close link between Antarctic temperature and CO<sub>2</sub> change during the deglaciation.


#### 4. Possible Mechanisms for In Situ CO<sub>2</sub> Production

[16] In some depth intervals the CO<sub>2</sub> concentrations in Siple Dome ice are than in the Vostok, Taylor

Dome and Dome C cores (Figures 2 and 3). These differences are up to 20 ppm greater (e.g., at around 11 kyr B.P.). One possible explanation for these differences is uncertainty in the age scales, particularly at times when CO<sub>2</sub> was changing. For example, during the two periods of rapid CO<sub>2</sub> increase during the last termination, an offset of 300 years between otherwise identical records would produce a CO<sub>2</sub> difference of 10 ppm. This effect may explain the differences between the Siple Dome versus the Dome C and Siple Dome versus the Taylor Dome cores (Figure 5c). The low CO<sub>2</sub> values of the Byrd ice core at about 39 kyr B.P. can be compared to low values in Taylor Dome at about 42 kyr B.P. (not shown in Figure 2) on the GISP2 age scale [Indermühle *et al.*, 2000]. However, elevated CO<sub>2</sub> levels in the Siple Dome at times of relatively little change in CO<sub>2</sub> (for example, during the Antarctic Cold Reversal or in the earliest Holocene) cannot be attributed to timescale uncertainties. Interestingly, most of the cold period (40 ~ 23 kyr B.P.) CO<sub>2</sub> data from the Siple Dome ice shows little difference in CO<sub>2</sub> when compared to other cores.

[17] As mentioned above, large variations of atmospheric CO<sub>2</sub> concentrations between Antarctic ice core sites are unlikely, prompting us to search for another explanation for the differences between the Siple Dome CO<sub>2</sub> and other records. To examine the possibility and the mechanisms of CO<sub>2</sub> production, we check the CO<sub>2</sub> difference between Siple Dome and Taylor Dome or Dome C (Figure 5c). We call this “excess CO<sub>2</sub>” although we cannot rule out the possibility of CO<sub>2</sub> consumption in other ice cores by the interaction between CO<sub>2</sub> in the bubbles and the carbonates in the ice (i.e.,  $\text{CO}_2 + \text{CO}_3^{2-} + \text{H}_2\text{O} \rightarrow 2\text{HCO}_3^-$ ) [Nefel *et al.*, 1982].

[18] The excess CO<sub>2</sub> is high at depth of 100 to 330 m (0.4 to 3.5 kyr B.P.) and 560 to 640 m (9.5 to 12.5 kyr B.P.) but is within the experimental uncertainty at depths >708 m



**Figure 5.** Comparison of possible processes of the excess CO<sub>2</sub> in Siple Dome ice cores on depth scale. (a) Siple Dome  $\delta D_{ice}$ . (b) Siple Dome CO<sub>2</sub>. (c) CO<sub>2</sub> difference between Siple Dome and Taylor Dome (solid line) or Dome C (shaded line). The dashed horizontal line represents 0. (d) Melt layers [Das, 2003]. The lengths of bars indicate the confidence level. 1, sure; 0.75, probable; 0.5, possible. (e) Potential CO<sub>2</sub> from acid-carbonate reaction. The dashed horizontal line represents 0. Non-sea-salt Ca (nssCa) was assumed to have been in the form of CaCO<sub>3</sub> and totally reacted with excess H<sup>+</sup>. Considering total gas content in the unit gram of ice [Severinghaus *et al.*, 2003], the CO<sub>2</sub> produced by the reaction was converted to a concentration change of CO<sub>2</sub>.

(17.5 kyr B.P.). The Siple Dome ice samples are more fractured than Taylor Dome samples below about 500 m (about 8 kyr). One possibility is that the CO<sub>2</sub> in the core might be compromised by air entering fractures in the core that reseal after the ice reaches the surface, during transportation and storage. However, there are high excess CO<sub>2</sub> values in the late Holocene Siple Dome ice where fracturing is minor, and low excess CO<sub>2</sub> in glacial ice where Siple Dome ice is highly fractured. Moreover, correlations between the scattering of data from adjacent samples and excess CO<sub>2</sub> does not support the possibility of ambient air contamination. The above observations imply the possibility that some of the artifacts in Siple Dome CO<sub>2</sub> records are related to in situ CO<sub>2</sub> production in the ice.

[19] Possible CO<sub>2</sub> production mechanisms include (1) carbonate-acid reaction [Delmas, 1993; Anklin *et al.*, 1995, 1997; Barnola *et al.*, 1995; Smith *et al.*, 1997a, 1997b], (2) oxidation of organic compounds abiologically [Tschumi and Stauffer, 2000], or biologically [Campen *et al.*, 2003], and (3) snowmelting-refreezing [Nefstel *et al.*, 1983; Stauffer *et al.*, 1985].

[20] We have calculated the potential CO<sub>2</sub> produced by the carbonate-acid reaction between CaCO<sub>3</sub> and H<sup>+</sup> [Nefstel

*et al.*, 1988], as shown in Figure 5e, using the non-sea-salt Ca (nssCa) ion content assuming all of the nssCa is from CaCO<sub>3</sub> and that all CaCO<sub>3</sub> reacts to produce CO<sub>2</sub>. This source of potential CO<sub>2</sub> is too low to produce the excess CO<sub>2</sub> observed during the Holocene and shows little correlation with the excess CO<sub>2</sub> data (Figure 5c). Moreover, electric conductivity measurements (ECM, proxy for H<sup>+</sup>) do not show any significant change at the depth intervals of high excess CO<sub>2</sub> in Figure 5c. Thus it is unlikely that a carbonate-acid reaction has affected the Siple Dome CO<sub>2</sub> record.

[21] Oxidation of organic compounds has been proposed to be at least as important as acid-carbonate reactions for CO<sub>2</sub> production (e.g., 2H<sub>2</sub>O<sub>2</sub> + HCHO → 3H<sub>2</sub>O + CO<sub>2</sub>) [Tschumi and Stauffer, 2000]. Organic acid concentrations in Siple Dome ice core have not yet been reported. H<sub>2</sub>O<sub>2</sub> is one of the important oxidants of organics in ice. The top 100 m of Siple Dome ice core shows concentrations near or below the detection limits of ~0.02 μM H<sub>2</sub>O<sub>2</sub> except at the 0 to 2.5 m depth [McConnell, 1997]. This H<sub>2</sub>O<sub>2</sub> concentration corresponds to a potential CO<sub>2</sub> production of less than 5 ppm assuming an adequate supply of organic acids, which does not easily explain the 20 ppm excess CO<sub>2</sub>. CH<sub>3</sub>COO<sup>-</sup> and HCOO<sup>-</sup> can act as other oxidants [Tschumi and Stauffer, 2000]. Another possible CO<sub>2</sub> production mechanism is respiration of microorganisms (consuming organic acids and producing CO<sub>2</sub>) within the ice [Campen *et al.*, 2003]. Natural organic acids and other organic materials have much lower values of  $\delta^{13}\text{CO}_2$ , approximately -25‰, compared to atmospheric  $\delta^{13}\text{CO}_2$  (preindustrial value of about -6.5‰) and CaCO<sub>3</sub> (0 ~ 4‰). Thus precise  $^{13}\text{CO}_2$  values could be used to investigate the sources of excess CO<sub>2</sub>. For example if the early Holocene CO<sub>2</sub> record has excess CO<sub>2</sub> values of 20 ppm derived from organic materials, it would reduce the  $\delta^{13}\text{CO}_2$  value by about 1‰. However,  $^{13}\text{CO}_2$  analyses in the Siple Dome ice core have been hampered by the contamination of the ice by drilling fluid, n-butyl acetate, which in the mass spectrometer lead to fragmentation interference with m/z = 45 [Ahn *et al.*, 2001].

[22] CO<sub>2</sub> can be considerably enriched in frozen melt layers due to the high solubility of CO<sub>2</sub> in meltwater [Nefstel *et al.*, 1983]. At Dye 3, Greenland, melt layers contribute about 7% by volume at today's climate conditions and show mean CO<sub>2</sub> concentration as much as 1500 ppm [Stauffer *et al.*, 1985]. The annual mean snow temperature of Siple Dome is -25.4°C, the highest among the Antarctic ice cores discussed in this paper [Severinghaus *et al.*, 2001], and occasional surface melting occurs during the summer, resulting in thin bubble free layers in the ice core. These melt layers can be detected visually, see arrows in Figures 3c and bars in Figure 5d [Das, 2003]. The ice samples measured for the Siple Dome CO<sub>2</sub> did not include visible melt layers. However, it is likely that small melt layers would not be detected visually, particularly deeper in the core where ice flow has thinned the layering. The mode of excess CO<sub>2</sub> is only weakly associated with that of melt layers (Figures 5c and 5d). Alternative possibilities beside warm temperature and the coastal location of the Siple Dome (Figure 1 and Table 1) include higher ambient temperatures during the austral summer (often being substantially higher than the mean annual temperature of



–25.4°C), combined with high sea salt content, resulting in CO<sub>2</sub> contamination.

## 5. Conclusion

[23] We present the CO<sub>2</sub> record of air occluded over the last 40,000 years in the Siple Dome ice core, Antarctica. The general time series of Siple Dome CO<sub>2</sub> concentration is similar to previous studies in other Antarctic ice core CO<sub>2</sub> records. Siple Dome ice also shows that surface temperature inferred from  $\delta D_{ice}$  correlates well with CO<sub>2</sub> concentration as shown in other Antarctic ice cores. During the last termination, it is likely that the change of the Siple Dome CO<sub>2</sub> concentrations lags the Siple Dome temperature change. Despite similarities with other Antarctic ice cores, the Siple Dome ice shows higher CO<sub>2</sub> concentrations than those in other Antarctic ice cores at some depth intervals, which may be due to in situ production in the ice. The cause of these elevated concentrations is not known with certainty. CO<sub>2</sub> production due to surface melting is the leading hypothesis, but the evidence is not definitive.

[24] **Acknowledgments.** We wish to thank Jeff Severinghaus, Sarah Das, Richard Alley, Joe McConnell, and Jihong Cole-Dai for advice and for providing unpublished data. Anonymous reviewers provided constructive comments. Staff at the National Ice Core Laboratory is thanked for the assistance with core processing. This work was supported by the Office of Polar Program of the NSF (NSF OPP 99-80619).

## References

- Ahn, J., M. Wahlen, and B. Deck (2001), Measurement of atmospheric CO<sub>2</sub> trapped in the Siple Dome ice core, *Eos Trans. AGU*, 82(47), Fall Meet. Suppl., Abstract A51F-0105.
- Anklin, M., J.-M. Barnola, J. Schwander, B. Stauffer, and D. Raynaud (1995), Processes affecting the CO<sub>2</sub> concentrations measured in Greenland ice, *Tellus, Ser. B*, 47, 461–470.
- Anklin, M., J. Schwander, B. Stauffer, J. Tschumi, A. Fuchs, J.-M. Barnola, and D. Raynaud (1997), CO<sub>2</sub> record between 40 and 8 kyr B.P. from the Greenland Ice Core Project ice core, *J. Geophys. Res.*, 102, 26,539–26,545.
- Barnola, J. M., M. Anklin, J. Porcheron, D. Raynaud, J. Schwander, and B. Stauffer (1995), CO<sub>2</sub> evolution during the last millennium as recorded by Antarctic and Greenland ice, *Tellus, Ser. B*, 47, 264–272.
- Blunier, T., and E. J. Brook (2001), Timing of millennial-scale climate change in Antarctica and Greenland during the last glacial period, *Science*, 291, 109–112.
- Blunier, T., et al. (1998), Asynchrony of Antarctic and Greenland climate change during the last glacial period, *Nature*, 394, 739–743.
- Brook, E. J., S. Harder, J. P. Severinghaus, E. J. Steig, and C. M. Suher (2000), On the origin and timing of rapid changes in atmospheric methane during the last glacial period, *Global Biogeochem. Cycles*, 14, 559–572.
- Caillon, N., J. P. Severinghaus, J. Jouzel, J.-M. Barnola, J. Kang, and V. Y. Lipenkov (2003), Timing of atmospheric CO<sub>2</sub> and Antarctic temperature changes across Termination III, *Science*, 299, 1728–1731.
- Campen, R. K., T. Sowers, and R. B. Alley (2003), Evidence of microbial consortia metabolizing within a low-latitude mountain glacier, *Geology*, 31, 231–234.
- Das, S. (2003), West Antarctic ice sheet surface melting and Holocene variability, Ph.D. thesis, Pa. State Univ., University Park.
- Delmas, R. J. (1993), A natural artifact in Greenland ice-core CO<sub>2</sub> measurement, *Tellus, Ser.*, 45, 391–396.
- Delmas, R. J., J. M. Ascencio, and M. Legrand (1980), Polar ice evidence that atmospheric CO<sub>2</sub> 20,000yr BP was 50% of present, *Nature*, 284, 155–157.
- Dome-F Deep Coring Group (1998), Deep ice-core drilling at Dome Fuji and glaciological studies in east Dronning Maud Land, Antarctica, *Ann. Glaciol.*, 27, 333–337.
- EPICA Dome C 2001–02 science and drilling teams (2002), Extending the ice core record beyond half a million years, *Eos Trans. AGU*, 83(45), 509, 517.
- Etheridge, D. M., and C. W. Wookey (1989), Ice core drilling at a high accumulation area of Law Antarctica, in *Ice Core Drilling: Proceedings of the Third International Workshop on Ice Core Drilling Technology*, edited by C. Rado and D. Beaudoin, pp. 86–96, Cent. Natl. de la Rech. Sci., Grenoble, France.
- Etheridge, D. M., L. P. Steele, R. L. Langenfelds, R. J. Francey, J.-M. Barnola, and V. I. Morgan (1996), Natural and anthropogenic changes in atmospheric CO<sub>2</sub> over the last 1000 years from air in Antarctic ice and firn, *J. Geophys. Res.*, 101, 4115–4128.
- Fischer, H., M. Wahlen, J. Smith, D. Mastroianni, and B. Deck (1999), Ice core records of atmospheric CO<sub>2</sub> around the last three glacial terminations, *Science*, 283, 1712–1714.
- Flückiger, J., E. Monnin, B. Stauffer, J. Schwander, and T. F. Stocker (2002), High-resolution Holocene N<sub>2</sub>O ice core record and its relationship with CH<sub>4</sub> and CO<sub>2</sub>, *Global Biogeochem. Cycles*, 16(1), 1010, doi:10.1029/2001GB001417.
- Goujon, C., J.-M. Barnola, and C. Ritz (2003), Modeling the densification of polar firn including heat diffusion: Application to close-off characteristics and gas isotopic fractionation for Antarctica and Greenland sites, *J. Geophys. Res.*, 108(D24), 4792, doi:10.1029/2002JD003319.
- Hamilton, G. S. (2002), Mass balance and accumulation rate across Siple Dome, West Antarctica, *Ann. Glaciol.*, 35, 102–106.
- Iglewicz, B., and D. C. Hoaglin (1993), *How to Detect and Handle Outliers*, ASQC Basic Ref. Qual. Control, vol. 16, 87 pp., Am. Soc. for Qual. Control, Milwaukee, Wis.
- Indermühle, A., et al. (1999), Holocene carbon-cycle dynamics based on CO<sub>2</sub> trapped in ice at Taylor Dome, Antarctica, *Nature*, 398, 121–126.
- Indermühle, A., E. Monnin, B. Stauffer, T. F. Stocker, and M. Wahlen (2000), Atmospheric CO<sub>2</sub> concentration from 60–20 kyr BP from the Taylor Dome ice core, Antarctica, *Geophys. Res. Lett.*, 27, 735–738.
- Kawamura, K., T. Nakazawa, S. Aoki, S. Sugawara, Y. Fujii, and O. Watanabe (2003), Atmospheric CO<sub>2</sub> variations over the last three glacial-interglacial climate cycles deduced from the Dome Fuji deep ice core, Antarctica using a wet extraction technique, *Tellus, Ser. B*, 55, 126–137.
- McConnell, J. R. (1997), Investigation of the atmosphere-snow transfer process for hydrogen peroxide, Ph.D. dissertation, Dep. of Hydrol. and Water Resour., Univ. of Ariz., Tucson.
- Monnin, E., A. Indermühle, A. Daellenbach, J. Flückiger, B. Stauffer, T. F. Stocker, D. Raynaud, and J.-M. Barnola (2001), Atmospheric CO<sub>2</sub> concentrations over the last glacial termination, *Science*, 291, 112–114.
- Nefel, A., H. Oeschger, J. Schwander, B. Stauffer, and R. Zumbunn (1982), Ice core sample measurements give atmospheric CO<sub>2</sub> content during the past 40,000 yr, *Nature*, 259, 220–223.
- Nefel, A., H. Oeschger, J. Achvander, and B. Stauffer (1983), Carbon dioxide concentration in bubbles of natural cold ice, *J. Phys. Chem.*, 87, 4116–4120.
- Nefel, A., H. Oeschger, T. Stauffelbach, and B. Stauffer (1988), CO<sub>2</sub> record in the Byrd ice core 50,000–5000 years BP, *Nature*, 331, 609–611.
- Petit, J. R., et al. (1999), Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, *Nature*, 399, 429–436.
- Raynaud, D., J. Jouzel, J. M. Barnola, J. Chappellaz, R. J. Delmas, and C. Lorius (1993), The ice record of greenhouse gases, *Science*, 259, 926–934.
- Schwander, J., J. Jouzel, C. U. Hammer, J.-R. Petit, R. Udisti, and E. Wolff (2001), A tentative chronology for the EPICA Dome Concordia ice core, *Geophys. Res. Lett.*, 28, 4243–4246.
- Severinghaus, J. P., A. Grachev, and M. Battle (2001), Thermal fractionation of air in polar firn by seasonal temperature gradients, *Geochem. Geophys. Geosyst.*, 2, Paper number 2000GC000146.
- Severinghaus, J. P., A. Grachev, B. Luz, and N. Caillon (2003), A method for precise measurement of argon 40/36 and krypton/argon ratios in trapped air in polar ice with applications to past firn thickness and abrupt climate change in Greenland and at Siple Dome, Antarctica, *Geochim. Cosmochim. Acta.*, 67, 325–343.
- Smith, H. J., M. Wahlen, D. Mastroianni, and K. Taylor (1997a), The CO<sub>2</sub> concentration of air trapped in GISP2 ice from the Last Glacial Maximum-Holocene transition, *Geophys. Res. Lett.*, 24, 1–4.
- Smith, H. J., M. Wahlen, D. Mastroianni, K. Taylor, and P. Mayewski (1997b), The CO<sub>2</sub> concentration of air trapped in Greenland Ice Sheet Project 2 ice formed during periods of rapid climate change, *J. Geophys. Res.*, 102, 26,577–26,582.
- Smith, H. J., H. Fischer, M. Wahlen, D. Mastroianni, and B. Deck (1999), Dual modes of the carbon cycle since the last glacial maximum, *Nature*, 400, 248–250.
- Staffelbach, T., B. Stauffer, A. Sigg, and H. Oeschger (1991), CO<sub>2</sub> measurements from polar ice cores: More data from different sites, *Tellus, Ser. B*, 43, 91–96.
- Stauffer, B., A. Nefel, H. Oeschger, and J. Schwander (1985), CO<sub>2</sub> concentration in air extracted from Greenland ice samples, in *Greenland Ice Core: Geophysics, Geochemistry and the Environment*, *Geophys.*

- Monogr. Ser.*, vol. 33, edited by C. C. Langway Jr. et al., pp. 85–89, AGU, Washington, D. C.
- Steig, E. J., D. L. Morse, E. D. Waddington, and P. J. Polissar (1998), Using the sunspot cycle to date ice cores, *Geophys. Res. Lett.*, *25*, 163–166.
- Tabacco, I. E., A. Passerini, F. Corbelli, and M. Gorman (1988), Determination of the surface and bed topography at Dome C, East Antarctica, *J. Glaciol.*, *44*, 185–191.
- Taylor, K. C., et al. (2004a), Dating the Siple Dome, Antarctic ice core by manual and computer interpretation of annual layering, *J. Glaciol.*, in press.
- Taylor, K. C., et al. (2004b), Abrupt climate change around 22 ka on the Siple Coast of Antarctica, *Quat. Sci. Rev.*, *23*, 7–15.
- Tschumi, J., and B. Stauffer (2000), Reconstructing past atmospheric CO<sub>2</sub> concentration based on ice-core analyses: Open questions due to in situ production of CO<sub>2</sub> in the ice, *J. Glaciol.*, *46*, 45–53.
- Waddington, E. D., and D. L. Morse (1994), Spatial variations of local climate at Taylor Dome, Antarctica: Implications for paleoclimate from ice cores, *Ann. Glaciol.*, *20*, 219–225.
- Wahlen, M., D. Allen, B. Deck, and A. Herchenroder (1991), Initial measurement of CO<sub>2</sub> concentrations (1530 to 1940 AD) in air occluded in the GISP2 ice core from central Greenland, *Geophys. Res. Lett.*, *18*, 1457–1460.
- 
- J. Ahn, B. L. Deck, and M. Wahlen, Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA 92093, USA. (jiahn@ucsd.edu)
- E. J. Brook, Department of Geology and Program in Environmental Science, Washington State University, Vancouver, WA 98686, USA.
- P. A. Mayewski, Climate Change Institute, University of Maine, Orono, ME 04469, USA.
- K. C. Taylor, Desert Research Institute, University of Nevada, Reno, NV 89512, USA.
- J. W. C. White, INSTAAR, University of Colorado, Boulder, CO 80309, USA.