

University of Wisconsin Milwaukee UWM Digital Commons

Theses and Dissertations

May 2014

Nonconvex Cases for Carpenter's Rulers

Ke Chen

University of Wisconsin-Milwaukee

Follow this and additional works at: <https://dc.uwm.edu/etd>

 Part of the [Computer Sciences Commons](#)

Recommended Citation

Chen, Ke, "Nonconvex Cases for Carpenter's Rulers" (2014). *Theses and Dissertations*. 577.
<https://dc.uwm.edu/etd/577>

This Thesis is brought to you for free and open access by UWM Digital Commons. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of UWM Digital Commons. For more information, please contact open-access@uwm.edu.

NONCONVEX CASES FOR
CARPENTER'S RULERS

by

Ke Chen

A Thesis Submitted in
Partial Fulfillment of the
Requirements for the Degree of

Master of Science
in Computer Science

at

The University of Wisconsin-Milwaukee

May 2014

ABSTRACT
NONCONVEX CASES FOR CARPENTER'S RULERS

by

Ke Chen

The University of Wisconsin-Milwaukee, 2014
Under the Supervision of Professor Adrian Dumitrescu

We consider the carpenter's ruler folding problem in the plane, i.e., finding a minimum area shape with diameter 1 that accommodates foldings of any ruler whose longest link has length 1. An upper bound of 0.614 and a lower bound of 0.476 are known for convex cases. We generalize the problem to simple nonconvex cases: in this setting we improve the upper bound to 0.583 and establish the first lower bound of 0.073. A variation is to consider rulers with at most k links. The current best convex upper bounds are 0.486 for $k = 3, 4$ and 0.523 for $k = 5, 6$. These bounds also apply to nonconvex cases. We derive a better nonconvex upper bound of 0.296 for $k = 3, 4$.

©Copyright by Ke Chen, 2014
All Rights Reserved

TABLE OF CONTENTS

LIST OF FIGURES	v
LIST OF TABLES	viii
1 Introduction	1
2 Upper Bound	8
2.1 Diameter of C	9
2.2 Algorithm for folding a ruler inside C	10
2.3 Minimum area of C	10
3 Lower Bound	11
3.1 Lower Bound with One 3-Link ruler	11
3.2 Lower Bound with One 5-link Ruler	14
4 k-universal Cases	20
4.1 Area of S_2	23
4.2 Area of C^2	24
4.3 C^2 is 4-universal	26
5 Summary and Future Directions	29
6 References	32

LIST OF FIGURES

1	A universal case (in bold lines) for Moser’s worm problem Left: a closed semidisk of unit diameter. Right: a plane curve of length 1 contained in the case.	1
2	Another universal case (in bold lines) for Moser’s worm problem. Left: a rhombus with a long diagonal of unit length and vertex angles of 60° and 120° . Right: a plane curve of length 1 contained in the case.	1
3	A universal case (in bold lines) for Lebesgue’s covering problem. Left: union of a unit Reuleaux triangle and a unit circle. Right: a Reuleaux triangle with unit diameter (shaded area) contained in the case.	2
4	Another universal case (in bold lines) for Lebesgue’s covering problem. Left: a regular hexagon circumscribed to a unit circle, with two corners cut off. Right: a Reuleaux triangle with unit diameter (shaded area) contained in the case.	3
5	A universal case (in bold lines) for Lebesgue’s covering problem under translation. Left: a unit square. Right: a Reuleaux triangle with unit diameter (shaded area) can rotate inside the square.	3
6	Smallest area translative universal case (in bold lines) for all open curves of unit length. Left: an equilateral triangle of unit height. Right: a unit open curve can rotate inside the equilateral triangle.	4
7	The minimum folding length of a ruler can be almost double the length of its longest link.	4
8	Left: convex universal case $R2$ (in bold lines). Right: folding a 6-link unit ruler $p_0p_1p_2p_3p_4p_5p_6$ into $R2$	5
9	Universal cases (in bold lines) are contained in a lens of radius 1. Left: convex universal case $R2$. Right: nonconvex universal case C for some $x \in [0, 0.5]$	6
10	A nonconvex case (in bold lines) with a folded ruler inside.	7
11	Left: nonconvex universal case C for some $x \in [0, 0.5]$; the shaded trapezoid can be discarded. Right: folding a 6-link unit ruler $p_0p_1p_2p_3p_4p_5p_6$ into C	8

12	Nonconvex universal case C (in bold lines).	9
13	A plot of $\text{area}(C)$ (left) and its derivative (right).	11
14	For a 3-link ruler $1, t, 1$, where t is fixed, the area of the shaded triangle is minimized when $ p_0p_3 = p_1p_3 = 1$	12
15	Legend for the 5-link ruler used (in bold lines).	15
16	Case 1 (left) and Case 2 (right): the lower bounds are given by the shaded areas in each case.	16
17	Case 3: area above (left) and area below (right). The lower bound is given by the sum of the two shaded areas.	17
18	Case 4: area above (left) and area below (right). The lower bound is given by the sum of the two shaded areas.	19
19	Left: 6-universal case $R1$ (in bold lines). Right: folding a 6-link unit ruler $p_0p_1p_2p_3p_4p_5p_6$ into $R1$	21
20	Left: 4-universal case $R1/2$ (in bold lines). Right: folding a 4-link unit ruler $p_0p_1p_2p_3p_4$ into $R1/2$	21
21	Left: convex 4-universal case $S2$ (in bold lines). Right: folding a 4-link unit ruler $p_0p_1p_2p_3p_4$ into $S2$	21
22	Left: nonconvex 4-universal case $C2$ (in bold lines). Right: folding a 4-link unit ruler $p_0p_1p_2p_3p_4$ into $C2$	22
23	A plot of $\text{area}(S2)$ (left) and its derivative (right).	23
24	Derive $C2$ from $S2$. Left: $S2$ (in bold lines) with the shaded triangle discarded. Right: $C2$ (in bold lines) has minimum area 0.295... when c lies in the middle of the circular arc bd	24
25	A plot of $\text{area}(C2)$ (left) and its derivative (right).	26
26	Two cases of folding a 3-link ruler into $C2$ (in bold lines). Left: the middle link is short. Right: the middle link is long.	26
27	Fold a 3-link ruler (in bold lines) into $C2$. Left: $t \leq x$. Right: $t > x$	27
28	Two cases of folding a 4-link ruler into $C2$ (in bold lines). Left: p_2p_3 is short. Right: p_2p_3 is long.	27

29 Fold a 4-link ruler (in bold lines) into $C2$. Left: $|p_2c| \leq x$. Right:
 $|p_2c| > x$ 28

LIST OF TABLES

1	Known bounds for convex cases.	7
2	Known and new (in bold) bounds for nonconvex cases.	7

1 Introduction

A *universal case* for a family of sets is a set that contains a copy (under congruence or translation) of each set in the family. Geometric questions on finding minimum universal cases for different families of sets have been studied for over a hundred years. Various measures of minimality such as smallest area and shortest diameter were also considered.

L. Moser's worm problem is one of the most famous problems of this type. It asks for a set in the plane with minimum area that can be used to cover any curve of length one. Meir showed that a closed semidisk of unit diameter (depicted in Figure 1 (left)) is a universal case for all curves of unit length. It has area $\pi/8 < 0.3927$.




Figure 1: A universal case (in bold lines) for Moser's worm problem Left: a closed semidisk of unit diameter. Right: a plane curve of length 1 contained in the case.

Poole and Gerriets [6] constructed a smaller universal case from a rhombus (see Figure 2 (left)) whose long diagonal is of unit length and the larger angles are 120° . It has area $1/(2\sqrt{3}) < 0.2887$.




Figure 2: Another universal case (in bold lines) for Moser's worm problem. Left: a rhombus with a long diagonal of unit length and vertex angles of 60° and 120° . Right: a plane curve of length 1 contained in the case.

Better upper bounds were gradually obtained over two decades since 1970s (see [11, 14, 15]). The current best upper bound for convex cases is 0.276, derived

by Norwood and Poole [10]. For nonconvex cases, a better upper bound of 0.246 was achieved by Hansen [7]. The current best convex lower bound 0.2194 is due to Wetzel [15] and dates back to 1973.

One of the oldest problems concerning universal cases was posed by Lebesgue. The problem asks for a smallest area convex universal case that contains a congruent copy of every planar set of unit diameter. A minimal universal case depicted in Figure 3 (left) is due to Eggleston [5]. It is the union of a Reuleaux triangle with diameter one and a unit circle where a pair of the triangle vertices is a diameter of the circle. It has area $1.0046\dots$




Figure 3: A universal case (in bold lines) for Lebesgue's covering problem. Left: union of a unit Reuleaux triangle and a unit circle. Right: a Reuleaux triangle with unit diameter (shaded area) contained in the case.

Pál [12] (see also [2, Problem 1, p. 457]) proved an upper bound of 0.8454 with his truncated hexagon (shown in Figure 4 (left)), a regular hexagon circumscribed to a unit circle, with two corners cut off. Pál also derived a lower bound of 0.8257 which was further increased to 0.832 by Braß and Sharifi [3].

Similar problems in which translation (but no rotation) is allowed were also studied. For the same problem, a universal case under translation is also a universal case under congruence. The direct analogue of Lebesgue's problem asks for a smallest universal case that contains a translate of every planar set of unit diameter. The unit square, shown in Figure 5 (left), is known to be a universal




Figure 4: Another universal case (in bold lines) for Lebesgue's covering problem. Left: a regular hexagon circumscribed to a unit circle, with two corners cut off. Right: a Reuleaux triangle with unit diameter (shaded area) contained in the case.

case under translation (and congruence). One of the oldest results for universal cases under translation is due to Pál [13] who proved that the smallest universal translative case for all open curves of unit length is an equilateral triangle with height one and area $1/\sqrt{3}$ (depicted in Figure 6 (left)).




Figure 5: A universal case (in bold lines) for Lebesgue's covering problem under translation. Left: a unit square. Right: a Reuleaux triangle with unit diameter (shaded area) can rotate inside the square.

Another problem of finding universal (one dimensional) cases asks to determine the shortest interval into which a chain of line segments can be folded. A chain of line segments is called a *carpenter's ruler* consists of links of different lengths that are hinged together making it possible to fold. The problem was posed by Hopcroft, Joseph and Whitesides in [8] where the authors proved it to be NP-complete and provided a factor 2-approximation algorithm. In [4], Călinescu and Dumitrescu improved this result by showing that there exists a fully polynomial-




Figure 6: Smallest area translative universal case (in bold lines) for all open curves of unit length. Left: an equilateral triangle of unit height. Right: a unit open curve can rotate inside the equilateral triangle.

time approximation scheme. Hopcroft et al. [8] also observed that for a ruler with its longest link having length 1, the minimum folding length can be at least $2 - \epsilon$ for all $1 \geq \epsilon > 0$. Consider a ruler with $2n - 1$ links as depicted in Figure 7, its links have lengths 1 and $1 - \epsilon$ alternatively in which $n = \lceil 1/\epsilon \rceil$. The minimum folding length of this ruler is $(n - 1)\epsilon + 1 \geq 2 - \epsilon$.




Figure 7: The minimum folding length of a ruler can be almost double the length of its longest link.

Călinescu and Dumitrescu [4] (see also [2, Problem 9, p. 461]) introduced a two-dimensional version of the carpenter's ruler folding problem in 2005: finding the minimum area universal case for carpenter's rulers in the plane whose diameter is of the same length as a ruler's longest link. More formally, a carpenter's ruler L of n links is a chain of n line segments with endpoints p_0, p_1, \dots, p_n , with consecutive segments connected by hinges. For $0 \leq i \leq n - 1$, the segment $p_i p_{i+1}$ is a *link* of the ruler. A ruler with its longest link having length 1 is called a *unit ruler*. A folding of a ruler L is represented by the $n - 2$ angles $\angle p_i p_{i+1} p_{i+2} \in [0, \pi]$ for all

$0 \leq i \leq n - 2$. A *case* is a planar shape whose boundary is a *simple* closed curve (i.e., with no self-intersections). In particular, a case has no interior holes.

Obviously a unit ruler requires a case whose diameter is at least one; on the other hand, there exist cases of unit diameter that allow folding of *any* unit ruler inside, e.g., a disk of unit diameter, regardless of the number of links in the ruler. A ruler L can be folded inside a case S if and only if there exists a point $p \in S$ and a folding of L such that all the points on L are in S when p_0 is placed at p . In a folded position of the ruler, its links may cross each other; an example is shown in Figure 8 (right).

A case is said to be *universal* if any unit ruler (or all unit rulers) can be folded inside it. The question asks for the minimum area of a convex universal case of unit diameter. A disk of unit diameter and the Reuleaux triangle with one arc removed (call it $R2$), were shown to be universal by the authors [4]. $R2$ is depicted in Figure 8, its area is $\frac{\pi}{3} - \frac{\sqrt{3}}{4} = 0.614\dots$; it is the current best upper bound for the area of a convex universal case. For any n -link unit ruler p_0, p_1, \dots, p_n , a folding of it inside $R2$ such that all p_i 's lie on the circular arcs can be computed in $O(n)$ time. The authors [4] also achieved a lower bound of 0.375 using 3-link rulers, and this was further improved by Klein and Lenz [9] to 0.476 using 5-link rulers.




Figure 8: Left: convex universal case $R2$ (in bold lines). Right: folding a 6-link unit ruler $p_0p_1p_2p_3p_4p_5p_6$ into $R2$.

A case is *k-universal* if any unit ruler with at most k links can be folded into it. In the problem of finding a universal case with minimum area, the number of links (as well as the total length) of the rulers is irrelevant. However, it is worth

study if fewer links in the rulers allow better bounds. In [1], Alt et al. studied convex universal cases for rulers with a small number of links for which better upper bounds were achieved.

Since a universal case has unit diameter, it must be contained in a *lens* of radius 1, namely the intersection of two disks of unit radius passing through the centers of each other (see Figure 9). It was shown in [9] that no subset of $R2$ with a smaller area is universal. All previous work has focused on convex cases; the lower bounds were derived using convex hull of the rulers used in the respective arguments.




Figure 9: Universal cases (in bold lines) are contained in a lens of radius 1. Left: convex universal case $R2$. Right: nonconvex universal case C for some $x \in [0, 0.5]$.

Călinescu and Dumitrescu [4] also asked whether the convexity of the case makes any difference. Here we deal with nonconvex cases, i.e., cases with spikes are allowed (see Figure 10), and give a first partial answer to this question. Our main result concerning nonconvex universal cases is summarized in the following theorem.

Theorem 1. *There exists a (simple) nonconvex universal case C of unit diameter and area at most 0.583. The folding of any unit ruler with n links inside C can*

be computed in $O(n)$ time. On the other hand, the area of any simple nonconvex universal case of unit diameter must be at least 0.073.

For the problem of finding k -universal cases, our main result for $k = 4$ is summarized in the following theorem.

Theorem 2. *There exists a (simple) nonconvex 4-universal case $C2$ of unit diameter and area at most 0.296. The folding of any unit ruler with at most 4 links inside $C2$ can be computed in $O(1)$ time.*




Figure 10: A nonconvex case (in bold lines) with a folded ruler inside.

Table 1 and Table 2 summarize the known and new bounds for convex and general (convex and nonconvex) cases respectively.

	Universal	3-universal	4-universal	5-universal	6-universal
Upper bounds	0.614	0.486	0.486	0.523	0.523
Lower bounds	0.476	0.375	0.375	0.476	0.476

Table 1: Known bounds for convex cases.

	Universal	3-universal	4-universal	5-universal	6-universal
Upper bounds	0.583	0.296	0.296	0.523	0.523
Lower bounds	0.073	0.038	0.038	0.073	0.073

Table 2: Known and new (in bold) bounds for nonconvex cases.

In Section 2, we prove that the nonconvex case C shown in bold lines in Figure 11 is a universal case for any $x \in [0, \frac{1}{2}]$. Its area is at most 0.583 (achieved




Figure 11: Left: nonconvex universal case C for some $x \in [0, 0.5]$; the shaded trapezoid can be discarded. Right: folding a 6-link unit ruler $p_0p_1p_2p_3p_4p_5p_6$ into C .

when $x = 0.165$), i.e., smaller than the area of $R2$. Notice that the case whose boundary is the convex hull of C is a convex universal case whose area is at least 0.694, larger than the area of $R2$.

In Section 3, lower bounds for nonconvex universal cases are considered, i.e., only areas required by the simplicity of the case boundary are taken into account. We first derive a lower bound of 0.038 using a suitable 3-link ruler, and then extend the calculation to a suitable 5-link ruler and improve the lower bound to 0.073.

In Section 4, the problem of finding k -universal cases for $k = 4$ is considered. We construct another nonconvex case $C2$ with unit diameter. It is proved to be 4-universal with an algorithm for folding unit rulers with at most 4 links inside it. $C2$ has area at most 0.296, smaller than 0.486 which is the current best upper bound for the area of a convex 4-universal case.

2 Upper Bound

The upper bound in Theorem 1 will be proved using the simple nonconvex shape C shown in Figure 12. C is constructed as follows.

- $|ac| = |af| = |bg| = 1$
- $|bd| = |cd| = |ef| = |eg| = x$, $x \in [0, \frac{1}{2}]$
- Arcs ab and gf are centered at e with radii $1 - x$ and x respectively

- Arcs ag and bc are centered at d with radii $1 - x$ and x respectively

Notice that when $x = \frac{1}{2}$, C becomes a disk with diameter 1; and when $x = 0$, C is identical to $R2$. We show below that for any $x \in [0, \frac{1}{2}]$, C is a universal case with diameter 1. Choosing $x = 0.165$ yields a universal case with area ≤ 0.583 ; notice that this area is smaller than $0.614\dots$, the area of $R2$, the current smallest convex universal case.




Figure 12: Nonconvex universal case C (in bold lines).

2.1 Diameter of C

We show that C has diameter 1 for any $x \in [0, \frac{1}{2}]$. The diameter is given by a pair of points on the convex hull, thus it suffices to consider points on arcs ab , bc , fg , ga and segment cf . Let p and p' be two points on the convex hull of C .

Fix p on arc ab . If p' is on arc ab , $|pp'| \leq |ab| < |ac| = 1$. If p' is on arc bc or segment cf , $|pp'| \leq |ac| = 1$. If p' is on arc fg , extend segment pe until it intersects arc fg at point p'' . If $p' = p''$, $|pp'| = |pe| + |ep''| = 1$; otherwise, segments pe, ep', pp' form a triangle, thus, $|pp'| < |pe| + |ep'| = |ae| + |ef| = 1$. If p' is on arc ga , $|pp'| \leq |bg| = 1$.

Fix p on arc bc . If p' is on segment cf , $|pp'| \leq |bf| < |bg| = 1$. If p' is on arc fg , $|pp'| \leq |bg| = 1$. By symmetry, C has diameter 1.

2.2 Algorithm for folding a ruler inside C

We show that the folding of any unit ruler with n links inside C can be computed in $O(n)$ time. We adapt the algorithm introduced in [4] to work with our case C . Fix the first free endpoint at some (arbitrary) point p on a circular arc. Iteratively fix the next point of the ruler at some intersection point between the arcs of C and the circle centered at p with radius the length of the current link.

Notice that for any point p on the circular arcs of C , and for any $t \in [0, 1]$, there exists at least one point p' on these arcs such that $|pp'| = t$. This guarantees the existence of the intersection points used in the iterative steps of the above algorithm.

2.3 Minimum area of C

The area of C is the sum of areas of the sectors dag , dbc , eab and efg minus the area of the triangle Δade . In the triangle Δade , we have $\angle ade = \arccos \frac{1-2x}{2-2x}$. The sectors dag and eab have the same area $\frac{(1-x)^2}{2} \arccos \frac{1-2x}{2-2x}$. The sectors dbc and efg have the same area $\frac{x^2}{2} \arccos \frac{1-2x}{2-2x}$.

The triangle Δade has area $\frac{1-2x}{4} \sqrt{3-4x}$. It follows that

$$\begin{aligned} \text{area}(C) &= 2\text{area}(\text{sector } dag) + 2\text{area}(\text{sector } dbc) - \text{area}(\Delta ade) \\ &= (1-x)^2 \arccos \frac{1-2x}{2-2x} + x^2 \arccos \frac{1-2x}{2-2x} - \frac{1-2x}{4} \sqrt{3-4x} \\ &= (1-2x+2x^2) \arccos \frac{1-2x}{2-2x} + \frac{2x-1}{4} \sqrt{3-4x}. \end{aligned}$$

Taking derivatives yields

$$\frac{d(\text{area}(C))}{dx} = (4x-2) \arccos \frac{1-2x}{2-2x} + \frac{3-7x+5x^2}{(1-x)\sqrt{3-4x}}.$$




Figure 13: A plot of $\text{area}(C)$ (left) and its derivative (right).

Solving for $\frac{d(\text{area}(C))}{dx} = 0$ yields a single root $x = 0.165\dots$, at which C has the smallest area, $\text{area}(C) \leq 0.583$ (see Figure 13).

3 Lower Bound

We start with Lemma 1 (in Subsection 3.1), which gives a lower bound of 0.038 for the area required by a suitable 3-link ruler. As it turns out, this lower bound is the best possible for all 3-link rulers. Lemma 1 will be reused when deriving a lower bound for 5-link rulers (in Subsection 3.2), improving this first bound to 0.073.

3.1 Lower Bound with One 3-Link ruler

For 3-link rulers, it is sufficient to consider the sequence of lengths $1, t, 1$ with $t \in (0, 1)$. Indeed, given a folding of ruler $1, t, 1$, and an arbitrary unit 3-link ruler with links a, t, b , make the t -links of the two rulers coincide, and fold the a - and b -links over the two unit links; the resulting folding is a valid one in the same case required by the $1, t, 1$ ruler.

For the 3-link ruler with link lengths $1, t, 1$, the two 1-links must intersect otherwise the diameter constraint will be violated, see Figure 14. The shaded triangle is the only area that counts for the nonconvex lower bound.




Figure 14: For a 3-link ruler $1, t, 1$, where t is fixed, the area of the shaded triangle is minimized when $|p_0p_3| = |p_1p_3| = 1$.

Lemma 1. *For any $t \in (0, 1)$, the shaded triangle in Figure 14 is minimized when $\alpha = \arccos \frac{t}{2} - \frac{\pi}{3}$ and $\beta = \arccos \frac{t}{2}$.*

Proof. By symmetry, we can assume that $\alpha \leq \beta$. Denote the area of the shaded triangle Δp_1p_2q by S . Since the triangle has base t , its height h determines the area. The height h is the distance between p_1p_2 and the intersection point between p_0p_1 and p_2p_3 . For any fixed $\alpha \in [\arccos \frac{t}{2} - \frac{\pi}{3}, \arccos \frac{t}{2}]$, the area is minimized when β is minimized without violating the diameter constraint $|p_0p_3| \leq 1$. Denote this angle by $\beta(\alpha)$; $\beta(\alpha)$ is a monotonically decreasing function that can be determined by computing the intersection of two circles of radius 1 centered at p_0 and p_2 . In the following discussion, we will refer to this angle by β .

It suffices to express the area S as a function of two parameters, t and α . In fact, $h \cot \alpha + h \cot \beta = t$ or

$$h = \frac{t}{\cot \alpha + \cot \beta}.$$

So

$$S(t, \alpha) = \frac{th}{2} = \frac{t^2}{2(\cot \alpha + \cot \beta)}.$$

Taking derivative with respect to α , we have

$$\frac{dS(t, \alpha)}{d\alpha} = \frac{t^2}{2(\cot \alpha + \cot \beta)^2} \left(\frac{1}{\sin^2 \alpha} + \frac{1}{\sin^2 \beta} \frac{d\beta}{d\alpha} \right).$$

To see that S is minimized when α is minimized, we need to show that $\frac{dS(t, \alpha)}{d\alpha} > 0$,

i.e.,

$$\begin{aligned} \frac{1}{\sin^2 \alpha} + \frac{1}{\sin^2 \beta} \frac{d\beta}{d\alpha} &> 0, \text{ or} \\ \frac{d\beta}{d\alpha} &> -\frac{b^2}{a^2}. \end{aligned}$$

Suppose p_1 is fixed at $(0, 0)$, p_2 is at $(t, 0)$, then p_0 has coordinates $(\cos \alpha, \sin \alpha)$, p_3 has coordinates $(t - \cos \beta, \sin \beta)$. Since $|p_0 p_3| = 1$, we have

$$(t - \cos \beta - \cos \alpha)^2 + (\sin \beta - \sin \alpha)^2 = 1, \text{ or}$$

$$t^2 + 1 - 2t \cos \beta - 2t \cos \alpha + 2 \cos(\alpha + \beta) = 0.$$

Taking derivative with respect to α , we have

$$2t \sin \beta \frac{d\beta}{d\alpha} + 2t \sin \alpha - 2 \sin(\alpha + \beta) \left(1 + \frac{d\beta}{d\alpha}\right) = 0, \text{ or}$$

$$\frac{d\beta}{d\alpha} = \frac{\sin(\alpha + \beta) - t \sin \alpha}{t \sin \beta - \sin(\alpha + \beta)}.$$

Notice that in the shaded triangle $\Delta p_1 p_2 q$,

$$\frac{a}{\sin \alpha} = \frac{b}{\sin \beta} = \frac{t}{\sin(\alpha + \beta)}.$$

So

$$\frac{d\beta}{d\alpha} = -\frac{1-a}{1-b}.$$

Thus $\frac{dS(t,\alpha)}{d\alpha} > 0$ is equivalent to

$$\frac{d\beta}{d\alpha} = -\frac{1-a}{1-b} > -\frac{b^2}{a^2}, \text{ or}$$

$$b^2 - b^3 > a^2 - a^3.$$

This inequality holds if $0 < a < 2/3$ and $a < b < (1-a)/2 + \sqrt{(1+3a)(1-a)}/2$. We know this is true since in triangle $\Delta p_0 p_3 q$, $(1-a) + (1-b) > 1$. Recall $\alpha \leq \beta$, so $a \leq b$. If $a > 1/2$, $(1-a) + (1-b) \leq 2(1-a) < 1$, so $a \leq 1/2$. And $b < 1-a < (1-a)/2 + \sqrt{(1-a)^2}/2$.

Observe that α and the correspondingly β are determined when $|p_1 p_3| = 1$ (see Figure 14). Moreover, this value of α is the minimum possible; indeed, if α is getting smaller, either $|p_0 p_3|$ or $|p_1 p_3|$ will violate the diameter constraint. In the isosceles triangle $\Delta p_1 p_2 p_3$, we have $\beta = \alpha + \angle p_0 p_1 p_3$ and $\cos \beta = \frac{t}{2}$. In the equilateral triangle $\Delta p_0 p_1 p_3$, we have $\angle p_0 p_1 p_3 = 60^\circ$. So $\beta = \alpha + \frac{\pi}{3} = \arccos \frac{t}{2}$. \square

Now we are ready to show our first lower bound on simple nonconvex cases. By Lemma 1,

$$S(t, \alpha) \geq U(t) := \frac{t^2}{2(\cot(\arccos \frac{t}{2} - \frac{\pi}{3}) + \cot \arccos \frac{t}{2})}. \quad (1)$$

It is easy to check that $U(0.676) \geq 0.038$, as desired. For $t \in (0, 1)$, $U(t)$ attains its maximum value for $t = 0.676\dots$, and this is the best possible bound for a single 3-link ruler.

3.2 Lower Bound with One 5-link Ruler

Consider a special ruler with 5 links of lengths 1, 0.6, 1, 0.6, 1 as shown in Figure 15. Recall that all the 1-links must pairwise intersect. Since the ruler is symmetric, w.l.o.g., we can assume that $\beta \geq \gamma$. The following lemma gives a

better lower bound using this ruler.




Figure 15: Legend for the 5-link ruler used (in bold lines).

Lemma 2. *The minimum area of a simple (nonconvex) case of unit diameter required by folding the ruler 1, 0.6, 1, 0.6, 1 inside it is at least 0.073.*

Proof. Put $t = 0.6$. The Cartesian coordinate is set up as follows: fix the origin at p_2 and let the x -axis pass through p_3 . We have $p_2 = (0, 0)$, $p_3 = (1, 0)$, $p_1 = (t \cos \beta, t \sin \beta)$ and $p_4 = (1 - t \cos \gamma, t \sin \gamma)$. Recall that the case is required to be simple, i.e., no self-intersections or holes are allowed. According to the analysis of 3-link rulers, $\beta, \gamma \in [\arccos \frac{t}{2} - \frac{\pi}{3}, \arccos \frac{t}{2}]$. We distinguish four cases according to the angles β and γ .

Case 1: The two t -links do not intersect. This case includes the situation that p_3p_4 is folded below p_2p_3 . As shown in Figure 16 (left), each shaded triangle is minimized using Lemma 1.

$$\beta = \gamma = \arccos \frac{t}{2} = 72.54\dots^\circ, \quad \alpha = \delta = \arccos \frac{t}{2} - \frac{\pi}{3} = 12.54\dots^\circ.$$

Observe that this is not a valid folding since the two 1-links p_0p_1 and p_4p_5 do not intersect. However, it gives a valid lower bound since for any fixed β and γ , increasing α or δ (to make the 1-links intersect) will increase the total area. By

(1), the lower bound for Case 1 is

$$2U(t) = \frac{t^2}{\cot(\arccos \frac{t}{2} - \frac{\pi}{3}) + \cot \arccos \frac{t}{2}} \geq 0.074.$$




Figure 16: Case 1 (left) and Case 2 (right): the lower bounds are given by the shaded areas in each case.

Case 2: The two t -links intersect and both β and γ are at least 16° . As shown in Figure 16 (right), increasing β or γ will enlarge the upper shaded area consisting of the triangles $\Delta_{q_0p_1p_2}$ and $\Delta_{q_0p_3q_1}$. The area of the triangle below p_2p_3 will decrease but we simply ignore it when computing the lower bound in this case. Similar to the case of 3-link rulers, when $\beta = \gamma = 16^\circ$, α should be minimized under the constraint $|p_0p_3| \leq 1$ otherwise the area of the upper right small triangle $\Delta_{q_1p_1q_2}$ will increase. In this configuration, triangle $\Delta_{q_0p_1p_2}$ has height $t \sin \beta$. Its base $|p_2q_0|$ is the difference between the projections of the segments p_2p_1 and q_0p_1 on the x -axis, and $\angle p_1q_0p_3 = \alpha + \beta$. It follows that

$$b = |p_2q_0| = t \cos \beta - \frac{t \sin \beta}{\tan(\alpha + \beta)}. \quad (2)$$

Triangle $\Delta q_0 p_3 q_1$ has base $1 - b$. Its height h equals to the y -coordinate of q_1 which is the intersection point of lines $p_0 p_1$ and $p_3 p_4$. The equation of line $p_0 p_1$ is

$$y = \tan(\alpha + \beta)(x - t \cos \beta) + t \sin \beta.$$

The equation of line $p_3 p_4$ is

$$y = (1 - x) \tan \gamma.$$

The y -coordinate of their intersection is

$$h = \frac{(t \sin \beta + (1 - t \cos \beta) \tan(\alpha + \beta)) \tan \gamma}{\tan \gamma + \tan(\alpha + \beta)}. \quad (3)$$

The total shaded area is the sum of the two areas of triangles $\Delta q_0 p_1 p_2$ and $\Delta q_0 p_3 q_1$, namely

$$\frac{bt \sin \beta + (1 - b)h}{2} \geq 0.073. \quad (4)$$

Case 3: The two t -links intersect and $\beta \geq 16^\circ, \gamma \leq 16^\circ$. In this case, the lower




Figure 17: Case 3: area above (left) and area below (right). The lower bound is given by the sum of the two shaded areas.

bound consists of two parts, the minimum shaded areas above and below p_2p_3 , denoted by S_a and S_b respectively.

As shown in Figure 17 (left), with a similar argument as in Case 2, the minimum shaded area above p_2p_3 is achieved when $\beta = 16^\circ$, $\gamma = \arccos \frac{t}{2} - \frac{\pi}{3}$ (which is the minimum value) and α is minimized under the constraint $|p_0p_3| \leq 1$. Plugging in these values into (2), (3) and (4) in Case 2 yields $S_a \geq 0.067$.

Observe that when β and γ increase, α and δ can take smaller values under the constraints $|p_0p_3| \leq 1$, $|p_2p_5| \leq 1$ and thus form a smaller triangle below p_2p_3 . So the area of triangle $\Delta q_0q_1q_2$ is minimized when both β and γ take the maximum values, i.e., $\gamma = 16^\circ$ and β is chosen such that p_4 lies on p_1p_2 (p_1p_2 and p_3p_4 need to intersect). Then, both α and δ are minimized under the diameter constraints. This configuration is shown in Figure 17 (right). Similar to (2), we have

$$\begin{aligned} |p_2q_0| &= t \cos \beta - \frac{t \sin \beta}{\tan(\alpha + \beta)}, \\ |q_1p_3| &= t \cos \gamma - \frac{t \sin \gamma}{\tan(\gamma + \delta)}. \end{aligned} \tag{5}$$

The base of triangle $\Delta q_0q_1q_2$ is $b = |p_2q_0| + |q_1p_3| - 1$. The height h of this triangle is the absolute value of the y -coordinate of q_2 , the intersection point of lines p_0p_1 and p_4p_5 . The equation of line p_0p_1 is

$$y = \tan(\alpha + \beta)(x - t \cos \beta) + t \sin \beta.$$

The equation of line p_4p_5 is

$$y = \tan(\gamma + \delta)(1 - t \cos \gamma - x) + t \sin \gamma.$$

Solving for their intersection point gives

$$h = \frac{\tan(\alpha + \beta) \tan(\gamma + \delta)(t \cos \beta + t \cos \gamma - 1)}{\tan(\alpha + \beta) + \tan(\gamma + \delta)} - \frac{t \tan(\gamma + \delta) \sin \beta + t \tan(\alpha + \beta) \sin \gamma}{\tan(\alpha + \beta) + \tan(\gamma + \delta)}. \quad (6)$$

It follows that $S_b = \frac{1}{2}hb \geq 0.006$, and consequently, the minimum total shaded area is $S_a + S_b \geq 0.073$.

Case 4: both β and γ are no more than 16° . Notice that since $t = 0.6$, the two t -links must intersect. Similar to Case 3, the lower bound is calculated as




Figure 18: Case 4: area above (left) and area below (right). The lower bound is given by the sum of the two shaded areas.

the sum of minimized areas of shaded triangles above and below p_2p_3 . For the triangle above p_2p_3 , recall that β and γ both have the minimum possible value, $\arccos \frac{t}{2} - \frac{\pi}{3}$, as shown in Figure 18 (left). The minimized isosceles triangle above p_2p_3 has base 1 and height $\frac{\tan \beta}{2}$. Its area is

$$S_a = \frac{\tan(\arccos \frac{t}{2} - \frac{\pi}{3})}{4} \geq 0.055.$$

The area of the triangle below p_2p_3 is minimized when both β and γ take the maximum value (16°). Using (5) and (6) in Case 3 and $\alpha = \delta$, $\beta = \gamma$, the triangle below p_2p_3 has base

$$b = 2 \left(t \cos \beta - \frac{t \sin \beta}{\tan(\alpha + \beta)} \right) - 1$$

and height

$$h = \frac{(2t \cos \beta - 1) \tan(\alpha + \beta)}{2} - t \sin \beta.$$

Its area is $S_b = \frac{hb}{2} \geq 0.019$. The minimum total shaded area is $S_a + S_b \geq 0.074$.

In summary, by Cases 2 and 3 of the analysis, the minimum nonconvex area required by folding the ruler 1, 0.6, 1, 0.6, 1 within a case of unit diameter is at least 0.073. □

4 k -universal Cases

In this section, we consider the problem of finding k -universal cases of minimum areas. Let A_k be the smallest area of a convex k -universal case and B_k be the smallest area of an arbitrary (convex or nonconvex) k -universal case. For any $k \in \mathbb{N}$, $B_k \leq A_k$. Additionally, $\forall i < j \in \mathbb{N}$, $A_i \leq A_j$, $B_i \leq B_j$. Our goal is to find better bounds for B_k . This problem was first studied by Alt et al. [1], in which the authors proved that $A_3 \leq A_4 < 0.486$ and $A_5 \leq A_6 \leq 0.523$. With a simple nonconvex case $C2$ (see Figure 22), we derive a better upper bound, $B_4 < 0.296$, which is smaller than the current best upper bound of $A_4 < 0.486$.

Replacing an circular arc in $R2$ (Figure 8) with its chord results in the case $R1$, depicted in Figure 19, a sector with radius one and center angle 60° , its area is $\pi/6$. Alt et al. [1] proved that $R1$ is 6-universal but not 7-universal. Thus, $A_6 \leq \text{area}(\text{sector } abc) = \pi/6 = 0.532\dots$

Further replacing half of the remaining circular arc in $R1$ with a line segment




Figure 19: Left: 6-universal case $R1$ (in bold lines). Right: folding a 6-link unit ruler $p_0p_1p_2p_3p_4p_5p_6$ into $R1$.




Figure 20: Left: 4-universal case $R1/2$ (in bold lines). Right: folding a 4-link unit ruler $p_0p_1p_2p_3p_4$ into $R1/2$.

produces the case $R1/2$ shown in Figure 20. $R1/2$ consists of the sector abc with radius one and center angle 30° and the isosceles triangle Δacd with base $|ac| = 1$ and height $|bd|/2 = 1/2$. $R1/2$ was shown to be 4-universal but not 5-universal in [1]. Thus, $A_4 \leq \text{area}(\text{sector } abc) + \text{area}(\Delta acd) = \pi/12 + 1/4 = 0.512\dots$




Figure 21: Left: convex 4-universal case $S2$ (in bold lines). Right: folding a 4-link unit ruler $p_0p_1p_2p_3p_4$ into $S2$.

Alt et al. [1] improved the upper bounds of A_4 with a better 4-universal case $S2$ depicted in Figure 21. $S2$ is constructed as follows. $|ab| = 1$, pick an arbitrary

point c on the circular arc ob , let $x = |bc|$. Draw a circle centered at c with radius x , let it intersect arc oa at point d . Notice that d exists only if $|oc| \leq |cb|$. So c must be at the middle point of arc ob or higher, i.e., $\angle cab \geq 30^\circ$. In $\triangle abc$, $|bc| = x$, $|ac| = |ab| = 1$. Thus $|bc| = |cd| = x \geq \sqrt{2 - \sqrt{3}} = 0.517\dots$. Similar to the universal case C introduced in Section 2, $S2$ is proved to be 4-universal with any value of $x \in [\sqrt{2 - \sqrt{3}}, 1]$. Notice that when $x = 1$, $S2$ becomes the 6-universal case $R1$; when c lies at the middle of the arc ob , $S2$ is identical to $R1/2$.

The minimum area of $S2$ was claimed (without proof) to be approximately 0.485. We confirm this result by providing full details in Section 4.1. However, we observe that the shaded triangle $\triangle ade$ in Figure 21 (left) is not necessary for the folding algorithm introduced in [1]. If we discard this triangle, a family of nonconvex cases $abcde$ with parameter x can be obtained. In Section 4.2, we




Figure 22: Left: nonconvex 4-universal case $C2$ (in bold lines). Right: folding a 4-link unit ruler $p_0p_1p_2p_3p_4$ into $C2$.

prove that for $x = \sqrt{2 - \sqrt{3}} = 0.517\dots$, i.e., c is at the middle point of arc ob , the nonconvex case $C2$ shown in Figure 22 has the smallest area 0.295... in this family. In Section 4.3, we show that $C2$ is a 4-universal but not 5-universal case. Thus, the upper bound of B_4 is improved from 0.486 to 0.296.

4.1 Area of S_2

In preparation for calculating the area of C_2 , we provide the missing details in [1] for calculating the minimum area of S_2 . In Section 4.2, the area of C_2 is derived based on the following calculations. The area of S_2 is the sum of the areas of its three parts: the isosceles triangles Δbcd , Δabd and the circular segment cb . In Δbcd , $\alpha = \arccos \frac{1}{2x}$ and its area is $\frac{\sqrt{4x^2-1}}{4}$. In isosceles triangle Δabc , $\beta = \arccos \frac{x}{2}$ and its area is $\frac{x}{4}\sqrt{4-x^2}$. The area of the circular segment cb is the area of the circular sector abc minus the area of Δabc , i.e., $\frac{\pi}{2} - \arccos \frac{x}{2} - \frac{x}{4}\sqrt{4-x^2}$. In Δabd , $\angle abd = \beta - \alpha$, its area is $\sin(\beta - \alpha)/2 = \sin\left(\arccos \frac{x}{2} - \arccos \frac{1}{2x}\right)/2$. In summary,

$$\begin{aligned} \text{area}(S_2) &= \text{area}(\Delta bcd) + \text{area}(\Delta abd) + \text{area}(\text{circular segment } cb) \\ &= \frac{\sqrt{4x^2-1}}{4} + \sin\left(\arccos \frac{x}{2} - \arccos \frac{1}{2x}\right)/2 \\ &\quad + \frac{\pi}{2} - \arccos \frac{x}{2} - \frac{x}{4}\sqrt{4-x^2}. \end{aligned}$$

Taking derivative yields,

$$\begin{aligned} \frac{d(\text{area}(S_2))}{dx} &= \frac{4+x^2}{4\sqrt{4-x^2}} - \frac{\sqrt{4-x^2}}{4} + \frac{x}{\sqrt{4x^2-1}} \\ &\quad - \left(\frac{1}{2x\sqrt{4x^2-1}} + \frac{1}{2\sqrt{4-x^2}}\right) \cos\left(\arccos \frac{x}{2} - \arccos \frac{1}{2x}\right). \end{aligned}$$




Figure 23: A plot of $\text{area}(S_2)$ (left) and its derivative (right).

Solving for $\frac{d(\text{area}(S2))}{dx} = 0$ yields two roots in $[0.5, 1]$ (see Figure 23). When $x = 0.743\dots$, $S2$ has the smallest area $\text{area}(S2) \leq 0.486$.

4.2 Area of $C2$




Figure 24: Derive $C2$ from $S2$. Left: $S2$ (in bold lines) with the shaded triangle discarded. Right: $C2$ (in bold lines) has minimum area $0.295\dots$ when c lies in the middle of the circular arc bd .

Due to the subtraction of triangle Δade , we need to calculate the area of triangle Δabe instead of Δabd which is used in the area formula of $S2$. In Δabc (see Figure 24 (left)), $\angle bac = \pi - 2\beta$. In Δabe , $\angle abe = \beta - \alpha$ and

$$\angle aeb = \pi - \angle bae - \angle abe = \pi - (\pi - 2\beta) - (\beta - \alpha) = \beta + \alpha.$$

So $\frac{|be|}{\sin(\pi - 2\beta)} = \frac{|ab|}{\sin(\beta + \alpha)} = \frac{1}{\sin(\beta + \alpha)}$, or $|be| = \frac{\sin(2\beta)}{\sin(\beta + \alpha)}$.

The area of Δabe is

$$\begin{aligned} \text{area}(\Delta abe) &= |ab| \cdot |be| \sin(\beta - \alpha) / 2 \\ &= \frac{\sin\left(2 \arccos \frac{x}{2}\right) \sin\left(\arccos \frac{x}{2} - \arccos \frac{1}{2x}\right)}{\sin\left(\arccos \frac{x}{2} + \arccos \frac{1}{2x}\right)}. \end{aligned}$$

Notice that

$$\begin{aligned}\sin\left(2\arccos\frac{x}{2}\right) &= \frac{x\sqrt{4-x^2}}{2}, \\ \sin\left(\arccos\frac{x}{2}-\arccos\frac{1}{2x}\right) &= \frac{\sqrt{4-x^2}-x\sqrt{4x^2-1}}{4x} \\ \sin\left(\arccos\frac{x}{2}+\arccos\frac{1}{2x}\right) &= \frac{\sqrt{4-x^2}+x\sqrt{4x^2-1}}{4x}.\end{aligned}$$

Thus, the area of the convex case $C2$ is

$$\begin{aligned}\text{area}(C2) &= \text{area}(\Delta bcd) + \text{area}(\text{circular segment } cb) + \text{area}(\Delta abe) \\ &= \frac{\sqrt{4x^2-1}}{4} + \frac{\pi}{2} - \arccos\frac{x}{2} - \frac{x}{4}\sqrt{4-x^2} \\ &\quad + \frac{x(4-x^2)-x^2\sqrt{(4-x^2)(4x^2-1)}}{4\sqrt{4-x^2}+4x\sqrt{4x^2-1}}.\end{aligned}$$

Taking derivative yields

$$\begin{aligned}\frac{d(\text{area}(C2))}{dx} &= \frac{4+x^2}{4\sqrt{4-x^2}} - \frac{\sqrt{4-x^2}}{4} + \frac{x}{\sqrt{4x^2-1}} \\ &\quad - \frac{4x(8x^2-1)(4-x^2)}{(4\sqrt{4-x^2}+4x\sqrt{4x^2-1})^2\sqrt{4x^2-1}} \\ &\quad - \frac{4x^3\sqrt{4x^2-1}-32x^4\sqrt{4-x^2}}{(4\sqrt{4-x^2}+4x\sqrt{4x^2-1})^2} \\ &\quad + \frac{4-3x^2-2x\sqrt{(4-x^2)(4x^2-1)}}{4\sqrt{4-x^2}+4x\sqrt{4x^2-1}} \\ &\quad - \frac{x^2(8x(4-x^2)-2x(4x^2-1))}{8(4-x^2)\sqrt{4x^2-1}+8x(4x^2-1)\sqrt{4-x^2}}.\end{aligned}$$

We verify that $\frac{d(\text{area}(C2))}{dx} = 0$ has no real root in $[\sqrt{2-\sqrt{3}}, 1]$. Indeed, as shown in Figure 25, $\text{area}(C2)$ is a monotonically increasing function with respect to x in this range. Thus, the area of $C2$ is minimized when $x = \sqrt{2-\sqrt{3}}$, as depicted in Figure 24 (right). Since c lies in the middle of the circular arc bd , line segment ce




Figure 25: A plot of $\text{area}(C2)$ (left) and its derivative (right).

is perpendicular to ed and $|ed| = 1/2$, $|ce| = \sqrt{7 - 4\sqrt{3}}/2$. The minimum area of $C2$ is $\text{area}(\text{sector } abc) + \text{area}(\Delta cde) = \pi/12 + \sqrt{7 - 4\sqrt{3}}/8 = 0.295\dots$

4.3 $C2$ is 4-universal

First we show that any 3-link unit ruler $p_0p_1p_2p_3$ can be folded into $C2$ (see Figure 26).




Figure 26: Two cases of folding a 3-link ruler into $C2$ (in bold lines). Left: the middle link is short. Right: the middle link is long.

Lemma 3. $C2$ is a nonconvex 3-universal case.

Proof. It is sufficient to consider rulers with links $1, t, 1$ for any $t \in (0, 1]$.

Case 1: t is small, i.e., $t \leq x = \sqrt{2 - \sqrt{3}}$. So both end points of the center link t can be placed on the circular arc bc (see Figure 27 (left)). Since arc bc is




Figure 27: Fold a 3-link ruler (in bold lines) into $C2$. Left: $t \leq x$. Right: $t > x$.

centered at a with radius 1, both links of length one can be folded from arc bc to a .

Case 2: t is large, i.e., $|cd| = x < t \leq 1 = |bd|$. As shown in Figure 27 (right), starting from point d , there exists a point p on arc bc such that $|dp| = t$. So the $1, t, 1$ ruler can be placed from b to d to p to a . \square

Now we prove that any 4-link unit ruler $p_0p_1p_2p_3p_4$ can be folded inside $C2$ (as illustrated in Figure 28).




Figure 28: Two cases of folding a 4-link ruler into $C2$ (in bold lines). Left: p_2p_3 is short. Right: p_2p_3 is long.

Lemma 4. $C2$ is a nonconvex 4-universal case.

Proof. It is sufficient to consider unit rulers with links $1, t, t', 1$.

Case 1: $t + t' \leq 1$. The folding problem of this ruler can be reduced to a folding problem of a 3-link ruler $1, t + t', 1$ which is already solved by Lemma 3.

Case 2: $t + t' > 1$. Without loss of generality, we assume $t \geq t'$. Label the end points of the ruler by p_i , $i = 0, 1, 2, 3, 4$, such that $|p_0p_1| = |p_3p_4| = 1$, $|p_1p_2| = t$ and $|p_2p_3| = t'$. Fold the ruler such that the first two links overlap each other, i.e., p_2 lies on p_0p_1 . As illustrated in Figure 29 (left), we place p_0p_1 at bd , then p_2 is on eb otherwise $t + t' \leq 1$. $|p_2d| = t \geq t'$ and $|p_2b| = 1 - t < t'$, so there exists a point p on the circular arc bcd (notice that the arc cd is imaginary) such that $|p_2p| = t'$.

Case 2a: p lies on arc bc . We can place p_3 at p then p_4 at a as illustrated in Figure 29 (left).

Case 2b: p lies on (the imaginary) arc cd . We flip the ruler around with respect to the axis ac . As shown in Figure 29 (right), now p_0 is at d , p_1 is at b and p_2 is at p'_2 . Point p is also flipped to point p' on arc bc . So we can place p_3 at p' , p_4 at a .

Thus, in both cases the 4-link ruler can be folded into $C2$. More formally, if p_0p_1 (or p_1p_0) is placed at bd and p_2 is folded on p_0p_1 , the distance $|p_2c| = \sqrt{(1/2 - t)^2 + (7/4 - \sqrt{3})}$. If $|p_2c| \geq t'$ (*Case 2a*), p_0 is placed at b . Otherwise $|p_2c| < t'$ (*Case 2b*), p_0 is placed at d . Then in both cases, there exists a point p on arc bc such that $|p_2p| = t'$. \square




Figure 29: Fold a 4-link ruler (in bold lines) into $C2$. Left: $|p_2c| \leq x$. Right: $|p_2c| > x$.

Hence Theorem 2 follows. Additionally we show that there exists a ruler with 5 links that cannot be folded inside $C2$.

Lemma 5. $C2$ is not a 5-universal case.

Proof. Consider folding the 5-link ruler 1, 0.6, 1, 0.6, 1 into $C2$. The 0.6 links are in between the 1 links so both their end points must lie on a , d or arc bc . But since $1 > 0.6 > \sqrt{2 - \sqrt{3}}$, between d and some point on arc bc is the only possible position among all the combinations. Suppose the first 0.6 link is placed in this position, the other end points of the two 1 links adjacent to it must be placed at b and a respectively. But the second 0.6 link must also be placed to this only position which is impossible. \square

5 Summary and Future Directions

For the problem of finding minimum nonconvex universal cases for carpenter's rulers, we have shown:

- The area of a smallest nonconvex universal case with unit diameter is at most 0.583.
- The area of any nonconvex universal case is at least 0.073.
- The area of a smallest nonconvex 3-universal case is at most 0.296 and at least 0.038.
- The area of a smallest nonconvex 4-universal case is at most 0.296.

In Section 3, the best possible lower bound given by one 3-link ruler is achieved, whereas the one given by a 5-link ruler is not. Computer experiments suggest that 5-link rulers require folding area at least 0.137; more precisely:

- The minimum folding of a 5-link ruler with lengths 1, 0.6, 1, 0.6, 1 has (nonconvex) area at least 0.092.
- The minimum folding of a 5-link (symmetric) ruler with lengths 1, t , 1, t , 1 has area at least 0.115 when $t = 0.8$.

- The minimum folding of a 5-link (asymmetric) ruler with lengths $1, t_1, 1, t_2, 1$ has area at least 0.137 when $t_1 = 0.7, t_2 = 0.4$.

The difficulty of approaching these better bounds lies in the complicated computations of nonconvex areas in many sub-cases. Note however that even the computational results were used, the resulting lower bounds would still be far away from the current upper bound of 0.583, which we believe is closer to the truth.

Based on these observations, possible future research directions are:

1. Derive better lower bounds for universal cases using rulers with more links.
2. Derive better lower bounds for universal cases using combinations of multiple rulers.

Recall that the area of $R1$ is $0.523\dots$ which is the current best upper bound for A_5, A_6 and B_5, B_6 . In Section 4, we improved the upper bound for B_4 (and B_3) to 0.296 with the nonconvex 4-universal case $C2$. But we also showed that $C2$ is not a 5-universal case. Based on these results, following questions can be asked:

3. Is there a 3-universal case with area smaller than the area of $C2$, namely 0.296?
4. Is there a 5-universal case with area smaller than the area of $R1$, namely 0.523?
5. What are the tight bounds for A_k and B_k when $k = 3, 4, 5$ and 6?
6. Are there convex k -universal cases for $k > 6$ with area smaller than 0.614, the area of the current best convex universal case $R2$?
7. Are there k -universal cases for $k > 6$ with area smaller than 0.583, the area of the current best (nonconvex) universal case C ?
8. Especially, we are interested in the connection between the problem of finding minimum area k -universal cases and the problem of finding minimum area

universal cases. More specifically, is there a k such that A_k (B_k) matches the universal convex (nonconvex) bound?

Additionally, in [9], Klein and Lenz showed that no subset of $R2$ with smaller area than $R2$ is a universal case. This is proved by using a ruler with n links where n goes to infinity. The authors showed that the only possible folding inside $R2$ covers the whole area of $R2$. Motivated by this result, we propose the following interesting open problems:

9. Does a similar result hold for the nonconvex universal case C ?
10. Does a similar result hold for the nonconvex 4-universal case $C2$? Notice that in this problem, the method with rulers having the number of links going to infinity (which was used in [9]) is not applicable. Rulers are restricted to have at most 4 links. However, a possible approach is to use a combination of multiple rulers and show that no matter how the rulers are folded, $C2$ will be covered.

6 References

- [1] H. Alt, K. Buchin, O. Cheong, F. Hurtado, C. Knauer, A. Schulz, and S. Whitesides, Small boxes for carpenter’s rules, manuscript, 2006.
<http://page.mi.fu-berlin.de/alt/papers/carpenter.pdf>
- [2] P. Braß, W. Moser, and J. Pach. *Research Problems in Discrete Geometry*, Springer, New York, 2005.
- [3] P. Braß, M. Sharifi, A lower bound for Lebesgue’s universal cover problem, *International Journal of Computational Geometry and Applications*, 15(5), 2005, pp. 537–544.
- [4] G. Călinescu and A. Dumitrescu, The carpenter’s ruler folding problem, in J. Goodman, J. Pach and E. Welzl (editors), *Combinatorial and Computational Geometry*, Mathematical Science Research Institute Publications, Cambridge University Press, 2005, pp. 155–166.
- [5] H. G. Eggleston, Minimal universal covers in E^n , *Israel Journal of Mathematics*, 1(3), 1963, pp. 149–155.
- [6] J. Gerriets and G. Poole, Convex regions which cover arcs of constant length, *The American Mathematical Monthly*, 81, 1974, pp. 36–41.
- [7] H. C. Hansen, The worm problem (in Danish), *Normat*, 40, 1992, pp. 119–123.
- [8] J. E. Hopcroft, D. Joseph and S. Whitesides, Movement Problems for 2-Dimensional Linkages. *SIAM Journal on Computing*, 13(3), 1984, pp. 610–629.
- [9] O. Klein and T. Lenz, Carpenters rule packings—a lower bound, *Abstracts of 23rd European Workshop on Computational Geometry*, 2007, pp. 34–37.
- [10] R. Norwood and G. Poole, An Improved Upper Bound for Leo Moser’s Worm Problem. *Discrete and Computational Geometry*, 29(3), 2003, pp. 409–417.
- [11] R. Norwood, G. Poole and M. Laidacker, The worm problem of Leo Moser, *Discrete and Computational Geometry*, 7(1), 1992, pp. 153–162.
- [12] J. Pál, Über ein elementares Variationsproblem, *Math.-fys. Medd., Danske Vid. Selsk.*, 3(2), 1920, pp. 2–35.
- [13] J. Pál, Ein Minimumproblem für Ovale, *Math. Annalen*, 83, 1921, pp. 311–319.
- [14] G. Poole and J. Gerriets, Minimum covers for arcs of constant length. *Bulletin of the American Mathematical Society*, 79(2), 1973, pp. 462–463.
- [15] J.E. Wetzel, Sectorial Covers for Curves of Constant Length. *Canadian Math. Bull.*, 16, 1973, pp. 367–375.