

1 **Cutaneous Leishmaniasis in North Lebanon: re-emergence of an important**
2 **neglected tropical disease**

3

4 Dima El Safadi^a, Sabah Merhabi^a, Rayane Rafei^a, Hassan Mallat^a, Monzer Hamze^{a*}, Alvaro
5 Acosta-Serrano^b

6

7 ^aHealth and Environment Microbiology Laboratory, Doctoral School of Sciences and
8 Technology, Faculty of Public Health, Lebanese University, Tripoli, Lebanon; ^bDepartment of
9 Vector Biology, Liverpool School of Tropical Medicine, Liverpool L3 5QA, England, UK

10

11 *Corresponding author: Tel: +961(3)228881; Email: mhamze@monzerhamze.com

12

13 **Abstract**

14 **Background:** Cutaneous leishmaniasis (CL) is the most prevalent neglected tropical disease
15 among externally displaced people in the Middle East. In recent years, the Lebanese population
16 has increased >30% mainly due to a mass influx of Syrian migrants, thousands of them carrying
17 CL among other infectious diseases. Here, we revisited the current CL prevalence among
18 refugees in Northern Lebanon.

19 **Methods:** This cohort study was conducted at the Al-Bachaer Medical Center in North Lebanon
20 between 01/2017 and 06/2017. Randomly selected, forty eight suspected CL patients were
21 clinically diagnosed by dermatologists, and samples were obtained for microscopic examination
22 and molecular identification by PCR-RFLP. The treatment response to antimonials was assessed
23 at each week and was followed for up 6 months.

24 **Results:** *Leishmania tropica* was the most predominant species (91.7%) followed by *L. major*
25 (8.3%). Confirmed cases were treated with 1-2 courses of antimonials, and healing was achieved
26 mainly after receiving a second course of treatment. Importantly, we show evidence of possible
27 local CL transmission by indigenous sand flies in three separate patients who had no history of
28 recent travel to Syria.

29 **Conclusions:** This highlights the urgent necessity to implement preventive disease strategies to
30 avoid further dispersion of *L. tropica* CL in North Lebanon.

31 **Keywords**

32 Cutaneous leishmaniasis, Glucantime, Lebanon, *Leishmania tropica*, PCR-RFLP, Syrian
33 refugees.

34

35 **Introduction**

36 Cutaneous leishmaniasis (CL) is the most prevalent neglected tropical disease (NTD) in
37 the conflict zone in the Middle East, which currently affects more than hundred thousands of
38 refugees annually.^{1,2} Ongoing civil war in Syria resulted in mass migration of five million
39 Syrians to neighboring countries.³ Lebanon, situated to the west of Syria, has the highest per-
40 capita of refugees in the world (232 per 1,000 inhabitants) by hosting more than one million
41 Syrian refugees, an enormous stress on a historically fragile country with a population of just 4
42 million.⁴ As reported by Oxfam T5, the area of North Lebanon, including districts of Tripoli,
43 Zgharta, Becharreh, Minieh-Dinieh, El-Koura hosts 283,728 Syrian refugees that counted as one
44 third of the total T5 population with 833,728 inhabitants. While Akkar district hosts 95,403
45 Syrian refugees counted as 33% of the total population of Akkar numbering around 293,577
46 persons.⁵

47 Historically, Lebanon has had a very low annual prevalence of CL. Only 6 CL cases were
48 recorded in Lebanon during 2000-2012,⁶ in addition to few cases of visceral leishmaniasis in
49 rural areas located at the Lebanese-Syrian borders.⁷ Recently, outbreaks of CL among Syrian
50 refugees started to be reported by hospitals and primary health care centers.⁸ Furthermore, mass
51 migration of Syrians fled from Aleppo and Edleb were reported several thousand cases of CL.^{9,10}
52 Moreover, the phlebotomine sand fly vectors *Phlebotomus tobbi*, *Ph. syriacus*, *Ph. alexandri* and
53 *Ph. sergenti* have been identified in high-altitude mountains of Lebanon.¹¹ Thus, the presence of
54 local sand fly vectors and a continual influx of CL-infected refugees into other areas of Lebanon
55 further increase the risk of disease emergence.

56 The clinical manifestations of CL vary widely from spontaneous self-healing ulcer to
57 debilitating chronic or disfiguring lesions.¹² The drugs used to treat CL are expensive with
58 significant toxicity and side effects. Meglumine antimoniate (Glucantime) remains the first-line
59 treatment for CL in most countries, including Lebanon.¹³ In many cases, CL ulcers develop
60 secondary infections, which according to the World Health Organization (WHO) are first treated
61 with a combination of antibiotics and antifungal creams followed by 1-2 courses of intralesional
62 (or intramuscular) meglumine antimonite depending on the number, size and location of the
63 lesions.¹³ The WHO has offered more than 10,000 ampoules of Glucantime for the treatment of
64 CL in Lebanon.

65 In this study, we report on the epidemiology of CL among refugees in North Lebanon
66 between January to June 2017, and showed that most patients were infected with *L. tropica*.
67 Furthermore, we present evidence of potential autochthonous CL transmission in North Lebanon,
68 which highlights the need of establishing a coordinate disease control program to avoid a further
69 CL dispersion in the region.

70

71 **Methods**

72 **Inclusion of patients**

73 This cohort study was conducted at the Al-Bachaer Medical Center in North Lebanon
74 between 01/2017 and 06/2017. Forty-eight CL patients were clinically diagnosed by well-trained
75 dermatologist according to symptoms (presence of characteristic sore or ulcer lesions). Samples
76 were taken from 18 females and 30 males, ranging in age from 1 to 60 years, with a median age
77 of 10 years (interquartile range, 5-17 years). The majority of patients (79%) were younger than

78 18 years. Case record studies and information sheets were obtained for all patients and re-
79 labelled with the appropriate study code. The recorded patient information included age, sex,
80 region of origin, date of entrance to Lebanon, address and all relevant clinical data (i.e. lesion
81 size, number(s) and location(s) on the body, clinical features and treatment response). All data
82 were kept in secure computer folders.

83 **Sample collection**

84 Needle aspiration was performed in each patient for microscopy analysis after Giemsa
85 staining. Isohelix (SK-2S, UK) swab samples were taken by rubbing the ulcer approximately 20
86 times in a clockwise direction after removing the covered crust or scabs; the surrounded area was
87 then cleaned with 70% ethanol. All swabs were then processed for molecular diagnosis as
88 indicated below.

90 **DNA extraction and molecular identification**

91 Total genomic DNA was extracted from all lesion swabs using Qiagen DNeasy Blood &
92 Tissue Kit (Qiagen, USA) according to the manufacturer's instructions. *Leishmania* species were
93 identified by PCR-RFLP (Polymerase chain reaction- Restriction fragment length
94 polymorphism) analysis of the ribosomal Internal Transcribed Spacer 1 (*ITS1*), as described
95 previously.¹⁴ Primers used were: LITSR (5'- CTGGATCATTTTCCGATG -3') and L5.8 (5'-
96 TGATACCACTTATCGCACTT -3'). The PCR mixture contained 4.0 mM MgCl₂, 200 μM
97 dNTPs, 500 nM primers, 2 U Taq polymerase (KAPA Biosystem) and 1.5 mM MgCl₂. After 5
98 minutes denaturation at 94°C, PCR was performed using 35 cycles of 30 seconds at 94°C, 30
99 seconds at 53°C, and 60 seconds at 72°C. There was a final extension step of 72°C for 8 mins.¹⁴
100 For swab samples negative by *ITS1*-PCR, DNA was extracted from smears and amplified using

101 the same PCR protocol.^{14,15} Subsequently, 10 µL of PCR product was digested with 1 unit of
102 *Hae*III digest (New England Biolabs) for 2 hours at 37°C. Restriction fragments were analysed
103 on 2% (w/v) agarose gels using ethidium bromide. The leishmaniasis infections were identified
104 by clinical assessment and confirmed by laboratory analysis.

105

106 **Drug Treatment**

107 Clinically diagnosed CL-infected patients were treated with 0.5-5ml (85 mg Sb/ml) of
108 intralesional antimonials twice a week, for 3-4 weeks until healing was completed. Systemic
109 administration of antimonials (20 mg Sb/kg/day for 20 consecutive days) was also used in
110 patients presenting multiple (>5), large (>5 cm) or disfiguring lesions. Treatment failure was
111 defined by the lack of re-epithelialization after receiving one course (12 doses) within 45 days of
112 treatment starting.¹³

113

114 **Data and statistical analysis**

115 Statistical analyses were performed using the IBM SPSS Statistics v22.0 software.
116 Results were presented as the median with interquartile range for quantitative variables (age) and
117 number (percentages) for qualitative variables. The association of the variable swab sample
118 technique along with PCR, in infected patients with suppurative lesions and dry lesions, was
119 determined using the Fisher's exact test and differences were considered to be statistically
120 significant at $p < 0.05$.

121

122 **Results**

123 **Origin and distribution of CL patients**

124 Confirmed patients mainly originated from CL endemic and non-endemic areas of North
125 Syria. Most of them migrated from Edleb (44%; 21/48), followed by Hama (25%; 12/48),
126 Aleppo (17%; 8/48), Rakkah (6%; 3/48), Homos (6%; 3/48) and Tartous (2%; 1/48) (Figure 1).
127 By the time this study was conducted, these patients were residing in North Lebanon, especially
128 in Tripoli district (29%; 15/48) followed by Miniyeh-Dannieh (24%; 10/48) and Zgharta districts
129 (24%; 10/48). They are localized primarily in Badawi camps and Tebbeneh region, which are
130 one of the poorest areas of Northern Lebanon. The level of poverty in these regions is quite high,
131 with insufficient hygiene, inadequate sanitation and infrastructure.

132

133 **Clinical presentations of CL-infected patients**

134 Forty-three patients developed newly active CL lesion while five had chronic lesions.
135 The duration of the lesions prior to sampling ranged from 4 weeks to 48 months. Furthermore,
136 the majority of the ulcers were reported to have appeared between 1 week up to 3 years after
137 their displacement to Lebanon. A higher proportion of the patients have lesions located in the
138 facial area (67%; 32/48), followed by the arms (31%; 15/48), legs (14.5%; 7/48), trunk (4%;
139 2/48) and shoulder (4%; 2/48) (Figure 2). Moreover, the lesion size varied from 0.5 to 7 cm with
140 an average mean of 1.3 cm. Nodular lesions were the most predominant clinical presentation
141 (67%; 32/48) followed by papular (23%; 11/48) and plaque lesions (10%; 5/48), and suppuration
142 was predominately present in nodular (66%; 21/32) and plaque forms (40%; 2/5).

143 Interestingly, CL was also diagnosed in three patients that have settled in Lebanon for
144 more than 32 months and with no recent history of traveling or diagnosed with CL. These
145 individuals lived in very poor areas of Tripoli (Abou Samra-chok), Akkar and Kalamoun, which
146 are characterized for accumulation high amount rubbish and organic matter from animals (well-

147 known sand fly breeding sites) (Figures 1 and 3). All three patients were diagnosed by both
148 microscopy examination and PCR, and presented ulcerative nodules bigger than 2 cm in size for
149 at least 4 months.

150

151 **Microscopy and molecular identification**

152 After DNA extraction from Isohelix ulcer swab and smear aspiration, the 48 samples
153 from CL suspect individuals were positive for *Leishmania* spp. by PCR. *L. tropica* was the most
154 predominant species (44/48; 91.7%) followed by *L. major* (4/48; 8.3%). Patients infected by *L.*
155 *tropica* originated mainly from Edleb (21/44; 48%), Hama (12/44; 27%), Aleppo (7/44; 16%),
156 Homes (3/44; 7%), and Tartous (1/44; 2%). Whereas, Al Rakka (3/4; 75%) and Aleppo (1/4;
157 25%) were the original cities for patients infected by *L. major* (Figure 1).

158 By microscopic examination (ME), 39 samples were positive by identifying at least one
159 *Leishmania* amastigote in smears. The sensitivity of ME for *L. tropica* was 73% (32/44) and
160 100% (4/4) for *L. major*. The majority of suppurative lesions (83%; 19/23) were successfully
161 genotyped by PCR after DNA extraction of Isohelix swabs whereas only 48% (12/25) of dry
162 lesion swabs gave positive results. This indicates that swab sampling method along with PCR is
163 significantly more suitable for the molecular diagnosis of suppurative lesions than dry lesions
164 (Fisher-test=0.016).

165

166 **Treatment response**

167 Among the 48 confirmed CL patients, we followed the treatment response for 43 patients
168 who developed new active lesions. Results showed that only 20 patients (46%) healed after one
169 course of intralesional Glucantime (12 doses within 45 days), whereas, 18 (42%) and 3 (7%) of

170 the patients required a second and third course of Glucantime, respectively. In addition, two
171 patients (5%) receiving systemic drug treatment were cured after one course of treatment over a
172 period of 20 days.

173

174 **Discussion**

175 Dispersion of CL continues to be one of the most important health consequences of the
176 current Syrian conflict, particularly in externally displaced Syrian migrants. In this work, we
177 present an update of CL epidemiology in the northern districts of Lebanon, which are in
178 geographical proximity to CL-endemic Syrian regions such as Homos, Hama and Edleb and
179 where ca. 75% of patients included in our study originated from. The molecular analysis showed
180 that *L. tropica* is the main parasite species identified in the above cities in addition to Aleppo and
181 Tartous. Interestingly, most of the patients infected with *L. major* originated from Al-Raqqah,
182 which was the capital of the so-called Islamic State terrorist group over 3 years of war. This
183 region is not historically endemic by CL, but it became one of the most affected by this disease
184 during the Syrian conflict.¹ However, *L. major* is also known to be prevalent in Deir Al-Zour and
185 Al-Hasakeh cities,¹⁶ which are located approx. 150 km from Al-Raqqah. These results support a
186 previous report¹ suggesting that a massive internal human displacement combined with
187 ecological disruption of *Ph. papatasi* (*L. major* vector) habitats in neighbor regions may have
188 contributed to the marked increase of CL in Al-Raqqah region. In June 2017, a new outbreak was
189 also recorded in the western countryside of Dara'a governate, mostly in the town tell Shihab.
190 This city had almost no incidence of CL in the past, but the number of confirmed CL cases has
191 currently increased suggesting that sandflies vector might have been carried by families moving
192 from the infected areas.¹⁷ In parallel, the effect of Syrian civil war on the epidemiology of CL in

193 other neighboring countries such as Turkey was also demonstrated in many recent studies.^{18,19}
194 Turkey is a CL-endemic country and it currently hosts around ~3 million Syrian refugees in the
195 south/southeaster part of the country, leading to a large increase in the number of CL cases.¹⁸

196 Most of the CL-infected individuals included in our study were young children (67%)
197 presenting permanent scars, which are likely to blight their social interactions and incorporation
198 into the Lebanese society. This creates a terrible social stigma due to the double impediment to
199 their mental health of being both refugees and carrying CL.²⁰ In fact, the presence of CL scars
200 alone can cause depression, anxiety and an overall decrease in the quality of life of infected
201 individuals.²¹ These psychological and socio-economic impacts will be substantially more
202 noticeable in refugee settings due to the conditions of some of these places. Therefore, future
203 studies aiming to assess the mental health burden of the CL patients in refugee settings are
204 needed in order to formulate recommendations about how simple interventions to address mental
205 health problems can be introduced.

206 During the first four years of the Syrian conflict, Lebanon maintained open borders with
207 Syria. While there are no formal refugee camps for Syrians were established in Lebanon, roughly
208 18% of Syrians live in informal tented settlements around the country. The others are left to
209 dwell in villages and cities without any centralized or regional plan. They are localized in the
210 Northern districts, one of the most deprived regions with severe poverty levels and a high
211 number of refugee migrants. This implies that approximately half of the population of Lebanon's
212 Northern districts (T5) are refugees living below the poverty line.²² The huge influx of refugees
213 into these regions has put a strain on the scanty local resources. It has increased the depletion of
214 water, sanitation and hygiene, the lack of basic infrastructure and the accumulation of rubbish,
215 which collectively promote the perfect breeding sites for *Ph. sergenti* vector.⁵ Thus, the

216 combination of this suitable environment, the presence of sand fly vectors and refugee migrants
217 presenting either active or inactive (asymptomatic carriers) CL, exhibit a higher threat for the re-
218 emergence of this disease in some areas in Lebanon. Furthermore, in our study, a possible local
219 CL transmission by indigenous sandflies was observed in the three separate cases, although we
220 cannot rule out that these patients were asymptomatic carriers before they migrated to Lebanon.
221 As the last entomological data concerning sandflies in Lebanon is related to a phlebotomine sand
222 fly collection realized in 1995,¹¹ the characterization of medically important local sand fly
223 species, including bloodmeal preferences and vectorial capacity, is a high priority to update
224 information on the current distribution and dispersion of sand fly fauna in North Lebanon and to
225 prevent future local outbreaks. In addition, xenodiagnosis of wild-caught sand flies will reveal
226 the possible existence of autochthonous transmission and the necessity to implement a local CL
227 preventive control strategy. Although the number of identified local cases is relatively low in our
228 study, other similar cases could be present but either underestimated, misdiagnosed or
229 underreported. For instance, studies showed that CL incidence in Jordan may be underestimated
230 by 40- to 47- fold in national surveillance data.²³ No empirical assessments of CL underreporting
231 are available from the Lebanese official surveillance data.

232 Patients treated in Lebanon receive 1-2 courses of intralesional (or intramuscular in case
233 of multiple lesions), highly toxic and painful pentavalent antimonials according to current CL
234 treatment regimens. This creates a problem because, as our data show, most of the confirmed CL
235 cases were infected with *L. tropica*, which is naturally more resistant to Glucantime²⁴ and in fact
236 most of the cases needed a second (and sometimes a third) course of treatment to achieve clinical
237 cure (49%). The four cases infected by *L. major* were healed with a second course of anti-
238 leishmanial treatment. However, we cannot conclude about species treatment response due to the

239 low number of *L. major* isolates. The variations to antimonial treatment may either be influenced
240 by different factors such as parasite species, the presence of secondary infections within/around
241 the lesion and the spread of drug-resistant parasites.²⁴⁻²⁵

242 As for the prevalence of visceral leishmaniasis (VL) among Syrian migrants, some cases
243 were recorded at the coast Syrian borders.⁷ *Ph. syriacus* and domestic dogs (*Canis lupus*
244 *familiaris*) were identified as the main implicated VL vector and reservoir, respectively.²⁶
245 Unfortunately, the burden of Syrian crisis on VL and their reservoir in Lebanon has not been yet
246 determined. During 2014–2017, five Syrian refugee children died of VL caused by *L. infantum*
247 because of a late diagnosis and lack of awareness of this disease in Lebanon.²⁷ Recently, four
248 other cases of VL from Syrian refugees were diagnosed in the Lebanese hospitals and were
249 treated by Amphotericin B.²⁸ The risk of introduction of *L. major* and/or *L. tropica* besides *L.*
250 *infantum* is highly regarded with the presence of their correspondent reservoirs and vectors. Cats
251 (*Felis catus*) were identified as susceptible reservoir for *L. tropica* and *L. major* in MENA
252 region.²⁹ On the other hand, humans are considered the main reservoir for the anthroponotic
253 transmission of *L. tropica*,²⁹ and *Ph. tobbi*, *Ph. sergenti*, *Ph. pappatasi*, *Ph. syriacus* and *Ph.*
254 *alexandri* sand fly species have been already identified in the Lebanese mountains.¹¹
255 Collectively, this suggests that Lebanon currently presents all the conditions for the
256 autochthonous transmission of both forms of Leishmaniasis if not preventive disease control
257 measures are put in place by local health authorities.

258 While the conflict in Syria approaches its ninth year, the military conflict has stopped in
259 some cities that pushed thousands of refugees to return amid fresh hope for last ceasefires.
260 According to the UNHCR, more than 440,000 internally displaced Syrian and about 22,000 of
261 whose fled abroad had come back to cities, which are partly or wholly controlled by the Syrian

262 government, including Damascus, Aleppo, Hama and Homos.³⁰ This recent situation is highly
263 dangerous because of the risks of returning to a destroyed and unsafe country lacking
264 international support. The deterioration of basic infrastructure, buildings and hospitals, the lack
265 of access to the primary health care and the increasing rate of many infectious diseases such as
266 leishmaniasis will introduce a high risk of morbidity for persons coming back. Sandflies vector
267 will be at high density in these adequate regions for multiplication and without control measures.
268 As a result, people can contract leishmaniasis and then become reservoirs that could contribute
269 with disease spreading on a large scale, especially with the non-sustainability of this return
270 without the social determinants of global health such as employment opportunities, adequate
271 food and water and health care infrastructure.

272 In Summary, *L. tropica* is the most predominant species among refugees in North
273 Lebanon. The majority of patients needed at least two courses of glucantime to achieved cured.
274 The identification of a possible local transmission in Lebanon highlights the urgent necessity to
275 perform entomological studies to characterize the local sand fly vectors. This will help develop
276 targeted vector control strategies in CL-endemic areas to specifically prevent local *L. tropica*
277 transmission.

278

279

280 **Authors' contributions:**

281 DES, MH and AAS designed the study. DS, SM, RR, HM performed the study fieldwork. DS, SM, RR,
282 HM carried out the molecular identification and analysed the data. DS and SM wrote the first draft of the
283 manuscript. DS, AAS, MM, RR and HM critically revised the manuscript for intellectual content. All
284 authors read and approved the final manuscript. MM and AAS are the guarantors of the paper.

285 **Acknowledgements**

286 The authors would like to thank all those who participated in the study, the staff and
287 dermatologists of Al-Bachaer Medical Center in Tripoli for their assistance in sample collection
288 and processing. We would like to thank Waleed Al-Salem (Saudi Ministry of Health) for
289 scientific advice and kind help and Yasser Alraey (Liverpool School of Tropical Medicine) for
290 PCR advice.

291

292 **Funding**

293 This work was supported by the Lebanese University and the Shefa Fund [to AAS].

294 **Competing interests:**

295 None declared.

296 **Ethical considerations**

297 Written informed consent was obtained from all the adults who participated in the study. Consent
298 for inclusion of young children was obtained from parents or guardians. The study was reviewed
299 and approved by the Azm Center for Research in Biotechnology and its application Institutional
300 Review Board (Ethic number: CE-EDST-02-2017) and patient data used in this study was
301 anonymized.

302

303 **References**

- 304 1. Al-Salem WS, Pigott DM, Subramaniam K et al. Cutaneous Leishmaniasis and Conflict in
305 Syria. *Emerg Infect Dis* 2016;22(5):931-3.
- 306 2. Du R, Hotez PJ, Al-Salem WS et al. Old World Cutaneous Leishmaniasis and Refugee
307 Crises in the Middle East and North Africa. *PLoS Negl Trop Dis* 2016;10(5):e0004545.
- 308 3. United Nations High Commissioner for Refugees. Situation Syria Regional Refugee
309 Response. 2018. <https://data2.unhcr.org/en/situations/syria> [accessed 2 July 2018].
- 310 4. Danish Refugee Council. Closing Borders, Shifting Routes: Summary of Regional
311 Migration Trends Middle East – May, 2016. [https://reliefweb.int/report/world/closing-](https://reliefweb.int/report/world/closing-borders-shifting-routes-summary-regional-migration-trends-middle-east-may-2016)
312 [borders-shifting-routes-summary-regional-migration-trends-middle-east-may-2016](https://reliefweb.int/report/world/closing-borders-shifting-routes-summary-regional-migration-trends-middle-east-may-2016)
313 [accessed 2 August 2018].
- 314 5. Oxfam. Local governance under pressure - Social Stability in T5 area, North Lebanon,
315 2016. <https://data2.unhcr.org/en/documents/details/43854> [3 August accessed 2018].
- 316 6. Ministry of Public Health. Epidemiologic Surveillance Department.
317 <http://www.moph.gov.lb> [accessed 14 October 2018].
- 318 7. Desjeux P. Information on the epidemiology and control of the leishmaniasis by country or
319 territory. Geneva : World Health Organization; 1991 Report No.: WHO/LEISH/91.30.
320 <http://apps.who.int/iris/handle/10665/62132> [accessed 10 July 2018].
- 321 8. Saroufim M, Charafeddine K, Issa G et al. Ongoing epidemic of cutaneous leishmaniasis
322 among Syrian refugees, Lebanon. *Emerg Infect Dis* 2014;20(10):1712-5.
- 323 9. Hayani K, Dandashli A, Weisshaar E. Cutaneous leishmaniasis in Syria: clinical features,
324 current status and the effects of war. *Acta Derm Venereol* 2015;95(1):62-6.

- 325 10. Haddad N, Saliba H, Altawil A et al. Cutaneous leishmaniasis in the central provinces of
326 Hama and Edlib in Syria: Vector identification and parasite typing. *Parasit Vectors*
327 2015;8:524.
- 328 11. Haddad N, Leger N, Sadek R. Sandflies of Lebanon: faunistic inventory. *Parasite*
329 2003;10:99-110.
- 330 12. de Vries HJ, Reedijk SH, Schallig HD. Cutaneous leishmaniasis: recent developments in
331 diagnosis and management. *Am J Clin Dermatol*. 2015;16(2):99-109.
- 332 13. World Health Organization, Manual for case management of cutaneous leishmaniasis in the
333 WHO Eastern Mediterranean Region. 2014.
334 http://applications.emro.who.int/dsaf/EMROPUB_2013_EN_1590.pdf [accessed 10 October
335 2018].
- 336 14. el Tai NO, Osman OF, el Fari M et al. Genetic heterogeneity of ribosomal internal
337 transcribed spacer in clinical samples of *Leishmania donovani* spotted on filter paper as
338 revealed by single-strand conformation polymorphisms and sequencing. *Trans R Soc Trop*
339 *Med Hyg*. 2000;94(5):575-9.
- 340 15. Ertabaklar H, Ertug S, Caliskan SO et al. Determination of *Leishmania* species by PCR-
341 RFLP in the smear samples taken from the lesions of cutaneous leishmaniasis cases.
342 *Mikrobiyol Bul* 2016;50(2):300-6.
- 343 16. Rioux JA, Dereure J, Khiami A et al. Ecoepidemiology of leishmaniasis in Syria.
344 *Leishmania major* Yakimoff and Schokhor (Kinetoplastida-Trypanosomatidae) infestation
345 of *Psammomys obesus* Cretzschmar (Rodentia-Gerbillidae). *Ann Parasitol Hum Comp*
346 1990;65(5-6):203-7.

- 347 17. Leishmaniasis, cutaneous - Syria (12): (Dara'a) increasing incidence. ProMED. 2017
348 October 29. <http://www.promedmail.org/post/20171029.5410639>. [Accessed 2 July 2018].
- 349 18. Inci R, Ozturk P, Mulayim MK et al. Effect of the Syrian Civil War on Prevalence of
350 Cutaneous Leishmaniasis in Southeastern Anatolia, Turkey. Med Sci Monit 2015;21:2100-
351 4.
- 352 19. Ozkeklikci A, Karakus M, Ozbel Y et al. The new situation of cutaneous leishmaniasis after
353 Syrian civil war in Gaziantep city, Southeastern region of Turkey. Acta Trop 2017;166:35-
354 8.
- 355 20. Hotez P, Ottesen E, Fenwick A et al. The neglected tropical diseases: the ancient afflictions
356 of stigma and poverty and the prospects for their control and elimination. Adv Exp Med
357 Biol 2006;582:23-33.
- 358 21. Yanik M, Gurel MS, Simsek Z et al. The psychological impact of cutaneous leishmaniasis.
359 Clin Exp Dermatol 2004;29(5):464-7.
- 360 22. United Nations Office for the Coordination of Humanitarian Affairs. North & Akkar
361 Governorates, Lebanon, 2016. [https://reliefweb.int/report/lebanon/lebanon-north-akkar-
362 governorates-profile-october-2018](https://reliefweb.int/report/lebanon/lebanon-north-akkar-governorates-profile-october-2018) [accessed 3 July 2018].
- 363 23. Mosleh IM, Geith E, Natsheh L et al. Cutaneous leishmaniasis in the Jordanian side of the
364 Jordan Valley: severe under-reporting and consequences on public health management.
365 Trop Med Int Health 2008;13(6):855-60.
- 366 24. Gourbal B, Sonuc N, Bhattacharjee H et al. Drug uptake and modulation of drug resistance
367 in *Leishmania* by an aquaglyceroporin. J Biol Chem 2004;279(30):31010-7.

368 25. Gimblet C, Meisel JS, Loesche MA et al. Cutaneous Leishmaniasis Induces a Transmissible
369 Dysbiotic Skin Microbiota that Promotes Skin Inflammation. *Cell Host Microbe*
370 2017;22(1):13-24 e4.

371 26. Nuwayri-Salti N, Baydoun E, el-Tawk R et al. The epidemiology of leishmaniasis in
372 Lebanon. *Trans R Soc Trop Med Hyg.* 2000;94:164–6.

373 27. El Hajj Rana , El Hajj Hiba , Ibrahim K. Fatal Visceral Leishmaniasis Caused by
374 *Leishmania infantum*, Lebanon. *Emerg Infect Dis.* 2018;24(5).

375 28. Gemayel G, Khalife I, Minari A et al. Visceral leishmaniasis in Lebanon a report of four
376 cases: case series and review. *The Journal of Infection in Developing Countries*
377 2018;12(02.1):31S.

378 29. Moriconi M, Rugna G, Calzolari M et al. Phlebotomine sand fly-borne pathogens in the
379 Mediterranean Basin: Human leishmaniasis and phlebovirus infections. *PLoS Negl Trop Dis*
380 2017;11(8):e0005660.

381 30. Hall S. Syria’s spontaneous returns - Syrian Arab Republic. ReliefWeb.
382 <https://reliefweb.int/report/syrian-arab-republic/syria-s-spontaneous-returns> [accessed 2 July
383 2018]

384

385

386


387

388

389

390

391



392

393

394 **Figure 1. Geographic distribution of Cutaneous Leishmaniasis (CL) cases identified among**
395 **Syrian refugees in Lebanon (Small box) and their relative region of origin (Round dots).**


396 The majority of patients are coming from the northern Syrian regions especially Edleb (44%) and
397 Hama (25%). Infected Syrian refugees are migrated mainly to Tripoli (31%). *L. tropica* (Red
398 dots) is the main genotype identified among CL cases followed by *L. major* (Yellow dots). The
399 green strikes indicated the location of the three probable local transmission CL cases in North
400 Lebanon.

401

402

403

404



405

406 **Figure 2. Patterns of cutaneous leishmaniasis among Syrian refugees in Lebanon, 2017. A**
407 **and B, Disfiguring facial lesions. C, Satellite lesions on the outer edge of CL primary lesion.**

408

409

410



411

412 **Figure 3. Unsanitary living conditions where CL patients live. Accumulated rubbish and**
413 **presence of animals create perfect sand fly habitats and promote local disease transmission.**