

2001

# The Genus *Striga* (Scrophulariaceae) in Africa

Kamal I. Mohamed

Lytton John Musselman  
*Old Dominion University*, lmusselm@odu.edu

Charles R. Riches

Follow this and additional works at: [https://digitalcommons.odu.edu/biology\\_fac\\_pubs](https://digitalcommons.odu.edu/biology_fac_pubs)



Part of the [Botany Commons](#), and the [Plant Biology Commons](#)

---

## Repository Citation

Mohamed, Kamal I.; Musselman, Lytton John; and Riches, Charles R., "The Genus *Striga* (Scrophulariaceae) in Africa" (2001). *Biological Sciences Faculty Publications*. 189.  
[https://digitalcommons.odu.edu/biology\\_fac\\_pubs/189](https://digitalcommons.odu.edu/biology_fac_pubs/189)

## Original Publication Citation

Mohamed, K. I., Musselman, L. J., & Riches, C. R. (2001). The genus *Striga* (scrophulariaceae) in Africa. *Annals of the Missouri Botanical Garden*, 88(1), 60-103. doi:10.2307/2666132

---

# THE GENUS *STRIGA* (SCROPHULARIACEAE) IN AFRICA<sup>1</sup>

---

Kamal I. Mohamed,<sup>2</sup> Lytton John  
Musselman,<sup>3</sup> and Charles R. Riches<sup>4</sup>

## ABSTRACT

Twenty-eight species and six subspecies of the hemiparasitic genus *Striga* Lour., witchweed, occur in Africa. Twenty-two species are endemic. Witchweeds occur throughout the continent with greatest diversity in the grasslands and savannas north of the equator. *Striga gesnerioides* (Willd.) Vatke and *S. hermonthica* (Del.) Benth. have developed host-specific strains that may have distinct morphotypes. We accord these no formal taxonomic status. *Striga hermonthica* and *S. asiatica* (L.) Kuntze, limited to agroecosystems, may be recently derived from non-ruderal species. *Striga hirsuta* Benth. and *S. lutea* Lour., sometimes treated as conspecific with *S. asiatica*, are recognized as distinct species restricted to natural ecosystems. *Striga linearifolia* Hepper, with its strongly bilabiate corolla and perennial habit, is considered as a subspecies of *S. bilabiata* (Thunb.) Kuntze. *Striga baumannii* Engl. and *S. fulgens* Hepper are excluded from the genus. *Striga baumannii* has tuberous roots and a ligneous calyx and may represent a distinct genus. *Striga fulgens*, with free petals, is aligned with *Chascanum* E. Mey. in the Verbenaceae.

**Key words:** Africa, Scrophulariaceae, *Striga*, witchweed.

---

Globally, witchweeds have a greater impact on human welfare than any other parasitic angiosperms because their hosts are subsistence crops in areas marginal for agriculture. They are also among the most specialized of all root-parasitic Scrophulariaceae (or Orobanchaceae, depending on how the families are circumscribed). Some members of the Scrophulariaceae are holoparasitic (without chlorophyll); most are hemiparasites (with chlorophyll). *Striga* combines life styles of both with a holoparasitic seedling and a green, chlorophyll-containing emergent plant. These furtive pathogens attack their hosts underground and, by the time the flowering stem of the parasite emerges and is evident, the crop is damaged. This occult behavior may be the source of the Latin name, "Striga," meaning "hag" or "witch." In this way, hosts are "bewitched" because the farmer is unaware of the parasite until it comes up. Due to the serious injury they inflict, witchweeds have been the targets of many control strategies. Helpful reviews of control measures and other aspects of

witchweed biology are found in Sand et al. (1990), Sauerborn (1991), Parker and Riches (1993), and Press and Graves (1995).

Because witchweeds are parasitic and not totally dependent upon leaves, no exclusively vegetative stems are produced. In other words, each stem bears an inflorescence. Flowers are usually showy, in some cases spectacular. Most species that have been studied are autogamous (e.g., Musselman & Hepper, 1986), although *S. hermonthica* and *S. aspera* (Willd.) Benth. are outcrossers (Aigbokhan et al., 1998). Seed production is prodigious. Conditioning, germination, parasitic contact, and penetration are mediated by elegant systems of chemical communication between host and parasite (Parker & Riches, 1993; Press & Graves, 1995; Maass, 1999).

Witchweeds tolerate a relatively wide range of climatic and soil conditions. They grow in areas with annual rainfall ranging from 25 to 150 cm per year with a decrease in the severity of infestation in areas with higher rainfall (Musselman & Ayensu,

---

<sup>1</sup> This work is based mainly on the dissertation research of K. I. Mohamed that was supported, in part, by grant USDA 58-319R-5-0006. Research conducted in Nigeria was supported by grant AID DHR 5600-G-00-1021-00. Many people have assisted in our studies of the genus over the past several years. We thank E. Aigbokhan, A. E. Awad, F. F. Bebwai, D. Berner, R. Eplee, H. Glen, F. N. Hepper, C. Jarvis, D. Kenfack, M. O. Khidir, R. Moberg, D. L. Nickrent, R. Norris, S. N. C. Okonkwo, C. Parker, A. Pieterse, K. V. Ramaiah, J. Ransom, A. Raynal-Roques, S. B. Safa (deceased), M. J. Vasudeva Rao, P. J. Terry, J. Thieret, M. Thulin, J. H. Visser (deceased), E. P. Weber, and A. Wolfe. Appreciation is extended to the curators and staff of BRE, BM, GH, K, MO, ODU, S, US, and UPS for the loan of specimens.

<sup>2</sup> Department of Biology, State University of New York College at Oswego, Oswego, New York 13126-3501, U.S.A.  
mohamed@oswego.edu.

<sup>3</sup> Mary Payne Hogan Professor of Botany, Old Dominion University, Norfolk, Virginia 23529-0266, U.S.A.  
lmusselman@odu.edu.

<sup>4</sup> Natural Resources Institute, University of Greenwich, Central Avenue, Chatham Maritime, Chatham, Kent ME4 4TB, U.K. charlie.riches@bbsrc.ac.uk.

1984). Optimum temperature for seed germination is between 30°C and 40°C, with no germination at 15°C or lower, or higher than 45°C (see review of literature in Parker & Riches, 1993).

Phenology among the different species of *Striga* varies. In a given area, there is a progression of appearances from the onset of the rains to the end of the rainy season. For example, in the Southern Kordofan region of Sudan, the first *Striga* to appear is *S. forbesii* Benth., followed by *S. passargei* Engl. → *S. aspera* → *S. gesnerioides* → *S. hirsuta* → and lastly, *S. hermonthica*.

In general, low soil fertility, nitrogen deficiency, well-drained soils, and water stress accentuate the severity of *Striga* damage to the host. These are typically the environmental conditions for *Striga*-hosts in the semiarid tropics (Sauerborn, 1991; Parker & Riches, 1993). *Striga* usually causes less damage to hosts growing on fertile, deep soils.

*Striga* is distinguished from other genera in the subfamily Rhinanthoideae of the Scrophulariaceae by bilabiate corollas with a pronounced bend in the corolla tube, and unilocular anthers. The corolla feature distinguishes *Striga* from related *Buchnera* L. in which nine *Striga* species were once included. Based on this feature, the genus *Striga* was erected by the Italian botanist Loureiro in 1790 as distinct from *Buchnera*, which has a salverform corolla. Little data are available on seed structure in *Buchnera*, but Musselman and Mann (1976) reported secondary ornamentation of *Buchnera americana* L. similar to that of *Striga asiatica*. *Buchnera* germination and seedling development are also much less specialized (Parker & Riches, 1993). Lastly, *Buchnera* is found in both the New and Old Worlds, while *Striga* is strictly Old World.

Recent molecular work supports the long-assumed relationship between *Striga* and *Buchnera*. Olmstead et al. (in press), using three plastid genes, show *Buchnera floridana* Gandoger (= *B. americana*) and *S. asiatica* as sister taxa. Further work is needed to determine the relationship between *Alectra* and *Striga*. Unlike *Buchnera*, *Alectra* species require a germination stimulant and have a holoparasitic seedling stage.

All witchweeds are African except the following: *S. curviflora* Benth. (Australia, New Guinea), *S. densiflora* (Benth.) Benth. (India), *S. masuria* (Buch.-Ham. ex Benth.) Benth. (= *S. flava* Miq.?, *S. esquirolia* Lév.?) (China, India, Indonesia), *S. multiflora* Benth. (Australia), *S. parviflora* Benth. (Australia), and *S. sulphurea* Dalz. ex Dalz. & Gibbs (India). Our understanding of the genus in Africa requires a better knowledge of these species.

Taxonomic studies at the genus and tribal level

are few. Wettstein (1891) divided *Striga* into two sections: the *Pentapleurae*, which have a 5-ribbed calyx, and the *Polypleurae*, characterized by approximately 10-calyx ribs. Later, Engler (1897) created the section *Tetrosepalum* to include *S. baumannii*. Saldanha (1963) noted that the number of ribs is inconsistent within species. In spite of this, rib number has remained as the only infrageneric character.

Figures 1 and 2 illustrate the general features of witchweeds.

Following is a summary of features helpful in determining *Striga* species.

1. *Duration.* *Striga* species are annuals or perennials. Duration may help identify plants in the field, but it is of limited value for herbarium specimens, especially when roots are lacking. All agriculturally important species are annuals, perhaps because their hosts are also annuals. The exception is sugarcane (*Saccharum officinarum* L.), which is perennial. In natural grasslands witchweed species do not seem to affect their hosts the same as cultivated crops; damage is not evident. Perennial species produce fleshy underground rhizomes or tubers that bear scale leaves from which new stems arise annually. Generally, perennial species have reduced or narrow stem leaves. See Raynal-Roques (1993) for a helpful discussion of duration in witchweeds.

2. *Stem shape.* Three stem types can be distinguished. Terete stems are round in cross section, obtusely square stems are square with blunt corners, and winged stems are square and acutely angled. A furrowed stem has grooves parallel to the long axis.

3. *Indumentum.* Indumentum types, both surface features and trichomes, are diverse. Many species have scabrous leaf and stem surfaces at maturity. The most common trichome types include: glandular trichomes; stiff hairs (hispid); long soft hairs (pubescent or pilose); long stiff hairs (ciliate) (Neumann et al., 1996).

4. *Leaf lobing, dentation.* Leaves of most species are unlobed, and only a few have teeth. Venation may extend to the tips of leaves in a few species.

5. *Inflorescence.* Inflorescence types include solitary flowers, spikes, and racemes. The length of the inflorescence compared to the vegetative part of the stem (longer or shorter); compactness of the flowers (lax or dense); and bract size and shape (similar to or different from stem leaves) are also helpful distinguishing features. Bracteoles are present just above the floral bract. Flowers can be opposite or alternate on the inflorescence. In almost




Figure 1. *Striga angolensis* K. I. Mohamed & L. J. Musselman. —A, B. Habit. Note the reduction of leaves acropetally and basipetally. —C. Calyx. Two bracteoles are evident, with the much larger bract on the left. This calyx has four lobes. —D, E. Corolla. —F. Seeds. —G. Capsule. —H. Pistil. (R. G. N. Young 1365, MO). Drawing by Karillos Karillos in Mohamed and Musselman, Brittonia 49: 118–121. 1997.




Figure 2. *Striga hirsuta* (Benth.) Benth. Note bract and bracteole, lower left. Musselman & Mansfield 5531 (ODU). Drawing by Daniel L. Nickrent from Musselman et al., Sida 8: 196–201. 1979. (Originally determined as *S. asiatica* (L.) Kuntze.) Scale = 1 cm.

all species the 2 to 4 pairs of lower bracts at the base of the inflorescence are longer than the upper bracts and are described separately.

6. *Calyx*. Calyx lobes can be equal or subequal. In some species the adaxial lobe is reduced.

7. *Corolla color and tube bend*. Most witchweed corollas are red or white, but they can also be salmon, orange, or yellow. The bend in the corolla tube can be positioned distally or proximally. If proximal, it may be within the calyx teeth. If distal, it is closer to the corolla limb. Fragrance has been reported only in *S. hermonthica* (Musselman & Hepper, 1986).

8. *Hosts*. Witchweeds have one of the narrower host ranges of any parasitic Scrophulariaceae. Not surprisingly, hosts of agronomically important species are better documented than hosts of native species. Most *Striga* species, however, have not been critically examined for host range and host specificity. Except for *S. gesnerioides*, which is restricted to dicots, these hosts are all Poaceae with the odd connection to non-grasses noted, disproportionately, by collectors. Great caution is advised when ascribing any form of host specificity to root parasites. It is essential that the host-parasite connection be located, not that hosts be inferred from proximity. Further, it is virtually impossible to determine how many different plants, perhaps different species, are attacked by a single witchweed.

Too few studies on other features of witchweeds

are available, including cytology. Reported chromosome numbers have varied widely, from  $n = 18$  to  $n = 40$  (reviewed in Aigbokhan et al., 1998). Careful cytological studies are needed to clarify the counts. Ultrastructural studies are also limited. Pollen has few useful features, at least in the species studied (DeLeonardis, 1986; Minkin & Eshbaugh, 1989). Current taxonomy is based almost exclusively on morphology.

Seed size and structure have been used to separate *S. angustifolia* from other species because of its larger seed size and its germination without a host stimulus (e.g., Parker & Riches, 1993). Secondary ornamentation of *S. angustifolia* is also distinct from other species (Musselman & Parker, 1981). However, in a survey of nine species, no other consistent seed ornamentation features were found to separate species (Musselman & Parker, 1981).

A conspectus of the genus on a continent-wide basis is important for agricultural researchers, among others, to enable the determination of new parasite problems. Certainly, future expansion of food crop species possibly susceptible to parasitism by witchweed should take into account indigenous *Striga* species that could potentially attack crops. There are several examples where this has occurred. These include maize planted in the KweKwe region of Zimbabwe (Knepper et al., 1991) where *S. forbesii* occurred. Another is the planting of sugarcane in Ethiopia, which was attacked by *S. latericea* Vatke (Parker, 1988). Moreover, meaningful quarantine measures depend on knowledge of the distribution of the parasite.

A study of *Striga* taxonomy invites comparison with the taxonomy of the other, agronomically significant genus in the family (assuming parasitic Scrophulariaceae and Orobanchaceae are monophyletic, e.g., Nickrent et al., 1998; Young et al., 1999). This is *Orobanche*, the genus of broomrapes.

Historically, broomrapes have suffered from extreme splitting and the establishment of varieties, forms, and even subforms based largely on herbarium studies with little consideration of host-induced variation. Thus, a plethora of names encumbers the taxonomy of the group. Unlike *Orobanche*, the cytology of witchweeds needs much additional work. Based on the limited data available, aneuploidy does not play a major role in witchweed speciation and variability. Some groups of broomrapes, on the other hand, exhibit aneuploidy, polyploidy, and parthenogenesis (Cubero, 1996). In short, the taxonomy of witchweeds is less daunting than that of broomrapes.

Despite the well-known impact of *Striga* on sub-

sistence crops in Africa, there is no single reference to the taxonomy and distribution of African taxa. The purpose of this paper is to describe all the reported taxa in Africa, map their distribution, record notes on their biology, and provide a key for their identification.

Maps of *Striga* collection sites were constructed using the *Britannica Atlas*. Collection sites were plotted on a large computer-generated map (Anonymous, 1996).

**Striga** Lour., Fl. Cochinchinensis 1: 22. 1790.

TYPE: *Striga lutea* Lour.

Annual or perennial, herbaceous hemiparasites. Stem: stiffly erect, square or terete, ridged in some species. Indumentum: hirsute or pilose, scabrid, glandular pubescent; rarely glabrate. Leaves: opposite or subopposite, rarely alternate, reduced to small scales at the base of the stem in all species. Inflorescence: dense or lax spike or raceme; rarely solitary. Calyx: tubular, with 4 to 8, usually 5, equal or subequal lobes. Corolla: divided into a tube and a limb, the tube cylindrical with an acute bend, the limb 2-lipped, with lobes fused at least at their bases to form a bilabiate corolla with 3 lower and 2 upper lips. Stamens: 4, didynamous, anthers dorsifix, unilocular, dehiscence loculicidal. Pollen: with 3–4 colpi, 15–30 µm with 2 nuclei at shedding. Style: terete, elongate, and persistent. Capsule: oblong or subovoid, placentation axillary, dehiscence loculicidal. Seeds [except in *S. angustifolia* (Don) Saldanha]: dust-like, 0.35 × 0.20 mm, with prominent ridges that are ornamented. Germination: hypogeal and cryptocotylar [except in *S. angustifolia*]. Chromosome numbers ranging from  $n = 18$  to  $n = 40$ . Hosts: usually Poaceae; few species restricted to dicots.

#### KEY TO THE AFRICAN *STRIGA* SPECIES

- |   |    |   |                              |
|---|----|---|------------------------------|
| 1. Calyx 4- to 5-ribbed (sect. <i>Pentapleurae</i> ) .....  | 2  | tic, shortly petiolate; corolla bright salmon-pink, tube 22–25 mm long .....  | 15. <i>S. hallaei</i>        |
| 1'. Calyx 10- or more ribbed (sect. <i>Polypleurae</i> ) .....  | 14 | Calyx lobes about as long as tube; leaf lanceolate, sessile; corolla mauve, tube 12–13 mm long .....  | 2. <i>S. angolensis</i>      |
| 2(1). Lobes of lower lip fused to >50% of their length forming a strongly bilabiate corolla .....   | 3  | Plant succulent, usually branched from base (tufted); leaf reduced, ≤10 mm long, appressed .....  | 7                            |
| 2'. Lobes of lower lip fused to <50% of their length forming a weakly bilabiate corolla .....   | 4  | Plant not succulent, unbranched or if branched then from middle; leaf large, > 10 mm long, ascending or spreading .....                       | 9                            |
| 3(2). Plant glabrate with few hairs on leaf margins and bracts; calyx ≤4 mm long .....  | 1  | Corolla densely pubescent, bent proximally within calyx teeth; lower corolla lobes lanceolate .....   | 21. <i>S. lepidagathidis</i> |
| 3'. Plant pubescent or hispid; calyx > 4 mm long .....  | 6  | Corolla sparsely pubescent or glandular-pubescent, bent distally above calyx; lower corolla lobes obovate .....                               | 8                            |
| 4(2). Leaf prominently 3-veined, leaf margin toothed .....  | 5  | Plant glabrate, about 6 cm tall, perennial; corolla tube up to 8 mm long .....  | 12. <i>S. gastonis</i>       |
| 4'. Leaf veins obscure, not apparent, leaf margin entire .....  | 6  | Plant sparsely to densely hispid or pubescent, > 10 cm tall, annual (except strain on <i>Euphorbia</i> ); corolla tube 8–14 mm long .....     | 13. <i>S. gesnerioides</i>   |
| 5(4). Calyx lobes deltate, < ½ the tube length, adaxial lobe reduced to < 1 mm .....  | 9  | Calyx lobes linear to lanceolate, > ½ the tube length, adaxial lobes > 1 mm .....   | 10                           |
| 5'. Calyx lobes linear to lanceolate, > ½ the tube length, adaxial lobes > 1 mm .....   | 12 | Herbage with dense retrorse hispid hairs; corolla retrorsely pubescent; leaf 1–2 mm wide .....  | 10                           |
| 6(4). Plant branched; corolla > 10 mm long, lobes of lower lip 7–15 mm long, 3–6 mm wide; calyx tube 5–10 mm long .....                             | 11 | Herbage with dense ascending hispid hairs; corolla with ascending hairs; leaf > 2 mm wide .....   | 11                           |
| 6(5). Plant usually unbranched; corolla < 10 mm long, lobes of lower lip 1–3 mm long, 1–1.5 mm wide; calyx tube 3–4 mm long .....                   | 11 | Plant branched; corolla > 10 mm long, lobes of lower lip 7–15 mm long, 3–6 mm wide; calyx tube 5–10 mm long .....                             | 16. <i>S. hermonthica</i>    |
| 7(6). Corolla white (rarely pink or yellow); upper bracts as long as or longer than calyx; flowers sessile, spike longer than vegetative stem ..... | 12 | Plant usually unbranched; corolla < 10 mm long, lobes of lower lip 1–3 mm long, 1–1.5 mm wide; calyx tube 3–4 mm long .....                   | 14. <i>S. gracillima</i>     |
| 7'. Corolla pinkish purple; upper bracts shorter than calyx; flowers shortly pedicellate, rame short than vegetative stem .....                     | 13 | Corolla white (rarely pink or yellow); upper bracts as long as or longer than calyx; flowers sessile, spike longer than vegetative stem ..... | 24. <i>S. passargei</i>      |
| 8(6). Calyx tube 4–6 mm long; lower corolla lobes 4–8 mm long, 2–5 mm wide .....  | 5  | Corolla pinkish purple; upper bracts shorter than calyx; flowers shortly pedicellate, rame short than vegetative stem .....                   | 13                           |
| 8'. Calyx tube 1.5–3 mm long; lower corolla lobes 2–3 mm long, 1–2 mm wide .....  | 7  | Calyx tube 4–6 mm long; lower corolla lobes 4–8 mm long, 2–5 mm wide .....  | 5. <i>S. aspera</i>          |
| 9(6). Leaf pinnatifid; flowers solitary, few, axillary .....  | 15 | Calyx tube 1.5–3 mm long; lower corolla lobes 2–3 mm long, 1–2 mm wide .....  | 7. <i>S. brachycalyx</i>     |
| 9'. Leaf not pinnatifid; flowers in terminal rame or spike .....  | 15 | Leaf pinnatifid; flowers solitary, few, axillary .....  | 25. <i>S. pinnatifida</i>    |
| 10(14). Leaf > 3 mm wide, with 2 to 5 prominent veins, margin toothed .....   | 16 | Leaf not pinnatifid; flowers in terminal rame or spike .....  | 15                           |
| 10(15). Leaf < 3 mm wide, veins obscure, margin entire .....  | 20 | Leaf > 3 mm wide, with 2 to 5 prominent veins, margin toothed .....   | 16                           |
| 11(16). Corolla salmon-pink; lower bracts > 2 cm long; inflorescence lax at base .....  | 17 | Leaf < 3 mm wide, veins obscure, margin entire .....  | 20                           |
| 11(17). Corolla pink or white; lower bracts < 2 cm long; inflorescence compact .....  | 18 | Corolla salmon-pink; lower bracts > 2 cm long; inflorescence lax at base .....  | 17                           |
| 11(18). Stem hairs stiff; lower bracts distinctly 3-veined, toothed; lower corolla lobes 6–9 mm long .....  | 11 | Corolla pink or white; lower bracts < 2 cm long; inflorescence compact .....  | 18                           |
| 11(19). Stem hairs stiff; lower bracts distinctly 3-veined, toothed; lower corolla lobes 6–9 mm long .....  | 11 | Stem hairs stiff; lower bracts distinctly 3-veined, toothed; lower corolla lobes 6–9 mm long .....  | 11. <i>S. forbesii</i>       |

- 17'. Stem hairs soft; lower bract veins obscure, entire; lower corolla lobes 10–15 mm long ..... 20. *S. latericea*
- 18(16). Leaf linear, base of hairs inconspicuous; bracts linear ..... 9. *S. dalzielii*
- 18'. Leaf elliptic or narrowly elliptic, base of hairs enlarged; bracts lanceolate ..... 19
- 19(18). Corolla pink, tube 10–13 mm long; lower corolla lobes 4–7 mm long, 2–4 mm wide; leaf shorter than internode ..... 19. *S. klingii*
- 19'. Corolla white, tube 17–24 mm long; lower corolla lobes 7–10 mm long, 4–8 mm wide; leaf longer than internode ..... 23. *S. macrantha*
- 20(15). Corolla brilliant scarlet with yellow throat; flowers opposite in dense raceme ..... 10. *S. elegans*
- 20'. Corolla not brilliant scarlet; flowers alternate in lax raceme or spike ..... 21
- 21(20). Corolla white or creamy white ..... 22
- 21'. Corolla pink, purple, yellow, or dark red ..... 23
- 22(21). Calyx 8–12 mm long; corolla tube 10–15 mm long, lower lobes 4–8 mm long, 2–4 mm wide ..... 3. *S. angustifolia*
- 22'. Calyx 13–21 mm long; corolla tube 27–35 mm long, lower lobes 14–20 mm long, 10–17 mm wide ..... 27. *S. pubiflora*
- 23(21). Calyx with at least 15 ribs; plant perennial ..... 24
- 23'. Calyx with 10 ribs; plant annual ..... 25
- 24(23). Corolla purple; leaf linear,  $\geq 1$  cm long; calyx 11–15 mm long, lobes 5, equal ..... 18. *S. junodii*
- 24'. Corolla yellow; leaf lanceolate,  $< 1$  cm long; calyx 8–9 mm long, lobes 5, unequal ..... 26. *S. primuloides*
- 25(23). Leaf  $\geq 15$  mm long, longer than internode; lower bracts  $> 15$  mm long ..... 4. *S. asiatica*
- 25'. Leaf  $< 15$  mm long, shorter than internodes; lower bracts  $< 15$  mm long ..... 26
- 26(25). Corolla tube 20–25 mm long; calyx 8–15 mm long ..... 8. *S. chrysanthra*
- 26'. Corolla tube  $< 15$  mm long; calyx up to 6 mm long ..... 27
- 27(26). Plant 6 cm tall (rarely 20); lower surface of leaf and bract with a single row of hispid hairs along margin and midrib; calyx lobes about equaling tube length ..... 17. *S. hirsuta*
- 27'. Plant up to 40 cm tall; lower surface of leaf and bract with two rows of hispid hairs along margin and midrib; calyx lobes  $\frac{1}{2}$  the tube length ..... 22. *S. lutea*

**1. *Striga aequinoctialis* A. Chev. ex Hutch. & Dalz., Expl. Bot. Afr. Oc. Fr. 1: 476. 1920.**  
TYPE: Guinea (Conakry). Montagne de Nzo, Chevalier 21035 (holotype, P!).

Perennial, to 50 cm tall, slender, glabrate, usually simple or with 2 or 3 branches from below middle, stem winged. Leaves 10–24  $\times$  1(–2) mm, filiform, sessile, decussate, ascending, shorter than internodes, hispid along margins, margin entire, veins obscure. Bracts 4–6(–12)  $\times$  1(–2) mm, linear, hispid along margins, incurved, longer than calyx. Flowers decussate in open spikes, spike shorter

than vegetative stem. Calyx 5-ribbed, 3–4 mm long; tube 2–2.5 mm long; lobes 5, unequal, lanceolate, 1–2 mm long, slightly shorter than tube. Corolla pale mauve, tube 10–12 mm long, bent and expanded distally above calyx, retrorsely pubescent; lobes of lower lip 3–7  $\times$  2–3 mm, ovate, tip acute, upper lip 3–5  $\times$  4–7 mm, slightly notched, tip obtuse.

Restricted to the moist montane grasslands of west tropical Africa between 5°16'N and 11°0'N (Fig. 3). Known from Guinea, Sierra Leone, Liberia, and Ivory Coast with only one collection from outside West Africa, Young 1300 (MO), from Angola. Like *S. bilabiata*, *S. aequinoctialis* has a bilabiate corolla, decussate leaves, slender stem, retrorsely pubescent corolla and shares part of the geographical range. *Striga aequinoctialis* is distinguished by its glabrous stem, filiform leaves, and small calyces. The calyx is the smallest in the genus.

*Selected specimens examined.* ANGOLA. Lunda South: Saurimo-Dala Road, 11°2'S, 20°12'E, Young 1300 (MO). GUINEA. Eastern: Kankan, 45 mi. NW of Beyla, 9°11'N, 9°9'W, Collenette 3 (K). Southern: N'Zerekore, 7°43'N, 8°50'W, Adam 5111, 24653, 24386, 26054, 26531 (all MO). Western: Fouta Djalon, near Bandakure, SW of Pita, 11°0'N, 12°16'W, Adames 284 (K). IVORY COAST. Northwestern: Odienne, Tienko, 10°5'N, 7°30'W, Morton & Gledhill 18771 (K). Southwestern: Danane, Nimba Mountains, 7°34'N, 8°24'W, Boughey 18052, 18067b, 180995 (all K). LIBERIA. Nimba: Mountain Nimba, 7°34'N, 8°40'W, Adam 21457 (K, MO, UPS), Vallah 65 (K); Grassfield-Nimba-Ideria, 6°45'N, 8°30'W, Adam 25550 (K, MO). Sinoe: Jedepo, 5°16'N, 8°20'W, Adam 24039, 27478 (MO). Liberia: Adam 25592 (K). SIERRA LEONE. Bintinani, Morton & Gledhill 1093 (K). Eastern: Sankanbiaiwa, 8°45'N, 11°0'W, Cole 199 (K). Northern: Bumbuna to Farangbaia, 9°1'N, 11°44'W, Deighton 5146 (K); Loma, 9°15'N, 11°2'W, Anonymous 9783 (K). Western: Freetown, Mendi (Njamumu), 8°29'N, 13°0'W, Burbridge 510 (K).

**2. *Striga angolensis* Mohamed & Musselman, Brittonia 49: 118–121. 1997.** TYPE: Angola. Vila Luzo, R. Luena, Young 1365 (holotype, MO!). Figure 1.

Perennial, to 70 cm tall, stiffly erect, simple or with 2 or 3 branches from below middle, densely glandular-pubescent, stem obtusely square. Leaves (10–)20–35  $\times$  2–8 mm, lanceolate, sessile, alternate, coarsely toothed, 3-veined, longer than internodes at the lower part of stem, becoming shorter above. Bracts 5–8  $\times$  2 mm, abruptly different from leaves, lanceolate, shorter than calyx. Flowers opposite, lax at the base of inflorescence, dense toward the apex, spike shorter than vegetative stem. Calyx 4- or 5-ribbed, 6–8 mm long; tube 2.5–4 mm long; lobes 4 or 5, subequal, lanceolate, 3–5 mm




Figure 3. Distribution based on specimens of *Striga aequinoctialis* (circle), *S. angolensis* (triangle), and *S. angustifolia* (diamond).

long, as long as tube. Corolla weakly bilabiate, mauve, tube 12–13 mm long, bent and expanded distally above calyx, densely glandular-pubescent; lobes of lower lip 7–8 × 2 mm, obovate, upper lip 5 × 3 mm, obovate, slightly notched.

*Striga angolensis* is restricted to boggy grassland and wet areas in Angola between 9°48'S and 12°50'S (Fig. 3). Because there are numerous collections, we assume that *S. angolensis* was fairly common in this region, but we are not aware of any recent collections.

While many witchweeds are plants of dry areas, *S. angolensis* is part of a guild of witchweeds that inhabit wetlands, including *S. latericea*, *S. forbesii*, and *S. junodii* Schinz.

A statistical analysis of the genus (Mohamed, 1994) suggested that *Striga angolensis* is closely related to *S. hallaei* A. Raynal, *S. dalzielii* Hutch., *S. klingii* (Engl.) Skan, and *S. macrantha* (Benth.) Benth. They can be distinguished, however, on leaf and bract characters. *Striga angolensis* and *S. hallaei* are the only two *Pentapleurae* with prominently veined leaves, a feature common in at least six species in section *Polypleurae*. *Striga dalzielii*, *S. macrantha*, and *S. klingii* (sect. *Polypleurae*) differ from *S. angolensis* in having dense leafy spikes and

leaf-like bracts that surmount the calyx, at least at the base of the inflorescence.

*Selected specimens examined.* ANGOLA. Benguella: Benguella, 12°36'S, 13°20'E, Gossweiler 3504 (K); Catumbela, 12°50'S, 14°40'E, Faulkner A376 (K). Lunda South: small river between River Luachi & River Lumeji, 10°10'S, 20°12'E, Young 1261 (ODU). Malange: Gando, 9°48'S, 16°10'E, Faulkner A376 (K). Moxico: boggy grassland by River Chibamba, 12°40'S, 20°45'E, Milne-Redhead 4161 (K); River Luena, Vila Luzo, 11°45'S, 19°50'E, Exell & Mendonca 1353 (BM), Young 1347 (BM), Young 1365 (MO).

### 3. *Striga angustifolia* (Don) Saldanha, Bull. Bot. Surv. India 5: 70. 1963. *Buchnera angustifolia* Don, Prodr. Fl. Nepal: 91. 1825. TYPE: Nepal. Wallich s.n. (holotype, K!).

Annual, 10–50 cm tall, stiffly erect, unbranched or with 2 or 3 branches from below middle, pubescent or sometimes densely hispid with short upward curved hairs, stem obtusely square. Leaves 10–20(–30) × 1–3 mm, linear, alternate, as long as internodes, entire, veins obscure. Lower bracts 10–25 × 1–2 mm, leaf-like, longer than calyx, upper bracts subulate, shorter than calyx. Flowers alternate in lax racemes, raceme shorter than vegetative stem. Calyx 15-ribbed; ribs prominent, pilose, 8–

12 mm long; tube 3–6 mm long; lobes 5, equal, linear to lanceolate, 3–6 mm long, equal or subequal to the tube. Corolla creamy white with greenish tube, tube 10–15 mm long, bent and expanded distally above calyx, densely pubescent; lobes of lower lip 4–8 × 2–4 mm, obovate, upper lip 3–4 × 3–4 mm, broadly obovate, emarginate or truncate.

In Africa, *S. angustifolia* grows in the seasonally wet grasslands of Tanzania, Malawi, Mozambique, Zambia, and Zimbabwe between 9°52'S and 22°05'S (Fig. 3). It does not attack crops in eastern Africa, although it has been cited as a pest of cereals and sugarcane in India (Ramaiah et al., 1983). The occurrence of *S. angustifolia* in native grasslands and its absence in agroecosystems suggest it is indigenous to East Africa. *Striga angustifolia* is sympatric with and apparently related to *S. pubiflora* Klotzsch. and *S. junodii* (Mohamed, 1994). The Indian species *S. masuria* superficially resembles *S. angustifolia*, but the relationship between these two species has not been investigated. *Striga angustifolia* is widespread in India and Sri Lanka, and is known from a single collection in Oman. See Hepper (1986) regarding typification.

*Selected specimens examined.* MALAWI. Central: 6 km N of Kasungu, 12°55'S, 33°29'E, Grosvenor & Renz 1038 (K); Kasungu Hills, 13°0'S, 33°15'E, Jackson 2282 (K); Kasungu National Park, 13°35'S, 33°32'E, Pawek 12015A (K); near Tamanda Mission, Robson 1094 (BM, K). Northern: Nyika Plateau, 10°30'S, 34°0'E, McCounie 178 (K). Southern: Mount Mlanje above Likhubula, 16°5'S, 35°40'E, Hepper 7368 (K); Saturi Stream, Likhubula Valley, Mulanj Mountain, 16°15'S, 35°27'E, Chapman & E. G. 7186, 9000 (K); Zomba Mountain, Chivunde Valley, 15°30'S, 35°19'E, La Croix 2735 (MO); Zomba, 15°30'S, 35°19'E, Hepper 7333 (K). MOZAMBIQUE. Inhambane: Mabote, 22°5'S, 34°5'E, de Koning 7703 (K). Niassa: Marrupa, 13°8'S, 37°30'E, Nuvunga 471 (K). TANZANIA. Ruvuma: just E Songea District Boundary, 10°42'S, 35°37'E, Milne-Redhead & Taylor 10577 (K); Songea, 3 km W of Gumbiro, 10°15'S, 35°42'E, Milne-Redhead & Taylor 10024 (K); S of Hanga River, 64 km from Songea on Njombe Road, also collected 120 km from Songea on Lindi road, Songea District, 9°52'S, 35°22'E, Gilbert 17889 (K). ZAMBIA. Southern: 17 mi. NE of Choma, Brachystegia Wood, 16°40'S, 27°18'E, Hutchison & Gilbert 3537 (K); Choma, 16°55'S, 26°59'E, Robinson 1214 (K). ZIMBABWE. Manicaland: Inyanga, Cheshire, 18°15'S, 32°48'E, Norlindh & Weimark 4878 (K); Inyanga, Ghikori Reserve, 18°0'S, 32°30'E, Masierson 217 (K); Umtali, Zimunya Reserve, 18°50'S, 32°37'E, Chase 6854 (K). Mashonaland North: River Ruya, 16°50'S, 31°20'E, Philcox & Drummond 9054 (K). Mashonaland South: Corononzi, Kopje Domboshawa, 17°40'S, 31°34'E, Philcox et al. 9093 (K); Goromonzi, Kopje Ngomakurira, 17°58'S, 31°22'E, Philcox et al. 9127 (K); Lower Sabi, Videw, 19°20'S, 31°25'E, Videw 8110 (K); Macheke, 18°0'S, 31°47'E, Eyles 8799

(K); Marandellas, Charter Estate, 18°15'S, 31°20'E, Cleghorn 2357 (K).

4. *Striga asiatica* (L.) Kuntze, Rev. Gen. Pl. 2: 466. 1891. *Buchnera asiatica* L., Sp. Pl.: 630. 1753. TYPE: Comoro Islands. St. Joanna, To-reen s.n. (LINN, photograph!).

Plant annual, up to 40 cm tall, stiffly erect, usually branched from middle, scabrous, hispid, stem obtusely square. Leaves (10–)20–50 × 1–3 mm, linear or narrowly elliptic, opposite, longer than internodes, margin entire, veins obscure. Lower bracts 15–35(–50) × 1–3 mm, leaf-like, upper bracts lanceolate, all bracts longer than calyx. Flowers alternate in lax terminal racemes, raceme longer than vegetative stem. Calyx 10-ribbed, 7–9 mm long; tube 5–7 mm long; lobes 5, equal, or 6 to 8 unequal, narrowly lanceolate, < ½ tube length. Corolla red with yellow throat, occasionally yellow, tube 11–14 mm long, bent and expanded distally above calyx, sparsely to densely pubescent; lobes of lower lip 3–5 × 2–3 mm, obovate, spreading, upper lip 3 × 3–4(–7) mm, emarginate, wider than long.

The red-flowered witchweed, which severely damages sorghum [*Sorghum bicolor* (L.) Moench.], maize (*Zea mays* L.), and other crops, is *S. asiatica*.

Hepper (1974) retained the name *S. asiatica* for the well-branched plant with dark-colored flowers over *S. lutea*, the name used for many years for the species of agronomic importance (e.g., in the United States, Sand et al., 1990). *Striga lutea* (see discussion below) and *S. hirsuta* may also have red corollas.

*Striga lutea* and *S. hirsuta* can also attack crops, but with no discernible damage. Because of confusion among these plants, reports in the literature of a red-flowered witchweed damaging a crop need to be verified.

Some workers, e.g., Musselman and Hepper (1986), have lumped *S. asiatica*, *S. lutea*, and *S. hirsuta* as a single species, *S. asiatica*. However, Mohamed (1994) reported consistent morphological, ecological, and phenological differences among these and concluded that three taxa should indeed be recognized.

The geographical range of *S. asiatica* includes southern and central Africa, where it is common on light sandy soils. It has also been collected from Abu Zaabal, Nile Delta, Egypt (Fig. 4), no doubt through contaminated grain seed, which is a major source of dispersal (Berner et al., 1992). In Madagascar and southern Africa, especially Botswana, *S. asiatica* does most damage to crops like sor-




Figure 4. Distribution based on specimens of *Striga asiatica* (circle) and *S. aspera* (triangle).

ghum, maize, and millet [*Pennisetum americanum* (L.) Schumann] (Ralston et al., 1987).

*Striga asiatica* is not recorded as a pest in Africa outside this region except in Kenya (Musselman, unpublished) and its introduction in the Republic of Togo (Agbobli, 1991). Recently, rice in Madagascar has been attacked by *S. asiatica* (Geiger et al., 1996).

The center of origin of *S. asiatica* is most likely African (Raynal-Roques, 1991); it probably evolved in southern Africa, Madagascar, or the Mascarenes. This assumption is supported by the presence of the very similar and putatively related (Mohamed, 1994) *S. elegans* Benth. only in southern and eastern Africa. If correct, then the *S. asiatica* population introduced to the Carolinas in the United States came from southern Africa. Allozyme studies indicate that the introduction occurred once from an autogamous population (Werth et al., 1984).

*Selected specimens examined.* ANGOLA. Cuanza North: Cassoalala, 9°25'S, 14°20'E, Anonymous 8347 (K). Huambo: 12°44'S, 15°5'E, Wellman 1769 (K). Loanda: Zenzado, 8°48'S, 13°14'E, Gossweiler 4466 (K). BOTSWANA. Central Mahalatswe Lands 200 km N of Gabarone, Musselman 7003 (ODU). CENTRAL: Swooning, near Serowe, 22°28'S, 26°47'E, Mitchison 1975 (K). Chobe: Formabachena, near Pandamatenga, 18°9'S, 24°24'E, Musselman et al. 9/4/86 (ODU). Ngamiland: Okavango, Moremi Wildlife, 19°10'S, 23°16'E, Smith 845 (K). SOUTHERN: Lobatsi, Magabane, 25°12'S, 25°38'E, De Bell 878 (K). BURUNDI. EASTERN: Astrida (Rubona Ineac), 2°33'S, 29°45'E, Michel 5367 (K). CONGO. KINSHASA: Kinshasa (Leopoldville), 4°20'S, 15°15'E, Baldwin s.n. (K). DEMOCRATIC REPUBLIC OF CONGO. Bas-Zaire: Boma, 5°48'S, 13°0'E, Soyaux 189 (K). Kasai Oriental: Gandajika, 6°45'S, 23°57'E, Sieber 2401 (GH). EGYPT. Delta: Abu Zaabal, 30°15'N, 31°21'E, Elhadi & Khattab 90/188 (S). KENYA. COAST: Kilifi, 3°35'S, 39°55'E, Moggridge 548 (K). EASTERN: Jakake Forest, Jeffery K242 (K). NAIROBI: Nairobi, 1°17'S, 36°48'E, Napier 51 (K). NORTH EASTERN: Chom Airship, 18 mi. NW Kiunga, 1°35'S, 41°22'E, Gillespie 186 (K). RIFT VALLEY: Isiolo to Mathews

Range,  $0^{\circ}24'N$ ,  $37^{\circ}33'E$ , and Mountain Nyiru,  $2^{\circ}10'N$ ,  $36^{\circ}50'E$ , *Nesbould* 1958 (K). LESOTHO. **Western:** Mafetieng,  $29^{\circ}50'S$ ,  $27^{\circ}10'E$ , *F.B.* 6431 (K). MADAGASCAR. **Southern:** Katoala, S of Ambovombe,  $25^{\circ}8'S$ ,  $46^{\circ}5'E$ , *Decary* 9014 (US). MALAWI. **Central:** Bunda Agriculture College Farm, 23 km S of Lilongwe,  $14^{\circ}8'S$ ,  $33^{\circ}47'E$ , *Brummit* 8661 (K). **Northern:** Rumpi, 1 mi. up Shelinda River,  $10^{\circ}55'S$ ,  $33^{\circ}53'E$ , *Pauw* 12566 (K). **Southern:** Lower Shire Valley,  $15^{\circ}45'S$ ,  $35^{\circ}0'E$ , *Lawrence* 15 (K). MOZAMBIQUE. **Cabo Delgado:** 15 km on Mocimboa-Mueda Road,  $10^{\circ}40'S$ ,  $40^{\circ}35'E$ , *Jasen* 8158 (K). **Gaza:** 3 km from Chibuto,  $24^{\circ}39'S$ ,  $33^{\circ}30'E$ , *De Lemes & Balsinhas* 92 (K). **Inhambane:** Govuro, 7 km from Banamana,  $24^{\circ}2'S$ ,  $34^{\circ}54'E$ , *Correia & Marques* 4142 (K). **Manica:** Odzi River Valley,  $18^{\circ}59'S$ ,  $32^{\circ}37'E$ , *Jeagu* 84 (K). **Maputo:** Namaacha,  $25^{\circ}59'S$ ,  $32^{\circ}05'E$ , *Sousa* 555 (K). **Niassa:** E Coast of Lake Nyasa, *Johnson* 101 (K). **Zambezian:** Lugela, Hamagru,  $16^{\circ}25'S$ ,  $36^{\circ}43'E$ , *Faulkner* 90 (K). NAMIBIA. **Owambo:** Olukonda  $18^{\circ}0'S$ ,  $16^{\circ}3'E$ , *Reed* 3/1888 (K). SOUTH AFRICA. **Natal:** Durban North, Prospect,  $29^{\circ}55'S$ ,  $30^{\circ}56'E$ , *Forbes* 1/1929 (K). **Nxikazi:** Kangwane Homeland, *Musselman* 7022 (ODU). **Orange Free State:** Kroonstad,  $27^{\circ}46'S$ ,  $27^{\circ}12'E$ , *Rogers* 96 (US). **Transki:** Crigualand East, Clydis-dall,  $30^{\circ}40'S$ ,  $29^{\circ}0'E$ , *Tyson* 1222 (GH). **Transvaal:** near Pretoria,  $25^{\circ}40'S$ ,  $28^{\circ}10'E$ , *Gkvile* 41 (K). SUDAN. **Al-jazirah:** Gir, *Anonymous* 2159 (K). **Bahr Al-Gabal:** Addai,  $9^{\circ}50'N$ ,  $24^{\circ}0'E$ , *Anonymous* 2180 (K). **Northern:** Ghattas, *Schweinfurth* 2285 (K). **Jungali:** Bor, Zandi Land, Li Atafu,  $6^{\circ}12'N$ ,  $31^{\circ}20'E$ , *Wyld* 225 (K). TANZANIA. **Arusha:** Endabash, Lake Manyara National Park,  $3^{\circ}40'S$ ,  $35^{\circ}50'E$ , *Greenway & Polhill* 11305 (K). **Dar es Salaam:** Dar es Salaam,  $6^{\circ}47'S$ ,  $39^{\circ}18'E$ , *Kuntze* 7/iv94 (US). **Iringa:** Ruaha National Park, at road to Idodi,  $7^{\circ}40'S$ ,  $35^{\circ}25'E$ , *Bjornstad* 1294 (K). **Lindi:** Nachingwea,  $10^{\circ}28'S$ ,  $38^{\circ}49'E$ , *Anderson* 846 (K). **Mafia Island:** Kikuni, *Greenway* 5097 (K). **Mara:** Kampi Ya Mawe, 25 mi. from Sronera Musoma,  $1^{\circ}30'S$ ,  $34^{\circ}6'E$ , *Greenway* 10519 (K). **Morogoro:** Nguru Mountains,  $6^{\circ}3'S$ ,  $37^{\circ}26'E$ , *Thulin & Mhoro* 3140 (K). **Mwanza:** Nyahanga,  $2^{\circ}20'S$ ,  $33^{\circ}3'E$ , *Tanner* 810 (K). **Pwani:** 23 km SE of Bagamoyo,  $6^{\circ}34'S$ ,  $39^{\circ}1'E$ , *Thulin & Mhoro* 488 (K). **Rukwa:** Tumba,  $7^{\circ}30'S$ ,  $31^{\circ}45'E$ , *Bullock* 3770 (K). **Ruvuma:** Tunduru, 60 mi. from Masasi,  $11^{\circ}7'S$ ,  $37^{\circ}2'E$ , *Richards* 17980 (K). **Shinyanga:** Shinyanga,  $3^{\circ}30'S$ ,  $33^{\circ}30'E$ , *Choner* 2005 (K). **Singinda:** Manyoni, S of Itigi Station,  $5^{\circ}37'S$ ,  $34^{\circ}27'E$ , *Greenway & Polhill* 11736 (K). **Tanga:** Dahil-Madanga,  $4^{\circ}45'S$ ,  $39^{\circ}E$ , *Tanner* 3018 (K). **Zanzibar:** Chaani, and Pemba, *Greenway* 1204 (K). UGANDA. **Eastern:** Amuria,  $2^{\circ}4'N$ ,  $33^{\circ}37'E$ , *Williams* 32 (K). ZAMBIA. **Central:** near Kapushi River, Lukanga Plains,  $14^{\circ}30'S$ ,  $27^{\circ}40'E$ , *Troprell* 2050 (K). **Eastern:** Luangwa Valley, Tunwe Munkanya,  $14^{\circ}2'S$ ,  $30^{\circ}30'E$ , *Phiri* 79 (K). **Northern:** Chisinoima Falls, Luombi River,  $10^{\circ}6'S$ ,  $31^{\circ}0'E$ , *Richards* 5230 (K). ZIMBABWE. **Mashonaland North:** Midlands, near Makava,  $17^{\circ}17'S$ ,  $32^{\circ}42'E$ , *Musselman* 4/17/86 (ODU). **Matabeleland North:** Bulawayo, Waterford,  $20^{\circ}7'S$ ,  $28^{\circ}32'E$ , *Norrgrann* 459 (K). **Midlands:** Gwelo, Msasa-Mnondo, Senka Area,  $19^{\circ}0'S$ ,  $29^{\circ}46'E$ , *Biegel* 1912 (K).

**5. *Striga aspera* (Willd.) Benth., Companion Bot. Mag. 1: 362. 1835. *Euphrasia aspera* Willd., Syst. 2: 776. 1801. TYPE: Ghana. *Isert* s.n. (holotype, BM!; isotype, C not seen).**

*Striga aspera* var. *schweinfurthii* Skan, Fl. Trop. Afr. 4(2): 403. 1906. TYPE: Sudan. *Schweinfurth* 1992 (holotype, K!).

*Striga aspera* var. *filiformis* Benth., in DC. Prodr. 10: 501. 1846. TYPE: Niger. *Vogel* 15 (holotype, K!).

Annual, to 53 cm tall, slender, unbranched or branched from the middle, densely hispid, stem obtusely square. Leaves (10–)20–35(–60) × 1(–2) mm, opposite, linear, ascending or spreading, longer than internodes, margin entire, veins obscure. Lower bracts 8–30(–40) mm, leaf-like, longer than calyx, upper bracts lanceolate, shorter than calyx. Flowers alternate in open racemes, raceme shorter than vegetative stem. Calyx 5-ribbed, 5–10 mm long; tube 4–6 mm long; lobes 5, almost equal, linear, 3–5 mm long, as long as tube. Corolla weakly bilabiate, pinkish purple (rarely white), tube 12–16 mm long, bent and expanded distally above calyx, sparsely glandular-pubescent; lobes of lower lip 4–8 × 2–5 mm, obovate, upper lip 3–7 × 4–7 mm, emarginate or subtruncate, as long as wide.

Superficially, *Striga hermonthica* and *S. aspera* resemble one another. There are three characters to distinguish between them: corolla tube bend, pubescence, and size of calyx lobes. In the field, *S. aspera* can be differentiated from *S. hermonthica* by its slender stem and the pinkish purple corolla tube bent distally above the calyx teeth. The tube of *S. hermonthica*, on the other hand, bends in the middle, that is, proximally within the calyx teeth. Unfortunately, there are plants with intermediate characters. *Striga aspera* is sparsely glandular-pubescent, while *S. hermonthica* lacks a glandular-pubescent corolla tube. The glandular pubescence of the corolla and the short calyx lobes are the most consistent characters separating these two species. Calyx lobes in *S. aspera* are almost equal to the tube length with the adaxial lobe more than 1 mm long, whereas the lobes of *S. hermonthica* are short, triangular. The adaxial lobe is less than 1 mm long in *S. hermonthica*.

Leaves of *S. hermonthica* are more than 2 mm wide, whereas leaves of *S. aspera* are rarely 2 mm wide. This and the smaller stems make the overall aspect of *S. aspera* more delicate.

*Striga aspera* and *S. hermonthica* interbreed; both are outbreeders. Aigbokhan et al. (1998, 2000) demonstrated a high rate of interfertility between the two species. Morphological differences between the two species, compared to within-species variation, and the low germination percentage of their hybrids' seeds (Aigbokhan et al., 1998), support the current taxonomic treatment of recognizing two species. Chromosome numbers are also

different, with *S. aspera* N = 18 and *S. hermonthica* N = 19 (Aigbokhan et al., 1998).

*Striga aspera*, in its more delicate forms, may resemble *S. brachycalyx* Skan and *S. passargei*, and the three species are sympatric in most of their range.

*Striga aspera* is in the Sahelian and Sudanian domains of Africa (Fig. 4), from Senegal to Sudan between 4°43'N and 15°18'N (Fig. 4). Together with *S. gesnerioides* and *S. hermonthica* it has the most northern distribution of all species. Unlike *S. hermonthica*, however, the range of *S. aspera* does not extend south of the equator. *Striga aspera* has been reported to attack maize but not sorghum in Ethiopia (Parker, 1988) and fonio [*Digitaria exilis* (Kippist) Stapf], maize, rice (*Oryza sativa* L. and other species), and sugarcane in West Africa (Parker & Riches, 1993). The extension of agriculture into natural grasslands may expose crops to this species, especially in West Africa where it is common.

Kenfack et al. (1996) have shown that *S. aspera* in northern Cameroon has the broadest host range (Poaceae and Cyperaceae, 18 genera, 24 species) of the eight species of witchweeds they studied.

*Selected specimens examined.* BURKINA FASO. **Center East:** 5 km N of Tenkodogo, 11°50'N, 0°19'W, *Safa & Musselman* 10/25/87 (ODU). **Eastern:** Fada N'Gourma, 12°12'N, 0°25'E, *Raynal* 22127 (K). **Haut Bassins:** Banfora, 10°50'N, 4°50'W, *Raynal* 22394, 22395 (K). **Northern:** 5 km E of Godi, 11°55'N, 2°24'E, *Safa & Musselman* 10/12/87 (ODU). **Volta:** Boromo, 11°50'N, 2°55'W, *Raynal* 22309 (K). CAMEROON. **Northern:** Champs/Piedmont, Maroua, 10°35'N, 14°22'E, *Meurillon* 1363 (K). CENTRAL AFRICAN REPUBLIC. **Vakaga:** Dongolo Plains, 9°33'N, 21°24'E, *Fay* 2794 (K). ETHIOPIA. **Sidamo:** Upper Birr State Farm, N of Bore, 4°43'N, 37°40'E, *Parker* 4065 (K). GAMBIA. **Central:** Denfi, Terry 3173 (K). **Eastern:** Basse, 13°25'N, 14°28'W, Terry 1942 (K). GHANA. **Eastern:** 2/3 of way between Adidame and Ho, 6°40'N, 0°30'E, *Morton s.n.* (K). **Northern:** just N of Buige Bridge on Kintampo to Tamale Road, 8°25'N, 1°30'W, *Hossain & Agyakwah* 37826 (K). **Upper:** Bolgatanga, 10°40'N, 0°50'W, *Hossain & Agyakwah* 37829 (K, US). GUINEA. **Eastern:** Kankan, 10°30'N, 9°20'W, *Pobeguin* 1020 (K). IVORY COAST. **Central:** CIDT Office at Kouto, 45 km N of Bouandougou, 9°0'N, 6°32'W, *Riches* 230, 233 (K). MALI. **Koulakoro:** 11 km N of Didienni, 13°55'N, 8°25'W, *Musselman & Mohamed* 10/6/88 (ODU). **Segou:** Famabougou, 17 km from Sokolo, 14°39'N, 6°15'W, *Raynal* 22581 (K). NIGER. Boundom to Chetimari, *Saadou* 1839 (K). **Dosso:** Dongondoutchi, 13°50'N, 4°8'E, *Parker* 2314 (ODU). **Maradi:** Dan Issa Road, 3 mi. from Maradi, 13°30'N, 7°7'E, *Hall* 452 (K). NIGERIA. **Bauchi:** Hassan Estate, Luna Road, 10°10'N, 9°20'E, *Knepper* 9/26/89 (ODU). **Kano:** 7 km S of Katsina on Kano Road (163 km N Kano), 12°57'N, 7°40'E, *Parker* 2046 (K). **Plateau:** Jos Plateau, Naroguta, 9°50'N, 4°35'E, *Lawlor & Hall* n 46508 (K). **Sokoto:** 10 mi. N of Bussa, 50 mi. S of Yelwa, 10°15'N, 4°35'E, *Cook* 358 (K). SENEGAL. **Oriental:** 41 km NE of Velingara,

13°26'N, 13°55'W, *Musselman* 7085 (K, ODU). SUDAN. **Bahr Al-Ghazal:** on River Sue, between Peili and Wau, 7°10'N, 28°0'E, *Meyers* 7329 (GH, K). **Jungali:** 12 km E of Nyany, 80 km N of Bor, 6°52'N, 31°25'E, *Lock* 81/132 (K). **South Kurdufan:** Dilling, Nuba Mountains, 12°5'N, 29°41'E, *Andrews* 6 (K). **Upper Nile:** Djur, Jeriba Ghafas, 8°39'N, 28°35'E, *Schweinfurt* 1992 (S).

## 6. *Striga bilabiata* (Thunb.) Kuntze, Rev. Gen. Pl. 3(2): 240. 1898. *Buchnera bilabiata* Thunb., Prodr. Pl. Cap 100. 1800. TYPE: South Africa. *Thunberg* s.n. (holotype, UPS!).

Perennial, 20–60 cm tall, stiffly erect, unbranched or with 2 to 3 branches from below middle, pilose or scabrous hispid. Stem either terete, square and furrowed, or winged. Leaves 2–6 × 1–2 mm, lanceolate, appressed, or 10–60 × 1–2(–5) mm, linear, ascending or spreading, shorter or longer than internodes, margin entire, veins obscure. Bracts 5–20 × 1–3 mm, linear or lanceolate, shorter or longer than calyx. Calyx tubular, 5-ribbed, 4–12 mm long; tube 2–6(–8) mm long; lobes 4 equal or 5 unequal, linear or deltate, 1–5 mm long. Corolla densely pubescent, tube 7–17 mm, bent and expanded proximally or distally; lobes of lower lip 1–4(–7) × 1–2 mm, upper lip 2–4(–7) × (1–)2–4(–7) mm, the three lobes of the lower lip and the two lobes of the upper lip are fused almost their entire length forming a strongly bilabiate corolla, tips obtuse or acute, short.

The lack of dependable characters separating *S. bilabiata*, *S. rowlandii* Engl., *S. barteri* Engl., *S. ledermannii* Pilger, and *S. welwitschii* Engl. led Hepper (1960) to reduce these species to subspecies of *S. bilabiata*. Statistical analysis of populations of each subspecies of *S. bilabiata* supported Hepper's treatment (Mohamed, 1994). All subspecies of *S. bilabiata* form a natural group with characters grading into one another. Like *S. bilabiata*, *S. linearifolia* also possesses a strongly bilabiate corolla, winged stem, perennial habit, and occurs in small populations. We consider *S. linearifolia* as a sixth subspecies of *S. bilabiata*; otherwise, we follow Hepper (1960).

The following is a key, followed by brief descriptions, of the subspecies of *S. bilabiata*:

### KEY TO SUBSPECIES OF *S. BILABIATA*

1. Bract broader than leaf; lower corolla lobes obtuse ..... 2
- 1'. Bract narrower than leaf, lower corolla lobes acute ..... 3
- 2(1). Calyx lobes subulate, about same length as tube, flowers opposite in spikes, spike as long as vegetative stem ..... 6b. subsp. *bilabiata*
- 2'. Calyx lobes lanceolate, < ½ tube length, flow-




Figure 5. Distribution based on specimens of *Striga bilabiata* subsp. *barteri* (diamond), *S. bilabiata* subsp. *bilabiata* (circle), and *S. bilabiata* subsp. *rowlandii* (triangle).

ers imbricate in strobilus-like spikes, spike much shorter than vegetative stem ..... 6c. subsp. *jaegeri*

3(1). Leaf scale-like, < 1 cm long, plant covered with appressed, hispid hairs; corolla bent proximally within calyx teeth .... 6e. subsp. *linearifolia*

3'. Leaf not scale-like, 3–8 mm long; plant covered with soft or hispid divergent hairs; corolla bent distally above calyx ..... 4

4(3). Plant ciliate; inflorescence strobilus-like; calyx 5 mm long ..... 6d. subsp. *ledermannii*

4'. Plant pilose; inflorescence not strobilus-like; calyx 6–12 mm long ..... 5

5(4). Stem winged; calyx lobes deltate,  $\frac{1}{2}$  tube length; spike shorter than vegetative stem ..... 6f. subsp. *rowlandii*

5'. Stem terete; calyx lobes linear, equaling tube length; spike longer than vegetative stem ..... 6a. subsp. *barteri*

**6a. *Striga bilabiata* subsp. *barteri* (Engl.).** Hepper, Kew Bull. 14: 414. 1960. *Striga barteri* Engl., Bot. Jahrb. Syst. 23: 514. 1897. TYPE: Nigeria. Nupe, Barter 1170 (holotype, K!).

*Striga glandulifera* Engl., Bot. Jahrb. Syst. 23: 514, fig. H-J. 1897. TYPE: Sudan. Schweinfurth 2931 (isotype, K!).

Plant to 18 cm tall, usually unbranched, densely pilose with divergent hairs, stem terete. Leaves 10–40  $\times$  1–3 mm, linear, spreading, tortuous when dry,

longer than internodes. Lower bracts 10–22(–45)  $\times$  1 mm, linear, longer than calyx, upper bracts lanceolate, shorter than calyx. Flowers alternate, the upper ones opposite, forming long open spikes, spike longer than vegetative stem. Calyx 6–12 mm long; tube 4(–8) mm long; lobes 5, unequal, linear, 3–4 mm long, about the same length as tube. Corolla lilac or pinkish, 12–14 mm long, bent and expanded distally above calyx, tube pubescent or glandular-pubescent; lobes of lower lip 2–3  $\times$  1 mm, tips acute, upper lip 2–4  $\times$  2 mm, tips acute.

The range of *S. bilabiata* subsp. *barteri* extends from Senegal to southern Sudan and south into Uganda and Tanzania between 14°N and 3°10'S (Fig. 5). Within this region, it occurs only in small scattered populations among grasses in rocky savanna woodland.

*Selected specimens examined.* BURUNDI. Ruyigi: Kigamba, 3°6'S, 30°33'E, Reekmans 7125, 7166 (K). CHAD. Logone-Oriental: Doba, 8°39'N, 16°51'E, Elliott s.n. (K). GUINEA. Northwest: Kouroussa, Banko, 10°45'N, 9°54'W, Adam 26896 (MO). GUINEA BISSAU. Southern: Madina do Boe, 11°45'N, 14°13'W, Epírito Santo 3199 (K); Kipkarren, Hill 755 (K). MALI. Kayes: Madina, 13°25'N, 8°48'W, Robert 16608 (K). NIGER. Barter 1170 (GH). NIGERIA. Kwara: slope near waterfall at head of Jebba Glen, 9°5'N, 4°50'E, Meckle 1011 (K). RWANDA. Southern: Mutara, Ngarama, 2°20'S,

30°10'E, *Alcool* 4116 (K). SENEGLA. **Oriental:** Tambacounda, Hasirik, 13°45'N, 13°40'W, *Adam* 17917, 17934 (MO), *Tutin* 162 (K). SUDAN. **Equatoria:** Zandi-land, Ibba, 4°50'N, 29°10'E, *Wyld* 725 (BM). TANZANIA. **Mwanza:** Nyanhangha, Bugufi, Ngara, 3°10'S, 33°8'E, *Tanner* S110 (K). **Ziwa Magharibi:** Bukoba, near Nddma, 1°30'S, 31°10'E, *Harrer* 2227 (K); W Lake Province, Lusahanga, Bihamulo, 2°52'S, 31°12'E, *Tanner* 5655 (K); W lake Prov., Murukurazo, Bugufi, Ngara, 2°25'S, 30°35'E, *Tanner* 5108 (K). UGANDA **Eastern:** Sorati, Kapiri Ferry, 1°43'N, 33°35'E, *Lind* 340 (K). **North Buganda:** Butemba-Singo, 1°8'N, 31°35'E, *Brown* 2027 (K); Kambuga, Kizezi, 1°0'S, 29°58'E, *Purseglove* 2711 (K). **Northern:** 2 km SE of Kikandwa, 3°38'N, 32°7'E, *Lester & Morrison* 2337 (K). **South Buganda:** Masaka, 0°21'S, 31°45'E, *Elliott* 1481 (K); Mubende, Lukwamnigo Hills, 1°0'N, 31°0'E, *Lanbrestreter* 18 (K). **Western:** Mubende, 1°5'N, 31°5'E, *Lind* 2027 (K), *Plant* 706 (K); Toro, Oruha Hill, 0°25'N, 30°30'E, *Eggeling* 4033 (K).

### 6b. *Striga bilabiata* Kuntze subsp. *bilabiata*

*Striga thunbergii* Benth., Companion Bot. Mag. 1: 363. 1835. TYPE: Botswana. (?)*Anonymous* 2494 (holotype, K!). *Striga thunbergii* var. *grandiflora* Benth., in DC. Prodr. 10: 502. 1846. TYPE: South Africa. Kat River Poort, Drége 2297b (holotype, K!). *Striga welwitschii* Engl., Bot. Jahrb. Syst. 23: 514, 12, fig. F, G. 1897. TYPE: Angola. *Welwitsch* 5821/5820 (holotype, K!); isotype, BM!. *Striga thunbergii* var. *grandiflora* Engl., in Baum Kunene-Samb. Exped. 369. 1903. TYPE: Angola. Longamindung, *Baum* 552 (holotype, K!).

Plant 20(–40) cm tall, usually unbranched or with 2 or 3 branches from base, scabrous, densely hispid with divergent hairs, stem square, furrowed. Leaves 10–20(–30) × 1(–2) mm, linear or lanceolate, ascending, decussate, longer than internodes. All bracts similar, 6–18 × 1–3 mm, lanceolate, ciliate, enclosing calyx, broader than leaves. Flowers opposite in short, dense spikes, spike as long as vegetative stem. Calyx 6–10 mm long; tube 3–5(–7) mm long; lobes 5, unequal, subulate, 3–5 mm long, about the same length as tube. Corolla white or mauve, tube 10–15 mm long, bent and expanded proximally within calyx teeth, densely glandular-pubescent; lobes of lower lip 2–4 × 1–2 mm, tips obtuse, upper lip 2–5(–7) × 2–4 mm, tips obtuse.

*Striga bilabiata* subsp. *bilabiata* is found only between 2°22'S and 34°02'S (Fig. 5) in southern Democratic Republic of Congo, Zambia, Angola, Zimbabwe, Botswana, and South Africa. It is less frequent to the north and unknown from the East African coast. This subspecies varies considerably in height but is typically about 20 cm tall. *Striga bilabiata* subsp. *bilabiata* is strongly autogamous (Ralston et al., 1987).

Selected specimens examined. ANGOLA. **Banguella:**

Banguella, 12°36'S, 13°20'E, *Gossweiler* 2234 (K). **Moxio:** 2 mi. N of Jimele River bridge on Caianda-Mwinilunga road, 11°17'S, 23°48'E, *Richards* 17147 (K). BOTSWANA. 3 mi. SW Nzinzi down Omuramba Mpungu, *Winter* 3995 (K). **Ngamiland:** banks of Okavango River at Runtu, 19°34'S, 23°15'E, *Winter & Marais* 4478, 4913 (K). **North District:** Tsetse Fly Camp, W of Parakarungu, *Lambrech* 264 (K). **Southern:** Digkatlong Ranch, 25°40'S, 25°3'E, *Hansen* 3016 (K, MO). DEMOCRATIC REPUBLIC OF CONGO. **Kivu:** 6.3 km NW of Katshungu, 2°22'S, 27°22'E, *Anonymous* 4863 (K). **Shaba:** Katanga, Biano Plateau, 7°0'S, 27°22'E, *Russell* 41 (K). LESOTHO. **Central:** N of Blue Mountains Pass, 29°30'S, 28°30'E, *Dieterlen* 527 (K). **Northern:** Leribe, 28°58'S, 28°0'E, *Dieterlen* 529 (K). MALAWI. **Central:** Lilongwe, 13°59'S, 33°47'E, *Jackson* 649 (K). MOZAMBIQUE. **Maputo:** Namaacha, Mt. Ponduine, 26°0'S, 32°25'E, *Schuter & Nuvunga* 6644 (K). NAMIBIA. **Windhoek:** 20 mi. SW of Windhoek, Farm Haris, 22°35'S, 17°4'E, *Winter* 2540 (K). SOUTH AFRICA. **Cape:** Stockenstrom, Nogsback, 32°40'S, 26°50'E, *Leighton* 2948 (MO). **Natal:** Bethlehem, Royal National Park, Tugela River, 28°45'S, 29°28'E, *Ross* 2240 (K). **Orange Free State:** Barkly West, Cristaaffrontein, 28°30'S, 24°26'E, *Acocks* 1481 (K). **Transkei:** Griqualand East, Clydesdale, 30°40'S, 29°0'E, *Tyson* 894 (GH). **Transvaal:** Albany, Alcedale Road, 26°6'S, 28°5'E, *Bayliss* 8112 (MO, US). SWAZILAND. **Hhohho:** 5 km NW of Mbabane, 26°18'S, 31°6'E, *Kemp* 701 (MO, US). **Matobo:** Farm Besua Kobila, *Miller* 7260 (K). ZAMBIA. **Central:** 5 mi. from Mumbwa on Road to Kafue Hoek, 15°0'S, 26°52'E, *Drummond & Cookson* 6185 (K). **Copperbelt:** 3 km NE of Mufulira, 12°30'S, 28°17'E, *Cruse* 372 (K). **Lusaka:** Broken Hill, 14°30'S, 28°29'E, *Mutimushi* 932 (K). **Mbala:** Lyopa Plain, *Sanane* 980 (K). **Northern:** 16 km N of Mbala, 8°40'S, 31°15'E, *Philcox et al.* 10170 (K). **Western:** 4 mi. N of Kalobo, 14°55'S, 22°30'E, *Drummond & Cookson* 6451 (K). ZIMBABWE. Farm Besna Kobila, *Miller* 2630, 2671, 5737 (K). **Lomagundi:** Great Dyke, *Philcox & Muller* 9080 (K). **Manicaland:** Inyanga, Cheshire, 18°15'S, 32°42'E, *Nordinh & Weimarck* 44347 (K). **Mashonaland South:** Macheka, 18°6'S, 31°47'E, *Eyles* 2015 (K). **Matabeleland South:** Matopo, 20°28'S, 28°29'E, *Miller* 1590 (MO).

### 6c. *Striga bilabiata* subsp. *jaegeri* Hepper, Kew Bull. 14: 415. 1960. TYPE: Mali. Massif de Kita, *Jaeger* k.2 (753/51) (holotype, K!).

*Striga brouilletii* Mielcarek, Bull. Jard. Bot. Belg. Bull. Nat. Plantentuin Belg. 58: 121–127. 1988. TYPE: Guinea (Conakry). Mt. Kankan: Koumbankourou, *Lisowski* 80579 (holotype, POZG, photo!; isotype, BR not seen).

Plant 30–40 cm tall, unbranched, densely hispid with divergent hairs, stem winged. Leaves 40–70 × 2–3 mm, linear to narrowly elliptic, spreading, decussate, alternate toward the apex, with prominent midrib, longer than internodes. All bracts similar, 9–11 × 3–4 mm, widely obovate, acuminate, ciliate, enclosing calyx, broader than leaves. Flowers imbricate in short, dense, strobilus-like spikes, spike much shorter than vegetative stem. Calyx 7–8 mm long; tube 5–6 mm long, lobes 4 equal, or 5




Figure 6. Distribution based on specimens of *Striga bilabiata* subsp. *jaegeri* (triangle), *S. bilabiata* subsp. *ledermannii* (diamond), and *S. bilabiata* subsp. *linearifolia* (circle).

unequal, lanceolate, 1–2 mm long, < ½ tube length. Corolla mauve, tube 15–17 mm long, bent and expanded distally above calyx, densely pubescent; lobes of lower lip 2–4 × 2 mm, tips obtuse, upper lip 1.5–2.5 × 1–2 mm, tips obtuse.

*Striga bilabiata* subsp. *jaegeri* is a rare taxon determined by its broad bracts and short compact spikes. It has been collected from savanna on rocky plateaus in Guinea, Senegal, and Mali between 10°18'N and 13°50'N (Fig. 6). *Striga bilabiata* subsp. *barteri* and *S. bilabiata* subsp. *jaegeri* have more northerly distributions than *S. bilabiata* subsp. *linearifolia* and *S. bilabiata* subsp. *rowlandii* (Figs. 5, 6).

*Selected specimens examined.* GUINEA. Northern: 5 km NW of Karifameriya, 11°55'N, 13°30'W, Lisowski 80551 (BR). MALI. Kayes: Kita, 13°10'N, 9°34'N, Jaeger 753/51, 2541 (K). SENEGAL. Oriental: Tambacounda, Gouloumbou, 14°0'N, 13°40'W, Adam 14861 (MO).

**6d. *Striga bilabiata* subsp. *ledermannii* (Pilger)**  
Hepper, Kew Bull. 14: 413. 1960. *Striga ledermannii* Pilger, Bot. Jahrb. 45: 215. 1910.  
TYPE: Cameroon. Garoua, *Ledermann* 4529 (lectotype, designated by Hepper (1960), BM!).

Plant 36 cm tall, unbranched, scabrous, with ciliate divergent hairs, stem winged. Leaves 10–23 × 2–3(–5) mm, lanceolate, decussate, decurrent, appressed, enclosing stem, longer than internodes. All bracts similar, 5–15 × 2–4 mm, imbricate, elliptic or lanceolate, ciliate, enclosing calyx. Flowers imbricate in short, dense, strobilus-like spikes, spike shorter than vegetative stem. Calyx 5 mm long; tube 2.5–3 mm long, lobes 5, unequal, linear, 2–3 mm long, about the same length as tube. Corolla pink, tube 12 mm long, bent and expanded distally above calyx, usually retrorsely pubescent; lobes of lower lip 1–2 × 0.5–1 mm, short, tips acute or obtuse, upper lip 3 × 1–2 mm, tips obtuse.

*Striga bilabiata* subsp. *ledermannii* is endemic to the northern province of Cameroon between 9°N and 10°N (Fig. 6). It is infrequent, like other subspecies of *S. bilabiata*, perhaps because it only parasitizes perennial hosts such as *Loudetia simplex* (Nees) C. E. Hubbard, *Schizachyrium sanguineum* (Retz.) Alston, and *Hyparrhenia* species (Kenfack et al., 1996). Although the hosts of other subspecies of *S. bilabiata* have not been studied, parasitism of perennial hosts may explain why *S. bilabiata* subsp. *barteri*, *S. bilabiata* subsp. *linearifolia*, and *S. bil-*

*abiata* subsp. *jaegeri* are infrequent and occur only in small populations.

*Selected specimens examined.* CAMEROON. **Northern:** 20 km SW of Dourbey, Biatomni, *Fotius* 2219 (K); 23 km S Garoua, 9°0'N, 13°25'E, *Musselman et al.* 10/16/88 (K, ODU); Bogalo by River Benue, 9°0'N, 13°50'E, *Talbot* 313 (BM); Buar to Buala 9°34'N, 13°46'E, *Michael* 9531 (K); Garoa, 9°16'N, 13°25'E, *Ledermann* 4529 (BM); Jaiya Buge, 6°20'N, 16°15'E, *Michael* 9706 (K); Kounden, 6°7'N, 14°38'E, *Saxer* 254 (K); near Koubout, 9°57'N, 13°45'E, *Thomas* 2452 (MO).

**6e. *Striga bilabiata* subsp. *linearifolia* (Schum. & Thonn.) Mohamed, comb. nov.** Basionym: *Buchnera linearifolia* Schum. & Thonn., Beskr. Guin. Pl.: 279. 1827. *Striga linearifolia* (Schum. & Thonn.) Hepper, Kew Bull. 14: 416. 1960. TYPE: Ghana. *Thonning* 284 (holotype, C, holotype negative # 2785, 2786; isotypes, K!, P-JU not seen).

*Striga strictissima* Skan, Fl. Trop. Afr. 4(2): 407. 1906. TYPE: Nigeria. *Barter* 1263 (holotype, K!).

Plant up to 62 cm tall, unbranched or with 1 to 2 branches from base, covered with dense, white, appressed, hispid hairs, stem winged. Leaves reduced, 3–8 × 1–2 mm, lanceolate, decussate, decurrent, appressed, shorter than internodes. All bracts similar, 5–7(–10) × 1–2 mm, lanceolate, slightly longer than calyx. Flowers opposite in dense spikes, spike shorter than vegetative stem. Calyx 4–7 mm long; tube 2–3(–4) mm long; lobes 5, unequal, linear, 1.5–3 mm long, about the same length as tube. Corolla whitish to bluish purple, tube 7–14 mm long, bent and expanded proximally within calyx teeth, densely retrorsely pubescent; lobes of lower lip 1–3(–7) × 0.5–1 mm, tips acute, upper lip 2–5 × 2–4 mm, slightly round.

*Striga bilabiata* subsp. *linearifolia* is widely distributed in Africa in wet grassy savannas. Its range extends from 13°N and reaches its southern limit in Malawi (10°55'S) (Fig. 6). Like other subspecies of *S. bilabiata*, it occurs in small populations often of only a few plants. For example, we gathered it once from populations of two plants in Mali, Ivory Coast, and Guinea (Conakry).

*Selected specimens examined.* ANGOLA. **Benguela:** Benguela, Sambo, 12°30'S, 13°26'E, *Teixerira* 303 (BM). BURKINA FASO. **Central:** S of Sapones, 12°0'N, 1°30'W, *Safa & Musselman* 10/21/87 (ODU). **Haut Bassins:** 5 km S of Banfora, 10°40'N, 5°50'W, *Safa & Musselman* 10/3/87 (ODU). BURUNDI. **Central:** Gitega, 3°20'S, 29°55'E, *Anonymous* 1829 (K). **Southern:** Danga (Moso), Bururi, 4°12'S, 30°0'E, *Reekmans* 6786 (K). CENTRAL AFRICAN REPUBLIC. **Bangoran:** 4 km N of Camp Koumbala, Gaumba Creek, 8°31'N, 21°13'E, *Fay* 7057 (MO). ETHIOPIA. **Gamu Gofa:** Gughe high-

lands, 6°3'N, 37°35'E, *Gilbert & Thulin* 415 (MO). **Ogaden Harerge:** Harar, Harrar-Jijega, 9°13'N, 42°29'E, *de Wilde* 6399 (MO). **Shewa:** Awash National Park, 8°55'N, 39°55'E, *Gilbert & S. B.* 1250 (K). GHANA. **Accra:** Accra Plains, 5°35'N, 0°6'W, *Morton* 6123 (K). **Eastern:** Keta Krachi, 5°55'N, 1°2'E, *Morton* 7171 (K). **Northern:** 2 mi. S Nasia on Tamale-Bolgatanga Road, 9°55'N, 0°50'W, *Adams* 4152 (K). **Upper:** 10 mi. from Lambusie, 10°52'N, 2°40'W, *Morton* 8794 (K). VOLTA: Akuse, 6°6'N, 0°8'E, *Vigne* 4359 (US). GUINEA. *Probegum* 364 (K). IVORY COAST. **Northeast:** Gawi-seye, along Iringou River, 9°14'N, 3°41'W, *Amshoff* 225 (MO). **Odienne:** Odienne, 9°31'N, 7°38'W, *Adam* 27125 (MO). **Seguela:** Seguela and Man, 8°0'N, 6°38'W, *Ake* 8755 (K). KENYA. **Central:** Lake Naivasha, 0°48'S, 36°20'E, *Mathew* 6126 (K). **Eastern:** Embu, Rumbia, 0°32'S, 37°28'E, *Graham* 2405 (K). **Rift Valley:** Chyulu Hills, 2°40'S, 37°52'E, *Gilbert & Kariuki* 18863 (K). **Shiyanga:** Kahawa, Kenyatta College, 3°50'S, 32°36'E, *Akestrid* 2527 (UPS). MALAWI. **Northern:** 27 mi. N of Rumpi, 10°55'S, 33°54'E, *Pawek* 12317 (K, MO). MALI. **Koulikoro:** Simidji, 11°40'N, 7°35'W, *Musselman & Mohamed* 9/30/88 (ODU). NIGER. *Barter* 1263 (GH). NIGERIA. **Kaduna:** Katsina, Maska, 13°0'N, 7°35'E, *Keay* 25896 (K). **Niger:** between Gawa–Abuja Road Junction, 9°15'N, 6°50'E, *Onochie* 18697 (K). **Northern:** Lemme, *Lely* 146 (K). **Plateau:** Shere Mountains, 9°10'N, 9°50'E, *Hall & Daramola* 67404 (MO). **Sokoto:** 10 km N of Bussa, 50 km S of Yelwa, 10°15'N, 4°35'E, *Cook* 445 (K). SUDAN. **Bahr Al-Ghazal:** Wau, 7°46'N, 28°2'E, *Anonymous* 1651 (K). TANZANIA. **Arusha:** 3 mi. E of Lake Manyara, 3°40'S, 35°53'E, *Lamprey* 515 (K). **Kigoma:** Buha, 56 mi. from Kasulu on Kigoma Road, 5°30'S, 30°0'E, *Verdcourt* 3330 (K). **Kilimanjaro:** E Kilimanjaro, Mran Rombo, 3°12'S, 37°20'E, *Haarer* 517 (K). TOGO. **Central:** 9 km N of Sokode, 9°7'N, 1°10'E, *Hakki et al.* 289 (K). UGANDA. **Bokora:** Iriri, *Nilson* 332 (K). **Central:** E of Mengo, Kakonge, 0°15'N, 32°28'E, *Rwaburindore* 580 (MO). **Eastern:** Busana, Bugisu, 1°15'N, 34°20'E, *Eggeling* 350 (K). **South Buganda:** near Sanje Rest Camp, 0°49'S, 31°30'E, *Ankole & Eggeling* 610 (K). **Southern:** Ankole, Ibanda, 0°8'S, 30°30'E, *Ragslane* 1251 (US). **Western:** Busoga, 8 mi. W of Kamuli, 0°58'N, 33°1'E, *Wood* 726 (K).

**6f. *Striga bilabiata* subsp. *rowlandii* (Engl.) Hepper, Kew Bull. 14: 415. 1960. *Striga rowlandii* Engl., Bot. Jahrb. Syst. 23: 513. 1897.** TYPE: Nigeria. Western Lagos, *Rowland* 1893 (holotype, K!).

Plant up to 34 cm tall, unbranched or with 2 or 3 branches from base, sparsely to densely pilose with divergent hairs, stem winged. Leaves 10–55 × 1–3 mm, linear, decussate, decurrent, ascending, longer than internodes. Lower bracts 7–10(–18) × 1–2 mm, lanceolate, longer than calyx, upper bracts shorter than calyx. Flowers alternate at base of inflorescence, opposite above in a somewhat dense spike, spike shorter than vegetative stem. Calyx 6–9 mm long; tube 5–6(–8) mm long; lobes 5, unequal, deltate, 1–2 mm long, ½ tube length. Corolla pink, mauve, or white, tube 9–12(–15) mm

long, bent and expanded distally above calyx, densely pubescent; lobe of lower lip 2–4 × 1 mm, tips acute, upper lip 2–4 × 2–4 mm, tips acute.

The subspecies is endemic to the relatively wet, rocky-plateau grasslands in West Africa between 13°N and 6°10'N (Fig. 5); its range extends from Senegal to Cameroon.

*Selected specimens examined.* BURKINA FASO. **Haut Bassins:** 18 km N of Banfora, 10°39'N, 4°50'W, *Leeuwenberg* 4357 (K, MO). CAMEROON. **Northern:** Mont Boulor, 5 km N of Maroua, 10°37'N, 14°22'E, *Letouzey* 624 (K). GAMBIA. *Hayes* 539 (K). GHANA. **Eastern:** 2 mi. NE of Kwahu-Tofo on road to Adawso, 6°15'N, 0°15'W, *Innes* 909 (K); 3 mi. N of Kwahu-Tafo, 6°17'N, 0°20'W, *Harris* 195 (K). **Northern:** 10 mi. from Lumbuga, 9°20'N, 2°5'E, *Morton* 8795 (K); 2 mi. beyond Kwaha-Tafo on Mankrong Road, 6°10'N, 0°18'W, *Morton* A663 (K); Afram Glams, 6°50'N, 0°25'W, *Onsow* 709 (K); Damongo, Yendi, 9°17'N, 0°22'E, *Hepper & Morton* A3110 (K); Kwaha-Tafo, 6°10'N, 1°0'W, *Hall* 131, 260 (K); Okroso to Otiso Ferry, 10°0'N, 0°25'E, *Hepper & Morton* A3039 (K); Onyaboni to Jaketi, *Morton* A3016 (K); Sesihamango, *Thomas* D138 (K). GUINEA. **Northern:** Seriba, Gaoual, 11°50'N, 13°10'W, *Adam* 14812 (MO). **Southern:** N'zerekore, 7°40'N, 8°50'W, *Adam* 26898, 27661 (MO). IVORY COAST. **Northeast:** Dabakala, 45 km NW Dabakala, 8°32'N, 4°9'W, *Geerling & Bokdam* 2029 (MO). NIGER. *Barter* 1169 (GH). NIGERIA. **Bauchi:** Bauchi Plateau, 10°25'N, 9°55'E, *Lely* 83 (K); 10°20'N, 9°45'E, *Lely* P83 (K, MO). **Kaduna:** 3 mi. N of Zaria, 11°8'N, 7°43'E, *Milne-Redhead* 5043 (K); Kaduna, 10°31'N, 7°25'E, *Hill* 51 (K); near Zaria, 11°7'N, 7°43'E, *Dalziel* 407 (K); Riruwa-Kano Hills, 11°15'N, 8°17'E, *Carpenter* 192 (K); Zaria, Mando, Gayam, 10°27'N, 7°25'E, *Keay* 25833 (K); Zaria, Samaru, 11°5'N, 7°43'E, *Harris* 11 (K); N Nigeria, *Pobeguin* 364 (K). **Oyo:** 8 mi. E of Igboho, Upper Ogun Forest Reserve, 8°47'N, 4°0'E, *Keay s.n.* (K); Kogosti, 27 mi. E of Ilesha, 7°35'N, 4°58'E, *Lowe* 3378 (K). **Plateau:** Vom, Bauchi Plateau, 9°40'N, 8°48'E, *Young* 185 (K). SENEGAL. **Casamance:** Ziguinchor, Bayotter, 12°35'N, 16°12'W, *Adam* 13724 (MO). **Oriental:** Niokolo-Koba, 13°0'N, 13°0'W, *Berhaut* 1630 (K); Tambacounda, Niokolo-Koba, 13°5'N, 13°0'W, *Adam* 14200, 26891, 26897 (MO). SIERRA LEONE. Kafari (Malal), *Bakshi* 146 (K); Lake Sonfon, *Gledhill* 285 (K). **Northern:** Buyabuya Scarcies, 9°45'N, 12°10'W, *Elliott* 4279 (GH); Falaba, 9°52'N, 11°20'W, *Elliott* 5207 (K, MO); Loma Manza, 9°15'N, 11°2'W, *Thomas* 155 (K); near Mongo, Bure Makonte Chiefdom, 9°40'N, 11°50'W, *Glanville* 241 (K), *Adames* 224 (K); Sumbuya, 7°39'N, 11°58'W, *Thomas* 603 (K); Rhombe Swamp, *Adames* 146 (K); Robis, *Jordan* 190 (K).

In summary, we recognize six subspecies of *S. bilabiata*, the five subspecies originally proposed by Hepper (1960) in addition to *S. bilabiata* subsp. *linearifolia*. These form a natural group: they are generally autogamous, perennials, unbranched, have strongly bilabiate corollas, winged stems, and occur in small populations. Many of the morphological features used to distinguish other species,

such as heights and sizes, overlap in this species complex, making them unreliable characters.

**7. *Striga brachycalyx* Skan, Fl. Trop. Afr. 4(2): 403. 1906. TYPE: Nigeria. Nupe, *Barter* 1858 (holotype, K!).**

*Striga warneckei* Engl. ex Skan, Fl. Trop. Afr. 4(2): 414. 1906. TYPE: Togo. *Warnecke* 201 (isotypes, BM!, K!).

Annual, to 70 cm tall, slender, branched from middle, scabrous, hispid, stem square, furrowed. Leaves (5–)20–37(–50) × 1(–2) mm, subopposite, linear, ascending or spreading, longer than internodes, margin entire, veins obscure. Lower bracts 5–20 × 1 mm, linear, as long as or longer than calyx, upper bracts subulate, shorter than calyx. Flowers alternate, forming somewhat open racemes, raceme shorter than the vegetative stem. Calyx 5-ribbed, 3–6 mm long; tube 1.5–3 mm long; lobes 5, almost equal, linear, 1–3.5 mm long, about the same length as tube. Corolla purple with white center, tube (6–)9–19 mm long, bent and expanded distally above calyx, sparsely glandular-pubescent; lobes of lower lip 2–3(–5) × 1–2 mm, tips obtuse, upper lip 1–2(–4) × 2–4 mm, obovate.

*Striga brachycalyx* is widespread and often common in dense stands in the savannas of west and central Africa between 13°12'N and 5°17'N (Fig. 7). Its range extends from Ivory Coast into Burkina Faso, Mali, Ghana, Togo, and Nigeria, reaching its eastern limits in Sudan. It is also known from Congo. This species is morphologically homogenous throughout its range. *Striga brachycalyx* resembles *S. aspera*, but can be distinguished by its smaller flowers and more slender stems. The two species are sympatric, found in relatively drier savannas, and frequently encountered in large populations. Unlike *S. aspera*, which occasionally invades agroecosystems, *S. brachycalyx* is not known to damage grains even though it is very common.

*Selected specimens examined.* BURKINA FASO. **Center East:** 5 km N of Tenkodogo, 12°0'N, 0°19'W, *Safa & Musselman* 10/9/87 (ODU). **Center West:** 5 km W of Godi, 11°55'N, 2°24'W, *Safa & Musselman* 10/12/87 (ODU). **Central:** 1 km S of Kamboiene, 12°25'N, 1°40'W, *Musselman* 6118 (ODU). **Haut Bassins:** 5 km S of Banfora, 10°37'N, 4°44'W, *Safa & Musselman* 10/13/87 (ODU). **Northern:** 5 km N of Gourey, 13°12'N, 2°22'W, *Safa & Musselman* 10/10/87 (ODU). **Volta:** near Boroma and Laba, 11°50'N, 2°55'W, *Raynal* 2282 (K). DEMOCRATIC REPUBLIC OF CONGO. **Kasai Oriental:** Tumba Camp, Kasumanyense Mbuga, Rukwa Valley, 3°7'S, 23°34'E, *Siame* 1444 (BM). GHANA. Ayafe, *Adams* 4693 (K). **Northern:** Bole, 9°5'N, 2°29'W, *Morton* A3295 (K). **Upper:** 4 mi. from Tono, Navrongo, 10°57'N, 1°40'W, *Irvine* 4668 (K). IVORY COAST. **Northern:** Kor-




Figure 7. Distribution based on specimens of *Striga brachycalyx* (triangle), *S. elegans* (circle), and *S. hallaei* (diamond).

hogo, 10 km SW of Korhogo, 9°27'N, 5°38'W, *Riches* 243 (K). MALI. **Koulakoro:** 12 km N of Oulessebougou, 12°0'N, 7°40'W, *Musselman & Mohamed* 9/30/88 (ODU). **Sikasso:** 6 km from Bougouni, 11°32'N, 7°30'W, *Raynal* 22196 (K). NIGERIA. **Bauchi:** Bauchi Plateau, 10°25'N, 9°55'E, *Lely* P567 (K). **Benue:** Abinsi, 7°20'N, 9°25'E, *Dalziel* s.n. (K). **Gongola:** Bogalo, close to River Benue, 8°45'N, 10°15'E, *Talbot* 312 (K). **Kaduna:** Zaria, Samaru, 11°5'N, 7°43'E, *Harris* 4, 7 (K). **Kwara:** Ilorin, 8°30'N, 4°32'E, *Parkinson* 9/17/86 (ODU). **Niger:** Bida/Mokwa Junction, 9°8'N, 5°40'E, *Mansfield* 9/4/78 (ODU). TOGO. **Southern:** Barani-Ideppé, 6°30'N, 1°45'E, *Kersting* 665 (K). **Southern:** Togo Forest, 6°29'N, 1°9'E, *Anonymous* s.n. (US).

#### 8. *Striga chrysanthia* A. Raynal, Bull. Jard. Bot. Belg. 39: 378. 1969. TYPE: Central African Republic. Zemio-Rafai, *Descoings* 12466 (holotype, Pl!).

Annual, small, < 12 cm tall, tufted, scabrous, densely pubescent, stem obtusely square. Leaves < 10 × 2(-3) mm, lanceolate, scale-like, shorter than internodes, margin entire, veins obscure. All bracts similar, 6 × 2 mm, lanceolate, shorter than calyx. Flowers alternate in rather dense spikes, spike as long as vegetative stem. Calyx 10-ribbed, 8–15 mm long; tube 6–13 mm long; lobes 5, subequal, deltate, 2 mm long, much shorter than tube. Corolla bright yellow, tube slender, 20–25 mm long, bent

and expanded distally above calyx, glandular pubescent; lobes of lower lip 15 × 8 mm, spreading, broadly round, upper lip 8 × 10 mm, reflexed backward, emarginate.

*Striga chrysanthia* is from the southern region of the Central African Republic and northern Democratic Republic of Congo (Raynal-Roques, 1969). It occurs between 3°40'N and 5°20'N (Fig. 8). The 10-calyx ribs, indumentum, and the yellow corolla suggest a relationship among this species, *S. lutea*, and *S. hirsuta*.

#### 9. *Striga dalzielii* Hutch., Fl. W. Afr., ed. 1(2): 226. 1931. TYPE: Nigeria. Zungeru, *Dalziel* 168 (holotype, K!).

Annual, 19–35 cm tall, stiffly erect, unbranched, scabrous, densely ciliate with long (1 mm), divergent hairs, stem square. Leaves 15–30 × 4–7 mm, 2-veined, opposite, linear, becoming reduced along the stem, as long as or longer than internodes, margin toothed. All bracts similar, 8–18 × 4 mm, linear, enclosing calyx, densely hispid along margins and midrib. Flowers sessile, imbricate, forming very dense spikes, spike shorter than vegetative stem. Calyx 10-ribbed, 5–8 mm long; tube 3–5 mm long; lobes 5, unequal, lanceolate, 2–3 mm long,




Figure 8. Distribution based on specimens of *Striga chrysanthia* (diamond), *S. gastonii* (triangle), *S. gesnerioides* (circle), and *S. lepidagathidis* (star).

shorter than tube. Corolla pink, tube 10–20 mm long, bent and expanded distally above calyx, glandular-pubescent; lobes of lower lip 3–8 × 3–5 mm, obtuse, upper lip 3–6 × 6 mm, emarginate.

*Striga dalzielii* occupies the wet Guinea savannas in Mali, Guinea, and Nigeria (Raynal-Roques, 1987). It resembles *S. klingii* and *S. macrantha*, but *S. dalzielii* has 10-calyx ribs whereas *S. klingii* and *S. macrantha* have 10–15 ribs. *Striga dalzielii* is characterized by its leaf-like bracts and long hispid hairs.

**10. *Striga elegans* Benth., Companion Bot. Mag. 1: 363. 1836. TYPE: South Africa. Burke 443/3591 (holotype, K!).**

Plant annual, 30(–50) cm tall, stiffly erect, unbranched or with 2 or 3 branches from base, sca-

brous, densely hispid, stem obtusely square, furrowed. Leaves 10–20(–30) × 2–3 mm, opposite, linear or lanceolate, shorter than internodes, margin entire, veins obscure. All bracts similar, 3–12 × 1–2 mm, lanceolate, shorter than calyx. Flowers opposite in dense racemes, raceme shorter than vegetative stem. Calyx 10–15-ribbed, 8–13 mm long; tube 6–7(–9) mm long; lobes 5 equal or 6 unequal, linear to lanceolate, 2–5 mm long, shorter than tube. Corolla brilliant scarlet, with yellow throat, tube 14–18 mm long, bent and expanded distally above calyx, densely glandular-pubescent; lobes of lower lip 5–10 × 3–5(–7) mm, ovate, upper lip 3–5 × 6–11 mm, emarginate.

*Striga elegans* is common in East and South Africa between 0°30'S and 33°30'S (Fig. 7). Its range extends from central Kenya south into Tan-

zania, Malawi, Zambia, Zimbabwe, Mozambique, Swaziland, Botswana, and Lesotho. *Striga elegans* shows considerable variation in corolla size and color that does not seem to correlate with geography. This beautiful species resembles *S. asiatica* and the two are sympatric in East Africa (see notes under *S. asiatica*), but it can be distinguished from *S. asiatica* by its relatively short, dense inflorescence with opposite flowers and large brilliant scarlet corolla lobes. Authentic crop loss from this parasite has not been reported.

*Selected specimens examined.* ANGOLA. **Huila:** Qui-lengues, Binde, 15°S, 13°30'E, de Menezes A.377 (K). **Mangle:** Pungo Andongo, 9°40'S, 15°30'E, Welwitsch 5851 (K). **Moxico:** between River Namavumba and River Lukaiia, 12°40'S, 20°45'E, Milne-Redhead 4018 (K). BOTSWANA. **Ngamiland:** Leshums Valley, 18°30'S, 23°20'E, Holub 1834 (K). **Southern:** Lobatse, 25°11'S, 25°40'E, Tapscott 2548 (K). KENYA. **Central:** Nairobi, 1°20'S, 36°48'E, Perceval 1901 (MO). **Coast:** Loitokitok, NE of Kilimanjaro 3°37'S, 37°20'E, Rogers 560 (K). **Rift Valley:** Rumuruti, Laikipia, 50 km N of Colchecio Lodge, 0°19'N, 36°34'E, Hepper & Jaeger 6618 (K, MO). LESOTHO. **Northern:** Mont-aux-Sources, 28°56'S, 28°50'E, Grant 2212387 (MO). MALAWI. **Central:** Kasungu, Chipala Hill, 4 mi. N of Kasungu, 12°55'S, 33°29'E, Robson & Jackson 1179 (K). MOZAMBIQUE. Paraia Sepulveda, Lavranos 4265 (MO). NAMIBIA. **Kavango:** 2.4 mi. E masari Camp on road to Nyangana, 18°0'S, 20°41'E, de Winter 4087 (K). SOUTH AFRICA. **Bechuanaland:** Little Klipbolikhonni, 27°0'S, 23°15'E, Burchell 2482 (K). **Cape:** Albany, Woest Hill, 33°0'S, 26°20'E, Bayliss 4526 (GH, MO). **Karasburg:** Orange River Colony, 28°45'S, 19°0'E, Walich 1278 (K). **Natal:** between Mooi River and Estcourt, 29°18'S, 29°56'E, Wood 394 (K). **Orange Free State:** Walton, 12 km S of Harrismith, 28°18'S, 29°3'E, Jocabsz 218 (K). **Transkei:** Griqualand, 30°40'S, 29°0'E, Tyson 1368 (MO). **Transvaal:** Randburg, Kelland, 26°6'S, 27°59'E, Liebenberg 8581 (MO). SWAZILAND. **Louwsburg:** 2 mi. E of Goedgegun, 27°20'S, 31°7'E, Ross 1743 (K, MO). TANZANIA. **Arusha:** Arusha National Park, 3°25'S, 36°37'E, Richards 25019 (K). **Iringa:** 8 km E of Kilolo, which is 30 km SE of Iringa, 7°40'S, 35°40'E, Peterson 833 (K). **Kilimanjaro:** E Kilimanjaro Mountain, 30°0'S, 37°38'E, Haarer 637 (K). **Tanzania. Mara:** Chamliho Hill, Ikiwu Musoma, Lake Province, 1°30'S, 33°58'E, Tanner 1690 (K). **Iringa:** Kyimbila, N of Lake Nyasa, 9°50'S, 34°50'E, Stoltz 2494 (GH, MO). ZAMBIA. **Lusaka:** 129 km E of Lusaka, 15°15'S, 28°40'E, Robinson 6174 (K). **Southern:** Kalomo, Kanchele, 17°0'S, 26°30'E, Astle 1571 (K). ZIMBABWE. **Maincaland:** 6 mi. N of Troutbeck, Inyangwa, 18°21'S, 32°15'E, Robinson 1961 (K). ZIMBABWE. **Mashonaland South:** Harare, Circular Drive, 17°50'S, 31°3'E, Hughes s.n. (ODU). **Matabeleland South:** Bellingwe (Mberengwa), Mount Buhwa, 20°37'S, 29°55'E, Wild 4315 (K). **Midlands:** Gokwe, 18°9'S, 29°0'E, Birmingham 1178 (MO).

## 11. *Striga forbesii* Benth., Companion Bot. Mag. 1: 364. 1836. TYPE: Mozambique. *Forbes s.n.* (holotype, K!).

Annual, 37–72 cm tall, stiffly erect, unbranched or with 2 to 4 branches below middle, scabrous to hispid, stem square, furrowed. Leaves 15–40(–90) × 4–12 mm, opposite, lanceolate, coarsely toothed, 3- to 5-veined, usually shorter than internodes. Lower bracts 20–50(–90) × 3–12 mm, leaf-like, longer than calyx, upper bracts shorter than calyx. Flowers opposite in open racemes, raceme shorter than vegetative stem. Calyx 10–15-ribbed, 9–14 (–20) mm long; tube 4–6 mm long; lobes 5, equal, broadly lanceolate, 5–9(–14) mm long, longer than tube. Corolla salmon-pink, tube 20–25 mm long, bent and expanded distally above calyx, densely glandular-pubescent; lobes of lower lip 6–9 × 3–6 mm, obovate, spreading, upper lip 3–6 × 6–9 mm, emarginate.

*Striga forbesii* ranges between 14°30'N and 28°30'S (Fig. 9). It is common in the Sudanian savannas from Senegal to Somalia and extends into East Africa reaching its southern limits in the Republic of South Africa. A plant of relatively wet open areas and swampy soils, *S. forbesii* can cause damage to irrigated crops such as sugar cane in Somalia (Musselman, unpublished).

Knepper et al. (1991) showed that some populations of *S. forbesii* are distylous. Flowers either had styles strongly exserted from the corolla tube exposing the stigma, or stigmas were among or just below the epipetalous stamens, with increasing possibilities of self-pollination.

*Striga forbesii* resembles *S. latericea*. Both are sympatric in East Africa, occupy wet habitats, and have unique salmon-pink corollas. No work has been done to demonstrate interfertility between these two species. *Striga forbesii* is distinguished from *S. latericea* by its annual habit, stiff hairs that give the plant a harsh texture, coarsely toothed and broader leaves, and shorter corollas with smaller lobes and a brighter salmon-pink color. The corolla color is darker in *S. latericea*.

*Selected specimens examined.* ANGOLA. **Cuanza North:** Cuanza, 8°50'S, 15°0'E, Johnston 83 (K). **Moxico:** between River Masanu and River Katetele, 13°0'S, 21°0'E, Milne-Redhead 4155 (BM, K). BOTSWANA. **Chobe:** near Pandamentenga, 18°9'S, 24°24'E, Musselman 4/9/86 (ODU). **Ngamiland:** NE Maun, 20°0'S, 23°26'E, Lambrecht 479 (K). **Bujumbura:** Karazi, 3°23'S, 29°22'E, Lewalle 335 (K). **Ruand:** Kigunee Territory, Biumba, 2°12'S, 30°32'E, Alcool 4142 (K). CAMEROON. **Northern:** 5 km W Marona, 10°35'N, 14°0'E, de Wilde et al. 2954 (K). DEMOCRATIC REPUBLIC OF CONGO. **Bandundu:** Kisambo, Kwango, 6°25'S, 18°10'E, Anonymous 3384 (K). **Equateur:** Olru-Uele, 4°0'N, 22°35'E, Lebrun 3557 (GH). **Haut Zaire:** Oriental, Dungu, Garamba National Park, 3°40'N, 28°30'E, Jaeger 1225 (K). **Katanga:** Elisabethville, 11°40'S, 27°28'E, Quarre 1451 (K). **Oriental:** Kasai, Gandajika, 6°42'S,




Figure 9. Distribution based on specimens of *Striga forbesii* (circle).

34°0' E, *Liben* 2038 (K). **Shaba:** River Lualaba, 50 km downstream from Bukama, 8°55'S, 25°55'E, *Paterson* 59 (K). **ETHIOPIA.** **Ihubabor:** Ihubabor, Gambela, Baro River, 8°16'N, 34°35'E, *Ash* 544 (K). **Welo:** Asaeta, Awash Valley, 33 km S of Tandah, 11°17'N, 40°54'E, *Ash* 2259 (K). **GAMBIA.** **Eastern:** Kuntaur, 13°49'N, 14°52'W, *Ruxton* 67, 149 (K). **Upper:** Bolgatanga, Crocodile Pool, 10°50'N, 0°50'W, *Hall* 558 (K). **GUINEA BISSAU.** **Northern:** between Farim and Bigene, 12°10'N, 15°15'W, *Anonymous* 3063 (K). **Western:** Sao, Fa, 12°27'N, 16°17'W, *Guerra* 3812 (K). **KENYA.** 30 mi. W of Bwa, *Wham* 26/24 (K). **Coast:** Kipini, Mlango Ya Simba, 2°30'S, 40°32'E, *Greenway & Rawlin* 9463 (K). **Narok:** Mara Plains, 1°0'S, 35°20'E, *Stewart* 327 (K). **Rift Valley:** Chyulu-Goohills, 1°0'N, 35°50'E, *Bally* 8072 (K). **Western:** Uasingishu Plateau and Trans Nzoia, 0°30'N, 34°35'E, *Douson* 674 (K). **MALAWI.** **Central:** 2 mi. N of Mzambazi, 13 mi. S of Mperembe, 11°58'S, 33°33'S, *Pawek* 13954 (K). **Kyimbila:** Nyasa Hochland-Station, *Stoltz* 213 (GH, US). **Northern:** 16 mi. N of Rumphi, 10°45'S, 33°54'E, *Pawek* 13913 (K). **Southern:** Lower Shire Valley, Elephant Marsh, 16°20'S, 35°0'E, *Robertson* 5 (K). **MALI.** **Massina:** Mopti, 14°30'N, 4°8'W, *Lean* 37 (K). **MOZAMBIQUE.** **Maputo:** Maputo (Lourenco Moazgues), Vila Luisa, 25°45'S, 32°35'E, *Grandraux & Delemos* 7913 (K). **Sofala:** near Massara, 18°20'S, 34°5'E, *Anonymous* 1858 (K). **Zambezia:** Morumbala Plains, 17°22'S, 35°36'E, *Kirk* 15/1/63 (K). **NIGERIA.** **Amambra:** Udi, upper Nau River, 6°18'N, 7°20'E, *Anderson* 1364 (K). **Bauchi:** Bauchi Plateau, 10°27'N, 9°57'E, *Lely* P369 (K). **Bendel:** 3 mi. N Asaba, 6°7'N, 6°47'E, *Killick* 234 (K). **Cross River:** Oban, 5°17'N, 8°35'E, *Talbot* 308 (BM). **Gongola:** Mambilla Plateau, 6°5'N, 11°14'E, *Chapman* 2935 (K). **Kaduna:** 2 mi. NE

of Kaduna, 10°32'N, 7°26'E, *Jackson* 254 (K). **Niger:** Zungeru, 9°48'N, 6°5'E, *Dalziel* 167 (K). **Plateau:** Jos Plateau, 9°50'N, 8°40'E, *Batten-Poole* 371 (K). **Sokoto:** 10 mi. N Bussa, 10°20'N, 4°35'E, *Cook* 386 (K). **SOMALIA.** **Jubba House:** Middle Juba, Mareeri, 0°25'N, 42°42'E, *Musselman* 7086 (ODU). **SOUTH AFRICA.** **Natal:** Inanda, 26°7'S, 28°3'E, *Wood* 440 (K). **Oriqualand:** N of Kaap, Barberton, 28°30'S, 23°45'E, *Thorneiop* 2206 (K). **Transvaal:** Crocodile River, beyond Brits, 25°42'S, 27°45'E, *Dyer & Merdoom* 3421 (K). **SUDAN.** **Bahr Al-Ghazal:** Wau, 7°46'N, 28°2'E, *McCintosh* 57k, 109k (K). **Equatoria:** Mount Nakbi between Yambio and Tambura, 5°0'N, 28°5'E, *Myers* 6997 (K). **Jungali:** 20 km N of Kongor, 7°10'N, 31°21'E, *Lock* 82/31 (K). **Kassala:** Metemma, Gallabat, 12°59'N, 36°0'E, *Schweinfurth* 179 (K). **North Darfur:** Darfur, Zalingei, 12°45'N, 23°29'E, *Lynes* 577 (BM). **South Kurdufan:** Kadugli, Nuba Mountains, 11°1'N, 29°41'E, *Bashir s.n.* (K). **SWAZILAND.** **Central:** Tshaneni, Swaziland Irrigation Scheme, 26°30'S, 31°30'E, *Parker* 1066 (ODU). **TANZANIA.** **Morogoro:** Kilosa, 6°50'S, 36°50'E, *Robertson* 676 (K). **Pwani:** Dar es Salaam, University College, 6°47'S, 39°18'E, *Mwasumbi* 10487 (K). **Rukwa:** Milepa, 8°3'S, 31°56'E, *Lea* LR20 (K). **Shinyanga:** Shinyanga, 3°45'S, 33°27'E, *Koritschoner* 2059 (K). **Tanga:** Handeni, 5°27'S, 28°2'E, *Ryae* H40/33 (K). **Ziwa:** Bukoba, Kakindu, 1°30'S, 31°10'E, *Haarer* 2307 (K). **Zanzibar:** Pemba, *Vaughan* 402, 1613, 1664 (K). **UGANDA.** **Busoga:** Bugerere, 0°45'N, 33°30'E, *Liebenberg* 1532 (K). **Eastern:** Elgon, Pemba, 1°10'N, 34°30'E, *Fames* 1905 (K). **Karamoja:** Honyili Ridge, 2°25'N, 33°40'E, *Wilson* 880 (K). **Nile:** Arua West Hill, 3°1'N, 31°10'E, *Hazel* 442 (K). **North Buganda:** Kakoge, Mengo, 1°0'N, 32°30'E, *Brown* 1732 (K). **Northern:** Lango, 2 km W of Ngunyoboke, 2°13'N, 32°34'E, *Lye* 4747

(K). **South Buganda:** Masaka Rest Camp, 0°20'S, 31°44'E, Chancellor 72 (K). **Southern:** Mbarara, 0°37'S, 30°39'E, Mailand 103 (K). **ZAMBIA.** **Copperbelt:** Kitwe, 12°54'S, 28°17'E, Fanshawe 10003 (K). **Eastern:** Luangwa Valley, 13°20'S, 31°35'E, Astle 4471, 4570 (K). **Luapula:** 5 km E of Lusaka, 15°28'S, 28°19'E, King 260 (K). **Shaba:** 3 mi. NE Ndola-Waterlogged Dambo, 12°58'S, 28°46'E, Draper 25 (K). **Southern:** Kafue River, at Lubalansuki, Namwala District, 15°50'S, 26°30'E, Mitchell 24/78 (K). **ZIMBABWE.** **Manicaland:** Melsetter, 6 mi. S Mutambara Village, 19°50'S, 32°47'E, Whiteside 107793 (K). **Mashonaland North:** Mazore, Shamva, near Lions Heads, 17°17'S, 31°30'E, Wild 599 (K). **Mashonaland South:** Makwiro, 18°0'S, 30°20'E, Smith 22673 (K). **Matabeleland North:** Urungwe National Park, 271 km from Salisbury on Chirundu Road, 18°57'S, 27°6'E, Philcox et al. 8796 (K). **Midlands:** Kwe Kwe, 18°55'S, 29°49'E, Musselman 124 (K).

**12. *Striga gastonii* A. Raynal, Bull. Mus. Natl. Hist. Nat., B, Adansonia, 3: 225–227. 1987.**  
TYPE: Chad. Danamadjji, Audru 1245 (holotype, P not seen; isotype, ALF not seen).

Perennial, 7 cm tall, succulent, tufted, glabrous, stem obtusely square. Leaves scale-like, 1–4 × 1–2 mm, lanceolate, appressed, longer than internodes, margin entire, veins obscure. All bracts similar, 3 × 1–2 mm, lanceolate, shorter than calyx. Spike dense, longer than vegetative stem. Calyx 5-ribbed, 3–5.5 mm long; tube 2–4 mm long; lobes 5, subequal, subulate, 1–2 mm long, shorter than tube. Corolla purple, tube 8 mm long, bent and expanded distally above calyx, sparsely glandular-pubescent, weakly bilabiate; lobes of lower lip 4 × 1–2 mm, obovate, upper lip 1 × 1–2 mm, emarginate, tips obtuse.

*Striga gastonii* is endemic to the lateritic plains in southern Chad and Central African Republic (Fig. 8) and is parasitic on *Lepidagathis* (Acanthaceae) species (Raynal-Roques, 1987). It is a small plant that resembles *S. gesnerioides* var. *minor* Santjuan in its overall appearance. The relationship of *S. gastonii* to the *S. gesnerioides* parasitizing *Lepidagathis* in India (e.g., Saldanha, 1963) should be investigated. *Striga gastonii* may be a host-specific strain of *S. gesnerioides*.

**13. *Striga gesnerioides* (Willd.) Vatke, Oesterr. Bot. Z. 25: 11. 1875. *Buchnera gesnerioides* Willd., Sp. Pl. 338. 1801. TYPE: India. *Koenig* s.n. (B 11573 not seen).**

*Buchnera orobanchoides* R. Br., Flora 2: 388, t.2. 1832.  
*Striga orobanchoides* (R. Br.) Benth., Companion Bot. Mag. 1: 361. 1836. TYPE: Ethiopia. *Salt* s.n. (holotype, BM!).  
*Striga orchidea* Hochst., Fl. XXIV. 43. 1841. TYPE: *Kotschy* 387 (holotype, K!).  
*Striga gesnerioides* var. *minor* Santapau, Kew Bull. 1948:

491. 1949. TYPE: *Santapau* 3072 (holotype, Blatter Herb., St. Xavier's College, Bombay not seen).

Annual or occasionally perennial, 12–30 cm tall, unbranched or branched from the base, usually succulent, drying black or brown, sparsely to densely hispid or pilose, stem terete or obtusely square. Leaves 4–10 × 1–3 mm, lanceolate, opposite/subopposite, scale-like, appressed, usually shorter than internodes, margin entire, veins obscure. All bracts similar, 2.5–6 × 2–3 mm, lanceolate, longer or shorter than calyx. Flowers opposite/subopposite in simple spikes, spike shorter or longer than vegetative stem. Calyx 5-ribbed, 4–9 mm long; tube 3–6 mm long; lobes 5, unequal or subequal, linear, deltate or lanceolate, 1–3 mm long, shorter than tube. Corolla creamy-white, blue, pink, purple, tube 8–14 mm long, bent and expanded distally above calyx, pubescent, with very few hairs or glandular hairs; lobes of lower lip 2–6 × 1.3–3 mm, obovate, spreading, upper lip 1–2 × 2–2.5 mm, indistinctly bilobed or emarginate, usually wider than long.

*Striga gesnerioides* is the most widely distributed of all witchweed species (Fig. 8) and is found between 33°10'N and 32°15'S, extending into Arabia and Asia. *Striga gesnerioides* was introduced into Florida in the United States (Sand et al., 1990) but has not become an agronomic problem.

*Striga gesnerioides* parasitizes only dicotyledons. Host selection, succulence, reduction of leaves, and tufted habit have been considered (see Boeshore, 1920) to represent a reduction in the direction of the holoparasitic Orobanchaceae. Thalouarn et al. (1991) found that *S. gesnerioides* has a weak capacity to fix CO<sub>2</sub>. These features of morphology, host selection, and physiology distinguish *S. gesnerioides* from other members of the genus.

*Selected specimens examined.* (All collections of *S. gesnerioides* are listed here. Under each strain we have listed specimens representative of those strains). ANGOLA. **Benguela:** Benguela, 12°36'S, 13°20'E, Gossweller 3540 (K). **Cunene:** Cahama, 16°19'S, 14°15'E, Pearson 2421 (K). **Luanda North:** Catete, 9°2'S, 13°42'E, Anonymous 9200 (K). **Lunda South:** Cazange, 11°2'S, 20°45'E, Gossweller 647 (K). BOTSWANA. **Central:** Boteti Delta Area, S of Mopipi, 20°50'S, 25°5'E, MacDonald 21 (K). **Desert:** 5 km NW of Bokspits, 26°51'S, 20°41'E, Skarpe 52 (K). **Ghanzi:** 13 mi. S of Ghanzi on Lobatsi Road, 21°50'S, 21°36'E, Brown 8282 (K). **Kweneng:** Artesia, 24°2'S, 23°26'E, Mitchison 1975 (K). NGAMILAND: Mboma Island, 19°12'S, 23°18'E, Smith 830 (K). **Southern:** Gaborone, Aedume, 24°42'S, 25°54'E, Hansen 3375 (K). BURKINA FASO. **Central:** Korsimoro, 30 km from Kaya, 12°0'N, 1°0'W, Raynal 22603 (K). **Northern:** 55 km from Ouahigouya, 13°0'N, 2°0'W, Raynal 22613 (K). BURUNDI. **Bujumbura:** Magari, 4°44'S, 29°18'E, Reekmans 9834 (K). **Rubindi:** Bururi, 3°56'S, 29°33'E, Reek-

**mans** 6926 (K). **Rusizi:** 14 km from Bujumbura, 3°20'S, 29°27'E, *Lewalle* 1474 (K). **CAMEROON.** **Northern:** 5 km W of Maroua, 10°37'N, 14°12'E, *de Wilde & de Wilder-Duyfjes* 29629 (K). **CENTRAL AFRICAN REPUBLIC.** **Bamingui-Bangoran:** Saline on Niao River, 2 km from confluence with Ground River, 8°42'N, 21°22'E, *Fay* 3354 (K). **CONGO.** **Shaba:** Lubanda, Kivu Province, 5°10'S, 26°38'E, *Germain* 5717 (K). **DEMOCRATIC REPUBLIC OF CONGO.** **Bandundu:** Tweya, 0°56'S, 18°58'E, *Robyn* 1980 (K). **Haut Zaire:** Faradje-Olru, 3°44'N, 29°43'E, *Lebrun* 3535 (MO). **Shaba:** Luvngu, 8°48'S, 25°17'E, *Linder* 1966 (K). **ETHIOPIA.** **Arsi:** Lake Langano, 7°45'N, 38°45'E, *Ash* 347 (K). **Gamo Gofa:** 12 km S of Arba Minch, 6°5'N, 37°40'E, *Gilbert et al.* 345 (K). **Gonder:** Tana, 12°30'N, 37°20'E, *Pichi-Sermolli* 1296 (K). **Ilubabor:** Gambela, 8°18'N, 34°37'E, *Ash* 557, 1089 (K). **Ogaden Harerge:** E of Grawa, 8°50'N, 42°25'E, *Burger* 2846 (K, US). **Shewa:** SW Shore of Lake Langano, 7°25'N, 38°25'E, *Gilbert & S.B.* 1078 (K). **Tigray:** Takaze Valley, 13°43'N, 39°11'E, *Gilbert & Getachew* 2958 (K). **GAMBIA.** **Eastern:** Kuntaur, 13°45'N, 14°40'W, *Ruxton* 53 (K). **GHANA.** **Northern:** Tamale, 9°20'N, 0°59'W, *Irvine* 4572, 4574 (K), *Hall* 2058 (K). **Upper:** Bolgatanga, 10°45'N, 0°55'W, *Parker* 2023 (K). **Volta:** Keta, Rest House, 5°55'N, 12°E, *Akpabla* 36 (K). **KENYA.** **Coast:** Arabuko, Kilifi, 3°40'S, 39°50'E, *Graham* 1675 (K). **Eastern:** Buna, 3°26'N, 39°54'E, *Gilbert* 13104 (K). **Nyanza:** Karungu, Homa Mount Foot, 0°49'S, 34°9'E, *Twiner* 6730 (K). **Rift Valley:** Chebloc, Eudo, Upper Kerio Valley, 0°22'S, 35°15'E, *Tweedie* 2448 (K). **MALAWI.** **Northern:** 19 mi. S of Chilumba, 11°0'S, 33°54'E, *Pawek* 9601 (K). **Southern:** Chiwembu, Mankhudue Hill, 15°38'S, 34°32'E, *Smith et al.* 5872 (K); Sombani Plateau, Mulanje Mountain, 16°10'S, 35°33'E, *Mouculie* 574 (K). **MALI.** **Koulakoro:** Sotuba, 12°50'N, 7°59'W, *Raynal* 22063 (K). **Massina:** Dioura, N Macina, 14°52'N, 5°16'W, *Davey s.n.* (K). **Segou:** between Dougouba and Markala, 13°0'N, 6°0'W, *Raynal* 22565 (K). **Sikasso:** 4 mi. from Kouialta, 12°30'N, 5°30'W, *Raynal* 22211 (K). **MOROCCO.** **Agadir:** Agadir, 30°30'N, 9°40'W, *Hettewy* 52, 179, 186 (K), *Lynes* 53 (BM). **Azilal:** near Demnate, 31°40'N, 7°3'W, *Balls* 3022 (K). **Beni-Mellal:** Kesbeh, 32°20'N, 6°20'W, *Lewalle* 10634 (BM). **Goulimime:** 1 km E of Tirhmi, 29°6'N, 9°5'W, *Miller et al.* 615 (BM). **Quaoqvizert:** Moyen Atlas Mountain, 33°10'N, 5°0'W, *Jahandiez* 149 (BM). **Southwest:** Oued Massa, between Agadir and Tiznit, 3°N, 9°35'W, *Davis* 53515 (BM). **MOZAMBIQUE.** Liben, *Kuntze* 94 (K). **Maputo:** Maputo (Lourenco Marques), 25°58'S, 32°32'E, *Balsinhas* 152 (K). **Niassa:** Marrupa, slope of Mountain Kuwanko, 15 km on road to Nungo, 13°8'S, 37°30'E, *Janssen* 8040 (K). **Sofala:** Chibabava, Lowu Buzi, 20°25'S, 33°35'E, *Swynnerton* 1413 (K). **Tete:** Tete, 16°13'S, 33°33'E, *Kirk* (K), *Peters* 8 (K). **NAMIBIA.** Along road to Ethosha Pans National Park, *Musselman & Visser* 4/2/86 (ODU). **Gobabis:** 10 km E of Gobabis, 22°28'S, 19°5'E, *Wanntop* 760 (K). **Great Namaqualand:** Jafelberg, 24°45'S, 17°15'E, *Smith* 26 (K). **Grootfontein:** 11 mi. from Otavi-Grootfontein on flat between mountains, 19°38'S, 17°30'E, *Tolken & Hardy* 919 (K). **Kaokoland:** Mopane, Ojivero, 18°0'S, 13°29'E, *Winter & Leistner* 5368 (K). **Karasburg:** Grot Karasberg, Naruda Sud Ravine, 27°29'S, 18°45'E, *Pearson* 8122 (K). **Karibib:** Damaraland Otisondu, 22°0'S, 15°56'E, *Seydel* 4491 (K, US). **Kavango:** bank of Okavango River at Runyu, 17°56'S, 19°44'E, *Winter & Marais* 4479 (K). **Keetmanshoop:** Keetmanshoop, 3 mi. W of Aroab, 26°51'S, 19°35'E, *Winter* 3372 (K). **Mariental:** E Aranos, Farm Mara on the Botswana border, 24°2'S, 19°10'E, *Vuuren & Giess* 1132 (K). **Omaruru:** Farm Okandjon North, 30 km NW of Omaruru, 21°26'S, 16°0'E, *Wanntop* 814 (K). **Windhoek:** 6 mi. E of Homestead on Nossob River, Gibeon, Farm Haruchas, 22°15'S, 17°48'E, *Leistner* 1817 (K). **Windhoek:** Avis Dam, 22°46'S, 17°7'E, *Liebenberg* A516 (K). **NIGER.** **Maradi:** 10 km NW of Maradi, 13°45'N, 7°0'E, *Parker* 2044 (K). **Niamey:** Niamey, 13°35'N, 2°10'E, *Olufsen* 479 (K). **NIGERIA.** **Bauchi:** 10 mi. from Gombe on Biu Road, 10°27'N, 11°14'E, *Daramola* 62346 (K). **Gongola:** Kilba Hills and Yola, 9°20'N, 12°35'E, *Dalziel* 300 (K). **Kwara:** Ilorin, 36 mi. from Ilorin, on Ilorin-Bacita Road, 8°30'N, 4°35'E, *Gbile et al.* 63394 (K). **Niger:** between Mokwa and Bida, 9°3'N, 5°30'E, *Lowe* A12 (K). **Oyo:** Ado Rock, 7°40'N, 5°15'E, *Keay* 25446 (K). **Plateau:** Jos Plateau, 9°40'N, 8°35'E, *Lely* 535 (K). **RUANDA.** **Southern:** Rumonge, along Lake Tanganyika, 3°58'S, 29°25'E, *Shantz & Tanner* 4196 (K). **SENEGAL.** Djembering, *Broadbent* 158 (K), *Farmer* 131 (K). **SIERRA LEONE.** **Western:** Buyabuya, 12°10'N, 7°50'W, *Elliott* 4279 (K). **SOMALIA.** **Bari:** Galgallo, 64 km SW of Bosaso, 10°58'N, 49°1'E, *Bally & Melville* 15841 (K). **Bay:** Buur Heybo (bur Eibi), 2°59'N, 44°16'E, *Thulin & Mahamed* 7088 (K). **Jubbada Hoose:** 16 km S of Badade, 1°4'S, 41°23'E, *Gillett et al.* 25136 (K). **Mogadisho:** Mogadishu International American School, 2°7'N, 45°32'E, *Kuchar* 15312 (K). **Nugaal:** Nugaal Valley, 8°5'N, 49°35'E, *Hemming & Watson* 30216 (K). **Sanaag:** Sugli, 10°58'N, 48°53'E, *Ballenetto* 277 (K). **Woqqooyi Gelbeed:** Borama, 10°2'N, 43°10'E, *Gilbert* 4723 (K). **SOUTH AFRICA.** **Cape:** bank of Kareiga River, Bathurst District, 32°15'S, 23°30'E, *Bayliss* 5212 (GH). **Natal:** Inomda, Amonopl Grass, *Wood* 4276 (K). **Orange Free State:** Kimberly, Dronfield, 28°54'S, 24°41'E, *William* 5975 (K). **Transvaal:** 9 mi. W of Krugersdorp on Farm Gladysvale, 26°5'S, 27°24'E, *Robin* 3839 (K). **SUDAN.** **Bahr Al-Gebel:** Terakekka, 5°25'N, 31°42'E, *N.D.H* 7361 (K). **Equatoria:** 10 km S of Bemmeira, 40 km S of Bor, 5°55'N, 31°36'E, *Lock* 82/44 (K). **Kassala:** Gedaref, Um Shagara, 14°2'N, 35°30'E, *Beshir* 151 (K). **Lake Province:** near Baidit, 35 km N of Bor by Kongor Road, 6°30'N, 31°30'E, *Lock* 81/77 (K). **North Kurdufan:** Barah, Hadeid Sherifi, 13°38'N, 30°36'E, *Wickens* 536 (K). **Red Sea:** Erkowit, 18°27'N, 36°32'E, *Jackson* 2722 (K). **South Darfur:** 13 km S Nyala, 11°58'N, 24°51'E, *Wickens* 3159 (K). **South Kurdufan:** G.T.Z. Farm, near Telo, 10°30'N, 30°E, *Musselman* 6174 (K, ODU). **Upper Nile:** Bahr el Jebel, Sheikh Jowbe, 8°45'N, 30°E, *Brown* 1644 (K), *Shellai* 837 (K). **TANZANIA.** 1.5 mi. NE Dangani, *Drummond & Hemsley* 3319 (K). **Arunsha:** Lake Tanganyika, Mbulu, 3°52'S, 35°33'E, *Richards* 4512 (K). **Dodoma:** Handeni, Kwa Mtono, 5°15'S, 35°25'E, *Archbold* 2776 (K). **Iringa:** 19 km E of Ibuguziwa Ferry, 7°55'S, 35°35'E, *Bjornstad* 1363 (K). **Kigoma:** Mahali Mountains, 6°0'S, 30°E, *Jefford et al.* 53 (K). **Kilimanjaro:** Mrau Rombo, E Kilimanjaro, 3°7'S, 37°30'E, *Haarer* 5127 (K). **Mara:** Lake Province, Musoma Town, 1°30'S, 33°48'E, *Tanner* 3996 (K). **Mbeya:** Galula, S Saza, 8°37'S, 33°2'E, *Carter et al.* 2479 (K). **Morogoro:** 10 km from base of Kitonga Gorge, 7°40'S, 36°15'E, *Lovett & Congdon* 1404 (K). **Mpana:** Kataki Plains, 6°20'S, 31°30'E, *Sanane* 1571 (K). **Mtwara:** Tunduru, 60 mi. from Masasi, 11°0'S, 39°0'E, *Richards* 17965 (K). **Rukwawa:** Rukwa Valley, 7°40'S, 32°20'E, *Pielou* 130 (K). **Ruvuma:** 6.5 km E of Gumbiro, 10°15'S, 35°52'E, *Milne-Redhead & Taylor* 10018 (K). **Shinyanga:** Shinyanga,

3°45'S, 33°27'E, *Burtt* 3770 (K). **Singinda:** Doma, 7 mi. on Bahi to Kilimatinde Road, 5°54'S, 35°14'E, *Bidgood et al.* 1204 (K). **Ufipa:** Kalambo Falls on Tanzania side of the river, 8°37'S, 31°15'E, *Richards* 12813 (K). **TOGO.** **Northern:** near Dapaon (Dapango), 10°53'N, 0°16'E, *Ern et al.* 1478 (K). **UGANDA.** **Busoga:** 2 mi. N of Nkondo-Kigindi Road, 0°50'N, 33°30'E, *Wood* 805 (K). **Eastern:** Soroti Rock, 1°43'N, 33°38'E, *Lye* 5379 (K). **Karamoja:** Lodoketeminit, near Moroto, 2°30'N, 34°36'E, *Kerfoot* 1313 (K). **Northern:** N of Mengo, 2°N, 32°28'E, *Brown* 2185 (K). **Western:** Butiaba, Lake Albert, 1°50'N, 31°20'E, *Thomas* 3458 (K). **ZAMBIA.** **Central:** Lunza Valley, Kafakula, 15°32'S, 27°0'E, *Richards* 4813 (K). **Lupula:** Mweru Lake, near Nchelengi, 8°45'S, 29°0'E, *Bullock* 3826 (K). **Lusaka:** Mazabuka, 4 mi. from Chirunda on Lusaka Road, 15°58'S, 28°48'E, *Drummond* 5503 (K). **Northern:** Abercorn, Cassawa Sands, 8°30'S, 31°17'E, *Richards* 10921 (K). **Southern:** Livingstone, 17°45'S, 25°52'E, *Rogers* 7023 (K). **Western:** Barotse Floodplains, Lizulu Island, 15°30'S, 23°30'E, *Verboom* 1039 (K). **ZIMBABWE.** **Binga District:** Mwenda Research Station, *Grosvenor* 144 (K). **Manicaland:** Umtali, Zimunya's Reserve, 18°50'S, 32°37'E, *Chase* 6853 (K). **Mashonaland North:** 20 km from Makuti on road to Kariba, Urungwe National Park, 16°50'S, 28°0'E, *Philcox & Leppard* 8718 (K). **Mashonaland South:** 311.5 km from Harare towards Chirundu, Zambezi Escarpment, 17°27'S, 30°19'E, *Philcox & Leppard* 8575 (K). **Matabeleland North:** 30 km from Makuti on road to Kariba, 17°0'S, 28°0'E, *Philcox et al.* 8748 (K). **Matabeleland South:** tributary of Nuanetsi River, near Mabalauta, Gona Game Reserve, 21°10'S, 30°15'E, *Ngoni* 160 (K).

Host-specific strains of *S. gesnerioides* differ in succulence, number of branches, pubescence and color of stem, bract length relative to calyx, length of calyx lobes relative to tube, and corolla color, pubescence, and size. Seed size is fairly uniform, 0.30 × 0.20 mm, for all strains of *S. gesnerioides* except for the *Euphorbia*-strain, which measures 0.39 × 0.24 mm.

#### Strains of *Striga gesnerioides*

While *S. gesnerioides* has evolved a strong host-specificity among its populations, these "morphs" do not have enough morphological differences to justify according them the level of species or subspecies (Mohamed, 1994). However, because of the potential for damage to crops and as a reference for further molecular work, we consider it important to record information on the host strains.

**Euphorbia (Euphorbiaceae) Strain:** Perennial, succulent, 12–25 cm tall, with 2 to 4 branches from base. Stem greenish yellow, drying brown, terete, thick, densely pilose with retrorse hairs. Internodes 1–3.5 cm long. All bracts similar, lanceolate, ciliate, incurved, shorter than calyx. Flowers opposite/subopposite in dense spikes, spike much longer than vegetative stem. Calyx 7–9 mm long, pubescent; lobes unequal, lanceolate, ciliate,  $\frac{1}{2}$  the

tube length. Corolla light blue or dark purple, tube 10–12 mm long, sparsely hairy, bent and expanded distally above calyx; lobes of lower lip obovate, obtuse; upper lip indistinctly bilobed, wider than long. Parasitic on arborescent *Euphorbia* species, i.e., *E. abyssinica* J. F. Gmel. (Musselman & Hopper, 1986).

*Selected specimens examined.* ETHIOPIA. **Welo:** 1 km NE of Desse, near Borkenna River, 11°7'N, 39°38'E, *Sutherland* 381 (MO). NAMIBIA. **Namaland:** Great Naquaaland, Great Karas Mountains, 25°10'S, 17°50'E, *Ortendahl* 487 (UPS). SOUTH AFRICA. **Cape:** Bathurst, bank of Kareiga River, 32°15'S, 23°30'E, *Bayliss* 5212 (GH). **Transvaal:** Albany, Douglas Heights, 26°12'S, 28°15'E, *Bayliss* 3158 (GH). SUDAN. **Red Sea:** Gabal Mawinja, *Musselman* 6180 (ODU).

**Indigofera (Fabaceae) Strain:** Plant annual, succulent, up to 16 cm tall, with 2 to 4 branches from base. Stem light green or purple, drying black, slender, obtusely square, furrowed, covered with dense, soft, divergent hairs. Internodes short, not exceeding 1 cm. All bracts similar, ciliate, shorter than calyx. Flowers opposite/subopposite in open spikes, spike longer than the vegetative stem. Calyx 5–7 mm long, ribs hispid, translucent between ribs; lobes equal or subequal, linear to lanceolate, ciliate, about the same length as tube. Corolla light pink to deep purple, tube 7–10 mm long, sparsely hairy, bent proximally within calyx teeth; lobes of lower lip obovate, obtuse, longitudinally rolled inside when dry; upper lip emarginate, as long as wide.

*Selected specimen examined.* SUDAN. **Blue Nile:** Abu Naama, Gabal Obeid, 12°15'N, 33°28'E, *Mohamed* 10/25/88 (ODU).

**Tephrosia (Fabaceae) Strain:** Plant annual, up to 23 cm tall, unbranched. Stem pale green, often red, drying brown, slender, obtusely square, furrowed, glabrate. Internodes 2.5 mm long. All bracts similar, lanceolate, ciliate, much shorter than calyx. Flowers opposite/subopposite forming somewhat open spikes, spike much longer than vegetative stem. Calyx 5–6.5 mm long, ribs hispid, translucent between ribs; lobes unequal, lanceolate, ciliate,  $< \frac{1}{2}$  tube length. Corolla pink to light purple, tube 8–10 mm long, sparsely pubescent, bent and expanded distally above calyx, lobes of lower lip linear-oval; upper lip emarginate, wider than long.

*Selected specimens examined.* BURKINA FASO. **Center East:** 5 km N of Tenkodogo, 11°50'N, 0°19'W, *Safa & Musselman* 10/9/87 (ODU). CAMEROON. **Northern:** 29 km S of Garoua, 9°5'N, 13°25'E, *Musselman et al.* 10/16/88 (ODU). NIGERIA. **Niger:** Mokwa, 9°18'N, 5°2'E, *Knepper* 9/19/89 (ODU).

**Vigna (Fabaceae) Strain:** Plant annual, succulent, 8–26 cm tall, branched from base. Stem green or purple, drying black, slender, obtusely square, furrowed, covered with sparse, soft, divergent hairs. Internodes 1–3 cm long. Bracts lanceolate, ciliate, incurved, lower ones as long as or longer than calyx, upper ones shorter than calyx. Flowers opposite forming somewhat dense spikes, spike usually shorter than vegetative stem. Calyx 6–9 mm long, ribs hispid, translucent between ribs; lobes unequal, acuminate, ciliate,  $\frac{1}{2}$  tube length. Corolla light pink or bluish, tube 8–10 mm long, glabrate, bent and expanded distally above calyx, lobes of lower lip obovate, spreading, slightly rolled inside; upper lip emarginate, as long as wide. Economically, this is the most important strain of *S. gesnerioides* because it attacks *Vigna unguiculata* (L.) Walp., cowpea, which is a staple in parts of Africa. Damage can be extensive to this and a few other crops (Parker & Riches, 1993).

*Selected specimens examined.* BURKINA FASO. Northern: 5 km S Gourcy, 13°9'N, 2°22'W, Safa & Musselman 10/10/87 (ODU). MALI. Koulikoro: Bamako, Sotuba, 12°50'N, 8°0'W, Musselman & Mohamed 10/1/88 (ODU). NIGER. Maradi: Maradi, Weed Research Organization Station, 13°32'N, 7°5'E, Parker 2264 (ODU). NIGERIA. Kano: near Kano Airport, 12°5'N, 8°35'E, Knepper 2/22/89, 8/24/89 (ODU).

**Ipomoea (Convolvulaceae) Strain:** Plant annual, succulent, 15–30 cm tall, usually unbranched. Stem purple, drying black, obtusely square, furrowed, somewhat thick, scabrous, very sparsely pilose, with soft, divergent, short hairs. Internodes 1–4 cm long. All bracts similar, slightly incurved, shorter than calyx. Flowers opposite in open spikes, spike as long as or longer than vegetative stem. Calyx 5–7 mm long, ribs hispid, translucent between ribs; lobes unequal, acuminate, ciliate, shorter than tube. Corolla white to light blue, tube 9–13 mm long, glabrate, bent and expanded distally above calyx, lobes of lower lip spreading, linear-oval; upper lip emarginate, about as long as wide.

*Selected specimens examined.* MALI. Koulikoro: 14 km N Koulokani, 13°38'N, 7°54'W, Musselman & Mohamed 10/6/88 (ODU). SENEGAL. Dakar: 10 km N Dakar, Musselman et al. 7066 (ODU). Diourbel: Kheur Serigne Sarr, 16 km E Diourbel, 14°39'N, 16°0'W, Musselman 7064 (ODU). SUDAN. South Kordofan: Balangya, 10°30'N, 29°48'E, Musselman 8/23/86 (ODU).

**Jacquemontia (Convolvulaceae) Strain:** Plant annual, succulent, up to 30 cm tall, unbranched or with 2–3 branches from base. Stem purple, drying black, slender, obtusely square, furrowed, sparsely to densely pilose with divergent hairs. Internodes

2–4 cm long. All bracts similar, incurved, ciliate, shorter than calyx. Flowers opposite in open spikes, spike very much longer than vegetative stem. Calyx 5–8 mm long, ribs hispid, translucent between ribs; lobes unequal, acuminate,  $< \frac{1}{2}$  tube length. Corolla pale blue or mauve, tube 8–11 cm long, glabrate, bent and expanded distally above calyx, lobes of lower lip obovate; upper lip widely obovate, indistinctly bilobed, wider than long.

*Selected specimens examined.* NIGER. Dosso: Kane, 224 km from Niamey on Maradi Road, 13°26'N, 3°53'E, Parker 2312 (ODU). Maradi: Maradi, 13°32'N, 7°5'E, Parker 2031 (K, ODU). NIGERIA. Plateau: Jos-Shendam Road, 9°50'N, 9°25'E, Knepper 8/26/89 (ODU). SOUTH AFRICA. Musselman 6269 (ODU).

**Merremia (Convolvulaceae) Strain:** Plant annual, succulent, up to 18 cm tall, unbranched or with 2–3 branches from base. Stem purple, drying black, slender, obtusely square, furrowed, sparsely to densely pilose, with divergent hairs. Internodes short, 2 cm long. All bracts similar, ciliate, incurved, shorter than calyx. Flowers imbricate forming dense spikes, spike very much longer than vegetative stem. Calyx 5–6 mm long, ribs hispid, translucent between ribs; lobes unequal, acuminate, about  $\frac{1}{2}$  tube length. Corolla purple, tube 8–10 mm long, glabrate, bent and expanded distally above calyx, lobes of lower lip linear-obovate, longitudinally rolled inside when dry; upper lip emarginate, wider than long.

*Selected specimens examined.* SUDAN. Bahr-Al-Ghazal: Djur River, Aymalar 27/51 (K). SUDAN. Blue Nile: Tozi, 12°54'N, 33°36'E, Mohamed 10/26/88 (ODU).

**Nicotiana (Solanaceae) Strain:** Plant annual, 15–22 cm tall, unbranched. Stem pale green, drying brown, slender, obtusely square, furrowed, covered with dense, short, soft, divergent hairs. Internodes 1–3 cm long. Bracts ciliate, incurved, lower ones as long as or longer than calyx, upper ones shorter than calyx. Flowers sparse, opposite in open spikes, spike usually shorter than vegetative stem. Calyx 4–7 mm long, ribs hispid, translucent between ribs; lobes unequal, acuminate, about  $\frac{1}{2}$  tube length. Corolla pink/mauve, tube 8–10 mm long, with few hairs or glandular hairs, bent and expanded distally above calyx, lobes of lower lip obovate; upper lip emarginate, as long as wide. Tobacco in Zimbabwe has been seriously damaged by this strain (Wild, 1948).

*Selected specimens examined.* NIGER. Maradi: Maradi, near the Weed Research Organization Station, 13°32'N, 7°5'E, Parker 1316, 2265 (ODU).




Figure 10. Distribution based on specimens of *Striga gracillima* (triangle) and *S. hermonthica* (circle).

- 14.** *Striga gracillima* H. Melchior, in Mildbraed, Notizbl. Bot. Gart. Berlin-Dahlem 11: 681–682. 1932. TYPE: Nyassa Hochland. Station Kyimbila, Stoltz 1304 (holotype, B?; isotypes, BM!, K!).

Annual, 25–55 cm tall, slender, usually unbranched, densely scabrous, hispid, hairs ascending, stem obtusely square. Leaves 18–45 × 2–5 mm, opposite, narrowly elliptic, longer than internodes, margin entire, veins obscure. Lower bracts 10–25 × 2 mm, leaf-like, longer than calyx, upper bracts lanceolate, as long as or just shorter than calyx. Flowers alternate, shortly pedicellate, axillary at the base of inflorescence, becoming imbricate above middle, forming strobilus-like inflorescences, raceme shorter than vegetative stem. Calyx 5-ribbed, 5–7 mm long; tube 3–4(–5) mm long; lobes 5, unequal (adaxial reduced to < 1 mm), deltate, 1–2 mm long, shorter than tube. Corolla pur-

ple, tube 6–8 mm long, bent and expanded proximally within calyx teeth, densely hairy, never glandular-pubescent; lobes of lower lip 1–3 × 1–1.5 mm, spreading, obovate, upper lip 4 × 5 mm, obovate, emarginate, as long as wide.

*Striga gracillima* is a poorly understood species and has mistakenly been identified as *S. aspera* (Hepper, 1984). It has been collected from Malawi, Tanzania, and Zambia between 8°41'S and 15°45' S (Fig. 10). *Striga gracillima* differs from *S. hermonthica* in having a slender, sparsely branched stem; narrower, erect, linear-elliptic leaves; shorter corolla tube with smaller lobes; and much smaller calyces than those of *S. hermonthica*. The glandular pubescent corolla and the equal calyx lobes are among the features that distinguish *S. aspera* from *S. hermonthica* and *S. gracillima*. *Striga gracillima* has not been reported to attack crops. Its host range has not been investigated.

*Selected specimens examined.* MALAWI. Central: Dedza, Dedza Mountain Forest, 14°22'S, 34°20'E, *Salubeni* 1330 (GH, K, MO); Dedza, near Kanjali, Chongoni Forest, 14°27'S, 34°18'E, *Salubeni* 1341, 1539 (K); Lilongwe, 25 mi. S of Lilongwe, 14°17'S, 33°47'E, *Pawek* 14332A (K); Kyimbila, *Stolz* 1304 (GH, MO); Nyassa Hochland, Station Kyimbila, *Stolz* 1304 (BM, K). SOUTHERN: Chiradzulu, Chiradzulu Mountain, 15°45'S, 34°18'E, *Brummit & Banda* 9838 (K). TANZANIA. MOROGORO: Mahenge, 8°41'S, 36°41'E, *Schlieben* 2108 (US). ZAMBIA. NORTHERN: Kasama, 40 km W of Kasama, 10°16'S, 30°54'E, *Philcox et al.* 10210 (K).

**15. *Striga hallaei* A. Raynal, Bull. Jard. Bot. Belg. 39: 374. 1969. TYPE: Gabon. Efout, Halle & Villiers 4914 (holotype, P!).**

Annual, 30–60 cm tall, erect, slender, unbranched, pilose, stem obtusely square. Leaves 40–70 × 12(–30) mm, opposite, elliptic, shortly petiolate, coarsely toothed, 3-veined, shorter than internodes. Lower bracts 8–20 × 5 mm, leaf-like, upper ones reduced, all bracts longer than calyx. Flowers alternate in open racemes, raceme shorter than vegetative stem. Calyx 5-ribbed, 6–7 mm long; tube 2 mm long; lobes 5, equal, linear to lanceolate, 4–5 mm long, twice as long as tube. Corolla salmon-pink, tube 22–25 mm long, bent and expanded distally above calyx, densely pilose; lobes of lower lip 17 × 30 mm, broadly round; upper lip 12 × 15 mm, emarginate, reflexed backward.

This species inhabits openings within rainforests (Raynal-Roques, 1969) of Gabon and Democratic Republic of Congo (Fig. 7). The large toothed and nerved leaves, long calyx lobes, salmon-pink corollas, and adaptation to wet habitats suggest a relationship between this species, *S. forbesii*, and *S. latericea*. Of these three species, only *S. hallaei* has a 5-ribbed calyx.

**16. *Striga hermonthica* (Del.) Benth., Companion Bot. Mag. 1: 365. 1836. *Buchnera hermonthica* Del., Fl. Egypte: 245, 34 fig. 3. 1813. TYPE: Egypt. *Delile* s.n. (holotype, P!).**

*Striga senegalensis* Benth., in DC. Prodr. 10: 502 (1846). TYPE: Senegal. *Heudelot* s.n. (syntype, K!).

Annual, to 90 cm tall, stiffly erect, branched from middle, scabrous, densely hispid, hairs ascending, stem square, furrowed. Leaves 15–80 (–110) × (4–)10–25(–40) mm, opposite, linear or narrowly elliptic, ascending or spreading, longer than internodes, margin entire, veins obscure. Lower bracts 12–50(–95) × 2–5(–10) mm, leaf-like, longer than calyx, upper bracts lanceolate, as long as or shorter than calyx. Flowers opposite, forming a lax spike at base of inflorescence, becoming dens-

er above middle, spike shorter than vegetative stem. Calyx 5-ribbed, 7–12 mm long; tube 5–10 mm long; lobes 5, unequal (adaxial reduced to < 1 mm), deltate, 2–4 mm long, shorter than tube. Corolla pink or mauve, rarely white, with ascending hairs, tube (13)16–20 mm long, bent and expanded proximally within calyx teeth, sparsely hairy, never glandular-pubescent; lobes of lower lip 7–15 × 3–6(–10) mm, obovate, upper lip 4–9 × 4–7 mm, emarginate.

*Striga hermonthica*, giant witchweed, is a serious pest of food crops (sorghum, maize, millet, rice) especially in the Sahel of Africa. In fact, in parts of the Sahel it causes more damage to millet and sorghum than any other factor, with the possible exception of birds. *Striga hermonthica* is not particular to soil type or pH and can be abundant on lighter soils in West Africa, as well as on heavy clay soils in East Africa.

Giant witchweed is an obligate outcrosser as is the related species, *S. aspera* (Musselman et al., 1991). Several insects are potential pollen vectors. Comparison of populations of *S. hermonthica* parasitizing similar hosts but growing in different geographical localities revealed a high level of morphological and floral variability. This high phenetic variability was accompanied by high genetic diversity (Bharathalakshmi et al., 1990; Olivier et al., 1998).

The sorghum- and millet-strains of *S. hermonthica* have similar vegetative characteristics but different corolla sizes and shades. The millet-strain has a smaller dull pink corolla and is common on sandy soil, whereas the sorghum-strain has a larger pink corolla and is associated with clay soils. Although host-specificity reduces the host range of the parasite, it may lead to successful attachment and better adaptability to a given host under specific conditions. Such floral differences could be expected to reduce cross pollination between the two strains when sympatric. Breeders developing grains for tolerance to the parasite need to be aware of genetic interchange in this wide-ranging species.

*Striga hermonthica* extends from Senegal to Ethiopia with southern limits in the Democratic Republic of Congo and Tanzania, between 22°3'N and 9°30'S (Fig. 10). There are a few collections from Angola and Namibia. Collections from the Nile Delta and parts of Yemen likely represent introductions with contaminated seed, a well-documented means of dispersal (Berner et al., 1992). *Striga hermonthica* has not been found in native grasslands except in the Nuba Mountains of the Sudan Republic (Musselman & Hepper, 1986).

South of 9°30'N, *S. hermonthica* is less frequent and smaller in size. *Striga hermonthica*, *S. asiatica*, and *S. gesnerioides* are the only species found north of the Sahara (Figs. 4, 8, 10).

In any large population of *S. hermonthica* it is possible to find albino plants. These are plants totally lacking chlorophyll yet capable of flowering because of their parasitic nature (Musselman & Hepper, 1986).


*Selected specimens examined.* ANGOLA. Luanda: Catete, 9°2'S, 13°42'E, Gossweiler 199, 14219 (K). BURKINA FASO. Central: 60 km E of Ouagadougou, 12°35'N, 0°18'W, Safa & Musselman 10/9/87 (ODU). Northern: Bogoya, 5 km NW of Ouahigouya, 13°38'N, 2°27'W, Raynal 22263 (K). Volta: Boromo, 11°50'N, 2°55'W, Raynal 22600 (K). CAMEROON. Center South: near Djikainke, 4 mi. from Foumban, 4°43'N, 10°55'E, Leeuwenberg 10189 (K). Northern: Adamawa, near Gangumi, 7°0'N, 13°0'E, Latilo & Daramola 28792 (K). CENTRAL AFRICAN REPUBLIC. Ouham: Bossango Region, 6°30'N, 17°20'E, Fay & Doka 5070 (K). DEMOCRATIC REPUBLIC OF CONGO. Equateur: Libenge, 3°39'N, 18°38'E, Lebrun 1759 (GH). Haba: Niambebe, 2°14'S, 27°44'E, Steyagi s.n. (US). Haut Zaire: between Mahagi and Mahagi Port, 2°16'N, 31°7'E, Boutique 308 (K). Kivu: between Gabiro and Kidebe, 1°30'S, 30°25'E, Lebrun 9631 (K). Shaba: Kundelungu, 9°30'S, 27°45'E, Kassner exped. 2786 (K). DJIBOUTI. Eastern: Arta Road, 11°32'N, 43°9'E, Curle 98 (K). EGYPT. Cairo: S of Al-Jizah, Helwan, 29°58'N, 31°16'E, Boland 743 (K). Delta: Rodah, 30°15'N, 31°15'E, Schweinfurth 1880 (US). Red Sea: Al-Fashn, 28°27'N, 30°55'E, Faad 1881 (K), Radjil 594 (GH). ERITREA. Southern: 4 km S of Adi Kuala, 14°38'N, 38°50'E, Ryding et al. 1548 (K). Teseney: 15°6'N, 36°40'E, Pichi-Sermoli 1295 (K). ETHIOPIA. Gojam: Shoa, 10°5'N, 38°16'E, Meyer s.n. (K). Ilubabor: Blue Nile, Gore, 8°9'N, 35°32'E, Parker E22 (K). Kefo: Omo National Park, 8°25'N, 37°15'E, Ash 3648 (K). Ogaden Harerge: at 8°44'N, 40°30'E, Puff et al. 820922 (K). Shewa: N Addis Ababa, 9°12'N, 38°42'E, Meyer 7594 (US). Tigray: Enda Sellassie, W of Axum, 14°10'N, 38°11'E, Scott 217 (K). Welega: Condor, 9°50'N, 36°31'E, Massey 47 (K). GAMBIA. Central: Sapu, 13°0'N, 15°0'W, Terry s.n. (K), Saunders 60, 62 (K), Brookes 19 (K). GHANA. Northern: 2.3 mi. NW of Bunkpurugu, 10°32'N, 0°3'E, Innes 30748 (K). Upper: Bongo, near Bolgatanga, 10°50'N, 0°50'W, Hossain & Agyakwah 37855 (K). GUINEA BISSAU. Central: Bafata, Arredores Region, 12°12'N, 14°42'W, Espírito Santo 2796 (K). Eastern: Gahee, Canquelifa, 12°40'N, 13°50'W, Espírito Santo 2960 (K). IVORY COAST. Bouna: Varale, 60 km N of Bouna, 9°39'N, 3°6'W, Geerling & Bokdam 934 (K). Boundiali: Zaguinaso, 64 km N of Boundiali, 10°2'N, 6°32'W, Riches 249 (K). Firkesedougou: 5 km S of Firkesedougou, 9°36'N, 5°8'W, Riches 226 (K). KENYA. Nyanza: 3 mi. S of Homa Bay, 0°34'S, 34°22'E, Bogdan 4535 (K). MALI. Kayes: Dionkoulane Fields, Yelimane Circle, 15°8'N, 10°34'W, Yeafes 88 (K). Koulikoro: Sekeson, 7 km from Ouelessebougou, 90 km S of Bamako, Raynal 22194 (K). Segou: 7 km from Diabaly, 14°0'N, 6°0'W, Raynal 22573 (K). SIERRA LEONE. Cunene: Oshikango, 17°9'S, 16°10'E, Rodin 9379 (K). Owambo: Olukonda, 18°0'S, 16°3'E, Reed 14

(K). NIGER. Maradi: 2 km W Maradi on Niamey Road, 13°32'N, 7°4'E, Parker 2316, 2317 (ODU). Niamey: Niamey, Galome, 13°35'N, 2°10'W, Hagerup 478 (K), Barter 136, 1821 (GH). NIGERIA. Bauchi: 1 mi. N of Naraguta, 10°40'N, 8°55'E, Hepper 1162 (K). Bornu: Biu, 10°35'N, 12°10'E, Noble 30, 31 (K). Gongola: 10 mi. W of Gombi, 10°12'N, 12°41'E, Lowe 1671 (K). Kaduna: between Zaria and Kano, site of new tollgate, 11°30'N, 8°10'E, Lowe 4828 (K). Kano: Gaya, near Kano, 11°55'N, 9°2'E, Shorland 179 (K). Kwara: Lokoja, 7°50'N, 6°40'E, Parsons L79 (K). Niger: Kontajora, 10°25'N, 5°25'E, Dalziel 171 (K). Ogun: Ibadan, Botany Department, 7°10'N, 3°50'E, Lowe 2760 (K). Plateau: Jos-Shendam Road, 9°50'N, 9°25'E, Knepper 8/26/89 (ODU). Sokoto: Guadabawa, 13°8'N, 5°15'E, Ryan 58 (K). SENEGAL. Oriental: Bakel, 14°55'N, 12°29'W, Hepper 3721 (K). SUDAN. Al-Jazirah: Madani, Barakat, 14°20'N, 33°48'E, Musselman 6139 (ODU, K). Bahr Al-Ghazal: Wau, 7°46'N, 28°2'E, MacCintosh 103 (K). Blue Nile: Abu Naama, 12°49'N, 33°40'E, Mohamed 10/27/88 (ODU). Kassala: Al-Qadarif, 14°4'E, 35°24'E, Schweinfurth 595 (K). Khartoum: at 15°38'N, 32°32'E, Sawyer 13698, 13699 (K). North Darfur: Gabal Kurdufan, 13°23'N, 23°32'E, Wickens 628 (K). Gabal Abu Sinun, 12°52'N, 23°30'E, Wickens 309 (K). North Kurdufan: El Obeid, Khartaget, 13°13'N, 30°32'E, Musselman 6162 (ODU). Northern: Wadi Halfa, 1 km NW Debeira Station, 22°3'N, 31°23'E, Petterson 16819 (K). South Kurdufan: near Kadugli, 11°1'N, 29°41'E, Musselman 6155 (K). Upper Nile: Sobat, 8°55'N, 32°25'E, Petherick 1863 (K). West Kurdufan: El Greo, Pfund 679 (K). TANZANIA. Mara: Lake Province, Ikoma, Musoma, 2°2'S, 34°35'E, Tanner 4142 (K). Shinyanga: near Shinyanga, 3°40'S, 33°27'E, Box 293 (K). Ziwa Magharibi: West Lake Province, Biharamulo, 2°37'S, 31°19'E, Tanner 5274 (K). CHAD. Chari-Baguirmi: Ba Ill, 10°32'N, 16°27'E, Audry B2 (K). TOGO. Northern: 7 km N of Mango, 10°26'N, 0°17'E, Pash 7916 (K). UGANDA. Eastern: Amuria, 2°4'N, 33°37'E, Willems 30 (K). North Buganda: Sempayo-Kiboga Road, Semlilei Flats, 0°40'N, 31°45'E, Liebenberg 1029 (K). Northern: between Nimule and Gondokoro, along Bahr El Gabal, 3°10'N, 32°5'E, Mearns 3005 (GH). Southern: W Ankole, 0°30'S, 30°18'E, Dawe 444 (K). Western: Bunyoro, 1°38'N, 31°30'E, Fyffe 32, 164 (K).

## 17. *Striga hirsuta* Benth., in DC. Prodr. 10: 502. 1846. TYPE: China. *Loureiro* s.n. (holotype, Pl!). Figure 2.

*Striga pusilla* Hochst, in DC. Prodr. 10: 503. 1846. TYPE: Ethiopia. Gafata, Schimper 1209 (holotype, K!).

Annual, small, 2–6(–20) cm tall, slender, unbranched or branched from middle, sparsely hispid to ciliate, hairs ascending, stem obtusely square, furrowed. Leaves reduced, 3–15 × 0.5–1 mm, opposite/subopposite, linear, shorter than internodes, margin entire, veins obscure. Lower bracts 6–15 × 1 mm, narrowly lanceolate, longer than calyx, upper bracts lanceolate, shorter than calyx. Lower surface of leaves and bracts with a row of hispid hairs along margins and midrib, upper surface hispid only along margins. Flowers alternate in open racemes, raceme longer than vegetative stem. Calyx

Figure 11. Distribution based on specimens of *Striga hirsuta* (circle).

10-ribbed, 4–5.5 mm long; tube 2–3 mm long; lobes 5, equal, subulate, 1.5–2.5 mm long, as long as tube. Corolla dark red with yellow throat, 8–10 (–12) mm long, bent and expanded distally above calyx, sparsely glandular-pubescent; lobes of lower lip 2–3(–5) × 1–2(–3) mm, obovate, upper lip 1–2 × 2–4 mm, emarginate.

*Striga hirsuta* is frequent in the Sudanian savanna. Its range extends from Mali to Ethiopia and south into Kenya, Tanzania, Malawi, and Mozambique between 13°N and 26°S (Fig. 11). *Striga hirsuta* and *S. lutea* may rarely attack crop plants such as sorghum and maize, but no noticeable damage occurs.

*Striga hirsuta* is typically a small plant, up to 6 cm, that emerges, flowers, and matures in a fairly short time. It flowers sooner after the commencement of the rains than the most damaging species, *S. hermonthica*. *Striga hirsuta* is less hispid than *S. asiatica* and *S. lutea*; however, it is more scabrous than both species. See comments under *S. asiatica*. Musselman and Hepper (1986) included *S. hirsuta* and *S. lutea* in *S. asiatica*. Further work is needed to elucidate the relationship of this species to *S. asiatica*.

*Selected specimens examined.* ANGOLA. Benguella: Benguella, 12°36'S, 13°20'E, Gossweiler 2916 (K). Huila:

Couceiro, Vila Parire, 15°10'S, 13°30'E, Henriquez 853 (K). BURKINA FASO. Eastern: Fada-N'gourma, 12°12'N, 0°25'E, Safa & Musselman 10/25/87 (ODU). BURUNDI. Northern: Bubanza, Imbo, 3°10'S, 29°58'E, Reekmans 10314 (K). CAMEROON. Eastern: Batouri, 4°30'N, 14°30'E, Taylor 1 (K). North-Western: Nanpe, 8 km W of Bangante, 8°5'N, 10°36'E, de Wilde & de Wilde-Duyffes 2397 (K). Northern: Jangla, foot of Vogel Peak, 8°22'N, 11°55'E, Hepper 2745 (K). Western: Mbaw Plain, 6°15'N, 11°0'E, Brun 404 (K). CENTRAL AFRICAN REPUBLIC. Bamingui-Bangoran: 2 km E of Camp Koumbala, 8°29'N, 21°24'E, Fay 2867 (K). Nana-Mambere: Bouar, 6°0'N, 15°35'E, Mildraed 9501 (K). Ouham-Pende: near Nana-Bakassa River, 7°0'N, 17°0'E, Fay 5068 (K). CONGO. Kinshasa: at 4°30'S, 15°30'E, Woolhouse 51/61/22 (K). DEMOCRATIC REPUBLIC OF CONGO. Bas-Zaire: Boma, 5°48'S, 13°0'E, Dacremont 128, 129 (K). Haut Zaire: Ibila-Bofea, 2°20'N, 26°0'E, Eviard 2231 (K). Katanga: Kayuba, 9°45'S, 27°58'E, de Witte 3332 (K). Kivu: Beni, 0°30'N, 29°28'E, Gille 39 (K). Shaba: Pweto, 8°25'S, 28°51'E, Robyn 1971 (K). ETHIOPIA. Bale: Dallo, 6°22'N, 39°50'E, Mooney 8427 (K). Gamo Gofa: 8 km S of Arba Minch, 5°58'N, 37°36'E, Gilbert et al. 320 (K). Gojam: Beles Farm, W of Gojam, 10°50'N, 35°47'E, Parker 4074 (K). Ilubabor: Gambela, 8°18'N, 34°37'E, Ash 518 (K). Ogaden Harerge: Awash National Park, 8°55'N, 39°55'E, Gilbert 1663 (K). Tigray: Gafata, 12°55'N, 39°40'E, Anonymous 1842 (GH, US). Welega: 9°3'N, 36°11'E, Gilbert & Thulin 708 (K). KENYA. Central: Zawadi, 0°24'S, 36°59'E, Faden et al. 74/572 (K). Coast: 30 km W of Buru, 3°45'S, 38°18'E, Durham 26/2/24 (K). Eastern: Emali Hill, 2°3'S, 37°27'E, Someren 250 (K).

**Nairobi:** 1°17'S, 36°48'E, *Johnstone 1551* (K), *Red 1424* (K). **Rift Valley:** 2 km S of Lake Naivasha, 0°50'S, 36°14'E, *Gilbert 4840* (K). **LIBERIA.** **Maryland:** Harper, 4°25'N, 7°43'W, *Baldwin 5988* (US). **MALAWI.** **Central:** Fort Manning, Bua River, 13°47'S, 32°58'E, *Robson 1083* (K). **Northern:** 37 mi. SW Mzuzu, 11°47'S, 33°38'E, *Pawek 10942* (K). **Southern:** Blantyre, 2 mi. N of Limbe, 15°49'S, 35°7'E, *Brummit 8428* (K). **MALI.** **Koulakoro:** Baguineda, 22 km N of Bamako, 12°51'N, 7°46'W, *Musselman & Mohamed 10/8/88* (ODU). **MOZAMBIQUE.** **Maputo:** Inhaca Island, 23 km E of Maputo (Lourenco Marques), 26°1'S, 32°57'E, *Mogg 30317* (K). **NIGER.** *Barter 530* (ODU). **NIGERIA.** **Plateau:** Jos-Plateau, 9°50'N, 8°40'E, *Knepper 8/25/89* (ODU). **RWANDA.** **Kaduna:** Tegina, *Musselman & Mansfield 5531* (ODU). **Northern:** Ruhengeri, 1°30'S, 29°36'E, *Reekmans 4257* (K). **SEYCHELLES ISLAND.** Vallee de Mai, *Belle 196* (GH). **SOMALIA.** **Coast:** 5 km S of Goba, 0°47'N, 41°41'E, *Gillet et al. 5045* (K). **SUDAN.** **Bahr Al-Ghazal:** Djur River, 8°40'N, 28°35'E, *Ayhuer GA27151* (K). **Darfur:** Kalwe, *Hynes 478* (K). **Equatoria:** Nakbi, between Yambio and Tambio, 5°1'N, 28°5'E, *Myers 7000* (K). **Jungali:** near Bairut, 45 km N of Bor, 6°32'N, 31°36'E, *Lock 81/247* (K). **North Darfur:** Gabal marra, Kibbi, 12°1'N, 23°32'E, *Wickens 2241* (K). **South Kurufutan:** between Belatama and Kadugli, 11°30'N, 29°30'E, *Sherif 3952* (K). **Upper Nile:** Adok, 8°10'N, 30°18'E, *Petherick 1862* (K). **TANZANIA.** **Arusha:** Mbulu, Jebogo Hill, Mgati Plains, 4°5'S, 35°30'E, *Burtt 2105* (K). **Iringa:** 80 km N of Muhanga, 8°10'S, 36°0'E, *Bidgood et al. 65* (K). **Kigoma:** Livingstone College, 5°30'S, 30°0'E, *Verdcourt 3390* (K). **Kilamanjaro:** Ngungu, Shimba Hills, 4°28'S, 38°4'E, *Jackson 415* (K). **Mafia:** Chungaruma, *Greenway 5371* (K). **Magharibi:** Biharimulo, on Nyamiremb Road, 2°39'S, 31°23'E, *Balslev 447* (MO). **Mbeya:** Chimala, 85 km E of Mbeya, 8°52'S, 34°0'E, *Bally & Carter 16448* (K). **Morogoro:** 25 mi. NE of Morogoro, 6°30'S, 37°40'E, *Bidgood et al. 429* (K). **Rukwa:** Tumba, 7°30'S, 31°45'E, *Bullock B14* (K). **Ruvuma:** 10 km SW of Songea, 10°44'S, 35°33'E, *Milner-Redhead & Taylor 882* (K). **Shinyanga:** near Shinyanga, 3°45'S, 33°27'E, *Box 291* (K). **Tanga:** Handeni, Nyongolo Hill, 5°27'S, 38°2'E, *Mgaza 434* (K). **Zanzibar:** *Kirk 2179* (K). **Ziwa Magharibi:** Muhutwe, Bukoba, 1°35'S, 31°45'E, *Panayotis 98* (K). **UGANDA.** 2 mi. S of Akilok, *Langdale-Brown 2421* (K). **Central:** Entebbe, 0°4'N, 32°28'E, *Brown 21, 238* (K). **Karamoja:** SE of Karamoja, NE of Kocemaluk, 2°18'N, 34°18'E, *Hudson 296* (K). **Nile:** Koboko, 3°25'N, 31°0'E, *Looe P1048* (K). **North Buganda:** 10 km NW of Katera, 0°46'N, 31°58'E, *Lye s.n.* (K). **South Buganda:** Lake Kauonje, Mountain Masak, 0°21'S, 31°45'E, *Rose 10253* (K). **Southern:** Kambugo, Kisizi, 0°plain 45'S, 29°50'E, *Purseglowe 2424* (K). **ZAMBIA.** **Eastern:** Fort Jameson, 13°37'S, 32°44'E, *Bush 23* (K). **Lusaka:** 1.5 km N of Marangora, Zambezi Escarpment, Urungwe National Park, 15°45'S, 29°0'E, *Philcox et al. 8563, 8610* (K). **North-Western:** Mwinilunga, near Matonchi, 11°41'S, 24°24'E, *Townsend 170* (K). **Northern:** 16 km E of Kasama, Nambkung Ridge, 10°16'S, 31°15'E, *Philcox et al. 10190* (K). **Western:** Sesheke, 17°30'S, 24°50'E, *Gairdner 412* (K). **ZIMBABWE.** **Inyanga:** Inyanga, 18°15'S, 32°42'E, *Norlindh & Weimarck 4956* (K). **Manicaland:** River Tandai, 19°32'S, 32°40'E, *Myres 294* (K). **Mashonaland South:** Salisbury Experimental Station, 17°48'S, 31°1'E, *Wild 5718* (K). **Matabeleland North:** near Wankie Prison, 18°22'S, 26°30'E, *Raymond 319* (K). **Matabeleland South:** Se-

lukwe, near Surprise Mine, 19°38'S, 30°0'E, *Cecil 128* (K). **Victoria:** Kyle National Park, 20°19'S, 31°1'E, *Philcox & Leppard 8870* (K).

**18. *Striga junodii* Schinz., Mem. Herb. Boiss 10: 62. 1896–1900. TYPE: Mozambique. Delagoa Bay, *Junod 183* (holotype, G not seen).**

Perennial, 30–60 cm tall, stiffly erect, unbranched, scabrous, hispid, stem winged. Leaves 10–25 × 1–2 mm, linear, opposite at base of stem, alternate above, shorter than internodes, margin entire, veins obscure. Lower and upper bracts similar, 2–5(–12) × 1 mm, linear, shorter than calyx. Flowers alternate in open racemes, raceme much shorter than vegetative stem. Calyx 15–18-ribbed, 11–15 mm long; tube 7–9 mm long; lobes 5, equal, linear to lanceolate, 4–7 mm long, slightly shorter than tube. Corolla purple, tube 20–22 mm long, bent and expanded distally above calyx, densely pubescent; lobes of lower lip 8–11 × 4–5 mm, spreading, obovate, slightly notched; upper lip 5–8 × 8–11 mm, emarginate.

*Striga junodii* is represented by only a few collections from swampy grounds in southern Mozambique and northeastern South Africa between 24°S and 26°S (Fig. 12). It does not extend north into Mozambique where the similar *S. pubiflora* occurs (Fig. 15). The two species may have different ecological requirements. *Striga junodii* is also morphologically similar to the Asian species, *S. mansuria*; however, *S. junodii* has a characteristic notch on the lower corolla lobes. The current status of *S. junodii* remains unknown. We are unaware of extant populations. Collectors in Mozambique and South Africa should be aware of its presence.

*Selected specimens examined.* **MOZAMBIQUE.** **Inhambane:** Inhambane, 23°54'S, 35°30'E, *Anonymous 12092* (MO, S, UPS). **Maputo:** Lourenco Marques, Rikatla, 25°58'S, 32°32'E, *Junod 276* (BRE). **SOUTH AFRICA.** **Transvaal:** Muzi Swamp, 2 km from Phelendaba Crossing on Mbazwana Road, 23°55'S, 31°13'E, *Germishuizen 3534* (K); Nelspruit, 25°30'S, 30°59'E, *Prosser 1784* (BRE, K), *Britton 4778* (BRE).

**19. *Striga klingii* (Engl.) Skan, Fl. Trop. Afr. 4(2): 413. 1906. *Buchnera klingii* Engl., Bot. Jahrb. 18: 69. 1893. TYPE: Togo. *Büttner 239/221* (B?).**

Annual, to 86 cm tall, coarse, stiffly erect, often 3-branched just below inflorescence, scabrous, hispid, stem obtusely square. Leaves 20–40(–70) × 4–7(–12) mm, opposite, narrowly elliptic, tips obtuse, 3-veined, coarsely toothed, shorter than internodes. Lower bracts 10–20 × 2–4 mm, lanceolate, longer than calyx, upper bracts lanceolate, as long




Figure 12. Distribution based on specimens of *Striga junodii* (triangle) and *S. macrantha* (circle).

as calyx. Flowers imbricate forming very compact spikes, spike much shorter than vegetative stem. Calyx 12–15-ribbed, 8–10 mm long; tube 5–6 mm long; lobes 4 equal or 5 unequal, linear, 3–5 mm long, slightly shorter than tube. Corolla pink, tube 10–13 mm long, bent and expanded proximally within calyx teeth, pubescent or glandular-pubescent; lobes of lower lip 4–7 × 2–4 mm, spreading, oval, upper lip 2–3 × 3–4 mm, obovate, emarginate.

A very distinct witchweed, stiffly erect with bright pink corollas, *S. klingii* is usually found in small populations parasitizing grasses in the wet savannas of west and central Africa. Its range extends from Senegal to southern Sudan between 4°0'N and 15°0'N with disjunct localities in Tanzania and Malawi (Fig. 13). *Striga klingii* is similar to *S. macrantha*; both species have three terminal branches just below the inflorescence. However, *S. klingii* has pink flowers whereas *S. macrantha* has white flowers and glandular hairs.

The vestiture of *S. klingii* is unique, as noted by Neumann et al. (1996) and Musselman (unpublished), with curved hairs that have very broad shield-like bases.

*Selected specimens examined.* CAMEROON. **North-**  
**ern:** 10 km S of Ngaoundere, Adamaua Plateau, 7°16'N,

13°35'E, Breteler 258 (K); 7°20'N, 13°32'E, Musselman et al. s.n. (K); Ngaouyangga, 29 km N of Dogba, 7°20'N, 13°32'E, Musselman 10/16/88 (ODU). CENTRAL AFRI-  
CAN REPUBLIC. **Bamingui Bangoran:** Camp Koum-  
bala, 8°29'N, 21°13'E, Fay 6021 (K, MO). GHANA.  
**Brong Ahafo:** N of Krachi, 7°55'N, 0°1'W, Morton 6292 (K). GUINEA. **Northern:** Youkounkoun, Sambailo,  
12°40'N, 13°22'W, Adam 13884 (MO). **Southern:** Ma-  
mou, 10°27'N, 12°2'W, Shuell 6785 (K). GUINEA BIS-  
SAU. **Eastern:** Madina, 11°55'N, 14°5'W, Espírito Santo 3754 (K). **Northern:** Bafata, Bambadine, 12°12'N,  
14°42'W, Espírito Santo 3808 (K, MO); Farim, 12°35'N,  
15°10'W, Espírito Santo 3622 (K, MO); Piche, 12°25'N,  
13°55'W, Pereira 2328, 2368 (K); São Domingos,  
12°30'N, 16°8'W, Espírito Santo 3668 (K, MO). MALAWI.  
Kyimbila, N Lake Nyasa, Stoltz 2500 (GH). NIGER. Barter  
158 (GH). NIGERIA. **Gongola:** Vogel Peak Area, 8°22'N,  
11°55'E, Hepper 1397 (K). **Kaduna:** Zaria, Burra,  
11°0'N, 9°0'E, Harris 10 (K). **Niger:** near Mokwa,  
9°18'N, 5°2'E, Musselman & Mansfield 5533 (ODU).  
**North Nigeria:** Lely 680 (K), Roger 758 (K). SENEGAL.  
**Oriental:** Kedougou, 12°44'N, 12°15'W, Adam 26881 (MO). SIERRA LEONE. **Eastern:** between Kainkordu  
and Sefadu, 8°34'N, 10°54'W, Morton 2527 (K). **North-  
ern:** Fiedwont, 8°45'N, 11°28'W, Towa 8166 (K); Seku-  
rela, Kabala, 9°33'N, 11°34'W, Adam 21995 (GH, MO);  
Tingi Mountains, 8°56'N, 10°48'W, Morton & Gledhill  
3035 (K). SUDAN. **Equatoria:** Ibikes, 40 mi. NW of Yei,  
4°17'N, 30°32'E, Meyers 7867 (GH).

**20. *Striga latericea*** Vatke, Linnaea 43: 311.  
1882. TYPE: Somalia. (?) *Anonymous* s.n. (ho-  
lotype, K!).




Figure 13. Distribution based on specimens of *Striga klingii* (circle) and *S. latericea* (triangle).

*Rhamphicarpa stricta* Engl., Abh. Preuss. Akad. Wiss. 24–39. 1894. TYPE: Tanzania. Holst 2532 (holotype, K!).

Perennial, to 60 cm tall, stiffly erect, unbranched or with 2 to 3 branches from base, pubescent, stem square, furrowed. Leaves 20–35 × 4–6 mm, opposite, linear to lanceolate, 3-veined, shorter than internodes, margins slightly toothed. Lower bracts 20–33 × 2–6 mm, linear, often shorter than calyx, upper bracts lanceolate, shorter than calyx. Flowers opposite in racemes, raceme lax at its base, dense above middle, shorter than vegetative stem. Calyx at least 15-ribbed, 10–24 mm long; tube 10–15 mm long; lobes 5, unequal, broadly lanceolate, 5–15 mm long, longer than tube, lobes of lower calyces about 2 cm long, 3-veined. Corolla salmon-pink, tube 22–30 mm long, bent and expanded distally above calyx, densely glandular-pubescent; lobes of lower lip 10–15(–20) × 7–9 mm, obovate, upper lip 7–13 × 13–14 mm, emarginate.

Unlike *S. forbesii*, which is widely distributed in Africa, *S. latericea* (its closest putative relative) is confined to East Africa. It is found in swampy areas in Somalia, Ethiopia, Kenya, and Tanzania between 9°37'N and 4°40'S (Fig. 13) where it is sympatric with *S. forbesii*. We consider these two species distinct (also see comment under *S. forbesii*). Parker

(1988) has shown that *S. latericea* is a perennial that produces aerial shoots from adventitious buds developing from the roots, unique in the genus (see also Raynal-Roques, 1993, for a discussion of roots in this species). The aerial parts also have a perennial habit with new shoots arising from the lower nodes after much of the stem matures and dies. *Striga latericea* is a pest of sugar cane in Somalia and Ethiopia (Parker & Riches, 1993).

*Selected specimens examined.* ETHIOPIA. **Kefa:** Maji, 6°8'N, 35°37'E, Kaessner 1472 (BM). **Ogaden Harerge:** Awash National Park, Metahara Plain, 8°59'N, 39°58'E, Gilbert 1122 (K); Harar Sugar Estate, Awash Valley, 9°4'N, 40°9'E, Parker 4034 (K, ODU). **Shewa:** Yerer and Kereya, 8°54'N, 39°59'E, Puff *et al.* 8209 (K). **Sidamo:** 15 km NE Yavello, 4°59'N, 38°13'E, Gilbert & Jefford 4594 (K); 20 km S of Negele, 5°15'N, 39°36'E, Tadesse & Vollesen 3073 (K); Sidamo, 6°2'N, 37°46'E, Putt & Kelbessa 921220 (K). KENYA. Between Samburu and Mackinnin, Drummond & Homsley 4086 (K). COAST: between Makindu and Mtito Andei, 2°25'S, 37°57'E, Hale 105 (K); Kilifi, near Ganze, 3°0'S, 39°35'E, Robertson 4121 (K); Kwale, Kaya Bombo, 4°15'S, 39°10'E, Luke 3489 (K); Mwatati-Voi, 3°10'S, 38°32'E, Napier 957 (K); Mwatati-Voi, 3°25'S, 38°32'E, Joanna 8928 (K); Saka-Garissa Road, 0°45'S, 39°40'E, Faden & R.B. 74/744 (K); Vanga, Tanzania-Kenya border, 4°35'S, 39°12'E, Smith 10/92 (K); Diwanya Dakebuko, Parker GM/522/H (K). EASTERN: Machakos, Sultan Hamud, Subalata, 2°0'S, 37°40'E, Verdcourt 3690 (K); Mjiya Chumvi Isiolo, 0°24'N, 37°33'E, Adamson 615 (K); Sultan Hamud, 2°8'S,

37°40'E, Kaessner 645 (BM, K); Mwachi, 3 mi. S of Mazeras, Drummond & Hornsley 4247 (K); Mwea Irrigation Scheme, Parker 2984 (ODU). **Nairobi:** Embakasi, near airport, 1°17'S, 36°48'E, Napper & Faden 1839 (K); 20 km E of Nairobi, 1°17'S, 37°4'E, Bogdan 3270A (K); outside Nairobi at Mombasa Road, 1°19'S, 36°49'E, Hensen 854 (K); near Magi Ghumor, Kaessner 474 (K). **Northern:** Jex-Blake 6880 (K). **Rift Valley:** Emali and Sultan Hamud, 2°50'S, 37°57'E, Bally 8601 (K); Emali Station, 2°5'S, 37°27'E, Bally 8176 (K); Majihi Chumki, 0°42'N, 37°33'E, Adanson 615 (K, S). **SOMALIA.** Goan Libah, Bally 10025 (K). **Jobba Hoose:** Juba Sugar Project, 0°15'N, 42°0'E, Terry 3384 (K, ODU). **Woqooyi Gelbeed:** 80 km E of Hergesia, 9°37'N, 44°47'E, Laevranos 7394 (K); Gombale Plains, 40 mi. E of Hergesia, 9°37'N, 44°38'E, Glover & Gilliland 1231 (BM, K). **TANZANIA.** **Arusha:** Lekuruki Area, 3°20'S, 36°40'E, Richards 24897, 26863 (K). **Kilimanjaro:** Mkomazi, 4°40'S, 38°7'E, Harris 9 (K).

**21. *Striga lepidagathidis* A. Raynal, Bull. Mus. Natl. Hist. Nat., B, Adansonia 224–225. 1987.**  
TYPE: Guinea. Mamou, Jacques-Felix 775 (holotype, P!).

Plant perennial, up to 12 cm tall, succulent, tufted, branched from base, glandular-pubescent, stem whitish or purple, square, furrowed. Leaves 8 × 2 mm, lanceolate, reduced, appressed, longer than internodes, margin entire, veins obscure. Bracts similar, 6 × 2 mm, lanceolate, slightly longer than calyx. Flowers imbricate in dense spikes, spike longer than vegetative stem. Calyx 5-ribbed, 6 mm long; tube 4 mm long; lobes 5, unequal, subulate, 1.5–2 mm long, about half tube length. Corolla purple, pink, or white, tube 10–12 mm long, bent and expanded proximally within calyx teeth, densely pubescent; lobes of lower lip 4–5 × 2 mm, spreading, lanceolate, longitudinally rolled in when dry, tips acute, upper lip 2–3 × 2–3 mm, emarginate, as long as wide.

*Striga lepidagathidis* grows in Senegal, Guinea, and Guinea Bissau (Raynal-Roques, 1987) between 13°N and 10°23'N (Fig. 8). It is parasitic on *Lepidagathis* (Acanthaceae) species and other dicotyledon plants (Raynal-Roques, 1987). *Striga lepidagathidis* resembles *S. gesnerioides* and *S. gastoni* in their overall features and is difficult to distinguish from them. (See comments under *S. gastoni*.)

Selected specimens examined. GUINEA. Touba-Kande, Roberty 16608 (K). **Western:** Mamou (type), 10°23'N, 12°5'W, Jacques-Felix 1241 (P). GUINEA BISSAU. **Eastern:** Boe-Madina, 11°45'N, 14°13'W, Espírito Santo 3199 (K), Pereira 2984 (K).

**22. *Striga lutea* Lour., Fl. Cochinc: 22. 1790.**  
TYPE: China. Canton, Loureiro s.n. (holotype, P!; holotype, K photo!).

*Striga lutea* var. *bicolor* Kuntze, Rev. Gen. Pl. 3, 2: 240. 1898. TYPE: Mozambique. Kuntze 7494 (holotype, K!).

Annual, 7–40 cm tall, slender, unbranched or with 2- to 3-branches from middle, densely hispid, stem obtusely square. Leaves reduced, 3–7 × 1 mm, alternate, narrowly lanceolate, appressed, shorter than internodes, margin entire, veins obscure. All bracts similar, 1–4 × 0.5–1 mm, lanceolate, shorter than calyx. Lower surface of leaf and bract with two rows of hispid hairs along margins and midrib, upper surface hispid only along margins. Flowers sparse, alternate in lax racemes, raceme shorter than vegetative stem. Calyx 10-ribbed, 4–6 mm long; tube 3–4 mm long; lobes 5, equal, subulate, 1–2 mm long, about half tube length. Corolla yellow or rarely red, tube 10–11 mm long, bent and expanded distally above calyx, sparsely to densely glandular-pubescent; lobes of lower lip 2–3(–5) × 1–2 mm, obovate or oval, upper lip 1–2 × 2–3 mm, emarginate.

*Striga lutea* occurs between 15°N and 15°S (Fig. 14). The plants are easily overlooked in the field because they are small, possess a slender unbranched stem, and have reduced leaves. We do not know the relationship between this species and the Indian *S. lutea*. See notes under *S. asiatica* and *S. hirsuta*.

Selected specimens examined. BURKINA FASO. **Haut Bassins:** 5 km S of Banfora, 10°37'N, 4°43'W, Safa & Musselman 10/13/87 (ODU); Niangokolo Agronomic Research Station, 10°16'N, 4°54'W, Safa & Musselman 10/25/87 (ODU). CAMEROON. **Western:** Bangwa, 15 km NW of Bangante, 5°12'N, 10°36'E, de Wilde & de Wilde-Duyffes 2347 (K); Bawwda, Buru, 5°10'N, 10°30'E, Hawaiduud 1599 (K). CENTRAL AFRICAN REPUBLIC. **Ouaka:** 5 km E of Bambari, 5°41'N, 20°41'E, Fay 1896B (K). CONGO. **Haut-Mbomou:** Route Gwwane-Zemio, 4°55'N, 25°30'E, Bousique 168 (K). DEMOCRATIC REPUBLIC OF CONGO. **Shabas:** Albertville, Lake Tanganyika, 5°56'S, 29°12'E, Linder 1927 (GH). KENYA. **Coast:** Kwale, Shimba Hills, Giriam Point, 4°15'S, 39°10'E, Magogo & Estes 1152, 1153 (K). **Eastern:** Embu, 0°32'S, 37°28'E, Graham 2154 (K); Mahadra, Graham 1701 (K). **Rift Valley:** Chyulu Hills, 2°32'S, 37°47'E, Bally 8068 (K), Gilbert 6215 (K). MALLI. **Koulikoro:** Dialakoro, 32 km S of Bamako, 12°20'N, 7°56'W, Musselman & Mohamed 10/4/88 (ODU); just N of Bamako, 12°40'N, 8°0'W, Parker 4001 (ODU); Kamesoba, between Bamako-Siby, 12°30'N, 8°16'W, Musselman & Mohamed 10/2/88 (ODU); Zantiebougou, S of Baguineda, 12°45'N, 7°46'W, Musselman & Mohamed 10/5/88 (ODU). NIGERIA. **Niger:** Bida/Mokwa Junction, National Grain Corporation, 9°20'N, 5°40'E, Parkinson 9/18/85 (ODU), Mansfield 9/4/78 (ODU). SIERRA LEONE. **Northern:** near Mapaki, 9°40'N, 11°50'W, Sighton 1298 (BM). **Western:** Waterloo, 8°25'N, 13°5'W, Sighton 2042 (BM). TANZANIA. **Tanga:** Pombwe, Madanga, 4°50'S, 39°0'E, Tanner 2954 (K). UGANDA. **Central:** Bugoye, 0°34'N, 33°45'E, Dummer 2629 (US), Curtis 4 (GH).




Figure 14. Distribution based on specimens of *Striga lutea* (circle).

- 23. *Striga macrantha* (Benth.) Benth., in DC. Prodr. 10: 503. 1846. *Buchnera macrantha* Benth., Companion Bot. Mag. 1: 366. 1836. TYPE: Sierra Leone. *Afzelius s.n.* (holotype, K!).**

*Buchnera buettneri* Engl., Chev. Bot. 475. 1894. TYPE: Angola. Welwitsch 5907 (holotype, K!).

Plant annual, up to 2.5 m tall, coarse, stiffly erect, often 3-branched just below inflorescence, covered with dense, hispid, glandular hairs, stem square, furrowed. Leaves 50–120(–160) × 4–15(–20) mm, opposite, narrowly elliptic, 3-veined, coarsely toothed, longer than internodes. All bracts similar, 10–20 × 2–5 mm, lanceolate, recurved, as long as or longer than calyx. Flowers imbricate forming very compact racemes, raceme much shorter than vegetative stem. Calyx 10-ribbed, 7–10 mm long; tube 4–7 mm long; lobes 5, unequal, lanceolate, 2–4 mm long, about half tube length. Corolla white, tube 17–24 mm long, bent and expanded distally above calyx, densely pubescent; lobes of lower lip 7–10 × 4–8 mm, obovate, spreading, upper lip 3–7 × 4–10 mm, emarginate.

One of the tallest of all witchweeds, *S. macrantha* is found in the grassland savannas in West and Central Africa between 13°55'N and 3°51'N (Fig. 12) and also from a few localities in Angola and Zambia. Its range overlaps with *S. klingii*, but *S. macrantha* has a more southerly distribution in West Africa (Figs. 12, 13). *Striga macrantha* has unique glandular trichomes composed of a 10–16-celled glandular head (Neumann et al., 1996). It resembles *S. klingii* and *S. angolensis*. See notes under *S. klingii*.

*Selected specimens examined.* ANGOLA. Luanda: Cazengo, 8°47'N, 13°15'E, Gossweiler 558 (BM). Lunda North: Luballa, 9°10'S, 19°15'E, Mouteiro 6143 (K). CAMEROON. Center South: Mountain Febem, 3 km NW Yaounde, 3°51'N, 11°35'E, Breteler 2565 (K). Litoral: Douala to Mountain Bambouto, 4°2'N, 9°48'E, Meurillon CNAD577 (K). Northern: 29 km N of Dogba at Ngaouyang, 7°20'N, 13°32'E, Musselman et al. 10/16/88 (ODU). CENTRAL AFRICAN REPUBLIC. Bambari-Bangoran: 8 km S of Koumbala, 8°26'N, 21°15'E, Fay 4083 (K). GAMBIA. Brookes 4 (K), Lester 34 (K). GHANA. Brong-Ahafo: Techiman, 7°38'N, 1°58'W, Adams & Akpabla 4509 (K). Northern: between River Oti and Atafie, 10°0'N, 0°25'E, Morton 94 (K). VOLTA: Ho, 6°35'N, 0°30'E, Glumptre 25 (K). GUINEA. Northwest: Youkounkoun, Sambailo, 12°40'N, 13°22'W, Adam 13885 (MO). GUINEA BISSAU. Central: Bafata, Sallinho, 12°12'N, 14°42'W, Espirito Santo 3636 (K). Northern: Farim, 12°35'N, 15°10'W, Toul Botanica Exped. 1699 (K). Western: Tussubi, 12°0'N, 15°35'W, Toul Botanical Exped. 1464 (K). LIBERIA. Lofa: Inpaka, near Kolahun Summit, 8°12'N, 10°4'W, Bequaert 35 (GH). Sinoe: Greenville, 5°30'N, 8°50'W, Adam 24394 (MO), Adam 29806 (US). MALI. Koulikoro: Bamako, Mandinguies, 12°40'N, 7°59'W, Adam 11339 (MO). NIGER. Parker 985 (GH). NIGERIA. Bauchi: Fuel Plantation, SE of Gombe, 10°12'N, 11°15'E, Lowe 1484 (K). Benue: 2–3 mi. SE Otukpo, 7°15'N, 8°10'E, Anonymous 2143 (BM). GONGOLA: Adamawa, Dakemi, 8°25'N, 11°50'E, Hepper 1392 (K). Kaduna: Zaria, near dam for Institution for Agriculture Research, 10°58'N, 7°58'E, Gbile & Daramola 245 (K). Kwara: 10 km W of Isanlu-Makutu, 8°15'N, 5°45'E, Lowe 4463 (K). Lagos: W of Lagos, 6°28'N, 3°30'E, Rowland 1893 (K). Niger: 45 mi. N of Bida, 9°29'N, 5°58'E, Harris 8 (K). Northern: Dalziel 169, 911 (K). Ogun: Abeokuta, 7°6'N, 3°25'E, Baldwin 13639 (K, MO, US). OYO: Oyo, University of Ife Camp, 7°58'N, 4°0'E, Suabad 6994 (MO). Plateau: Jos Plateau, 9°50'N, 8°40'E, Batten-Poole 226 (K). Southern: southern Guinea Savana, Musselman & Mansfield 5522 (ODU). SENEGAL. Oriental: Kedougou, 12°44'N, 12°15'W, Adam 20029 (MO). SIERRA LEONE. Freetown: rifle range, near Waterloo,

8°22'N, 13°3'W, F.A.M. & T.H. 474 (K). **Northern:** Bisawa, above Bumbuna, 9°6'N, 11°44'W, King 175 (K). SUDAN. **North Darfur:** Ngertete, Gabal Marra, 12°50'N, 24°30'E, Wickens 2807 (K). CHAD. **Logone-Oriental:** Lere to Ham, at 9°50'N, 14°40'E, Talbot 324 (BM, K). ZAMBIA. **Southern:** Kalomo, 17°0'S, 26°30'E, Fanshawe 4689 (K).

**24. *Striga passargei* Engl., Bot. Jahrb. 23: 515, 12 figs. M, N. 1897.** TYPE: Cameroon. Jola, Passarge 48 (holotype, K!).

Annual, to 46 cm tall, erect, unbranched or branched from middle, densely hispid, stem obtusely square. Leaves 10–45 × 1–3 mm, narrowly elliptic, opposite/subopposite, ascending, longer than internodes, margin entire, veins obscure. Lower bracts 12–60 × 1–3 mm, linear, much longer than calyx, upper bracts as long as or longer than calyx. Flowers alternate in long, lax spikes, spike longer than vegetative stem. Calyx 5-ribbed, 8–12 mm long; tube 3–8 mm long; lobes 5, equal, linear to lanceolate, 3–8 mm long, about same length as tube. Corolla cream-white, rarely pink or yellow, tube 6–8 mm long, bent proximally within calyx teeth, glandular-pubescent; lobes of lower lip 2–6 × 2–4 mm, round, upper lip 1–3 × 2–5 mm, emarginate.

*Striga passargei* is found in the wet areas within the Sudanian domain in Africa. Its range extends from Senegal to Sudan between 14°44'N and 6°35'N (Fig. 15). In this area it is sympatric with *S. aspera* and *S. brachycalyx*. Unlike these two species, *S. passargei* always occurs in smaller populations. South of the equator, it occurs only in Tanzania, Zambia, and Namibia (Fig. 15). There is considerable variation in populations of *S. passargei* particularly in stem height, leaf and bract size, calyx lobes relative to tubes, and corolla color.

*Selected specimens examined.* BURKINA FASO. **Central:** Zongo, 69 km E of Ouagadougou, 12°24'N, 0°35'W, Raynal 22277 (K). **Center East:** 5 km S of Ouada, 11°30'N, 0°20'W, Safa & Musselman 10/9/87 (ODU). **Center North:** 55 km N of Ouagadougou, 12°51'N, 1°39'W, Safa & Musselman 10/19/87 (ODU). CAMEROON. **Northern:** Bogo, 35 km from Maroua, 10°44'N, 14°36'E, Bounougou 126 (K). GHANA. **Brong-Afao:** Bambo, 8°10'N, 2°12'W, Hall 2028 (K). GUINEA **Northern:** Jiripa, 10°55'N, 1°58'W, Thorold 251 (BM). **Central:** Pita, 11°6'N, 12°24'W, Adames 341 (K). MALI. **Koulikoro:** Dialakoro, 32 km S of Bamako, 12°18'N, 7°56'W, Musselman & Mohamed 10/4/88 (ODU). NAMIBIA. **Ovamboland:** Ondangwa, 17°56'S, 16°0'E, Kers 1399 (S). NIGERIA. **Bauchi:** Panshanu Pass, 10°5'N, 9°3'E, Lawlor & Hall 461 (K). **Bornu:** Lake Tchad Area, 13°5'N, 13°51'E, Parkinson A1 (K). **Gongola:** Yola, 9°10'N, 12°25'E, Dalziel 110 (K). **Kaduna:** Panshanu, 30 mi. from Jos on Jos-Bauchi Road, 10°8'N, 8°0'E, Lawlor & Hall 73 (K). **Ogun:** Ibadan, IITA, 7°17'N, 3°30'E, Par-

kinson 1985 (ODU). **Plateau:** Jos, Naraguta Hills, SE Naraguta, 9°50'N, 8°40'E, Lawlor & Hall 34 (K). **Sokoto:** Sokoto, 13°8'N, 5°15'E, Dalziel 355 (K). SENEGAL. **Oriental:** Dialakoto, 13°27'N, 13°18'W, Musselman et al. 7061, 7063 (ODU). SUDAN. **Blue Nile:** Abu Naama, Gabal Tozi, 12°54'N, 33°36'E, Lea 58 (K, ODU), Lea 180 (K). **Jungali:** near Baidit, 45 km N of Bor, 6°35'N, 31°36'E, Lock 81/248 (K). **North Darfur:** Kutum, 14°10'N, 24°42'E, Lynes 578 (BM). **North Kurdufan:** Al-Obied, 13°15'N, 30°12'E, Pfund 86 (K). **South Kurdufan:** Kadugli, Saref Research Station, 11°3'N, 29°42'E, Musselman 198 (K). TANZANIA. **Eastern:** 2 mi. W of Chibembe River, Luangwa Valley Game Reserve, 14°25'S, 30°30'E, Prince 200, 401 (K). **Lindi:** Kingupira Forest, 8°30'S, 38°34'E, Vollesen 3311 (K, UPS). **Mwanza:** Mwanza, 2°30'S, 32°58'E, Davis 255 (K). **Rukwa:** Tumba, Rukwa Valley, 4°15'S, 33°14'E, Siame 144 (K). **Shinyanga:** near Shinyanga, 3°47'S, 33°27'E, Rax 292 (K). **Tabora:** Rukwa-Nzega Plain, 4°10'S, 33°12'E, Whellar 1188 (K). TOGO. **Northern:** Dapaon, (Dapango), 10°44'N, 0°12'E, Hakki et al. 1504 (K). ZAMBIA. **Mashonaland North:** Kariba Airport, 16°30'S, 28°35'E, Philcox & Leppard 8677 (K).

**25. *Striga pinnatifida* Getachew Aweke, Kew Bull. 47: 293–294. 1992.** TYPE: Ethiopia. Sidamo Prov., Friis, Gilbert, Rasmussen & Vollesen 918 (holotype, K!; isotype, ETH not seen).

Annual, to 45 cm tall, erect, unbranched or with 2 to 3 branches from middle, covered with dense, hispid hairs, stem obtusely square. Leaves 15–30 × 10 mm, alternate, pinnatifid with linear segments, shorter than internodes. Lower bracts 10–25 × 8 mm, leaf-like, longer than calyx, upper bracts shorter than calyx. Flowers shortly pedicellate, alternate, axillary, solitary, sparse. Calyx 10-ribbed, 10–12 mm long; tube 6–7 mm long; lobes 5, unequal, lanceolate, 3–5 mm long, half tube length. Corolla white, tube 20 mm long, bent and expanded distally above calyx, pubescent; lobes of lower lip 6 × 5 mm, spreading, obovate, upper lip 3 × 6 mm, emarginate.

*Striga pinnatifida* is known solely from the type collection. It is the only witchweed with pinnatifid leaves and solitary flowers.

**26. *Striga primuloides* Chev., Bull. Soc. Bot. Fr. Mem. 8: 185. 1912.** TYPE: Ivory Coast. Anoumaba, Chevalier B.22.404 (holotype, Pl!; isotype, K!).

Perennial, 30–70 cm tall, erect, unbranched, appearing leafless, pilose, stem obtusely square. Leaves scale-like, 2–5 × 1–2 mm, lanceolate, alternate, appressed, shorter than internodes, margin entire, veins obscure. All bracts similar, 4–7 × 1–2 mm, lanceolate, shorter than calyx. Flowers sparse, alternate in lax spikes, spike much shorter




Figure 15. Distribution based on specimens of *Striga passargei* (triangle), *S. primuloides* (circle), *S. pubiflora* (diamond), and *S. yemenica* (star).

than vegetative stem. Calyx approximately 15-ribbed, 8–9 mm long; tube 5 mm long; lobes 5, unequal, lanceolate, 3–4 mm long, slightly shorter than tube. Corolla pale yellow, tube 22–30 mm long, bent and expanded distally above calyx, densely glandular-pubescent; lobes of lower lip 9–10 × 8–9 mm, round, spreading, as long as wide, upper lip 5 × 10 mm, broadly obovate, notched.

*Striga primuloides* is restricted to the southern parts of West Africa between 5°10'N and 10°37'N (Fig. 15). It is a rare species with only a few collections from Mali, Ivory Coast, Ghana, and southern Nigeria.

*Selected specimens examined.* GHANA. Brong-Ahafo: Bambo Ravine, 8°10'N, 2°8'W, Hall 1998 (K), Morton A4241 (K); Bambo, W of Wenchi, 8°15'N, 2°12'W, Morton GC8601 (K). Northeastern: along road from Gawi to Seye, in Gallery Forest of Iringou River, 9°14'N, 3°41'W, Amshoff 193 (MO). MALI. Mopti: Massina, Diou, 14°52'N, 5°16'W, Laferrere 112 (K). NIGERIA. South Nigeria: Awkaka, Birkett-Smith 49 (K).

**27. *Striga pubiflora* Klotzsch., in Peters Reise Mossamb. Bot.: 227. 1861. TYPE: Mozambique. Dayser s.n. (holotype, B not seen).**

*Striga zanzibarensis* Vatke, Linnaea 43: 310. 1882. TYPE: Mafia Isl. Hildebrandt 1904 (holotype, K!).

Perennial, 35–75 cm tall, stiffly erect, unbranched or branched from base, densely pilose or hispid, stem obtusely square, furrowed. Leaves 10–20(–50) × 1–2(–4) mm, linear, opposite on the lower part of the stem, alternate above, shorter than internodes, margin entire, rarely toothed, veins obscure. Lower bracts 22–27 × 2 mm, linear, longer than calyx, upper bracts shorter than calyx. Flowers alternate in lax racemes, raceme shorter than vegetative stem. Calyx 15-ribbed, 13–21 mm long; tube 8–12 mm long; lobes 5, equal, lanceolate, 4–12 mm long, as long as or shorter than tube. Corolla white, tube 27–35 mm long, bent and expanded distally above calyx, densely pubescent; lobes of lower lip 14–20 × 10–17 mm, obovate, spreading, upper lip 9–12 × 10–20 mm, widely obovate.

*Striga pubiflora* is common in the seasonally flooded savannas of southeastern Africa between 3°10'S and 23°54'S (Fig. 15). Its range extends from Kenya south into Tanzania, Zanzibar, Malawi, and Mozambique. It may be related to *S. junodii* and *S. angustifolia* (Mohamed, 1994). *Striga pubiflora* and *S. angustifolia* are sympatric and also have white flowers, but the corolla of *S. pubiflora* is much larger.

*Selected specimens examined.* KENYA. Coast: Kwale,

Shimba Hills, Mombasa Road between Vanga and Tiwi turn-off, 4°25'S, 39°8'E, *Magogo & Clover* 1050 (K). **MA-LAWI.** **Central:** Dowa, 25 km S of Kasungu, 13°14'S, 33°40'E, *Pawek* 4112 (K). **Southern:** Zomba Mountain, Chivunde Valley Road, 15°20'S, 35°19'E, *Croix* 2735 (MO). **MOZAMBIQUE.** 5 mi. NE of Lunga, *Moomaw* 953 (K). **Inhambane:** Inhambane, 23°54'S, 35°30'E, *Anonymous s.n.* (BM). **Manica:** 25 mi. S of Muda, 19°51'S, 33°50'E, *Leach* 11232 (K, MO). **Niassa:** 47 mi. W of Nova Ureixo on Mandimba Road, 12°40'S, 35°40'E, *Leach & Smith* 11011 (K), *Allen* 84 (K), *Stocks* 55 (K). **Sofala:** 41 mi. N of Sofala, 19°50'S, 34°34'E, *Methew* 257 (K). **Zambia:** Quelimanje, 17°53'S, 36°58'E, *Faulkner* K118 (K). **TANZANIA.** **Dar es Salaam:** 20 mi. S of Dar es Salaam, between Mbwa and Kimbiji, 7°0'S, 39°18'E, *Batty* 870 (K). **Iringa:** Mahenge Plateau, 7°37'S, 36°12'E, *Schlieben* 2281 (K). **Lindi:** 3 km W of Nainokwe/Njinjio-Nangurukuru Road junction on road to Kilwa, 9°0'S, 39°0'E, *Magogo & Innes* 373 (K). **Mafia:** Mji Mkuu, Kipumbwe, Mwera, *Tanner* 2850 (K). **Morogoro:** 3 km N of Mlahi, 8°16'S, 37°5'E, *Kesollesen* 2817 (K). **Pwani:** Kibaha, N of Bush, 6°46'S, 38°55'E, *Harris* 786 (S). **Rukwa:** Mlimba, 7°51'S, 31°0'E, *Paget-Wilkes* 700 (MO). **Ruvuma:** 5 km E of Gumbiro, 10°15'S, 35°48'E, *Milne-Redhead & Taylor* 8421 (K). **Tanga:** Aboni, Mgaza 4°50'S, 39°0'E, *Tanner* 89 (K). **Wani:** Kisarawe Mogo Forest Reserve, 7°3'S, 39°0'E, *Poulo* 105 (K, MO). **Zanzibar:** Fessland bec Lamu, *Hildebrandt* 1907 (UPS).

**28. *Striga yemenica* Musselman & Hepper,** Notes Roy. Bot. Gard. Edinburgh 45(1): 43–50. 1988. TYPE: Yemen Arab Republic. Jiblah, *Miller* 546 (holotype, E!; isotype, K!).

Annual, 18–22 cm tall, erect, usually unbranched, covered with dense, retrorse, hispid hairs, stem square, furrowed. Leaves 10–22 × 1–2 mm, lanceolate, opposite, ascending, shorter than internodes, margin entire, veins obscure. Lower bracts 8–12 × 1(–2) mm, lanceolate, ciliate, longer than calyx, upper bracts shorter than calyx. Flowers opposite in lax spikes, spike as long as vegetative stem. Calyx 5-ribbed, 5–7 mm long; tube 4–5 mm long; lobes 5, unequal, (adaxial reduced to < 1 mm) deltate, 1(–2) mm long, much shorter than tube. Corolla red or mauve, bent distally above calyx, densely retrorsely pubescent; lobes of lower lip 0.5–1 × 2 mm, round, upper lip small, 0.5 × 2 mm, truncate.

In Africa, only in Ethiopia between 9°6'N, 9°38'N and 41°48'E, 42°49'E (Fig. 15). It bears some morphological similarities to *S. hermonthica* and *S. gracillima*. In these three species, the calyx lobes are short and deltate with the adaxial lobe reduced to less than 1 mm in length; corollas are pink or mauve with simple hairs and never glandular. *Striga yemenica*, on the other hand, has retrorse hairs in contrast to the antrorse hairs of *S. hermonthica* and *S. gracillima*.

*Selected specimens examined.* ETHIOPIA. 31 km from

Alemaya on road to Grawa and Bedeno, *Gilbert* 4072 (K). **Ogaden Harerge:** 4 km W of China Hasen, 31 km NW of Jijiga, 9°26'N, 42°42'E, *Boulos* 10250 (K); Gures, 18 km NW of Jijiga, 9°20'N, 42°42'E, *Boulos* 10509 (K); Harrar, 9°18'N, 42°8'E, *Ellis* 272 (K); Mountain Hakim, 9°17'N, 42°6'E, *de Wilde* 7223 (K); Mountain Hakim, 9°16'N, 42°7'E, *Gilbert & S.B.* 1408 (K); 1 km W of Abu Yonis, 30 km NW Jijiga, 9°24'N, 42°45'E, *Boulos* 10316 (K, MO).

#### EXCLUDED SPECIES

*Striga baumannii* has tuberous roots and a lignous calyx, features not found in any other *Striga* species. The foliage is also unique with two large leaves formed during the growing season. In *Striga* the lowermost leaves are reduced to small scales. We consider *S. baumannii* distinct from any witchweed and therefore do not include it in the genus. It is thought to represent a new genus (Mohamed, unpublished).

*Striga fulgens* (= *Buchnera fulgens* Engl.) was named by Hepper (1984). However, Raynal-Roques (1993) presented evidence that it is a member of the Verbenaceae, likely in the genus *Chascanum* E. Mey. It has free lobes of the upper corolla lip as well as a glabrous calyx with characteristic white bands between the ribs and short calyx lobes (1–2 mm). Accordingly, we exclude it from *Striga*.

#### Literature Cited

- Agbobli, C. A. 1991. Effect of nitrogen rates on *Striga asiatica* emergence in maize culture in Togo. Pp. 28–30 in J. K. Ransom, L. J. Musselman, A. D. Worsham & C. Parker (editors), Proceedings of the Fifth International Symposium on Parasitic Weeds. CIMMYT, Nairobi.
- Aigbokhan, E. I., D. K. Berner & L. J. Musselman. 1998. Reproductive ability of hybrids of *S. aspera* and *S. hermonthica*. *Phytopathology* 88: 563–567.
- \_\_\_\_\_, \_\_\_\_\_ & H. D. Mignouna. 2000. Evaluation of variability in *Striga aspera*, *Striga hermonthica* and their hybrids using morphological characters and random amplified polymorphic DNA markers. *Weed Res.* 40: 375–386.
- Anonymous. 1996. MapInfo Professional, 2nd ed. Map-Info, Troy, New York.
- Berner, D. K., K. F. Cardwell & B. O. Fatuoti. 1992. *Striga hermonthica* distribution mechanisms and their implications in control. *Phytopathology* 82: 1164.
- Bharathalakshmi, C. R. Werth & L. J. Musselman. 1990. A study of genetic diversity among host-specific populations of the witchweed *Striga hermonthica* (Scrophulariaceae) in Africa. *Pl. Syst. Evol.* 172: 1–12.
- Boesshore, I. 1920. The morphological continuity of Scrophulariaceae and Orobanchaceae. *Contr. Bot. Lab. Morris Arbor. Univ. Pennsylvania* 5: 139–177.
- Cubero, J. I. 1996. Cytogenetics in Orobanchaceae: A review. Pp. 75–96 in M. T. Moreno & J. I. Cubero (editors), Advances in Parasitic Plant Research. Junta de Andalucía, Córdoba.
- DeLeonardis, M. A. 1986. The Pollen Exostructure of

- Striga* (Scrophulariaceae). M.S. Thesis, Old Dominion University, Norfolk.
- Engler, A. 1897. Scrophulariaceae africanae. II. Bot. Jahrb. 23: 497–517.
- Geiger, U., J. Kroschel & J. Sauerborn. 1996. *Striga asiatica*, a problem in the middle west of Madagascar. Pp. 479–486 in M. T. Moreno & J. I. Cubero (editors), Advances in Parasitic Plant Research. Junta de Andalucía, Córdoba.
- Hepper, F. N. 1960. The *Striga bilabiata* (Thunb.) Kuntze complex in Africa. Kew Bull. 14: 411–416.
- \_\_\_\_\_. 1974. Parasitic witchweed: *Striga asiatica* versus *S. lutea* (Scrophulariaceae). Rhodora 76: 45–47.
- \_\_\_\_\_. 1984. The species of *Striga* in East Africa. Pp. 262–265 in C. Parker, L. J. Musselman, R. M. Polhill & A. K. Wilson (editors), Proceedings of the Third International Symposium on Parasitic Weeds. ICARDA/International Parasitic Seed Plants Research Group, Aleppo.
- \_\_\_\_\_. 1986. Proposal to reject the name *Buchnera euphrasiooides* Vahl (*Striga euphrasiooides* (Vahl) Benth.) (Scrophulariaceae). Taxon: 35: 390–391.
- Kenfack, D., L. J. Musselman & H. J. Hovers. 1996. Hosts of eight *Striga* species (Scrophulariaceae) in Cameroon. Pp. 465–470 in M. T. Moreno & J. I. Cubero (editors), Advances in Parasitic Plant Research. Junta de Andalucía, Córdoba.
- Knepper, D. A., A. T. Obilana & L. J. Musselman. 1991. Morphology of *Striga forbesii* and preliminary screening for resistance in sorghum. Pp. 241–246 in J. K. Ransom, L. J. Musselman, A. D. Worsham & C. Parker (editors), Proceedings of the Fifth International Symposium on Parasitic Weeds. CIMMYT, Nairobi.
- Maass, E. 1999. A Comparative Study on the Germination Requirements of Some Economically Important *Striga* Species. Ph.D. Dissertation, University of Stellenbosch.
- Minkin, J. P. & W. H. Eshbaugh. 1989. Pollen morphology of the Orobanchaceae and rhianthoid Scrophulariaceae. Grana 28: 1–18.
- Mohamed, K. I. 1994. Biosystematics and Diversification in the Genus *Striga* Lour. (Scrophulariaceae) in Africa. Ph.D. Dissertation, Old Dominion University, Norfolk.
- Musselman, L. J. & E. S. Ayensu. 1984. Taxonomy and biosystematics of *Striga*. Pp. 37–45 in E. S. Ayensu, H. Doggett, R. D. Keynes, J. Marton Le-Fevre, L. J. Musselman, C. Parker & A. Pickering (editors), *Striga* Biology and Control. International Council of Scientific Unions Press, Paris.
- \_\_\_\_\_. & F. N. Hepper. 1986. The witchweeds (*Striga*, Scrophulariaceae) of the Sudan Republic. Kew Bull. 41: 205–221.
- \_\_\_\_\_. & W. F. Mann, Jr. 1976. A survey of surface characteristics of seeds of Scrophulariaceae and Orobanchaceae using scanning electron microscopy. Phytomorphology 26: 370–378.
- \_\_\_\_\_. & C. Parker. 1981. Surface features of *Striga* seeds (Scrophulariaceae). Adansonia, series 2 20: 431–437.
- \_\_\_\_\_, Bharathalakshmi, S. B. Safa, D. A. Knepper, K. I. Mohamed & C. L. White. 1991. Recent research on the biology of *Striga asiatica*, *Striga gesnerioides* and *Striga hermonthica*. Pp. 31–41 in S. K. Kim (editor), Combating *Striga* in Africa. Proceedings, International Workshops Organized by IITA, ICRISAT and IDRC. IITA, Ibadan.
- Neumann, U., J. Paré, A. Raynal-Roques, G. Sallé & H. Weber. 1996. Characteristic trichomes observed in some African parasitic Scrophulariaceae. Pp. 264–272 in M. T. Moreno & J. I. Cubero (editors), Advances in Parasitic Plant Research. Junta de Andalucía, Córdoba.
- Nickrent, D. L., R. J. Duff, A. E. Colwell, A. D. Wolfe, N. D. Young, K. E. Steiner & C. W. dePamphilis. 1998. Molecular phylogenetic and evolutionary studies of parasitic plants. Pp. 211–241 in D. E. Soltis, P. S. Soltis & J. J. Doyle (editors), Molecular Systematics of Plants II. DNA Sequencing. Kulwer Academic Publishers, Boston.
- Olivier, A., J. C. Glaszmann, C. Lanauud & G. D. Leroux. 1998. Population structure, genetic diversity, and host specificity of the parasitic weed *S. hermonthica* (Scrophulariaceae) in Sahel. Pl. Syst. Evol. 209: 33–45.
- Olmstead, R. G., C. W. dePamphilis, A. D. Wolfe, N. D. Young, W. J. Elisons & P. A. Reeves. Disintegration of the Scrophulariaceae. Amer. J. Bot. (in press).
- Parker, C. 1988. Parasitic plants in Ethiopia. Walia 11: 21–27.
- \_\_\_\_\_. & C. R. Riches. 1993. Parasitic Weeds of the World. CABI, Oxford.
- Press, M. C. & J. D. Graves (editors). 1995. Parasitic Flowering Plants. Chapman and Hall, London.
- Ralston, D. M., C. R. Riches & L. J. Musselman. 1987. Morphology and hosts of three *Striga* species (Scrophulariaceae) in Botswana. Bull. Mus. Natl. Hist. Nat., B, Adansonia 2: 195–215.
- Ramaiah, K. V., C. Parker, M. J. Vasudeva Rao & L. J. Musselman. 1983. *Striga* Identification and Control Handbook. ICRISAT Information Bulletin #15, Patancheru, India.
- Raynal-Roques, A. 1969. Deux nouveaux *Striga* africains (Scrophulariaceae). Bull. Jard. Bot. Belg. 39: 373–382.
- \_\_\_\_\_. 1987. The genus *Striga* (Scrophulariaceae) in western and central Africa—A survey. Pp. 675–689 in H. C. Weber & W. Forstreuter (editors), Proceedings of the Fourth International Symposium on Parasitic Flowering Plants. ISPPEP, Marburg.
- \_\_\_\_\_. 1991. Diversification in the genus *Striga*. Pp. 251–261 in J. K. Ransom, L. J. Musselman, A. D. Worsham & C. Parker (editors), Proceedings of the Fifth International Symposium on Parasitic Weeds. CIMMYT, Nairobi.
- \_\_\_\_\_. 1993. Contribution à la connaissance de la biologie des *Striga* (Scrophulariaceae): Types biologiques et phénologie. Bull. Mus. Natl. Hist. Nat., B, Adansonia 3–21.
- Saldanha, C. J. 1963. The genus *Striga* Lour. in western India. Bull. Bot. Surv. India 5: 67–70.
- Sand, P. F., R. E. Eplee & R. G. Westbrooks (Editors). 1990. Witchweed Research and Control in the United States of America. Weed Science Society of America, Champaign.
- Sauerborn, J. 1991. Parasitic Flowering Plants. Ecology and Management. Verlag Josef Margraf, Weikersheim.
- Thalouarn, P., M. C. Arnaud, C. Theodet & L. Rey. 1991. Cytological, biochemical and genetic aspects of carbon fixation in *Striga hermonthica* and *Striga gesnerioides*. Pp. 51–57 in J. K. Ransom, L. J. Musselman, A. D. Worsham & C. Parker (editors), Proceedings of the Fifth International Symposium on Parasitic Weeds. CIMMYT, Nairobi.
- Werth, C. R., J. L. Riopel & N. W. Gillespie. 1984. Genetic uniformity in an introduced population of witchweed (*Striga asiatica*) in the United States. Weed Sci. 32: 645–648.

- Wettstein, R. von. 1891. Scrophulariaceae. In A. Engler & K. Prantl (editors), Die Natürlichen Pflanzenfamilien IV (3b): 39–107.
- Wild, H. 1948. A suggestion for the control of tobacco witchweed [*Striga gesnerioides* (Willd.) Vatke] by leguminous trap crops. Rhodesia Agric. J. 45: 208–215.
- Young, N. D., K. E. Steiner & C. W. dePamphilis. 1999. The evolution of parasitism in Scrophulariaceae/Orobanchaceae: Plastid gene sequences refute an evolutionary transition series. Ann. Missouri Bot. Gard. 86: 876–893.

#### LIST OF SPECIES

1. *Striga aequinoctialis* A. Chev. ex Hutch. & Dalz.
2. *Striga angolensis* Mohamed & Musselman
3. *Striga angustifolia* (Don) Saldanha
4. *Striga asiatica* (L.) Kuntze
5. *Striga aspera* (Willd.) Benth.
6. *Striga bilabiata* (Thunb.) Kuntze
  - 6a. *Striga bilabiata* subsp. *barteri* (Engl.) Hepper
  - 6b. *Striga bilabiata* Kuntze subsp. *bilabiata*
  - 6c. *Striga bilabiata* subsp. *jaegeri* Hepper
  - 6d. *Striga bilabiata* subsp. *ledermannii* (Pilger) Hepper
  - 6e. *Striga bilabiata* subsp. *linearifolia* (Schum. & Thonn.) Mohamed
  - 6f. *Striga bilabiata* subsp. *rowlandii* (Engl.) Hepper
7. *Striga brachycalyx* Skan
8. *Striga chrysanthra* A. Raynal
9. *Striga dalzielii* Hutch.
10. *Striga elegans* Benth.
11. *Striga forbesii* Benth.
12. *Striga gastonii* A. Raynal
13. *Striga gesnerioides* (Willd.) Vatke
14. *Striga gracillima* H. Melchior
15. *Striga halliae* A. Raynal
16. *Striga hermonthica* (Del.) Benth.
17. *Striga hirsuta* Benth.
18. *Striga junodii* Schinz.
19. *Striga klingii* (Engl.) Skan
20. *Striga latericea* Vatke
21. *Striga lepidagathidis* A. Raynal
22. *Striga lutea* Lour.
23. *Striga macrantha* (Benth.) Benth.
24. *Striga passargei* Engl.
25. *Striga pinnatifida* Getachew Aweke
26. *Striga primuloides* Chev.
27. *Striga pubiflora* Klotzsch.
28. *Striga yemenica* Musselman & Hepper

#### INDEX TO EXSICCATAE

Specimens examined are listed alphabetically by collector; followed by collection numbers (or dates of collections for specimens with no numbers), herbarium, and the species number corresponding to the number in the List of Species.

- Acocks, J. P. 1481 (K) (6b), 1491 (K) (10), 2012 (K) (13); Adam, J. G. 5111 (MO) (1), 11248 (MO) (23), 11339 (MO) (23), 13724 (MO) (6f), 13884 (MO) (19), 13885 (MO) (23), 14200 (MO) (6f), 14812 (MO) (6f), 14861 (MO) (6c), 17917 (MO) (6a), 17934 (MO) (6a), 20029 (MO) (23), 21457 (K, MO, UPS) (1), 21957 (MO) (23), 21995 (GH, MO) (19), 24039 (MO) (1), 24386 (MO) (1), 24394 (MO) (23), 24653 (MO) (1), 25117 (MO) (23), 25550 (K, MO) (1), 25592 (K) (1), 26054 (MO) (1), 26531 (MO) (1), 26881 (MO) (19), 26891 (MO) (6f), 26896 (MO) (6a),

- 26897 (MO) (6f), 26898 (MO) (6f), 27125 (MO) (6e), 27478 (MO) (1), 27661 (MO) (6f), 29806 (US) (23); Adams, P. 146 (K) (6f), 193 (K) (23), 224 (K) (6f), 284 (K) (1), 341 (K) (24); Adams, C. D. 3548 (K) (6e), 4152 (K) (6e), 4419 (K) (16), 4472 (K) (23), 4693 (K) (7); Adams, C. D. & G. K. Akpabla 4107 (K) (16), 4272 (K) (5), 4509 (K) (23); Adanson, G. 615 (K, S) (20); Adamson, J. 527 (K) (4), 615 (K) (20), 3562 (K) (13); Agnew, A. 8700 (MO) (6e); Ajayi, M. O. 19271 (K) (23); Ake, L. 8755 (K) (6e); Akestrid 2527 (UPS) (6e); Akpabla, G. K. 36 (K) (13); Alcool, M. 963 (K) (16), 4116 (K) (6a), 4142 (K) (11); Allen, C. E. 124 (K) (13); Allen, E. F. 84 (K) (27); Allen, J. 10470 (K) (13); Amshoff, G. J. 193 (MO) (26), 225 (MO) (6e); Anderson, B. 846 (K) (4), 885 (K) (11), 1364 (K) (11); Andrews, F. W. 1018 (K) (17); Andrews, J. W. 6 (K) (5); Angus, A. 1066 (K) (6b); Ankole, E. & W. J. Eggeling 610 (K) (6e); Ankrah, J. O. 20183 (K) (6e); Anonymous 189 (K) (13), 782 (K) (13), 837 (K) (13), 1036 (K) (16), 1126 (K) (5), 1651 (K) (6e), 1784 (K) (18), 1829 (K) (6e), 1842 (GH, US) (17), 1858 (K) (11), 2069 (GH) (10), 2143 (BM) (23), 2159 (K) (4), 2180 (K) (4), 2381 (K) (16), 2482 (GH) (10), 3063 (K) (11), 3384 (K) (11), 4863 (K) (6b), 5749 (GH) (4), 8347 (K) (4), 9199 (US) (16), 9200 (K) (13), 9783 (K) (1), 11762 (K) (11), 12092 (MO, S, UPS) (18), 12092 (BM, K, US) (27), s.n. (BM) (27), s/ss (K) (16); Appelsbosch 1889 (UPS) (4); Archbold, A. E. 694 (K) (4), 732 (K) (4); Archbold, M. E. 2776 (K) (13); Ascheimeier, C. R. 11 (US) (17); Ash, J. W. 347 (K) (13), 518 (K) (17), 557 (K) (13), 1089 (K) (13), 3648 (K) (16); Ash, M. J. 544 (K) (11), 2259 (K) (11); Astle, W. L. 1023 (K) (6b), 1556 (K) (11), 1571 (K) (10), 4471 (K) (11), 4570 (K) (11), 5507 (K) (4); Audru, 1245 (P) (12); Audry, P. B2 (K) (16); Awaikiwe, et al. 148 (K) (11); Awbeya, 1430 (MO) (10); Aylmar, G. 27/50 (K) (11), 27/51 (K) (13e), 236 (K) (13).

- Baikie, W. B. 1865 (GH) (16); Baitha, R. 36 (K) (16); Bakshi, T. S. 146 (K) (6f); Baldwin, J. T. 5988 (US) (17), 13639 (K, MO, US) (23), s.n. (K) (4); Ballenetto 277 (K) (13); Balls, E. K. 3022 (BM, K) (13); Bally, M. 283 (K) (17); Bally, P. R. O. 4596 (K) (13), 7329 (K) (13), 8064 (K) (6e), 8068 (K) (22), 8072 (K) (11), 8176 (K) (20), 8601 (K) (20), 9675 (K) (13), 10025 (K) (20); Bally, P. R. & Carter 16448 (K) (17); Bally, P. R. O. & R. Melville 15841 (K) (13); Balsinhas, A. 152 (K) (13); Balsinhas, A. & L. Marrime 379 (K) (4); Balslev, H. 447 (MO) (17); Barbey 1051 (US) (13); Barbosa, G. & R. Corriss 8937 (K) (4); Barter, C. 136 (GH) (16), 158 (GH) (19), 530 (ODU) (17), 1169 (GH) (6f), 1170 (GH) (6a), 1263 (GH) (6e), 1821 (GH) (16); Basera, C. 343 (K) (13); Bashir, M.O. s.n. (K) (11); Batten-Poole, W.H. 226 (K) (23), 371 (K) (11); Battiscombe, E. 495 (K) (16); Batty, M. 131 (K) (27), 870 (K) (27); Baum, H. 490 (K) (10); Baur, R. 34 (K) (10); Bayliss, L. P. 11866 (K) (27); Bayliss, R. D. 3080 (US) (10), 3158 (GH) (13a), 4526 (GH, MO) (10), 5212 (GH) (13a), 7468 (MO) (10), 8112 (MO, US) (6b); Beckett, J. J. 747 (K) (13); Beesley, J. S. S. 252 (K) (10); Belle, E. R. 196 (GH) (17); Ben, V. D. 2441 (K) (6e); Bequaert, C. 35 (GH) (23); Berhaut, R. P. 1630 (K) (6f); Beshir, B. 113 (K) (16), 151 (K) (13), 507 (K) (16), 508 (K) (16), 509 (K) (16), 510 (K) (16), 511 (K) (16); Best, E. B. 52 (K) (11), 74 (K) (17); Bidgood, & J. Lovet 38 (K) (17); Bidgood, et al. 65 (K) (17), 429 (K) (17), 644 (K) (17), 1204 (K) (13); Biegel, H. M. 1912 (K) (4); Bigger, M. 1923 (K) (10); Bingham, M.G. 1178 (MO) (10); Birkett-Smith 49 (K) (26); Bittrimiwx 40 (K) (4); Bjronstad, A. 1294 (K) (4), 1363 (K) (13), 1605 (K) (17); Blackburn, 1863 (GH) (4); Blackmore, et al. 453 (K) (10); Blake, M. 2282 (K) (13); Boaler, S. B. 527 (K)

- (17); Bogdan, 3270A (K) (20), 4535 (K) (16), 4600 (K) (11), 4827 (K) (11); Boicher, E. 1841 (GH) (16); Boland, B. G. 743 (K) (16); Boughey, A. S. 14298 (K) (7), 18052 (K) (1), 18067b (K) (1), 180995 (K) (1); Boulos, L. 10250 (K) (28), 10316 (K, MO) (28), 10509 (K) (28); Bouner, E. 1846 (K) (16); Bounougou, E. 126 (K) (24); Boutique, 308 (K) (16); Bousique, R. 168 (K) (22); Bowson, W. J. 452 (K) (16); Box, R. D. 291 (K) (17), 293 (K) (16); Breteler, F. J. 258 (K) (19), 2565 (K) (23), 7110 (K) (7); Britton, L. L. 4778 (BRE) (18); Broadbent, J. A. 158 (K) (13); Brockman, D. 321 (K) (13); Brod 10503 (K) (11); Brodhurst-Hill 176 (K) (17); Brookes, A. J. 4 (K) (23), 19 (K) (16); Brown, A. J. 1644 (K) (13); Brown, E. 21 (K) (17), 238 (K) (17); Brown, F. 91 (K) (13); Brown, L. 1732 (K) (11), 2027 (K) (6a), 2066 (K) (11), 2185 (K) (13); Brown, R. C. 8282 (K) (13); Brown, & Shapiro 4112 (K) (6b); Bruce, E. M. 841 (K) (4); Brummit R. K. 8428 (K) (17), 8661 (K) (4), 18348 (K) (13); Brummit, R. K. & E. A. Banda 8527 (K) (4), 9838 (K) (14); Brummit, R. K. et al. 13988 (K) (6b), 14239 (K) (10), 14911 (K) (4), 16199 (K) (6e); Brunt, M. 404 (K) (17), 1389 (K) (16); Buchanan 1141 (US) (13), 1174 (K) (27); Buhrmann, A. E. 14 (K) (6b); Bullock, A. A. 2212 (K) (6b), 2293 (K) (4), 2391 (K) (4), 3748 (K) (11), 3770 (K) (4), 3826 (K) (13), B14 (K) (17); Bunke, R. 369 (GH) (10); Burbidge, K. 510 (K) (1), 534 (K) (23); Burchell, W. J. 396 (K) (6b), 2482 (K) (10); Burger, W. 914 (K, US) (13), 2846 (K, US) (13), 3204 (K) (13), 3229 (K) (13), 3489 (K) (13); Burke 443/3591 (K) (10); Burnett, G. F. s.n. (K) (11); Burtt, B. D. 1799 (K) (10), 2105 (K) (17), 2423 (K) (16), 2443 (K) (4), 2448 (K) (16), 2732 (K) (6e), 3770 (K) (13), 5095 (K) (13); Bush, R. P. 23 (K) (17). Carmichael, 390 (K) (10); Carpenter, S. W. 192 (K) (6f); Carter, S. 864 (K) (13), 870 (K) (13); Carter, S. et al. 2251 (K) (13), 2479 (K) (13); Cathorold, 210 (K) (13); Cecil, E. 128 (K) (17), 159 (K) (6b), 247 (K) (27), 259 (K) (10); Chafer, P. A. & A. Nuvunga 6644 (MO) (6b); Champion, A. M. T187 (K) (13); Chancellor, R. J. 5 (K) (17), 56 (K) (16), 72 (K) (11); Chandler, 106 (K) (16); Chapman, J. D. 2935 (K) (11), 3198 (K) (16), 70986 (K) (16); Chapman, J. P. H/579 (K) (13); Chapman, J. W.F. 64 (K) (11); Chapman, & E. G. 7186 (K) (3), 9000 (K) (3); Chase, N.C. 666 (K) (10), 1423 (K) (6b), 6382 (K) (10), 6819 (K) (13), 6853 (K) (13), 6854 (K) (3), 7925 (K) (6b); Cheesman, E. 16 (BM) (10); Chevalier, B. 22.404 (K) (26); Chikawa, I. 676 (K) (13); Choner, H. K. 2005 (K) (4); Cleghorn, W.B. 2357 (K) (3); Codd, L. E. 3729 (K) (13), 9661 (K) (6b); Cole, N. H. 199 (K) (1); Collenette, C. N. 3 (K) (1); Cook, C. D. 358 (K) (5), 386 (K) (11), 445 (K) (6e), 477 (K) (11); Cooper, T. 1349 (K) (10), 2829 (K) (6b); Correia, F. & A. Marques, 4142 (K) (4); Croat, T. B. 53417 (K, MO) (16); Croix, I. F. 2735 (MO) (27); Cruse, A. W. 121 (K) (6b), 372 (K) (6b); Cullett, J. B. 3243 (K) (10); Cunningham, T. F. 18 (K) (10); Curle, N. 98 (K) (16); Curtis, A. G. 707 (GH) (4), 860 (GH) (4), s.n. (GH) (22).
- Dacremont 128 (K) (17), 129 (K) (17); Dale, J. R. 252 (K) (13); Dalziel, J. M. 109 (K) (11), 110 (K) (24), 167 (K) (11), 168 (K) (9), 169 (K) (23), 171 (K) (16), 300 (K) (13), 355 (K) (24), 357 (K) (13), 407 (K) (6f), 711 (K) (23), 911 (K) (23), 1912 (BM) (23), s.n. (K) (7); Damwell et al. 252 (K) (13); Daramola, B. O. 62346 (K) (13); Davey, 27126 (K) (16); Davey, J. T. s.n. (K) (13); Davies, R. M. 2890 (K) (13); Davis, P. 53515 (BM) (13), 55245 (BM) (13); Davis, P. & J. Davis 48448 (BM) (13); Davis, R. L. 2 (K, US) (16), 6 (K) (4), 255 (K) (24); Davy, J. B. 932 (K) (6b); Dawe 120 (K) (16), 444 (K) (16); Dawson, W. J. 533 (K) (6e); De Bell, G. S. 878 (K) (4); de Koning, J. 7703 (K) (3); De Lemes, F. & A. Balsinhas 92 (K) (4); de Menezes, A. 377 (K) (10); de Wilde, J. J. 6399 (MO) (6e), 7223 (K) (28); de Wilde, W. J. 4585 (MO) (23); de Wilde, W. J. & B. E. de Wilde-Duyfjes 2347 (K) (22), 2397 (K) (17), 2954 (K) (11), 3471 (K) (13), 9693 (K) (16), 29629 (K) (13); de Winter, B. 3994 (K) (10), 4087 (K) (10); de Witte, G. F. 3332 (K) (17); Decary, R. 9014 (US) (4); Deighton, F. C. 4494 (K) (23), 5146 (K) (1), 6096 (K) (23); Descoings 12466 (P) (8); Dieterlen, M. 527 (K) (6b), 529 (K) (6b); Dinter, K. 5749 (GH) (4); Dowson, W. J. 674 (K) (11); Draper, 25 (K) (11); Drege, J. F. 92713 (MO) (10), 832608 (MO) (6b), 832649 (MO) (6b); Drummond, R. B. 5503 (K) (13); Drummond, R. B. & A. J. Cookson 6185 (K) (6b), 6451 (K) (6b); Drummond, R. B. & J. H. Hemmings 3319 (K) (13), 3721 (K) (13), 3839 (K) (4), 4086 (K) (20), 4247 (K) (20); Duben, W.V. 1845 (S) (4); Dummer, 2629 (US) (22); Durham, F. R. 26/2/24 (K) (17); Dyer, & Merdmo 3421 (K) (11); Dynes, H. 120A (K) (17).
- Easterhuper, E. 972 (K) (10); Ecklon, & Zeyher 3.10 (MO) (66), 10.11 (GH, MO, US) (6b); Edwards, L. C. 1923 (K) (16); Eggeling, W. J. 350 (K) (6e), 785 (K) (16), 4033 (K) (6a); Ejiofor, M. C. 30240 (K) (16), 30811 (K) (16); Elhadi, N. & A. Khattab 90/188 (S) (4); Elliott, G. S. 128 (K) (16), 1481 (K) (6a), 4174 (K) (23), 4279 (GH) (6f), 5207 (K, MO) (6f), s.n. (K) (6a); Ellis, P. E. 272 (K) (28); Ern et al. 406 (K) (16), 1477 (K) (16), 1478 (K) (13), 1634 (K) (5); Espirito Santo 2796 (K) (16), 2960 (K) (16), 3199 (K) (21), 3520 (K) (16), 3622 (K, MO) (19), 3636 (K) (23), 3643 (MO) (23), 3668 (K, MO) (19), 3754 (K) (19), 3808 (K, MO) (19), 3824 (K, MO) (23); Esterhiner, E. E. 970 (K) (6b); Evan, I. & J. Erens 1645 (K) (16); Evans-Pritchard, E. E. 27 (K) (16); Eviard, C. 2231 (K) (17); Exell, A. & F. Mendonca 1353 (BM) (2); Exell, A. et al. 214 (BM) (10), 1241 (K) (6b); Eyles F. 985 (K) (13), 1274 (K) (13), 2015 (K) (6b), 8799 (K) (3).
- F.A.M. & Hooker, T. 474 (K) (23), 287A (K, MO) (23); F.B. 6431 (K) (4); Faad, G. A. 1881 (K) (16); Faden, A. J. 77/309 (US) (13); Faden, A. J. & R. B. 74/744 (K) (20), 74/1122 (K) (13); Faden, A. J. et al. 74/572 (K) (17); Fames, E. 1905 (K) (11); Fanshawe, D. B. 615 (K) (6b), 2083 (K) (13), 4689 (K) (23), 4920 (K) (6b), 5439 (K) (13), 5543 (K) (13), 5819 (K) (6b), 10003 (K) (11); Farmer, L. 131 (K) (13); Farris, J. 1736 (K) (17); Fauckner, W. 3721 (K) (11), 4877 (K) (13); Faucknor, D. 844 (K) (4); Faucknor, H. 3605 (K) (4); Faucknor 2494 (K) (4); Faulkner, H. G. 90 (K) (4), K118 (K) (27), A376 (K) (2), 661 (K) (27); Fay, J. M. 1896B (K) (22), 2787 (K) (17), 2794 (K) (5), 2867 (K) (17), 2877 (K) (17), 3354 (K) (13), 4083 (K) (23), 5066 (K) (17), 5067 (K) (17), 5068 (K) (17), 5069 (K) (17), 5479 (MO) (6e), 6021 (K, MO) (19), 7057 (MO) (6e); Fay, J. M. & J. Doka 5070 (K) (16); Fitzgerald, D. V. 3235 (K) (17); Forbes, H. W. 1/1929 (K) (4); Fotius 2177 (K) (24), 2219 (K) (6d); Friis et al. 918 (K) (25); Frith, A. C. 147 (K) (5); Froment, D. 588 (K) (16); Fyffe, R. 32 (K) (16), 164 (K) (16).
- Gaepin 1709 (US) (11); Gairdner, A. E. 194 (K) (13), 412 (K) (17); Galpin, E. E. 9768 (K) (6b), 13131 (K) (6b); Gande, P. 39/74 (K) (11); Gardner, H. M. 3517 (K) (13); Gaughan, J. H. 1614 (K) (11); Gbesofssié 3724 (GH, K) (16); Gbile et al. 63394 (K) (13), 64088 (K) (16), 72899 (K) (16); Gbile & B. O. Daramola 245 (K) (23); Geering, C. & J. Bokdam 934 (K) (16), 2029 (MO) (6f); Geilingen 58 (K) (27), 499 (K) (27), 585 (K) (4), 2420 (K) (17), 4788 (K) (17); Germain, R. 4269 (K) (17), 5717 (K) (13), 6666 (K) (11), 6747 (K) (11); Germishuizen, G. 1018 (K) (6b), 3534 (K) (18); Gerrad, W. T. 424 (K) (11); Gerstner, J. 7151 (K) (4); Gilbert, J. B. 4253 (K) (13), 4347 (K) (13), 4722 (K) (13), 4723 (K) (13), 4918 (K) (13), 4938 (K)

- (13), 13104 (K) (13), 17889 (K) (3); Gilbert, J. B. & F. Kariuki 18863 (K) (6e); Gilbert, J. B. & S. P. Kibwua 19932 (K) (11); Gilbert, M. G. 1663 (K) (17), 2412 (K) (13), 4071 (K) (13), 4072 (K) (28), 4840 (K) (17), 6215 (K) (22), 6268 (K) (13); Gilbert, M. G. & A. Getachew 2958 (K) (13); Gilbert, M. G. & S. B. Gilbert 1078 (K) (13), 1250 (K) (6e), 1408 (K) (28), 1438 (K) (13); Gilbert, M. G. & T. G. Jefford 4594 (K) (20); Gilbert, M. G. & M. Thulin 415 (MO) (6e), 708 (K) (17); Gilbert, M. G. et al. 24 (K) (13), 304 (K) (13), 320 (K) (17), 345 (K) (13), 417 (K) (13), 524 (K) (13), 525 (K) (13); Gilbert, S. B. 1122 (K) (20); Gilger, W. 710 (K) (11); Gille 39 (K) (17); Gillespie, J. B. 186 (K) (4), 246 (K) (4); Gillett, J. B. 16369 (K) (16), 17889 (MO) (27); Gillett, J. B. & R. Faden 18152 (K) (13); Gillett, J. B. et al. 5045 (K) (17), 25031 (K) (13), 25136 (K) (13); Gilliland, H.B. k1398 (BM, K) (10); Giville, J. W. 41 (K) (4); Glanville, R. R. 241 (K) (6f); Gledhill, D. 285 (K) (6f), 328 (K) (23), 942 (K) (23); Glover, P. E. & H. B. Gilliland 1231 (BM, K) (20); Glover, P. E. & Samuel 2722 (K) (17); Glumptre, E. D. 25 (K) (23); Goodnny, H.V. 101 (K) (13); Goodies, R. 69 (K) (13); Gossweller, J. 199 (K) (16), 329 (K) (13), 558 (BM) (23), 647 (K) (13), 1466 (K) (4), 2234 (K) (6b), 2916 (K) (17), 3504 (K) (2), 3540 (K) (13), 4545 (K) (23), 14219 (K) (1); Goyder, D. J. & A. J. Paton 3515 (K) (11); Goyder, D. J. & A. R. Smith 3216 (K) (13); Goyder et al. 3631 (K) (17); Graham, M. A. 2405 (K) (6e); Graham, M. D. 3031 (K) (13); Graham, R. M. 272 (K) (4), 497 (K) (13), 1577 (K) (27), 1675 (K) (13), 1701 (K) (22), 2154 (K) (22); Grandraux, L. A. & F. Delemos 7913 (K) (11); Grant, A. L. 3141 (MO) (6b), 3505 (MO) (10), 3570 (MO) (6b), 2212387 (MO) (10); Greenstock, W. 1613834 (MO) (10); Greenway, P. J. 1204 (K) (4), 2274 (K) (17), 4358 (K) (13), 4462 (K) (6e), 5097 (K) (4), 5330 (K) (27), 5371 (K) (17), 7392 (K, US) (16), 7473 (K) (6e), 7840 (K, S) (10), 10519 (K) (4), s.n. (K) (13); Greenway, P. J. & J. M. Brennan 8095 (K) (6b), 8213 (K) (6b); Greenway, P. J. & Kanuri 11245 (K) (13), 11870 (K) (13), 14220 (K) (4); Greenway, P. J. & R. M. Polhill 11305 (K) (4), 11736 (K) (4); Greenway, P. J. & S. P. Rawlins 9463 (K) (11); Grosvenor, R. K. 144 (K) (13); Greenway, P. J. et al. 13160 (K) (13), 14082 (K) (13); Grosvenor, R. & J. Renz 1038 (K) (3); Guerra, J. A. 3812 (K) (11); Guile, D. P. 1662 (MO) (23).
- Haarer, Q. E. 177 (K) (13), 338 (K) (17), 517 (K) (6e), 596 (K) (17), 637 (K) (10), 1767 (K) (17), 2307 (K) (11), 5127 (K) (13), 5128 (K) (13), s.n. (K) (17); Hagerup, O. 478 (K) (16); Hakki, et al. 289 (K) (6e), 309 (K) (6e), 1504 (K) (24); Hale, M. 105 (K) (20); Hall, J. B. 90 (K) (11), 131 (K) (6f), 260 (K) (6f), 400 (K) (16), 452 (K) (5), 558 (K) (11), 1392 (K) (6e), 1998 (K) (26), 2028 (K) (24), 2058 (K) (13), 3079 (K) (5), s.n. (K) (7); Hall, J. B. & B. O. Daramola 67404 (MO) (6e); Hall, J. B. & Swaine GC46220 (US) (24); Halle, N. & Villiers 4914 (P) (15); Hambler, D. J. 768 (K) (23); Hansen, O. J. 854 (K) (20), 3016 (K, MO) (6b), 3375 (K) (13); Hardy, D. S. 961 (K) (13); Hardy, D. S. & P. R. O. Bally 10826 (K) (13); Harrison, 2 (K) (11); Harlan, H. V. 1029870 (US) (16); Harold, J. M. 903 (K) (10); Harrer, A. E. 2227 (K) (6a); Harris, B. J. 1774 (K) (13), 2523 (K) (11); Harris, B. J. & A. McCusker BJH2568 (MO) (27); Harris, E. 4 (K) (7), 5 (K) (16), 6 (K) (23), 7 (K) (7), 8 (K) (23), 9 (K) (16), 10 (K) (19), 11 (K) (6f); Harris, L. D. 9 (K) (20); Harris, T. M. 195 (K) (7), 786 (S) (27); Hayes, T. R. 539 (K) (6f); Haygarth, W. 92714 (MO) (10); Hazel, C. 289 (K) (16); Hazel, G. 442 (K) (11); Hazel, O. 521 (K) (17); Hedbert, O. 7022 (UPS) (4); Hemming, C. F. & R. M. Watson 30216 (K) (13); Henriques, C. 853 (K) (17); Hepper, F. N. 1162 (K) (16), 1218 (K) (16), 1278 (K) (16), 1392 (K) (23), 1397 (K) (19), 1847 (K) (16), 2745 (K) (17), 3721 (K) (16), 3759 (K) (16), 3915 (K) (23), 7333 (K) (3), 7368 (K) (3), 7373 (K) (4), 7377 (K) (13); Hepper, F. N. & D. V. Field 5218 (K) (4); Hepper, F. N. & A. Jaeger 6618 (K, MO) (10); Hepper, F. N. & J. K. Morton 2314 (K) (6e), A3039 (K) (6f), A3050 (K) (6e), A3110 (K) (6f), A3160 (K) (16); Het-tawy, A. W. 52 (K) (13), 179 (K) (13), 186 (K) (13); Hildebrandt, J. M. 1907 (UPS) (27); Hill, A. W. 51 (K) (6f); Hill, B. 755 (K) (6a); Hilliard, O. M. & B. L. Burtt 7861 (MO) (10), 10012 (MO) (6b), 12139 (K) (10); Hindrot 818 (K) (16); Holub, E. 1834 (K) (10), 1883 (K) (11), 1993 (K) (13); Hooper, S. S. & C. C. Townsend 503 (K) (6b); Hopkins, J. C. 9712 (K) (11), 14555 (K) (13); Hornby, R. M. 2898 (K) (11); Hossain, & Agyakwah 37826 (K) (5), 37829 (K, S) (5), 37831 (K) (13), 37954 (K) (16), 37855 (K) (16), 37957 (K, US) (16), s.n. (K) (5); Howaidou 1599 (K) (22); Hudson, D. 64 (K) (17), 296 (K) (17), 2566 (K) (17); Hughes, N. K. s.n. (ODU) (10); Huntley, B. J. 192 (MO) (6b); 2033 (MO) (10); Hutchinson, J. 2381 (K) (6b), 2586 (K) (6b), 2854 (K) (10); Hutchinson, J. & J. B. Gilbert 3537 (K) (3); Hynes, A. 478 (K) (17), 576 (K) (16).
- Innes, R. R. 373 (K) (27), 909 (K) (6f), 30116 (K) (16), 30748 (K) (16); Irvine, R. 146 (K) (6e), 466 (K) (6e), 1426 (K) (6e), 1620 (K) (6e), 4534 (K) (16), 4572 (K) (13), 4574 (K) (13), 4581 (K) (16), 4667 (K) (5), 4668 (K) (7), 4669 (K) (7), 4732 (K) (7), 4914 (K) (5), 4915 (K) (6e), s.n. (K) (6e); Ithwa, R. 1370 (US) (16); Iverson, et al. 87612 (K) (13).
- Jackson, G. 649 (K) (6b), 1656 (K) (4), 2282 (K) (3), U103 (K) (17); Jackson, J. A. D. 119 (K) (16), 254 (K) (11); Jackson, J. K. 2722 (K) (13), 3196 (K) (17); Jackson, W. B. 415 (K) (17); Jacques-Felix, 1241 (P) (21); Jaeger, H. D. 1225 (K) (11); Jaeger, P. 753/51 (K) (6c), 790 (K) (23), 2541 (K) (6c); Jahandiez, E. 149 (BM) (13), 202 (BM) (13); James, E. 1905 (K) (11), 1905 (K) (16), 1905 (K) (17); Jansen, P. C. 8040 (K) (13); Jarrett, T. 103 (K) (6e); Jasen, P. C. 8158 (K) (4); Jeagu, A. J. 84 (K) (4); Jeffery, G. M. K242 (K) (4); Jefford, et al. 53 (K) (13); Jex-Blake, M. 6880 (K) (20); Joanna, B. 8928 (K) (20), 8947 (K) (13); Jacobsz, M. L. 218 (K) (10), 2011 (K) (6b); Johannes, 26–1–22 (US) (10); Johnson, W. P. 32 (K) (13), 101 (K) (4); Johnston, A. H. 83 (K) (11); Johnstone, M. 1551 (K) (17); Jones, A. 3660 (K) (16), 7164 (BM) (23); Jones, K. W. 41 (K) (4), 501 (K) (16), 502 (K) (16), 503 (K) (16), 504 (K) (16), 505 (K) (16), 506 (K) (16), 512 (K) (16), 513 (K) (16), 514 (K) (16), 515 (K) (16), 516 (K) (16), 517 (K) (16), 518 (K) (16), 519 (K) (16), 521 (K) (16), 522 (K) (16), 523 (K) (16), 524 (K) (16), 525 (K) (16), 526 (K) (16), 527 (K) (16), 528 (K) (16), 529 (K) (16), 530 (K) (16), 531 (K) (16), 532 (K) (16), 533 (K) (16), 534 (K) (16), 535 (K) (16); Jones, V. 3 (K) (11); Jordan, H. D. 152 (K) (23), 190 (K) (6f); Junod, H. A. 276 (BRE) (18).
- Kaessner, T. 474 (K) (20), 645 (BM, K) (20), 1474 (BM) (20); Kassner Exped., 190 (K) (17), 2397 (K) (6b), 2786 (K) (16), 3148 (K) (16); Keay, R. W. J. 21674 (K) (16), 25446 (K) (13), 25833 (K) (6f), 25896 (K) (6e), 25977 (K) (24), s.n. (K) (6f); Kelly, R. D. 80 (K) (10); Kemp, S. 701 (MO, US) (6b); Kennedy, J. D. 2894 (K) (16); Kenyon, E. 1901 (K) (13); Kerfoot, O. 1313 (K) (13), 3815 (K) (13); Keritochonorag, H. 2166 (K) (16); Kers, L. E. 1399 (S) (24); Kersting, 665 (K) (7); Kesollesen, 2817 (K) (27); Kew, J. 19911 (MO) (27); Kilian, N. 1833 (K) (13); Killick, D. J. 3857 (K) (6b); Killick, H. J. 234 (K) (11); Killick & Leisner 3415 (K) (11); Kimani, Z. J. 186 (K) (13); King, A. E. 260 (K) (11); King, E. L. 175 (K) (23),

- 175B (K) (23); Kirk, J. 15/1/63 (K) (11), 2179 (K) (17), 9/68 (K) (4), 1 (K) (13); Knepper, D. K. 8/22/89 (ODU) (13h), 8/24/89 (ODU) (13h), 8/25/89 (ODU) (17), 8/26/89 (ODU) (13d), 8/26/89 (ODU) (16), 8/26/89 (ODU) (5), 8/25/89 (ODU) (5), 9/20/89 (ODU) (16), 9/22/89 (ODU) (16), 9/23/89 (ODU) (16), 9/23/89 (ODU) (5), 9/24/89 (ODU) (16), 9/26/89 (ODU) (5), 9/19/89 (ODU) (13g), 9/21/89 (ODU) (5), 9/23/89 (ODU) (5); Konging, J. de & D. Zunguze 8038 (K) (6b); Korans, J. 1508 (K) (10), 1596 (K) (13), 3060 (K) (6b), 3299 (K) (6b), 3364 (K) (13); Krauss, J. 213 (MO) (6b); Koritschoner, H. 2059 (K) (11); Kuchar, P. 15312 (K) (13); Kuntze, O. 7/iv94 (US) (4), 94 (K) (13), 1913 (K) (4).
- La Croiz, I. F. 2735 (MO) (3); Laferrere, M. 112 (K) (26); Lamb, P. H. 4 (K) (16); Lambrecht, D. 479 (K) (11); Lambrecht, F. L. 31 (K) (13), 264 (K) (6b), 440 (K) (6b); Lamprey, H. F. 515 (K) (6e); Lanbrestre, C. H. 18 (K) (6a); Langdale-Brown, I. 2392 (K) (16), 2421 (K) (17); Last, J. T. 1887 (K, US) (4), 1908 (K) (17); Latilo, M. G. 18214 (K) (16); Latilo, M. G. & B. O. Daramola 28792 (K) (16); Lavranos, J. J. 4265 (MO) (10), 7394 (K) (20); Lavranos, J. J. & Cater 23192 (K) (13); Lavranos, J. et al. 24938 (K) (13); Lawlor, D. W. & B. J. Hall 28 (K) (11), 34 (K) (24), 73 (K) (24), 461 (K) (24), 468 (K) (13), 508 (K) (7), 46508 (K) (5); Lawrence, E. 15 (K) (4), 28 (K) (4), 163 (K) (4); Lea, A. LR20 (K) (11); Lea, J. D. 10 (K) (27), 17 (K) (16), 58 (K, ODU) (24), 180 (K) (24); Leach, L. C. 9189 (K) (27), 11232 (K, MO) (27); Leach, L. C. & R. F. Smith 11011 (K) (27); Lean, O. B. 37 (K) (11); Lebrun, J. 1759 (GH) (16), 3374 (GH, K) (16), 3535 (MO) (13), 3557 (GH) (11), 9631 (K) (16); Lecard, F. s.n. (K) (16); Ledermann, C. 4529 (BM) (6d); Lee, J. 16 (K) (11); Leeuwenberg, A. J. 4357 (K, MO) (6f), 10189 (K) (16); Leighton, J. M. 2948 (MO) (6b); Leistner, O. A. 1817 (K) (13); Lehchase, N. 3673 (MO) (10), 3764 (MO) (10); Lely, H. V. 146 (K) (6e), 255 (K) (11), 489 (K) (13), 535 (K) (13), 590 (K) (16), 680 (K) (19), P83 (K, MO) (6f), P369 (K) (11), P567 (K) (7), P785 (K, MO) (23); Lester, J. B. 34 (K) (23); Lester, R. N. & M. Morrison, 2337 (K) (6a); Letouzey, R. 624 (K) (6f), 6299 (K) (11), 6406 (K) (16), 7209 (K) (16); Levy, B. 1177 (K) (13); Lewalle, H. J. 335 (K) (11), 1474 (K) (13), 2402 (K) (17), 10634 (BM) (13); Liben, L. 976 (K) (13), 2038 (K) (11), 2401 (K) (4), 3705 (K) (11); Liebenberg, L. C. C. 1029 (K) (16), 1532 (K) (11), 4942 (K) (13), 6904 (K) (6b), 6986 (K) (10), 8581 (MO) (10), 8748 (MO) (10), A516 (K) (13); Lind, E. 340 (K) (6a), 387 (K) (16), 2027 (K) (6a); Linder, D. H. 1927 (GH) (22), 1966 (K) (13); Linoblom, K. G. 1911 (S) (10); Lisowski 80551 (BR) (6c), 80579 (BR) (6c); Lloyd, H. 80 (K) (10); Lobin, W. 6687 (K) (13); Lock, J. M. 81/77 (K) (13), 81/132 (K) (5), 81/247 (K) (17), 81/248 (K) (24), 81/2658 (K) (13), 82/31 (K) (11), 82/44 (K) (13); Longnot & Verdcourt 3200 (K) (6e); Loope, P. P1048 (K) (17); Lord, J. K. 8169 (K) (13); Lovett, J. 1612 (K) (13), 1637 (K) (4); Lovett, J. & C. Congdon 1404 (K) (13), 1931 (K) (13); Lowe, J. 1379 (K) (16), 1484 (K) (23), 1671 (K) (16), 2760 (K) (16), 3378 (K) (6f), 4463 (K) (23), 4828 (K) (16), A12 (K) (13); Lugard, E. J. 127 (K) (13), 164 (K) (13); Luke, W. R. 2459 (K) (11), 3054 (K) (13), 3390 (K) (13), 3489 (K) (20); Lye, K. A. 4747 (K) (11), 5379 (K) (13), 9078 (K) (13), s.n. (K) (17); Lynes, H. 18 (K) (10), 53 (BM) (13), 577 (BM) (11), 578 (BM) (24).
- Mabberley & McCall 86 (K) (17); Macaulay 867 (K) (6b), 927 (K) (6b); MacCintosh, D. G. 103 (K) (16); MacDonald, W. 21 (K) (13); MacOwan, L. 426 (GH) (10); Magajgi, S.O. M6660 (K) (16); Magogo, F. C. 373 (K) (27); Magogo, F. C. & R. Estes, 1152 (K) (17), 1152 (K) (22), 1153 (K) (17), 1153 (K) (22); Magogo, F. C. & P. Glover 1050 (K) (27), 1089 (K) (4); Magogo, F. C. & R. R. Innes 849 (K) (27); Maitland, J. 103 (K) (11), 1926 (K) (16), 1946 (K) (16), s.n. (K) (11), s.n. (K) (16); Mansfield, R. A. 9/4/78 (ODU) (22), 9/4/78 (ODU) (7); Marmo, V. 36 (K) (23); Masierson, G. 217 (K) (3); Massey, R. E. 47 (K) (16); Mathew, B. 6126 (K) (6e); Mauteiro 6193 (K) (13); Mavi, S. 626 (K) (6b); McCintosh, D. G. H. 3k (K) (17), 57k (K) (11), 109k (K) (11); McClean, A. 595 (K) (6b); McClean, B. 209 (K) (6b); McClounie, T. 60 (K) (4), 178 (K) (3); McDonald, J. 910 (K) (16); McGillett 1007 (K) (13); Mearns, E. A. 2885 (GH) (11), 3005 (GH) (16); Meel, L. V. 1432 (K) (17); Meikle, R. D. 746 (K) (16), 768 (K) (23), 1011 (K) (6a); Melville, F. A. & T. Hooker 287a (K, MO) (23); Mesfin et al. 4058 (K) (6e); Methew, L. 162 (K) (6b), 257 (K) (27); Meurillon, A. 1363 (K) (5), CNAD577 (K) (23); Meurillon, A. et al. 1976 (K) (16); Meyer, F. G. 7594 (K, US) (16), s.n. (K) (16); Meyers, J. G. 7329 (GH, K) (5), 7867 (GH) (19); Mgaza, C. D. 434 (K) (17); Michael 9531 (K) (6d), 9706 (K) (6d); Michel, B. 13 (K) (11); Michel, G. 5367 (K) (4); Migeod, F. W. 13 (BM) (27), 35 (BM) (27), 50 (BM) (27), 730 (BM) (27); Mildraed, 1-232 (K) (17), 9501 (K) (17); Miller, H. 3/1896 (K) (23); Miller, O. B. 1590 (MO) (6b), 1939 (K) (13), 2630 (K) (6b), 2671 (K) (6b), 4827 (K) (6b), 5737 (K) (6b), 7260 (K) (6b); Miller et al. 615 (BM) (13); Milne-Redhead, E. 3577 (K) (6b), 3756 (K) (17), 4018 (K) (10), 4155 (BM, K) (11), 4161 (K) (2), 5043 (K) (6f); Milne-Redhead, E. & B. Mhroro 9693 (K) (4); Milne-Redhead, E. & P. Taylor 882 (K) (17), 8421 (K) (27), 9524 (K) (13), 10018 (K) (13), 10024 (K) (3), 10577 (K) (3); Mitchell, B. L. 24/78 (K) (11); Mitchison, N. 1975 (K) (13), 1975 (K) (4); Mitte 314 (US) (16); Mogg, A. O. 5935 (MO) (6b), 30317 (K) (17); Moggridge, F. Y. 548 (K) (4), 1868 (US) (16); Mohamed, K. I. 10/25/88 (ODU) (13b), 10/26/88 (ODU) (16), 10/26/88 (ODU) (13e), 10/27/88 (ODU) (16); Moiser, B. 63 (K) (13), 9 (K) (16); Moll, E. J. 4723 (K) (6b); Moowam, J. C. 953 (K) (27); Mooney, H. F. 8427 (K) (17); Moriarty, A. 441 (K) (17), 469 (K) (27); Morton, J. K. 94 (K) (23), 1104 (K) (23), 1677 (K) (13), 2527 (K) (19), 6123 (K) (6e), 6214 (K) (16), 6224 (K) (16), 6292 (K) (19), 7171 (K) (6e), 8794 (K) (6e), 8795 (K) (6f), 9370 (K) (23), A663 (K) (6f), A1394 (K) (16), A2045 (K) (6e), A3016 (K) (6f), A3295 (K) (7), A3488 (K) (6e), A3926 (K) (6e), A4241 (K) (26), GC8601 (K) (26), s.n. (K) (5), s.n. (K) (23); Morton, J. K. & J. M. Dalziel 145 (K) (6e); Morton, J. K. & D. Gledhill 1093 (K) (1), 3035 (K) (19), 18771 (K) (1); Moucoulie, A. 574 (K) (13); Mouteiro 6143 (K) (23); Mugwedi, E. 1371 (K) (4); Muller, P. J. 2205 (K) (6b); Murray, A. L. 426 (K) (10); Musselman, L. J. 124 (K) (11), 198 (K) (24), 6118 (ODU) (7), 6122 (ODU) (5), 6123 (ODU) (5), 6124 (ODU) (5), 6131 (ODU) (7), 6139 (K, ODU) (16), 6140 (ODU) (16), 6143 (ODU) (16), 6144 (K, ODU) (16), 6145 (K, ODU) (16), 6152 (ODU) (16), 6154 (K) (16), 6155 (K) (16), 6162 (K, ODU) (16), 6174 (ODU) (13), 6180 (ODU) (13a), 6212 (K) (16), 6231 (K) (24), 6256 (K) (24), 6260 (K) (17), 6269 (ODU) (13d), 6279 (K, ODU) (16), 7003 (ODU) (4), 7011 (ODU) (6b), 7015 (ODU) (4), 7022 (ODU) (4), 7030 (ODU) (10), 7032 (ODU) (6b), 7038 (ODU) (10), 7039 (ODU) (10), 7040 (ODU) (6b), 7060 (K, ODU) (5), 7061 (ODU) (24), 7064 (ODU) (13c), 7085 (K, ODU) (5), 7086 (ODU) (11), 4/10/86 (ODU) (11), 4/9/86 (ODU) (11), 8/23/86 (ODU) (17), 8/23/86 (ODU) (13c), 8/23/86 (ODU) (24), 8/23/86 (K) (24), 10/13/88 (ODU) (5), 10/13/88 (ODU) (11), 10/16/88 (ODU) (19), s.n. (K) (24), s.n. (K) (17); Musselman, L. J. & R. A. Mansfield 5522 (ODU) (23), 5524 (ODU) (5), 5527 (ODU) (16), 5533 (ODU) (19),

- 5535 (ODU) (16); Musselman, L. J. & K. I. Mohamed 9/30/88 (ODU) (6e), 9/30/88 (ODU) (7), 10/1/88 (ODU) (5), 10/1/88 (ODU) (7), 10/1/88 (ODU) (13h), 10/1/88 (ODU) (16), 10/1/88 (ODU) (17), 10/2/88 (ODU) (5), 10/2/88 (ODU) (7), 10/2/88 (ODU) (17), 10/2/88 (ODU) (22), 10/4/88 (ODU) (5), 10/4/88 (ODU) (7), 10/4/88 (ODU) (17), 10/4/88 (ODU) (22), 10/4/88 (ODU) (24), 10/5/88 (ODU) (5), 10/5/88 (ODU) (22), 10/6/88 (ODU) (5), 10/6/88 (ODU) (7), 10/6/88 (ODU) (13c), 10/8/88 (ODU) (5), 10/8/88 (ODU) (17); Musselman, L. J. & J. H. Visser 4/2/86 (ODU) (13); Musselman, L. J. et al. 7 (K) (19), 064 (ODU) (13c), 7063 (ODU) (24), 7066 (ODU) (13c), 4/9/86 (ODU) (11), 4/17/86 (ODU) (4), 9/4/86 (ODU) (4), 10/13/88 (ODU) (17), 10/13/88 (ODU) (5), 10/16/88 (K, ODU) (6d), 10/16/88 (ODU) (23), 10/16/88 (ODU) (13g), s.n. (K) (19), s.n. (K) (23); Mutch, M. I. 27952 (K) (16); Mutimushi, J. M. 537 (K) (10), 932 (K) (6b); Mwasumbi, L. B. 10487 (K) (11); Myers, J. G. 6656 (K) (17), 6997 (K) (11), 7000 (K) (17), 7329 (K) (5), 7336 (K) (17); Myres, R. F. 294 (K) (17), 303 (K) (6b).
- N.D.H 7361 (K) (13); Nafuer, E. R. 5343 (K) (16); Napier, E. R. 51 (K) (4), 957 (K) (20); Napper, D. & R. Faden 1839 (K) (20); Nation, O. 29 (K) (10), 143 (K) (13), 313 (K) (6b); Ncube, M. 75 (K) (11); Nelson, W. 394 (K) (10); Nesbould, J. G. 1958 (K) (4); Newberre, R. J. 6 (K) (23); Newbould, J. G. B. 1958 (K) (13); Ngoni, J. F. 160 (K) (13), 295 (K) (13), 1249 (K) (6b); Niamir, M. 17 (K) (13); Nicholas, A. & V. D. Berg 1736 (K) (6b); Nicholas, A. & D. Neave 2109 (K) (6b); Nichols, E. H. 11 (K) (23); Niger 495 (UPS) (16); Nilson, J. 332 (K) (6e); Noak, S. L. 311 (K) (10); Noble, H. L. 30 (K) (16), 31 (K) (16); Norlindh, T. & H. Weimarck 4878 (K) (3), 4956 (K) (17), 44347 (K) (6b); Norman, E. M. 294 (K) (16); Norrgrann, G. 459 (K) (4); Nuvunga, A. 471 (K) (3).
- Okafor, J. C. 46885 (K) (23); Olmuntus, B. 2507 (K) (6e); Olorunfemi, J. 55046 (K) (16); Olufsen, O. 479 (K) (13), 705 (K) (23), 1927 (K) (23); Onochie, C. F. A. 18697 (K) (6e); Onsow, M. P. 709 (K) (6f); Onwudinjo, D. C. 24005 (K) (11); Ortendahl, V. 487 (UPS) (13a); Ossent, J. 478 (K) (13).
- Padva, J. H. 326 (K) (17); Paget-Wilkes, C. F. 700 (MO) (27); Panayotis, L. J. 98 (K) (17); Parker, C. 985 (GH) (23), 1066 (ODU) (11), 1316 (ODU) (13f), 1976 (K, ODU) (5), 1977 (ODU) (5), 1978 (K) (5), 1979 (K) (16), 1994 (ODU) (16), 2023 (K) (13), 2031 (K, ODU) (13d), 2031 (K) (13), 2044 (K) (13), 2046 (K) (5), 2210 (ODU) (11), 2264 (ODU) (13h), 2265 (ODU) (13f), 2306 (ODU) (16), 2312 (ODU) (13d), 2314 (ODU) (5), 2316 (ODU) (16), 2317 (ODU) (16), 2318 (ODU) (5), 2321 (ODU) (16), 2332 (ODU) (16), 2443 (ODU) (4), 2682 (K) (7), 2984 (ODU) (20), 3215 (ODU) (5), 4001 (ODU) (22), 4034 (K, ODU) (20), 4041 (ODU) (17), 4065 (K) (5), 4074 (K) (17), 4075 (ODU) (16), (K) (16), E22 (K) (16), GM/522/H (K) (20); Parkinson, V. 1985 (ODU) (24), 9/18/85 (ODU) (22), 9/17/86 (ODU) (7), A1 (K) (24), A3 (K) (16), A4 (K) (16); Parsons, D. A. L79 (K) (16); Pasch 7846 (K) (16), 7916 (K) (16); Patel, I. H. & W. Nachamba 1775 (K) (11); Patterson, A. 59 (K) (11); Paulo, S. 105 (K) (27); Pawek, J. 1649 (K) (17), 3349 (K) (4), 4112 (K) (27), 5135 (K) (13), 5266 (K) (13), 7799 (K) (6e), 8301 (K) (17), 9499 (K) (17), 9575 (K) (17), 9601 (K) (13), 10942 (K) (17), a12015 (MO) (27), b12015 (K) (3), 12317 (K, MO) (6e), 12566 (K) (4), 13856 (K) (17), 13913 (K) (11), 13954 (K, MO) (11), 14332A (K) (14), 14332B (K) (11); Pearson, H. H. 2038 (K) (13), 2421 (K) (13), 8122 (K) (13); Perceval, A. B. 1901 (MO) (10); Pereira, J. A. 2328 (K) (19), 2368 (K) (19), 2984 (K) (21), 2984 (K) (6a), 3580 (K) (16); Peters,
- 8 (K) (13); Peterson, C. 823 (K) (17), 833 (K) (10); Petherick 1862 (K) (17), 1863 (K) (16); Petterson, B. 16819 (K) (16); Pfund, J. 86 (K) (24), 679 (K) (16); Philcox, D. & R. B. Drummond 070 (K) (13), 9050 (K) (17), 9054 (K) (3), 9094 (K) (13); Philcox, D. & M. J. Leppard 8574 (K) (13), 8575 (K) (13), 8677 (K) (24), 8718 (K) (13), 8870 (K) (17); Philcox, D. & T. Muller 9080 (K) (6b), 9083 (K) (13); Philcox, D. et al. 8563 (K) (17), 8610 (K) (17), 8748 (K) (13), 8796 (K) (11), 8799 (K) (13), 8933 (K) (10), 9017 (K) (6b), 9019 (K) (10), 9093 (K) (3), 9127 (K) (3), 9873 (K) (4), 9908 (K) (17), 10170 (K) (6b), 10190 (K) (17), 10210 (K) (14); Phillips, E. 661 (GH, MO) (6e), 832 (K, MO) (6e), 1215A (MO) (6e); Phillipson, P. B. 138 (MO) (6b); Phipps, J. B. 342 (K) (6b), 2594 (K) (4); Phiri, R. 79 (K) (4); Pichi-Sermolli, R. 1295 (K) (16), 1296 (K) (13); Pielou, D. P. 130 (K) (13), s.n. (K) (17); Plant, H. 706 (K) (6a); Plowes, D. C. 32013 (K) (13); Pobeguin, M. 364 (K) (6f), 1020 (K) (5); Polhill, C. 496 (K) (17); Polhill, R. & S. Paulo 579 (K) (13), 950 (K) (13); Poulo, S. 105 (K, MO) (27); Powell, J. H. 876 (K) (4); Powys, 197 (K) (10); Prince, S. D. 116 (K) (4), 200 (K) (24), 215 (K) (11), 401 (K) (24); Probegum, M. 364 (K) (6e); Prosser, D. U. 1643 (K) (6b), 1784 (BRE, K) (18); Puff, C. & E. Kelbessa 921220 (K) (20); Puff, C. et al. 8209 (K) (20), 820922 (K) (16); Purseglove, J. W. 487 (K) (16), 560 (K) (16), 1609 (K) (6e), 2053 (K) (11), 2711 (K) (6a), P2424 (K) (17); Pusch, 2689 (K) (4).
- Quarre, 1451 (K) (11).
- Radjil 594 (GH) (16); Ragslane 1251 (US) (6e); Ramaiyah, K.V. 2973 (K) (24); Ranger, G. A. 304 (K) (10); Rax, R. D. 292 (K) (24); Raymond, R. J. 319 (K) (17); Raynal, A. R. 2282 (K) (7), 22057 (K) (7), 22063 (K) (13), 22127 (K) (5), 22139 (K) (7), 22149 (K) (5), 22181 (K) (7), 22194 (K) (16), 22196 (K) (7), 22200 (K) (16), 22210 (K) (16), 22211 (K) (13), 22220 (K) (16), 22221 (K) (16), 22244 (K) (13), 22263 (K) (16), 22277 (K) (24), 22290 (K) (5), 22309 (K) (5), 22394 (K) (5), 22395 (K) (5), 22435 (K) (7), 22526 (K) (16), 22565 (K) (13), 22573 (K) (16), 22581 (K) (5), 22602 (K) (16), 22603 (K) (13), 22613 (K) (13), 22636 (K) (13), 22640 (K) (7), 22641 (K) (24); Read, J. G. 10 (BM, K) (10), 27 (BM, K) (10); Reading University/BM 292 (BM) (13); Red, E. 1424 (K) (17); Reed, S. W. 16290 (K) (11); Reed 14 (K) (16), 3/1888 (K) (4); Reekmans, M. 4257 (K) (17), 6786 (K) (6e), 6926 (K) (13), 7125 (K) (6a), 7166 (K) (6a), 9834 (K) (13), 10314 (K) (17), 23/275 (K) (11); Rehmann, A. 4671 (K) (13); Rensburg, H. J. 1516 (K) (13), 2772 (K) (11), 23107 (K) (13); Revell, R. J. 321 (K) (17); Richards, H. M. 657 (K) (13), 668 (K) (11), 676 (K) (6b), 1169 (K) (6b), 1771 (K) (27), 4205 (K) (6b), 4512 (K) (13), 4699 (K) (6b), 4780 (K) (11), 4813 (K) (13), 5230 (K) (4), 5423 (K) (13), 7425 (K) (6b), 9026 (K) (13), 9103 (K) (11), 10542 (K) (6e), 10763 (K) (6b), 10921 (K) (13), 10953 (K) (6b), 12222 (K) (11), 12486 (K) (13), 12570 (K) (6b), 12813 (K) (13), 14824 (K) (11), 14858 (K) (13), 15025 (K) (6b), 16275 (K) (6b), 17147 (K) (6b), 17965 (K) (13), 17980 (K) (4), 19831 (K) (13), 24360 (K) (4), 24897 (K) (20), 25019 (K) (10), 25389 (K) (6e), 25465 (K) (13), 26863 (K) (20), 28461 (K) (13); Richie, A. H. 1924 (K) (4); Riches, C. R. 226 (K) (16), 228 (K) (16), 230 (K) (5), 233 (K) (5), 234 (K) (16), 242 (K) (16), 243 (K) (7), 249 (K) (16), s.n. (K) (5); Robert, G. 16608 (K) (6a); Robertson, S. A. 108 (K) (4), 676 (K) (11), 4121 (K) (20); Robertson, V. C. 5 (K) (11), 31 (K) (16); Roberty 16608 (K) (21), 16621 (K) (23); Robin, R. J. 3839 (K) (13), 4339 (K) (13), 4513 (K) (13); Robins, S. 902 (K) (13); Robinson, E. A. 637 (K) (23), 1214 (K) (3), 1847 (K) (6b), 1961 (K) (10), 2176 (K) (13), 3330 (K)

- (6b), 4024 (K) (6b), 6174 (K) (10), 6403 (K) (11), 6753 (K) (6b), 6897 (K) (13); Robson, N. K. B. 1083 (K) (17), 1094 (BM, K) (3), 1169 (BM) (10), 1487 (K) (13); Robson & Jackson 1179 (K) (10), 1199 (BM, K) (11); Robyn, L. 1971 (K) (17), 1980 (K) (13), 2216 (K) (16); Rodin, R. J. 9379 (K) (16); Roger 758 (K) (19); Rogers 1821 (GH) (16), 20943 (US) (10); Rogers, C. G. 560 (K) (10), 709 (K) (11), 740 (K) (11); Rogers, D. 96 (US) (4), 1493 (US) (6b); Rogers, F. A. 7023 (K) (13), 8215 (K) (13), 18297 (K) (6b); Rose, F. 10117 (K) (17), 10253 (K) (17); Ross, J. H. 1743 (K, MO) (10), 2240 (K) (6b); Rounce, N. V. 288 (K) (4), 355 (K) (11), 358 (K) (13), 405 (K) (13); Rowland 1893 (K) (23); Rudatis, F. 543 (US) (6b), 1829 (S) (6b); Rushworth, J. E. 1489 (K) (17); Russell, J. 41 (K) (6b); Russell, T. 4 (K) (17); Rutherford-Smith, R. O. 542 (K) (4), 565 (MO) (6b); Ruxton, J. P. 53 (K) (13), 67 (K) (11), 149 (K) (11), 152 (K) (5); Rwaburindore, P. K. 580 (MO) (6e), 817 (MO) (6e), 1813 (US) (11); Ryan, J. B. H40/33 (K) (11); Ryan, J. D. 28 (K) (16), 58 (K) (16); Ryder, E. F. 102 (K) (10); Ryding, O. 120 (UPS) (16); Ryding, O. et al. 1548 (K) (16), 1567 (K) (16).
- Saadou, M. 1839 (K) (5); Safa, S. B. & L. J. Musselman 10/3/87 (ODU) (6e), 10/9/87 (ODU) (13g), 10/9/87 (ODU) (16), 10/9/87 (ODU) (24), 10/9/87 (ODU) (7), 10/10/87 (ODU) (13h), 10/10/87 (ODU) (5), 10/10/87 (ODU) (7), 10/12/87 (ODU) (5), 10/12/87 (ODU) (7), 10/13/87 (ODU) (22), 10/13/87 (ODU) (5), 10/13/87 (ODU) (7), 10/19/87 (ODU) (24), 10/21/87 (ODU) (6e), 10/21/87 (ODU) (7), 10/25/87 (ODU) (17), 10/25/87 (ODU) (22); Sallinho Botanical Exped. 3824 (K) (23); Salubeni, A. J. 278 (K) (13), 416 (K) (17), 424 (K) (11), 1330 (GH, K, MO) (14), 1341 (K) (14), 1453 (K) (17), 1539 (K) (14); Sanane, M. 980 (K) (6b), 1571 (K) (13); Sanders, J. R. 54 (K) (10), 62 (K) (4); Sanderson, J. 428 (K) (6b); Saunders, D. H. 7 (K) (16), 60 (K) (16), 62 (K) (16); Sawer, E. R. 13698 (K) (16), 13699 (K) (16); Saxon, A. 254 (K) (6d), 294 (K) (17), 308 (K) (16), 309 (K) (11), 398A (K) (16), 588 (K) (16); Schenpera, J. C. 1048 (K, MO) (6b); Schimpler, M. W. 319 (K) (13); Schinz, H. 35 (K) (13), 42 (K) (16); Schlechter, R. 3814 (GH, K, MO) (6b), 3979 (GH, MO) (10), 4342 (K) (13), 4541 (GH) (4); Schlieben, H. J. 2108 (US) (14), 2281 (K, S) (27), 3874 (K) (11), 5791 (K) (13), 6021 (S) (27), 7662 (US) (10), 7668 (US) (6b); Scholes, J. S9 (K) (6e); Schuter, P. A. & A. Nuvunga 6644 (K) (6b); Schweinfurth, G. 179 (K) (11), 595 (K) (16), 1880 (US) (16), 1992 (S) (5), 2285 (K) (4); Schyff, H. 1897 (K) (11); Scott, H. 217 (K) (16); Semsei, S. R. 1720 (K) (27), A9/48 (K) (4); Seydel, R. 4491 (K, US) (13); Shannon, R. K. 88–39 (US) (5), 88–44 (US) (13); Shantz, H. L. 930 (K) (16); Shantz & Tanner 4196 (K) (13); Sherif, M. I. 3952 (K) (17); Shillito, E. M. 119 (K) (11); Shorland 179 (K) (16); Shuell 6785 (K) (19); Siame, W. 95 (K) (4), 112 (K) (11), 144 (K) (24), 144A (BM) (7), 164 (K) (13), 525 (MO) (6b); Sidey, J. L. 1287 (UPS) (6b), 1589 (MO) (6b); Sieber 2401 (GH, K) (4); Sington, F. C. 1298 (BM) (22), 2042 (BM) (22); Sillitoe, F. 457 (K) (17); Simfsuz, N. D. 2166 (K) (16); Simon, B. K. & G. Williamson 1403 (K) (6b); Simpson, N. D. 7302 (K) (17), 7406 (K) (16), 7606 (K) (16); Simson, H. E. 348 (K) (16); Skarpe, C. 52 (K) (13); Smith, L. C. 10/92 (K) (20); Smith, P. A. 26 (K) (13), 390 (K, MO) (6b), 830 (K) (13), 845 (K) (4), 887 (K) (13), 1369 (MO) (10), 1376 (K) (13), 1970 (K, MO) (6b), 2259 (K, MO) (10), 3114 (K) (13); Smith, P. A. et al. 5872 (K) (13); Smith, S. H. 22673 (K) (11); Smuts, J. C. & M. C. Gillett 3399 (K) (6b), 4127 (K) (6b); Snowden, J. D. 1160 (K) (16); Someren, V. 250 (K) (17), 7651 (K) (13); Sommerville, S. S. 726A (BM) (11); Sousa, A. F. 445 (K) (6b), 555 (K) (4); Soyaux, H. 189 (K) (4); Speke & Grant 572 (K) (16); Spinagi, C. A. 361 (K) (17); Stee, J. V. 3163 (K) (4); Stewart, D. R. M. 327 (K) (11); Stewart, M. M. 1/1911 (K) (6b); Steyagi, M. F. s.n. (US) (16); Stirton, C. H. 401 (K, MO) (6b), 559 (K) (6b), 1458 (K) (10), 1725 (K) (13); Stocks, J. 55 (K) (27); Stoltz, A. 213 (GH, US) (11), 540 (K) (17), 753 (K) (13), 1304 (BM, K, GH, MO) (14), 2494 (GH, MO) (10), 2500 (GH) (19); Streys, R. G. 6187 (K) (6b), 9297 (K) (6b), 9987 (K) (6b); Suabad, M. M. 6994 (MO) (23); Sutherland, D. 1856 (K) (10); Sutherland, J. 381 (MO) (13a); Swynnerton, C. F. 1413 (K) (13), 1451 (K) (27), 1951 (K) (27); Symes, Y. E. 112 (K) (11).
- T.C.D. Biol. Exped. 210 (BM) (13); Tadesse, F. M. & K. Vollesen 3073 (K) (20); Tadesse, M. T. & G. Y. Yagnew 1594 (K) (16), 1600 (K) (16); Talbot, P. A. 308 (BM) (11), 310 (K) (16), 312 (K) (7), 313 (BM) (6d), 324 (BM, K) (23), 1920 (BM) (24); Tanner, R. 89 (K) (27), 129 (K) (4), 747 (K) (16), 810 (K) (4), 1690 (K) (10), 1979 (K) (4), 2197 (K) (27), 2658 (K) (27), 2850 (K) (27), 2879 (K) (11), 2904 (K) (11), 2947 (K) (4), 2954 (K) (22), 3018 (K) (4), 3996 (K) (13), 4142 (K) (16), 4290 (K) (16), 4461 (K) (10), 5108 (K) (6a), 5274 (K) (16), 5655 (K) (6a), S110 (K) (6a); Tapscott, S. 2548 (K) (10); Taylor, W. W. 1885 (BM) (11); Taylor, T. W. 1 (K) (17); Teague, A. J. 33 (K) (13); Teixerira, B. 303 (BM) (6e); Terry, P. J. 1879 (K) (5), 1942 (K) (5), 3104 (K) (5), 3135 (K) (5), 3173 (K) (5), 3384 (K, ODU) (20), s.n. (K) (16); Thomas 4429 (K) (6e); Thomas, A. S. 183 (K) (17), 1031 (K) (16), 1532 (K) (16), 2015 (K) (11), 2015 (K) (11), 2893 (K) (11), 3458 (K) (13), 3554 (K) (13), D138 (K) (6f); Thomas, D. 2452 (MO) (6d); Thomas, N. W. 155 (K) (6f), 603 (K) (6f), 3164 (K) (23), 3208 (K) (23), 5829 (K) (23), 6665 (K) (23); Thompson 10 (K) (23); Thomson, B. s.n. (K) (13); Thorneiop, G. 2206 (K) (11); Thorold, C. A. 209 (K) (6e), 251 (BM) (24); Thulin, M. & B. A. Mahamed 7088 (K) (13); Thulin, M. & B. Mhoro 488 (K) (4), 3140 (K) (4); Timberlake, J. R. 459 (K) (13); Tolken, H. & D. S. Hardy 919 (K) (13); Toms, K. E. 5 (K) (4); Toul Botanical Exped. 1464 (K) (23), 1699 (K) (23); Towa, E. 8166 (K) (19); Townsend, C. C. 170 (K) (17), 2162 (K) (13); Troprell, C. G. 2050 (K) (4); Troupin, G. 621 (K) (11); Tutin, C. E. 162 (K) (6a); Tweedie 638 (K) (4), 2448 (K) (13), 3314 (K) (13), 3360 (K) (13), s.n. (K) (4); Twuner, A. 6730 (K) (13); Tyson, W. 894 (GH) (6b), 1222 (GH) (4), 1368 (MO) (10), 1504 (GH) (4).
- Vallance, G. B. 699 (K) (4), 700 (K) (4); Vallah 65 (K) (1); Vaughan, J. H. 402 (K) (11), 1613 (K) (11), 1614 (K) (17), 1664 (K) (11), 2328 (BM) (27); Verboom, W. C. 1039 (K) (13); Verdcourt, B. 581 (K) (13), 3330 (K) (6e), 3390 (K) (17), 3690 (K) (20); Vigne, C. 4359 (US) (6e); Visser, J. H. 248 (K) (10), 315 (K) (6b), 316 (K) (6b), 2528 (K) (13); Vollesen, K. 3311 (K, UPS) (24); Vrgine, G. 3355 (K) (6e); Vuuren, D. A. & W. Giess 1132 (K) (13).
- Waitland 838 (K) (6e); Wallich, Z. 1278 (K) (10); Wallace, G. B. 713 (K) (27); Walter, E. G. 10 (K) (13); Wannop, H. E. 515 (K) (13), 760 (K) (13), 814 (K) (13); Welch, J. R. 174 (K) (16); Wellby, M. S. 1901 (K) (16); Wellman, J. C. 1769 (K) (4); Welwitsch, A. P. 779 (K) (13), 5851 (K) (10); Werdermann & Oberdieck 2236a (US) (4); Wham, J. R. D. 26/2/24 (K) (11); Whellar, J. A. 1188 (K) (24), 1214 (K) (11); White, F. 2048 (K) (13), 6213 (K) (10), 7574 (K) (11); White, M. 142 (K) (13); Whiteside, J. O. 107793 (K) (11), 130476 (K) (11); Whyte, A. 1898 (K) (16), 1898 (K) (17), 1902 (K) (13), 1912 (K) (13); Wickens, G. E. 309 (K) (16), 536 (K) (13), 568 (K) (16), 587 (K) (13), 628 (K) (16), 1562 (K) (23), 2063 (K) (24), 2099 (K) (17), 2241 (K) (17), 2306 (K) (24), 2332 (K)

- (24), 2345 (K) (24), 2492 (K) (16), 2807 (K) (23), 3159 (K) (13), Wiehe, P. O. 28294 (K) (4), N/118 (K) (11), N/426 (K) (4); Wild, H. 599 (K) (11), 1631 (K) (13), 4315 (K) (10), 5718 (K) (17); William, M. 5975 (K) (13); Willemse, R. H. 30 (K) (16); Williams, G. R. 220 (K) (11); Williams, J. G. 5115 (K) (13), 12546 (K) (13), B5113 (K) (4); Williams, R. 545 (K) (16); Williams, R. H. 32 (K) (4); Williams, T. L. 534 (K) (6e); Williamson, D. J. 90 (K) (4); Wilson 349 (K) (16); Wilson, J. 373 (K) (13), 880 (K) (11); Winter, B. D. 2540 (K) (6b), 3372 (K) (13), 3995 (K) (6b), 4428 (K) (6b); Winter, B. D. & O. A. Leistner 5368 (K) (13); Winter, B. D. & W. Marais 4478 (K) (6b), 4479 (K) (13), 4567 (K) (10), 4731 (K) (13), 4913 (K) (6b); Winter, B. D. & H. J. Wise 4452 (K) (13); Wit et al. 166671 (K) (11); Wittle, D. 314 (K) (16); Wood, G. H. S. 726 (K) (6e), 805 (K) (13); Wood, J. M. 4/1908 (US) (4), 16 (K) (4), 20 (US) (4), 113 (K) (6b), 394 (K) (10), 440 (K) (11), 911 (US) (6b), 4276 (K) (13), 4934 (MO) (6b), 7252 (US) (6b), 7500 (US) (11), 9212 (GH) (6b), 10495 (MO) (6b), 10873 (US) (4), 10887 (US) (6b), 92715 (MO) (10); Woolhouse, H. W. 51/61/22 (K) (17); Worsdell, W. O. 12/1909 (K) (6b); Wright, J. M. 302 (K) (13); Wyld, J. W. 225 (K) (4), 725 (BM) (6a); Wylie, F. 1895 (K) (6b).  
Yeafes, M. 88 (K) (16); Yong, J. D. 184 (K) (11); Young, J. 185 (K) (6f); Young, R. G. 1261 (ODU) (2), 1300 (MO) (1), 1347 (BM) (2), 1365 (MO) (2).  
Zimmer, F. 102 (K) (27); Zunguze, D. & X. Boan 49 (K) (11).