

Old Dominion University ODU Digital Commons

Physics Faculty Publications

Physics

7-2017

Higher-Twist Corrections to Gluon TMD Factorization

I. Balitsky

Old Dominion University, ibalitsk@odu.edu

A. Tarasov

Follow this and additional works at: https://digitalcommons.odu.edu/physics_fac_pubs

 Part of the [Elementary Particles and Fields and String Theory Commons](#)

Repository Citation

Balitsky, I. and Tarasov, A., "Higher-Twist Corrections to Gluon TMD Factorization" (2017). *Physics Faculty Publications*. 67.
https://digitalcommons.odu.edu/physics_fac_pubs/67

Original Publication Citation

Balitsky, I., & Tarasov, A. (2017). Higher-twist corrections to gluon TMD factorization. *Journal of High Energy Physics*, 2017(7), 95.
doi:10.1007/jhep07(2017)095

This Article is brought to you for free and open access by the Physics at ODU Digital Commons. It has been accepted for inclusion in Physics Faculty Publications by an authorized administrator of ODU Digital Commons. For more information, please contact digitalcommons@odu.edu.

Higher-twist corrections to gluon TMD factorization

I. Balitsky^{a,b} and A. Tarasov^c

^a*Physics Department, Old Dominion University,
4600 Elkhorn Ave, Norfolk, VA 23529, U.S.A.*

^b*Thomas Jefferson National Accelerator Facility,
12000 Jefferson Ave, Newport News, VA 23606, U.S.A.*

^c*Physics Department, Brookhaven National Laboratory,
Bldg. 510A, Upton, NY 11973, U.S.A.*

E-mail: balitsky@jlab.org, atarasov@bnl.gov

ABSTRACT: We calculate power corrections to TMD factorization for particle production by gluon-gluon fusion in hadron-hadron collisions.

KEYWORDS: QCD Phenomenology

ARXIV EPRINT: [1706.01415](https://arxiv.org/abs/1706.01415)

Contents

1	Introduction	1
2	TMD factorization from functional integral	2
3	Power corrections and solution of classical YM equations	7
3.1	Power counting for background fields	7
3.2	Approximate solution of classical equations	9
3.3	Twist expansion of classical gluon fields	14
4	Leading higher-twist correction at $s \gg Q^2 \gg Q_\perp^2 \gg m^2$	16
5	Small-x limit and scattering of shock waves	19
6	Conclusions and outlook	21
A	Diagrams with retarded propagators	22
B	Solution of Yang-Mills equations in two dimensions	25

1 Introduction

Particle production in hadron-hadron scattering with transverse momentum of produced particle much smaller than its invariant mass is described in the framework of TMD factorization [1–5]. The typical example is the Higgs production at LHC through gluon-gluon fusion. Factorization formula for particle production in hadron-hadron scattering looks like [1, 6]

$$\frac{d\sigma}{d\eta d^2q_\perp} = \sum_f \int d^2b_\perp e^{i(q,b)_\perp} \mathcal{D}_{f/A}(x_A, b_\perp, \eta) \mathcal{D}_{f/B}(x_B, b_\perp, \eta) \sigma(ff \rightarrow H) + \text{power corrections} + \text{Y - terms} \quad (1.1)$$

where η is the rapidity, $\mathcal{D}_{f/A}(x, z_\perp, \eta)$ is the TMD density of a parton f in hadron A , and $\sigma(ff \rightarrow H)$ is the cross section of production of particle H of invariant mass $m_H^2 = Q^2$ in the scattering of two partons. (For simplicity, we consider the scattering of unpolarized hadrons.)

In this paper we calculate the first power corrections $\sim \frac{q_\perp^2}{Q^2}$ in a sense that we represent them as a TMD-like matrix elements of higher-twist operators. It should be noted that our method works for arbitrary relation between s and Q^2 and between q_\perp^2 and hadron mass m^2 (provided that pQCD is applicable), but in this paper we only present the result for the physically interesting region $s \gg Q^2 \gg q_\perp^2 \gg m^2$.




Figure 1. Particle production by gluon-gluon fusion.

To obtain formula (1.1) with first corrections we use factorization in rapidity [7]. We denote quarks and gluons with rapidity close to the rapidity of the projectile and target protons as A -fields and B -fields, respectively. We call the remaining fields in the central region of rapidity by the name C -fields and integrate over them in the corresponding functional integral. At this step, we get the effective action depending on A and B fields. The subsequent integration over A fields gives matrix elements of some TMD-like operators switched between projectile proton states and integration over B fields will give matrix elements between target states.¹

The paper is organized as follows. In section 2 we derive the TMD factorization from the double functional integral for the cross section of particle production. In section 3, which is central to our approach, we explain the method of calculation of higher-twist power corrections based on a solution of classical Yang-Mills equations. In section 4 we find the leading higher-twist correction to particle production in the region $s \gg Q^2 \gg q_\perp^2$. Finally, in section 5 we compare our calculations in the small- x limit to the classical field resulting from the scattering of two shock waves. The appendices contain proofs of some necessary technical statements.

2 TMD factorization from functional integral

We consider production of an (imaginary) scalar particle Φ in proton-proton scattering. This particle is connected to gluons by the vertex

$$\mathcal{L}_\Phi = g_\Phi \int d^4x \Phi(x) g^2 F^2(x), \quad F^2(x) \equiv F_{\mu\nu}^a(x) F^{a\mu\nu}(x) \quad (2.1)$$

¹It should be noted that due to the kinematics $Q^2 \gg Q_\perp^2, m^2$ we will not need the explicit form of the high-energy effective action which is much sought after in the small- x physics but not known up to now except a couple of first perturbative terms [7–11].

This is a $\frac{m_H}{m_t} \ll 1$ approximation [12, 13] for Higgs production via gluon fusion at LHC with

$$g_H = \frac{1}{48\pi^2 v} \left(1 + \frac{11}{4\pi} \alpha_s + \dots \right)$$

where $\alpha_s = \frac{g^2}{4\pi}$ as usual.² The differential cross section of Φ production has the form

$$d\sigma = \frac{d^3 q}{2E_q (2\pi)^3} \frac{g_\Phi^2}{2s} W(p_A, p_B, q) \quad (2.2)$$

where we defined the “hadronic tensor” $W(p_A, p_B, q)$ as

$$\begin{aligned} W(p_A, p_B, q) &\stackrel{\text{def}}{=} \sum_X \int d^4 x e^{-iqx} \langle p_A, p_B | g^2 F^2(x) | X \rangle \langle X | g^2 F^2(0) | p_A, p_B \rangle \\ &= \int d^4 x e^{-iqx} \langle p_A, p_B | g^4 F^2(x) F^2(0) | p_A, p_B \rangle \end{aligned} \quad (2.3)$$

As usual, \sum_X denotes the sum over full set of “out” states. It can be represented by double functional integral

$$\begin{aligned} W(p_A, p_B, q) &= \sum_X \int d^4 x e^{-iqx} \langle p_A, p_B | g^2 F^2(x) | X \rangle \langle X | g^2 F^2(0) | p_A, p_B \rangle \\ &= \lim_{t_i \rightarrow -\infty}^{t_f \rightarrow \infty} g^4 \int d^4 x e^{-iqx} \int^{\vec{A}(t_f)=A(t_f)} D\tilde{A}_\mu D A_\mu \int^{\vec{\psi}(t_f)=\psi(t_f)} D\tilde{\psi} D\tilde{\bar{\psi}} D\bar{\psi} D\psi \Psi_{p_A}^* (\vec{A}(t_i), \vec{\psi}(t_i)) \\ &\quad \times \Psi_{p_B}^* (\vec{A}(t_i), \vec{\psi}(t_i)) e^{-iS_{\text{QCD}}(\tilde{A}, \tilde{\psi})} e^{iS_{\text{QCD}}(A, \psi)} \tilde{F}^2(x) F^2(0) \Psi_{p_A} (\vec{A}(t_i), \psi(t_i)) \Psi_{p_B} (\vec{A}(t_i), \psi(t_i)) \end{aligned} \quad (2.4)$$

Here the fields A, ψ correspond to the amplitude $\langle X | F^2(0) | p_A, p_B \rangle$, fields $\tilde{A}, \tilde{\psi}$ correspond to complex conjugate amplitude $\langle p_A, p_B | F^2(x) | X \rangle$ and $\Psi_p(\vec{A}(t_i), \psi(t_i))$ denote the proton wave function at the initial time t_i . The boundary conditions $\vec{A}(t_f) = A(t_f)$ and $\vec{\psi}(t_f) = \psi(t_f)$ reflect the sum over all states X , cf. refs. [15–17].

We use Sudakov variables $p = \alpha p_1 + \beta p_2 + p_\perp$ and the notations $x_\bullet \equiv x_\mu p_1^\mu$ and $x_* \equiv x_\mu p_2^\mu$ for the dimensionless light-cone coordinates ($x_* = \sqrt{\frac{s}{2}} x_+$ and $x_\bullet = \sqrt{\frac{s}{2}} x_-$). Our metric is $g^{\mu\nu} = (1, -1, -1, -1)$ so that $p \cdot q = (\alpha_p \beta_q + \alpha_q \beta_p) \frac{s}{2} - (p, q)_\perp$ where $(p, q)_\perp \equiv -p_i q^i$. Throughout the paper, the sum over the Latin indices $i, j \dots$ runs over the two transverse components while the sum over Greek indices runs over the four components as usual.

To derive the factorization formula, we separate the (quark and gluon) fields in the functional integral (2.4) into three sectors: “projectile” fields A_μ, ψ_a with $|\beta| < \sigma_a$, “target”

²For finite m_t the constant g_H should be multiplied by $\frac{3\tau}{2} [1 + (1 - \tau) \arcsin^2 \frac{1}{\sqrt{\tau}}]$ with $\tau = \frac{4m_t^2}{m_H^2}$ [14].




Figure 2. Rapidity factorization for particle production.

fields with $|\alpha| < \sigma_b$ and “central rapidity” fields C_μ, ψ with $|\alpha| > \sigma_b$ and $|\beta| > \sigma_a$.³

$$\begin{aligned}
 W(p_A, p_B, q) &= g^4 \int d^4x e^{-iqx} \int^{\vec{A}(t_f)=A(t_f)} D\vec{A}_\mu D A_\mu \int^{\vec{\psi}_a(t_f)=\psi_a(t_f)} D\bar{\psi}_a D\psi_a D\tilde{\psi}_a D\tilde{\psi}_a \\
 &\times e^{-iS_{\text{QCD}}(\vec{A}, \vec{\psi}_a)} e^{iS_{\text{QCD}}(A, \psi_a)} \Psi_{p_A}^*(\vec{A}(t_i), \vec{\psi}_a(t_i)) \Psi_{p_A}(\vec{A}(t_i), \psi(t_i)) \\
 &\times \int^{\vec{B}(t_f)=B(t_f)} D\vec{B}_\mu D B_\mu \int^{\vec{\psi}_b(t_f)=\psi_b(t_f)} D\bar{\psi}_b D\psi_b D\tilde{\psi}_b D\tilde{\psi}_b \\
 &\times e^{-iS_{\text{QCD}}(\vec{B}, \vec{\psi}_b)} e^{iS_{\text{QCD}}(B, \psi_b)} \Psi_{p_B}^*(\vec{B}(t_i), \vec{\psi}_b(t_i)) \Psi_{p_B}(\vec{B}(t_i), \psi_b(t_i)) \\
 &\times \int DC_\mu \int^{\vec{C}(t_f)=C(t_f)} D\vec{C}_\mu \int D\bar{\psi}_C D\psi_C \int^{\vec{\psi}_c(t_f)=\psi_c(t_f)} D\tilde{\psi}_C D\tilde{\psi}_C \tilde{F}_C^2(x) F_C^2(0) e^{-i\tilde{S}_C + iS_C}
 \end{aligned} \tag{2.5}$$

where $S_C = S_{\text{QCD}}(A + B + C) - S_{\text{QCD}}(A) - S_{\text{QCD}}(B)$.

Our goal is to integrate over central fields and get the amplitude in the factorized form, as a (sum of) products of functional integrals over A fields representing projectile matrix elements (TMDs) and functional integrals over B fields representing target matrix elements. In the spirit of background-field method, we “freeze” projectile and target fields (and denote them the \vec{A} , $\vec{\xi}_a$, ξ_a and \vec{B} , $\vec{\xi}_b$, ξ_b respectively) and get a sum of diagrams in these external fields. Since $|\beta| < \sigma_a$ in the projectile fields and $|\alpha| < \sigma_b$ in the target fields, at the tree-level one can set with power accuracy $\beta = 0$ for the projectile fields and $\alpha = 0$ for the target fields - the corrections will be $O(\frac{m^2}{\sigma_a s})$ and $O(\frac{m^2}{\sigma_b s})$. Beyond the tree level, one should expect that the integration over C fields will produce the logarithms of the cutoffs σ_a and σ_b which will cancel with the corresponding logs in gluon TMDs of the projectile and the target.

³The standard factorization scheme for particle production in hadron-hadron scattering is splitting the diagrams in collinear to projectile part, collinear to target part, hard factor, and soft factor [1]. Here we factorize only in rapidity. For our purpose of calculation of power corrections in the tree approximation it is sufficient; however, we hope to treat possible logs of transverse scales in loop corrections in the same way as it was done in our rapidity evolution equations for gluon TMDs in refs. [18, 19].

As usual, diagrams disconnected from the vertices $F^2(x)$ and $F^2(0)$ (“vacuum bubbles” in external fields) exponentiate so the result has the schematic form

$$\int DC_\mu \int^{\tilde{C}(t_f)=C(t_f)} D\tilde{C}_\mu \int D\bar{\psi}_C D\psi_C \int^{\tilde{\psi}_c(t_f)=\psi_c(t_f)} D\tilde{\bar{\psi}}_C D\tilde{\psi}_C g^4 \tilde{F}_C^2(x) F_C^2(0) e^{-i\tilde{S}_C+iS_C} = e^{S_{\text{eff}}(U,V,\tilde{U},\tilde{V})} \mathcal{O}(q,x;A,\tilde{A},\psi_a,\tilde{\psi}_a;B,\tilde{B},\psi_b,\tilde{\psi}_b) \quad (2.6)$$

where $\mathcal{O}^{\mu\nu}(q,x;A,\psi_A;B,\psi_B)$ is a sum of diagrams connected to $\tilde{F}^2(x)F^2(0)$. Since rapidities of central fields and A, B fields are very different, one should expect the result of integration over C -fields to be represented in terms of Wilson-line operators constructed from A and B fields.

The effective action has the form

$$S_{\text{eff}}(U,V,\tilde{U},\tilde{V}) = 2\text{Tr} \int d^2x_\perp [-i\tilde{U}_i\tilde{V}^i + iU_iV^i + (\tilde{\mathcal{L}}_i(\tilde{U},\tilde{V})\tilde{\mathcal{L}}^i(\tilde{U},\tilde{V}) - 2\tilde{\mathcal{L}}_i(\tilde{U},\tilde{V})\mathcal{L}^i(U,V) + \mathcal{L}_i(U,V)\mathcal{L}^i(U,V)) \ln \sigma_a\sigma_b s + O(\ln \sigma_a\sigma_b s)^2] \quad (2.7)$$

where Wilson lines U are made from projectile fields

$$U(x_\perp) = [\infty p_2 + x_\perp, -\infty p_2 + x_\perp]^{A*}, \quad U_i = U^\dagger i\partial_i U$$

and Wilson lines V from target fields

$$V(x_\perp) = [\infty p_1 + x_\perp, -\infty p_1 + x_\perp]^{B\bullet}, \quad V_i = V^\dagger i\partial_i V$$

and similarly for \tilde{U} and \tilde{V} in the left sector. The explicit form of “Lipatov vertices” $L_i(U,V)$ is presented in [20]. Unfortunately, the effective action beyond the first two terms in (2.7) is unknown, but we will demonstrate below that for our purposes we do not need the explicit form of the effective action.

After integration over C fields the amplitude (2.4) can be rewritten as

$$\begin{aligned} W(p_A,p_B,q) &= \int d^4x e^{-iqx} \int^{\tilde{A}(t_f)=A(t_f)} D\tilde{A}_\mu D A_\mu \int^{\tilde{\psi}_a(t_f)=\psi_a(t_f)} D\tilde{\bar{\psi}}_a D\psi_a D\tilde{\bar{\psi}}_a D\tilde{\psi}_a \\ &\times e^{-iS_{\text{QCD}}(\tilde{A},\tilde{\psi}_a)} e^{iS_{\text{QCD}}(A,\psi_a)} \Psi_{p_A}^*(\vec{\tilde{A}}(t_i),\tilde{\psi}_a(t_i)) \Psi_{p_A}(\vec{A}(t_i),\psi(t_i)) \\ &\times \int^{\tilde{B}(t_f)=B(t_f)} D\tilde{B}_\mu D B_\mu \int^{\tilde{\psi}_b(t_f)=\psi_b(t_f)} D\tilde{\bar{\psi}}_b D\psi_b D\tilde{\bar{\psi}}_b D\tilde{\psi}_b \\ &\times e^{-iS_{\text{QCD}}(\tilde{B},\tilde{\psi}_b)} e^{iS_{\text{QCD}}(B,\psi_b)} \Psi_{p_B}^*(\vec{\tilde{B}}(t_i),\tilde{\psi}_b(t_i)) \Psi_{p_B}(\vec{B}(t_i),\psi_b(t_i)) \\ &\times e^{S_{\text{eff}}(U,V,\tilde{U},\tilde{V})} \mathcal{O}(q,x;A,\psi_a,\tilde{A},\tilde{\psi}_a;B,\psi_b,\tilde{B},\tilde{\psi}_b) \end{aligned} \quad (2.8)$$

Note that due to boundary conditions at t_f in the above integral, the functional integral over C fields in eq. (2.6) should be done in the background of the A and B fields satisfying

$$\tilde{A}(t_f) = A(t_f), \quad \tilde{\psi}_a(t_f) = \psi_a(t_f) \quad \text{and} \quad \tilde{B}(t_f) = B(t_f), \quad \tilde{\psi}_b(t_f) = \psi_b(t_f) \quad (2.9)$$

Our approximation at the tree level is that $\beta = 0$ for A, \tilde{A} fields and $\alpha = 0$ for B, \tilde{B} fields which corresponds to $A = A(x_\bullet, x_\perp)$, $\tilde{A} = \tilde{A}(x_\bullet, x_\perp)$ and $B = B(x_*, x_\perp)$, $\tilde{B} = \tilde{B}(x_*, x_\perp)$.

Now comes the important point: because of boundary conditions (2.9), for the purpose of calculating the integral (2.6) over central fields one can set

$$\begin{aligned}
 A(x_\bullet, x_\perp) &= \tilde{A}(x_\bullet, x_\perp), & \psi_a(x_\bullet, x_\perp) &= \tilde{\psi}_a(x_\bullet, x_\perp) \\
 \text{and} \\
 B(x_*, x_\perp) &= \tilde{B}(x_*, x_\perp), & \psi_b(x_*, x_\perp) &= \tilde{\psi}_b(x_*, x_\perp)
 \end{aligned}
 \tag{2.10}$$

Indeed, because A, ψ and $\tilde{A}, \tilde{\psi}$ do not depend on x_* , if they coincide at $x_* = \infty$ they should coincide everywhere. Similarly, if B, ψ_b and $\tilde{B}, \tilde{\psi}_b$ do not depend on x_\bullet , if they coincide at $x_\bullet = \infty$ they should be equal.

It should be emphasized that the boundary conditions (2.9) mean the summation over all intermediate states in corresponding projectile and target matrix elements in the functional integrals over projectile and target fields. Without the sum over all intermediate states the conditions (2.10) are no longer true. For example, if we would like to measure another particle or jet in the fragmentation region of the projectile, the second condition in eq. (2.10) breaks down.

Next important observation is that due to eqs. (2.10) the effective action (2.7) vanishes for background fields satisfying conditions (2.9). For the first two terms displayed in (2.7) it is evident, but it is easy to see that the effective action in the background fields satisfying (2.10) should vanish due to unitarity. Indeed, let us consider the functional integral (2.4) without sources $\tilde{F}^2(x)F^2(0)$. It describes the matrix element (2.11) without Φ production, that is

$$\sum_X \langle p_A, p_B | X \rangle \langle X | p_A, p_B \rangle = 1
 \tag{2.11}$$

(modulo appropriate normalization of $|p_A\rangle$ and $|p_B\rangle$ states). If we perform the same decomposition into A, B , and C fields as in eq. (2.4) we will see integral (2.8) without $\mathcal{O}^{\mu\nu}(q, x, y; A, \psi_a, \tilde{A}, \tilde{\psi}_a; B, \psi_b, \tilde{B}, \tilde{\psi}_b)$ which can be represented as

$$\langle p_A, p_B | e^{S_{\text{eff}}(U, V, \tilde{U}, \tilde{V})} | p_A, p_B \rangle = 1
 \tag{2.12}$$

which means that the effective action should vanish for the Wilson-line operators constructed from the fields satisfying eqs. (2.10). Summarizing, we see that at the tree level in our approximation

$$\begin{aligned}
 & \int DC_\mu \int^{\tilde{C}(t_f)=C(t_f)} D\tilde{C}_\mu \int D\tilde{\psi}_C D\psi_C \int^{\tilde{\psi}_c(t_f)=\psi_c(t_f)} D\tilde{\psi}_C D\psi_C g^4 \tilde{F}_C^2(x) F_C^2(0) e^{-i\tilde{S}_C + iS_C} \\
 &= \mathcal{O}(q, x; A, \psi_a; B, \psi_b)
 \end{aligned}
 \tag{2.13}$$

where now $S_C = S_{\text{QCD}}(C + A + B) - S_{\text{QCD}}(A) - S_{\text{QCD}}(B)$ and $\tilde{S}_C = S_{\text{QCD}}(\tilde{C} + A + B) - S_{\text{QCD}}(A) - S_{\text{QCD}}(B)$. It is known that in the tree approximation the double functional integral (2.13) is given by a set of retarded Green functions in the background fields [21–23] (see also appendix A for the proof). Since the double functional integral (2.13) is given by a set of retarded Green functions (in the background field $A + B$), the calculation of tree-level contributions to, say, $F^2(x)$ in the r.h.s. of eq. (2.13) is equivalent to solving YM

equation for $A_\mu(x)$ (and $\psi(x)$) with boundary conditions that the solution has the same asymptotics at $t \rightarrow -\infty$ as the superposition of incoming projectile and target background fields.

The hadronic tensor (2.8) can now be represented as

$$W(p_A, p_B, q) = \int d^4x e^{-iqx} \langle p_A | \langle p_B | \hat{\mathcal{O}}(q, x; \hat{A}, \hat{\psi}_a; \hat{B}, \hat{\psi}_b) | p_A \rangle | p_B \rangle \quad (2.14)$$

where $\hat{\mathcal{O}}(q, x; \hat{A}, \hat{\psi}_a; \hat{B}, \hat{\psi}_b)$ should be expanded in a series in $\hat{A}, \hat{\psi}_a; \hat{B}, \hat{\psi}_b$ operators and evaluated between the corresponding (projectile or target) states: if

$$\hat{\mathcal{O}}(q, x; \hat{A}, \hat{\psi}_a; \hat{B}, \hat{\psi}_b) = \sum_{m,n} \int dz_m dz'_n c_{m,n}^{\mu\nu}(q, x) \hat{\Phi}_A(z_m) \hat{\Phi}_B(z'_n) \quad (2.15)$$

(where $c_{m,n}^{\mu\nu}$ are coefficients and Φ can be any of A_μ, ψ or $\bar{\psi}$) then⁴

$$W = \int d^4x e^{-iqx} \sum_{m,n} \int dz_m c_{m,n}^{\mu\nu}(q, x) \langle p_A | \hat{\Phi}_A(z_m) | p_A \rangle \int dz'_n \langle p_B | \hat{\Phi}_B(z'_n) | p_B \rangle \quad (2.16)$$

As we will demonstrate below, the relevant operators are quark and gluon fields with Wilson-line type gauge links collinear to either p_2 for A fields or p_1 for B fields.

3 Power corrections and solution of classical YM equations

3.1 Power counting for background fields

As we discussed in previous section, to get the hadronic tensor in the form (2.14) we need to calculate the functional integral (2.13) in the background of the fields (2.10). To understand the relative strength of Lorentz components of these fields, let us compare the typical term in the leading contribution to W

$$\frac{64/s^2}{N_c^2 - 1} \int d^4x e^{-iqx} \langle p_A | \hat{U}_*^{mi}(x_\bullet, x_\perp) \hat{U}_*^{mj}(0) | p_A \rangle \langle p_B | \hat{V}_{\bullet i}^n(x_*, x_\perp) \hat{V}_{\bullet j}^n(0) | p_B \rangle \quad (3.1)$$

where

$$\hat{U}_{*i}^a(z_\bullet, z_\perp) \equiv [-\infty_\bullet, z_\bullet]_z^{ab} g \hat{F}_{*i}^b(z_\bullet, z_\perp), \quad \hat{V}_{\bullet i}^a(z_*, z_\perp) \equiv [-\infty_*, z_*]_z^{ab} g \hat{F}_{\bullet i}^b(z_*, z_\perp) \quad (3.2)$$

and some typical higher-twist terms. As we mentioned, we consider $W(p_A, p_B, q)$ in the region where $s, Q^2 \gg Q_\perp^2, m^2$ while the relation between Q_\perp^2 and m^2 and between Q^2 and s may be arbitrary. So, for the purpose of counting of powers of s , we will not distinguish

⁴Our logic here is the following: to get the expression for $\hat{\mathcal{O}}$ in eq. (2.13) we calculate \mathcal{O} in the background of two external fields $\Phi_A = (A_\mu, \psi_a)$ and $\Phi_B = (B_\mu, \psi_b)$ and then promote them to operators $\hat{\Phi}_A$ and $\hat{\Phi}_B$ in the obtained expressions for \mathcal{O} . However, there is a subtle point in the promotion of background fields to operators. When we are calculating \mathcal{O} as the r.h.s. of eq. (2.13) the fields Φ_A and Φ_B are c -numbers; on the other hand, after functional integration in eq. (2.4) they become operators which must be time-ordered in the right sector and anti-time-ordered in the left sector. Fortunately, as we shall see below, all these operators are separated either by space-like distances or light-cone distances so all of them (anti) commute and thus can be treated as c -numbers.

between s and Q^2 (although at the final step we will be able to tell the difference since our final expressions for higher-twist corrections will have either s or Q^2 in denominators). Similarly, for the purpose of power counting we will not distinguish between m and Q_\perp and will introduce m_\perp which may be of order of m or Q_\perp depending on matrix element.

The estimate of the leading-twist matrix element between projectile states is

$$\langle p_A | \hat{U}_{*i}^a(x_\bullet, x_\perp) \hat{U}_{*j}^a(0) | p_A \rangle = p_2^\mu p_2^\nu \langle p_A | \hat{U}_{\mu i}^a(x_\bullet, x_\perp) \hat{U}_{\nu j}^a(0) | p_A \rangle \sim s^2 (m_\perp^2 g_{ij}^\perp + m_\perp^4 x_i^\perp x_j^\perp) \quad (3.3)$$

(here we assume normalization $\langle p_A | p_A \rangle = 1$ for simplicity).

The typical higher-twist correction is proportional to (see e.g. eq. (4.4))

$$\begin{aligned} & d^{abc} \langle p_A | \hat{U}_{*i}^a(x_\bullet, x_\perp) \hat{U}_{*k}^b(x'_\bullet, x_\perp) \hat{U}_{*j}^c(0) | p_A \rangle \\ &= d^{abc} p_2^\mu p_2^\nu p_2^\lambda \langle p_A | \hat{U}_{\mu i}^a(x_\bullet, x_\perp) \hat{U}_{\nu k}^b(x'_\bullet, x_\perp) \hat{U}_{\lambda j}^c(0) | p_A \rangle \\ &\sim s^3 m_\perp^4 (g_{ij}^\perp x_k + g_{ik}^\perp x_j + g_{jk}^\perp x_i) + s^3 m_\perp^6 x_i x_j x_k \end{aligned} \quad (3.4)$$

Since $x_i^\perp \sim \frac{q_i^\perp}{q_\perp^2} \sim \frac{1}{m_\perp}$ we see that an extra $\hat{F}_{\mu i}$ in the matrix element between projectile states brings $p_{1\mu} m_\perp$ which means that $\hat{U}_{*i} \sim s m_\perp$.

Next, some of the higher-twist matrix elements have an extra U_{kl} like

$$d^{abc} \langle p_A | \hat{U}_{*i}^{ai}(x_\bullet, x_\perp) \hat{U}_{kl}^b(x'_\bullet, x_\perp) \hat{U}_{*j}^{cj}(0) | p_A \rangle \quad (3.5)$$

where

$$\hat{U}_{kl}(x_\bullet, x_\perp) \equiv [-\infty_\bullet, x_\bullet]_x g \hat{F}_{kl}(x_\bullet, x_\perp) [x_\bullet, -\infty_\bullet]_x \quad (3.6)$$

Since we consider only unpolarized projectile and target hadrons

$$\begin{aligned} & d^{abc} \langle p_A | \hat{U}_{*i}^{ai}(x_\bullet, x_\perp) \hat{U}_{kl}^b(x'_\bullet, x_\perp) \hat{U}_{*j}^{cj}(0) | p_A \rangle \\ &\sim s^2 (m_\perp^4 g_{ik}^\perp g_{jl}^\perp + m_\perp^6 g_{ik}^\perp x_j x_l + m_\perp^6 g_{jl}^\perp x_i x_k - k \leftrightarrow l) \end{aligned} \quad (3.7)$$

and, comparing this to eq. (3.3), we see that an extra \hat{F}_{kl} can bring an extra m_\perp^2 . Combining this with an estimate $U_{*i} \sim s m_\perp$ we see that the typical field \bar{A}_* is of order s while $\bar{A}_i \sim m_\perp$. Similarly, for the target fields we get $\bar{B}_\bullet \sim s$, $\bar{B}_i \sim m_\perp$.

Some of the power corrections involve matrix elements like

$$d^{abc} \langle p_A | \hat{U}_{*i}^{ai}(x_\bullet, x_\perp) \hat{U}_{*\bullet}^b(x'_\bullet, x_\perp) \hat{U}_{*j}^{cj}(0) | p_A \rangle \quad (3.8)$$

where

$$\hat{U}_{*\bullet}(x_\bullet, x_\perp) \equiv [-\infty_\bullet, x_\bullet]_x g \hat{F}_{*\bullet}(x_\bullet, x_\perp) [x_\bullet, -\infty_\bullet]_x \quad (3.9)$$

An extra field strength operator $\hat{F}^{\mu\nu}$ between the projectile states can bring $\frac{p_A^\mu p_2^\nu}{p_A \cdot p_2} - \mu \leftrightarrow \nu$ so that $\hat{F}_{*\bullet} \sim s m^2$.⁵ Since $\bar{A}_* \sim s$ we see that $\bar{A}_\bullet \sim m_\perp^2$. Similarly, for the target we get $\bar{B}_* \sim m_\perp^2$.

⁵The denominator $p_A \cdot p_2$ is due to the fact that p_2 enters only through the direction of Wilson line and therefore the matrix element should not change under rescaling $p_2 \rightarrow \lambda p_2$.




Figure 3. Typical diagram for the classical field with projectile/target sources. The Green functions of the central fields are given by retarded propagators.

Summarizing, the relative strength of the background gluon fields in projectile and target is

$$\begin{aligned}
 \bar{A}_*(x_\bullet, x_\perp) &\sim s, & \bar{A}_\bullet(x_\bullet, x_\perp) &\sim m_\perp^2, & \bar{A}_i(x_\bullet, x_\perp) &\sim m_\perp \\
 \bar{B}_*(x_*, x_\perp) &\sim m_\perp^2, & \bar{B}_\bullet(x_*, x_\perp) &\sim s, & \bar{B}_i(x_*, x_\perp) &\sim m_\perp
 \end{aligned} \quad (3.10)$$

To finish power counting, we need also the relative strength of quark background fields ψ_a and ψ_b . From classical equations for projectile and target

$$\begin{aligned}
 \bar{D}^\mu \bar{A}_{\mu\bullet}^a &= -g\bar{\psi}_a \gamma_\bullet t^a \psi_a, & \bar{D}^\mu \bar{A}_{\mu i}^a &= -g\bar{\psi}_a \gamma_i t^a \psi_a, & \bar{D}^\mu \bar{A}_{\mu*}^a &= -g\bar{\psi}_a \gamma_* t^a \psi_a \\
 \left[\frac{2}{s}(i\partial_* + g\bar{A}_*)\hat{p}_1 + \frac{2g}{s}\bar{A}_\bullet \hat{p}_2 + (i\partial_i + g\bar{A}_i)\gamma^i \right] \psi_a &= 0 \\
 \bar{D}^\mu \bar{B}_{\mu\bullet}^a &= -g\bar{\psi}_b \gamma_\bullet t^a \psi_b, & \bar{D}^\mu \bar{B}_{\mu i}^a &= -g\bar{\psi}_b \gamma_i t^a \psi_b, & \bar{D}^\mu \bar{B}_{\mu*}^a &= -g\bar{\psi}_b \gamma_* t^a \psi_b \\
 \left[\frac{2}{s}(i\partial_\bullet + g\bar{B}_\bullet)\hat{p}_2 + \frac{2g}{s}\bar{B}_* \hat{p}_1 + (i\partial_i + g\bar{B}_i)\gamma^i \right] \psi_b &= 0
 \end{aligned} \quad (3.11)$$

we get

$$\begin{aligned}
 \hat{p}_1 \psi_a(x_\bullet, x_\perp) &\sim m_\perp^{5/2}, & \gamma_i \psi_a(x_\bullet, x_\perp) &\sim m_\perp^{3/2}, & \hat{p}_2 \psi_a(x_\bullet, x_\perp) &\sim s\sqrt{m_\perp} \\
 \hat{p}_1 \psi_b(x_*, x_\perp) &\sim s\sqrt{m_\perp}, & \gamma_i \psi_b(x_*, x_\perp) &\sim m_\perp^{3/2}, & \hat{p}_2 \psi_b(x_*, x_\perp) &\sim m_\perp^{5/2}
 \end{aligned} \quad (3.12)$$

Thus, to find TMD factorization at the tree level (with higher-twist corrections) we need to calculate the functional integral (2.4) in the background fields of the strength given by eqs. (3.10) and (3.12).

3.2 Approximate solution of classical equations

As we discussed in Sect 2, the calculation of the functional integral (2.13) over C -fields in the tree approximation reduces to finding fields C_μ and ψ_c as solutions of Yang-Mills equations for the action $S_C = S_{\text{QCD}}(C + A + B) - S_{\text{QCD}}(A) - S_{\text{QCD}}(B)$

$$\begin{aligned}
 D^\nu F_{\mu\nu}^a(\bar{A} + \bar{B} + C) &= g \sum_f (\bar{\psi}_a^f + \bar{\psi}_b^f + \bar{\psi}_c^f) \gamma_\mu t^a (\psi_a^f + \psi_b^f + \psi_c^f) \\
 (i\not{\partial} + g\bar{A} + g\bar{B} + gC)(\psi_a^f + \psi_b^f + \psi_c^f) &= m(\psi_a^f + \psi_b^f + \psi_c^f)
 \end{aligned} \quad (3.13)$$

As we discussed above (see also appendix A) the solution of eq. (3.13) which we need corresponds to the sum of set of diagrams in background field $\bar{A} + \bar{B}$ with *retarded* Green functions (see figure 3). The retarded Green functions (in the background-Feynman gauge) are defined as

$$\begin{aligned} (x|\frac{1}{\bar{P}^2 g^{\mu\nu} + 2ig\bar{F}^{\mu\nu} + i\epsilon p_0}|y) &\equiv (x|\frac{1}{p^2 + i\epsilon p_0}|y) - g(x|\frac{1}{p^2 + i\epsilon p_0}\mathcal{O}_{\mu\nu}\frac{1}{p^2 + i\epsilon p_0}|y) \\ &+ g^2(x|\frac{1}{p^2 + i\epsilon p_0}\mathcal{O}_{\mu\xi}\frac{1}{p^2 + i\epsilon p_0}\mathcal{O}_{\nu\xi}\frac{1}{p^2 + i\epsilon p_0}|y) + \dots \end{aligned} \quad (3.14)$$

where

$$\begin{aligned} \bar{P}_\mu &\equiv i\partial_\mu + g\bar{A}_\mu + g\bar{B}_\mu, \quad \bar{F}_{\mu\nu} = \partial_\mu(\bar{A} + \bar{B})_\nu - \mu \leftrightarrow \nu - ig[\bar{A}_\mu + \bar{B}_\mu, \bar{A}_\nu + \bar{B}_\nu] \\ \mathcal{O}_{\mu\nu} &\equiv (\{p^\xi, \bar{A}_\xi + \bar{B}_\xi\} + g(\bar{A} + \bar{B})^2)g_{\mu\nu} + 2i\bar{F}_{\mu\nu} \end{aligned} \quad (3.15)$$

and similarly for quarks.

The solutions of eqs. (3.13) in terms of retarded Green functions give fields C_μ and ψ_c that vanish at $t \rightarrow -\infty$. Thus, we are solving the usual classical YM equations

$$D^\nu F_{\mu\nu}^a = \sum_f g\bar{\psi}^f t^a \gamma_\mu \psi^f, \quad (\not{P} - m_f)\psi^f = 0 \quad (3.16)$$

with boundary conditions

$$\begin{aligned} A_\mu(x) &\stackrel{x_* \rightarrow -\infty}{\equiv} \bar{A}_\mu(x_\bullet, x_\perp), & \psi(x) &\stackrel{x_* \rightarrow -\infty}{\equiv} \psi_a(x_\bullet, x_\perp) \\ A_\mu(x) &\stackrel{x_* \rightarrow -\infty}{\equiv} \bar{B}_\mu(x_*, x_\perp), & \psi(x) &\stackrel{x_* \rightarrow -\infty}{\equiv} \psi_b(x_*, x_\perp) \end{aligned} \quad (3.17)$$

following from $C_\mu, \psi_c \stackrel{t \rightarrow -\infty}{\rightarrow} 0$. These boundary conditions reflect the fact that at $t \rightarrow -\infty$ we have only incoming hadrons with ‘‘A’’ and ‘‘B’’ fields.

The solution of YM equations (3.16) in general case is yet unsolved problem, especially important for scattering of two heavy nuclei in semiclassical approximation. Fortunately, for our case of particle production with $\frac{q_\perp}{Q} \ll 1$ we can construct the approximate solution of (3.16) as a series in this small parameter. However, before doing this, it is convenient to perform a gauge transformation so that the incoming projectile and target fields will no longer have large components $\sim s$ as \bar{A}_* and \bar{B}_\bullet in eq. (3.10). Let us perform the gauge transformation of eq. (3.16) and initial conditions (3.17) with the gauge matrix $\Omega(x)$ such that

$$\Omega(x_*, x_\bullet, x_\perp) \stackrel{x_* \rightarrow -\infty}{\rightarrow} [x_\bullet, -\infty_\bullet]_{x_\perp}^{\bar{A}_*}, \quad \Omega(x_*, x_\bullet, x_\perp) \stackrel{x_* \rightarrow -\infty}{\rightarrow} [x_*, -\infty_*]_{x_\perp}^{\bar{B}_\bullet} \quad (3.18)$$

The existence of such matrix is proved in appendix B by explicit construction. After such gauge transformation, the YM equation of course stays the same but the initial conditions (3.17) turn to

$$\begin{aligned} gA_\mu(x) &\stackrel{x_* \rightarrow -\infty}{\equiv} U_\mu(x_\bullet, x_\perp), & \psi(x) &\stackrel{x_* \rightarrow -\infty}{\equiv} \Sigma_a(x_\bullet, x_\perp) \\ gA_\mu(x) &\stackrel{x_* \rightarrow -\infty}{\equiv} V_\mu(x_*, x_\perp), & \psi(x) &\stackrel{x_* \rightarrow -\infty}{\equiv} \Sigma_b(x_*, x_\perp) \end{aligned} \quad (3.19)$$

where

$$\begin{aligned}
U_\mu(x_\bullet, x_\perp) &\equiv \frac{2}{s} p_{2\mu} U_\bullet(x_\bullet, x_\perp) + U_{\mu\perp}(x_\bullet, x_\perp) \\
V_\mu(x_*, x_\perp) &\equiv \frac{2}{s} p_{1\mu} V_*(x_*, x_\perp) + V_{\mu\perp}(x_*, x_\perp) \\
U_i(x_\bullet, x_\perp) &\equiv \frac{2}{s} \int_{-\infty}^{x_\bullet} dx'_\bullet U_{*i}(x'_\bullet, x_\perp), & V_i(x_*, x_\perp) &\equiv \frac{2}{s} \int_{-\infty}^{x_*} dx'_* V_{\bullet i}(x'_*, x_\perp) \\
U_\bullet(x_\bullet, x_\perp) &\equiv \frac{2}{s} \int_{-\infty}^{x_\bullet} dx'_\bullet U_{*\bullet}(x'_\bullet, x_\perp), & V_*(x_*, x_\perp) &\equiv -\frac{2}{s} \int_{-\infty}^{x_*} dx'_* V_{*\bullet}(x'_*, x_\perp)
\end{aligned} \tag{3.20}$$

and Σ_a, Σ_b are defined as

$$\Sigma_a(z_\bullet, z_\perp) \equiv [-\infty_\bullet, z_\bullet]_z \psi_a(z_\bullet, z_\perp), \quad \Sigma_b(z_*, z_\perp) \equiv [-\infty_*, z_*]_z \psi_b(z_*, z_\perp) \tag{3.21}$$

The initial conditions (3.19) look like the projectile fields in the light-like gauge $p_2^\mu A_\mu = 0$ and target fields in the light-like gauge $p_1^\mu A_\mu = 0$ so our construction of matrix Ω in a way proves that we can take the sum of projectile fields in one gauge and target fields in another gauge as a zero-order approximation for iterative solution of the YM equations. Note also that our power counting discussed in previous section means that

$$U_\bullet \sim V_* \sim m_\perp^2, \quad U_i \sim V_i \sim m_\perp \tag{3.22}$$

so we do not have large background fields $\sim s$ after this gauge transformation. Finally, the classical equations for projectile and target fields in this gauge read:⁶

$$\begin{aligned}
D_U^\nu U_{\mu\nu}^a &= g^2 \sum_f \bar{\Sigma}_a^f \gamma_\mu t^a \Sigma_a^f, & i\cancel{D}_U \Sigma_a &= 0 \\
D_V^\nu V_{\mu\nu}^a &= g^2 \sum_f \bar{\Sigma}_b^f \gamma_\mu t^a \Sigma_b^f, & i\cancel{D}_V \Sigma_b &= 0
\end{aligned} \tag{3.23}$$

where $U_{\mu\nu} \equiv \partial_\mu U_\nu - \partial_\nu U_\mu - i[U_\mu, U_\nu]$, $D_U^\mu \equiv (\partial^\mu - i[U^\mu, \cdot])$ and similarly for V fields.

We will solve eqs. (3.16) iteratively, order by order in perturbation theory, starting from the zero-order approximation in the form of the sum of projectile and target fields

$$\begin{aligned}
g\mathcal{A}_\mu^{[0]}(x) &= U_\mu(x_\bullet, x_\perp) + V_\mu(x_*, x_\perp) \\
\Psi^{[0]}(x) &= \Sigma_a(x_\bullet, x_\perp) + \Sigma_b(x_*, x_\perp)
\end{aligned} \tag{3.24}$$

and improving it by calculation of Feynman diagrams with retarded propagators in the background fields (3.24).

The first step is the calculation of the linear term for the trial configuration (3.24). We rewrite field strength components as

$$\begin{aligned}
g\mathcal{F}_{\bullet i}^{[0]} &= U_{\bullet i} + V_{\bullet i} - i[U_\bullet, V_i], & g\mathcal{F}_{*i}^{[0]} &= U_{*i} + V_{*i} - i[V_*, U_i] \\
g\mathcal{F}_{*\bullet}^{[0]} &= U_{*\bullet} + V_{*\bullet} + i[U_\bullet, V_*], & g\mathcal{F}_{ij}^{[0]} &= U_{ij} + V_{ij} - i[U_i, V_j] + i[U_j, V_i]
\end{aligned} \tag{3.25}$$

Note that $U_{*i} \sim V_{\bullet i} \sim sm_\perp$, $U_{*\bullet} \sim V_{*\bullet} \sim sm_\perp^2$ while all other components are not large.

⁶Here we consider only u, d , and s quarks which can be regarded as massless.

The linear term has the form

$$\begin{aligned}
 L_i^a &\equiv \mathcal{D}^\mu \mathcal{F}_{\mu i}^{[0]a} + g \bar{\Psi}^{[0]} \gamma_i t^a \Psi^{[0]} = L_i^{(0)a} + L_i^{(1)a} \\
 L_i^{(0)a} &= -\frac{i}{g} [U^{jab} V_{ji}^b + V^{jab} U_{ji}^b + \mathcal{D}_j^{ab} (U^{jbc} V_i^c + V^{jbc} U_i^c)] \\
 &\quad - \frac{2i}{gs} (U_{*\bullet}^{ab} V_i^b - V_{*\bullet}^{ab} U_i^b) + g \bar{\Sigma}_a t^a \gamma_i \Sigma_b + g \bar{\Sigma}_b t^a \gamma_i \Sigma_a \\
 L_i^{(1)a} &= -\frac{2i}{gs} [U_{\bullet*}^{ab} V_{*i}^b + V_{*i}^{ab} U_{\bullet*}^b - i\{U_{\bullet}, V_{*}\}^{ab} U_i^b - i\{V_{*}, U_{\bullet}\}^{ab} V_i^b] \\
 L_{\bullet}^a &\equiv \mathcal{D}^\mu \mathcal{F}_{\mu\bullet}^{[0]a} + g \bar{\Psi}^{[0]} \gamma_{\bullet} t^a \Psi^{[0]} = L_{\bullet}^{(-1)a} + L_{\bullet}^{(0)a} + L_{\bullet}^{(1)a}, \quad L_{\bullet}^{(-1)a} = \frac{i}{g} U^{jab} V_{\bullet j}^b \\
 L_{\bullet}^{(0)a} &= \frac{i}{g} V^{jab} U_{\bullet j}^b + \frac{i}{g} \mathcal{D}^{jab} U_{\bullet}^{bc} V_j^c + g \bar{\Sigma}_a t^a \gamma_{\bullet} \Sigma_b + g \bar{\Sigma}_b t^a \gamma_{\bullet} \Sigma_a - \frac{4i}{gs} U_{\bullet}^{ab} V_{*\bullet}^b \\
 L_{\bullet}^{(1)a} &= \frac{2}{gs} (U_{\bullet} U_{\bullet})^{ab} V_{*}^b \\
 L_{*}^a &\equiv \mathcal{D}^\mu \mathcal{F}_{\mu*}^{[0]a} + g \bar{\Psi}^{[0]} \gamma_{*} t^a \Psi^{[0]} = L_{*}^{(-1)a} + L_{*}^{(0)a} + L_{*}^{(1)a}, \quad L_{*}^{(-1)a} = \frac{i}{g} V^{jab} U_{*j}^b \\
 L_{*}^{(0)a} &= \frac{i}{g} U^{jab} V_{*j}^b + \frac{i}{g} \mathcal{D}^{jab} V_{*}^{bc} U_j^c + g \bar{\Sigma}_a t^a \gamma_{*} \Sigma_b + g \bar{\Sigma}_b t^a \gamma_{*} \Sigma_a + \frac{4i}{gs} V_{*}^{ab} U_{*\bullet}^b \\
 L_{*}^{(1)a} &= \frac{2}{gs} (V_{*} V_{*})^{ab} U_{\bullet}^b \\
 L_{\psi} &\equiv \not{P} \Psi^{[0]} = L_{\psi}^{(0)} + L_{\psi}^{(1)} \\
 L_{\psi}^{(0)} &= \gamma^i U_i \Sigma_b + \gamma^i V_i \Sigma_a, \quad L_{\psi}^{(1)} = \frac{2}{s} \hat{p}_2 U_{\bullet} \Sigma_b + \frac{2}{s} \hat{p}_1 V_{*} \Sigma_a
 \end{aligned} \tag{3.26}$$

where $\mathcal{D}^j \equiv \partial^j - iU^j - iV^j$, $\mathcal{D}_{\bullet} = \partial_{\bullet} - iU_{\bullet}$, and $\mathcal{D}_{*} = \partial_{*} - iV_{*}$. The power-counting estimates for linear terms in eq. (3.26) are

$$\begin{aligned}
 L_i^{(0)} &\sim m_{\perp}^3, & L_i^{(1)} &\sim \frac{m_{\perp}^5}{s} \\
 L_{\bullet}^{(-1)} &\sim L_{*}^{(-1)} \sim sm_{\perp}^2, & L_{\bullet}^{(0)} &\sim L_{*}^{(0)} \sim m_{\perp}^4, & L_{\bullet}^{(1)} &\sim L_{*}^{(1)} \sim \frac{m_{\perp}^6}{s} \\
 L_{\psi}^{(0)} &\sim m_{\perp}^{5/2}, & L_{\psi}^{(1)} &\sim \frac{m_{\perp}^{9/2}}{s}
 \end{aligned} \tag{3.27}$$

Note that the order of perturbation theory is labeled by $(\dots)^{[n]}$ and the order of expansion in the parameter $\frac{m_{\perp}^2}{s}$ by $(\dots)^{(n)}$.

With the linear term (3.26), a couple of first terms in perturbative series are

$$\begin{aligned}
 A_{\mu}^{[1]a}(x) &= \int d^4 z (x | \frac{1}{\mathcal{P}^2 g^{\mu\nu} + 2ig\mathcal{F}^{[0]\mu\nu}} | z)^{ab} L^{b\nu}(z) \\
 A_{\mu}^{[2]a}(x) &= g \int d^4 z \left[-i(x | \frac{1}{\mathcal{P}^2 g^{\mu\eta} + 2ig\mathcal{F}^{[0]\mu\eta}} \mathcal{P}^{\xi} | z)^{aa'} f^{a'bc} A_{\xi}^{[1]b} A^{[1]cn} \right. \\
 &\quad \left. + (x | \frac{1}{\mathcal{P}^2 g^{\mu\eta} + 2ig\mathcal{F}^{[0]\mu\eta}} | z)^{aa'} f^{a'bc} A^{[1]b\xi} (\mathcal{D}_{\xi} A^{[1]c\eta} - \mathcal{D}^{\eta} A_{\xi}^{[1]c}) \right]
 \end{aligned} \tag{3.28}$$

for gluon fields (in the background-Feynman gauge) and

$$\Psi_f^{[1]}(x) = -\int d^4z (x|\frac{1}{\not{P}}|z)L_\psi(z), \quad \Psi_f^{[2]}(x) = -g\int d^4z (x|\frac{1}{\not{P}}|z)A^{[1]}(z)\Psi_f^{[0]}(z) \quad (3.29)$$

for quarks where

$$\mathcal{P}_\bullet = i\partial_\bullet + U_\bullet, \quad \mathcal{P}_* = i\partial_* + V_*, \quad \mathcal{P}_i = i\partial_i + U_i + V_i \quad (3.30)$$

are operators in external zero-order fields (3.24). Hereafter we use Schwinger's notations for propagators in external fields normalized according to $(x|F(p)|y) \equiv \int \bar{d}^4p e^{-ip(x-y)}F(p)$. Moreover, when it will not lead to a confusion, we will use short-hand notation $\frac{1}{\mathcal{O}}\mathcal{O}'(x) \equiv \int d^4z (x|\frac{1}{\mathcal{O}}|z)\mathcal{O}'(z)$. Next iterations will give us a set of tree-level Feynman diagrams in the background field $U_\mu + V_\mu$ and $\Sigma_a + \Sigma_b$.

Let us consider the fields in the first order in perturbation theory:

$$\begin{aligned} A_\mu^{[1]} &= \frac{1}{\mathcal{P}^2 g^{\mu\nu} + 2ig\mathcal{F}^{[0]\mu\nu}}L^\nu \quad (3.31) \\ &= \frac{1}{[\{\alpha + \frac{2}{s}V_*, \beta + \frac{2}{s}U_\bullet\}_2^s - (p+U+V)_\perp^2]g^{\mu\nu} + 2ig\mathcal{F}^{[0]\mu\nu} + i\epsilon p_0}L^\nu \\ \Psi_f^{[1]}(x) &= -\frac{1}{\not{P}}L_\psi = -\frac{(\alpha + \frac{2}{s}V_*)\not{p}_1 + (\beta + \frac{2}{s}U_\bullet)\not{p}_2 + \not{P}_\perp}{\{\alpha + \frac{2}{s}V_*, \beta + \frac{2}{s}U_\bullet\}_2^s - (p+U+V)_\perp^2 + i\epsilon p_0}L_\psi \end{aligned}$$

Here α , β , and p_\perp are understood as differential operators $\alpha = i\frac{\partial}{\partial x_\bullet}$, $\beta = i\frac{\partial}{\partial x_*}$ and $p_i = i\frac{\partial}{\partial x^i}$.

Now comes the central point of our approach. Let us expand quark and gluon propagators in powers of background fields, then we get a set of diagrams shown in figure 3. The typical bare gluon propagator in figure 3 is

$$\frac{1}{p^2 + i\epsilon p_0} = \frac{1}{\alpha\beta s - p_\perp^2 + i\epsilon(\alpha + \beta)} \quad (3.32)$$

Since we do not consider loops of C -fields in this paper, the transverse momenta in tree diagrams are determined by further integration over projectile (“A”) and target (“B”) fields in eq. (2.8) which converge on either q_\perp or m . On the other hand, the integrals over α converge on either α_q or $\alpha \sim 1$ and similarly the characteristic β 's are either β_q or ~ 1 . Since $\alpha_q\beta_qs = Q_\parallel^2 \gg Q_\perp^2$, one can expand gluon and quark propagators in powers of $\frac{p_\perp^2}{\alpha\beta s}$

$$\begin{aligned} \frac{1}{p^2 + i\epsilon p_0} &= \frac{1}{s(\alpha + i\epsilon)(\beta + i\epsilon)} \left(1 + \frac{p_\perp^2/s}{(\alpha + i\epsilon)(\beta + i\epsilon)} + \dots \right) \quad (3.33) \\ \frac{\not{p}}{p^2 + i\epsilon p_0} &= \frac{1}{s} \left(\frac{\not{p}_1}{\beta + i\epsilon} + \frac{\not{p}_2}{\alpha + i\epsilon} + \frac{\not{p}_\perp}{(\alpha + i\epsilon)(\beta + i\epsilon)} \right) \left(1 + \frac{p_\perp^2/s}{(\alpha + i\epsilon)(\beta + i\epsilon)} + \dots \right) \end{aligned}$$

The explicit form of operators $\frac{1}{\alpha+i\epsilon}$, $\frac{1}{\beta+i\epsilon}$, and $\frac{1}{(\alpha+i\epsilon)(\beta+i\epsilon)}$ is

$$\begin{aligned}
 (x|\frac{1}{\alpha+i\epsilon}|y) &= \frac{s}{2} \int \bar{d}^2 p_{\perp} \int \frac{\bar{d}\alpha}{\alpha+i\epsilon} \bar{d}\beta e^{-i\alpha(x-y)_{\bullet}-i\beta(x-y)_{*}+i(p,x-y)_{\perp}} \\
 &= -i\frac{s}{2}(2\pi)^2 \delta^{(2)}(x_{\perp}-y_{\perp}) \theta(x_{\bullet}-y_{\bullet}) \delta(x_{*}-y_{*}) \\
 (x|\frac{1}{\beta+i\epsilon}|y) &= \frac{s}{2} \int \bar{d}^2 p_{\perp} \int \bar{d}\alpha \frac{\bar{d}\beta}{\beta+i\epsilon} e^{-i\alpha(x-y)_{\bullet}-i\beta(x-y)_{*}+i(p,x-y)_{\perp}} \\
 &= -i\frac{s}{2}(2\pi)^2 \delta^{(2)}(x_{\perp}-y_{\perp}) \theta(x_{*}-y_{*}) \delta(x_{\bullet}-y_{\bullet}) \\
 (x|\frac{1}{(\alpha+i\epsilon)(\beta+i\epsilon)}|y) &= \frac{s}{2} \int \bar{d}^2 p_{\perp} \int \frac{\bar{d}\alpha}{\alpha+i\epsilon} \frac{\bar{d}\beta}{\beta+i\epsilon} e^{-i\alpha(x-y)_{\bullet}-i\beta(x-y)_{*}+i(p,x-y)_{\perp}} \\
 &= -\frac{s}{2}(2\pi)^2 \delta^{(2)}(x_{\perp}-y_{\perp}) \theta(x_{*}-y_{*}) \theta(x_{\bullet}-y_{\bullet}) \tag{3.34}
 \end{aligned}$$

After the expansion (3.33), the dynamics in the transverse space effectively becomes trivial: all background fields stand either at x or at 0 . (This validates the reasoning in the footnote on page 3).

One may wonder why we do not cut the integrals in eq. (3.34) to $|\alpha| > \sigma_b$ and $|\beta| > \sigma_a$ according to the definition of C fields in section 2.7. The reason is that in the diagrams like figure 3 with retarded propagators (3.34) one can shift the contour of integration over α and/or β to the complex plane away to avoid the region of small α or β .⁸

Note that the background fields are also smaller than typical $p_{\parallel}^2 \sim s$. Indeed, from eq. (3.22) we see that $p_{\bullet} = \frac{s}{2}\beta \gg U_{\bullet} \sim m^2$ (because $\alpha \geq \alpha_q \gg \frac{m^2}{s}$) and similarly $p_{*} \gg V_{*}$. Also $(p_i + U_i + V_i)^2 \sim q_{\perp}^2 \ll p_{\parallel}^2$. The only exception is the fields $V_{\bullet i}$ or U_{*i} which are of order of sm_{\perp} but we will see that effectively the expansion in powers of these fields is cut at the second term with our accuracy.

3.3 Twist expansion of classical gluon fields

Now we expand the classical gluon fields in powers of $\frac{p_{\perp}^2}{p_{\parallel}^2} \sim \frac{m_{\perp}^2}{s}$. It is clear that for the leading higher-twist correction we need to take into account only the first two terms (3.28) of the perturbative expansion of classical field. The expansion (3.28) of gluon field A_{\bullet} takes

⁷Such cutoffs for integrals over C fields are introduced explicitly in the framework of soft-collinear effective theory, see the review [24].

⁸This may be wrong if there is pinching of poles in the integrals over α or β but we will see that in our integrals for the tree-level power corrections the pinching of poles never occurs. In the higher orders in perturbation theory the pinching does occur so one needs to formulate a subtraction program to avoid double counting.

the form

$$\begin{aligned}
 A_{\bullet}^{[0]} + A_{\bullet}^{[1]} &= A_{\bullet}^{(0)} + A_{\bullet}^{(1)} + O\left(\frac{m_{\perp}^6}{s^2}\right) \quad (3.35) \\
 A_{\bullet}^{(0)a} &= A_{\bullet}^{([1]0)a} + \frac{1}{g}U_{\bullet}^a = \frac{1}{p_{\parallel}^2}L_{\bullet}^{(-1)a} + \frac{1}{g}U_{\bullet}^a = \frac{1}{g}U_{\bullet}^a + \frac{1}{2g\alpha}U_j^{ab}V^{jb} \\
 A_{\bullet}^{(1)a} &= \frac{1}{p_{\parallel}^2}L_{\bullet}^{(0)a} + \frac{1}{2gp_{\parallel}^2}((\{\alpha, U_{\bullet}\} + \{\beta, V_{\bullet}\} - \mathcal{P}_{\perp}^2)V^j)^{ab}\frac{1}{\alpha}U_j^b - 2i\frac{1}{p_{\parallel}^2}(V_{\bullet}^i)^{ab}A_i^{(1)b} \\
 &\quad + \frac{4i}{s}\frac{1}{p_{\parallel}^2}(U_{\bullet\bullet} + V_{\bullet\bullet})^{ab}\frac{1}{p_{\parallel}^2}L_{\bullet}^{(-1)b} - \frac{igf^{abc}}{\alpha s}A_{\bullet}^{([1]0)b}A_{\bullet}^{([1]0)c} - \frac{1}{p_{\parallel}^2}A_{\bullet}^{([1]0)ab}U_j^{bc}V^{cj}
 \end{aligned}$$

where

$$\begin{aligned}
 A_{\bullet}^{([1]0)a} &\equiv \frac{1}{p_{\parallel}^2}L_{\bullet}^{(-1)a} = \frac{i}{2\alpha g}f^{abc}U_j^bV^{cj}, \quad A_{\bullet}^{([1]0)a} \equiv \frac{1}{p_{\parallel}^2}L_{\bullet}^{(-1)a} = -\frac{i}{2\beta g}f^{abc}U_j^bV^{cj} \\
 \Rightarrow \mathcal{D}_{\bullet}A_{\bullet}^{([1]0)a} - \mathcal{D}_{\bullet}A_{\bullet}^{([1]0)a} &= \frac{s}{2g}f^{abc}U_j^bV^{cj} + O(m_{\perp}^2) \quad (3.36)
 \end{aligned}$$

Similarly, from eq. (3.28) one obtains

$$\begin{aligned}
 A_{\bullet}^{[0]} + A_{\bullet}^{[1]} &= A_{\bullet}^{(0)} + A_{\bullet}^{(1)} + O\left(\frac{m_{\perp}^6}{s^2}\right) \quad (3.37) \\
 A_{\bullet}^{(0)a} &= A_{\bullet}^{([1]0)a} + \frac{1}{g}V_{\bullet}^a = \frac{1}{p_{\parallel}^2}L_{\bullet}^{(-1)a} + \frac{1}{g}V_{\bullet}^a = \frac{1}{g}V_{\bullet}^a - \frac{1}{2g\beta}U_j^{ab}V^{jb} \\
 A_{\bullet}^{(1)a} &= \frac{1}{p_{\parallel}^2}L_{\bullet}^{(0)a} + \frac{1}{2gp_{\parallel}^2}((\{\alpha, U_{\bullet}\} + \{\beta, V_{\bullet}\} - \mathcal{P}_{\perp}^2)U^j)^{ab}\frac{1}{\beta}V_j^b - 2i\frac{1}{p_{\parallel}^2}(U_{\bullet}^i)^{ab}A_i^{(1)b} \\
 &\quad - \frac{4i}{s}\frac{1}{p_{\parallel}^2}(U_{\bullet\bullet} + V_{\bullet\bullet})^{ab}A_{\bullet}^{([1]0)b} + \frac{igf^{abc}}{\beta s}A_{\bullet}^{([1]0)b}A_{\bullet}^{([1]0)c} - \frac{1}{p_{\parallel}^2}A_{\bullet}^{([1]0)ab}V_j^{bc}U^{cj}
 \end{aligned}$$

and

$$\begin{aligned}
 A_i^{[0]} &= A_i^{(0)} = \frac{1}{g}(U_i + V_i) \quad (3.38) \\
 A_i^{[1]} + A_i^{[2]} &= A_i^{(1)} + A_i^{(2)} + O\left(\frac{m_{\perp}^7}{s^3}\right), \quad A_i^{(1)} = \frac{1}{p_{\parallel}^2}\tilde{L}_i^{(0)} \sim \frac{m_{\perp}^3}{s} \\
 A_i^{(2)a} &= \frac{1}{p_{\parallel}^2}\tilde{L}_i^{(1)a} + \frac{1}{p_{\parallel}^2}(\mathcal{P}_{\perp}^2 - \{\alpha, U_{\bullet}\} - \{\beta, V_{\bullet}\})^{ab}A_i^{(1)b} - 2i\frac{1}{p_{\parallel}^2}(\mathcal{F}_i^{[0]k})^{ab}A_k^{(1)b} + \dots
 \end{aligned}$$

where ($n = 1, 2$)

$$\tilde{L}_i^{(0)} = L_i^{(0)} + \frac{4i}{s}\left(V_{\bullet i}\frac{1}{p_{\parallel}^2}L_{\bullet}^{(-1)} + U_{\bullet i}\frac{1}{p_{\parallel}^2}L_{\bullet}^{(-1)}\right) = L_i^{(0)} - \frac{2i}{gs}(V_{\bullet i}U^j)^{ab}\frac{1}{\beta}V_j^b - \frac{2i}{gs}(U_{\bullet i}V^j)^{ab}\frac{1}{\alpha}U_j^b \quad (3.39)$$

In these formulas the singularity in $\frac{1}{\alpha}$ is always causal $\frac{1}{\alpha+i\epsilon}$ and similarly for $\frac{1}{\beta} \equiv \frac{1}{\beta+i\epsilon}$ and $\frac{1}{p_{\parallel}^2} \equiv \frac{1/s}{(\alpha+i\epsilon)(\beta+i\epsilon)}$, see eq. (3.34).

The corresponding expansion of field strengths reads

$$\begin{aligned}
 gF_{\bullet i}^{(-1)a}(x) &= V_{\bullet i}^a(x), & gF_{*i}^{(-1)a}(x) &= U_{*i}^a(x) \\
 gF_{\bullet i}^{(0)a}(x) &= U_{\bullet i}^a(x) - iU_{\bullet}^{ab}(x)V_i^b(x) - \frac{ig}{2\alpha}\tilde{L}_i^{(0)a}(x) + \mathcal{D}_i^{ab}V_j^{bc}(x)\frac{1}{2\alpha}U^{cj}(x) \\
 gF_{*i}^{(0)a}(x) &= V_{*i}^a(x) - iV_{*}^{ab}(x)U_i^b(x) - \frac{ig}{2\beta}\tilde{L}_i^{(0)a}(x) + \mathcal{D}_i^{ab}U_j^{bc}(x)\frac{1}{2\beta}V^{cj}(x) \\
 gF_{*\bullet}^{(-1)a}(x) &= U_{*\bullet}^a(x) + V_{*\bullet}^a(x) - \frac{is}{2}U_j^{ab}(x)V^{bj}(x) \\
 gF_{ik}^{(0)a}(x) &= U_{ik}^a(x) + V_{ik}^a(x) - i(U_i^{ab}(x)V_k^b(x) - i \leftrightarrow k)
 \end{aligned} \tag{3.40}$$

Power corrections to hadronic tensor are proportional to

$$F^2(x) \equiv F_{\mu\nu}^a(x)F^{a\mu\nu}(x) = \frac{8}{s}F_{\bullet i}^a(x)F_*^{ai}(x) + F_{ik}^a(x)F^{aik}(x) - \frac{8}{s^2}F_{*\bullet}^a(x)F_{*\bullet}^a(x) \tag{3.41}$$

so

$$\begin{aligned}
 (F^2(x))^{(-1)} &= \frac{8}{sg^2}U_{*i}^aV_{\bullet}^{ai} \\
 (F^2(x))^{(0)} &= F_{ik}^{(0)a}(x)F^{(0)aik} - \frac{8}{s^2}F_{*\bullet}^{(-1)a}(x)F_{*\bullet}^{(-1)a}(x) \\
 &\quad + \frac{8}{sg}V_{\bullet}^{ai}(x)F_{*i}^{(0)a}(x) + \frac{8}{sg}U_*^{ai}(x)F_{\bullet i}^{(0)a}(x)
 \end{aligned} \tag{3.42}$$

and the leading higher-twist correction is proportional to

$$\begin{aligned}
 (F^2(x))^{(0)}(F^2(0))^{(-1)} + (x \leftrightarrow 0) &= \left[F_{ik}^{(0)a}(x)F^{(0)aik}(x) - \frac{8}{s^2}F_{*\bullet}^{(-1)a}(x)F_{*\bullet}^{(-1)a}(x) \right. \\
 &\quad \left. + \frac{8}{sg}V_{\bullet}^{ai}(x)F_{*i}^{(0)a}(x) + \frac{8}{sg}U_*^{ai}(x)F_{\bullet i}^{(0)a}(x) \right] \frac{8}{sg^2}U_{*i}^a(0)V_{\bullet}^{ai}(0) + (x \leftrightarrow 0)
 \end{aligned} \tag{3.43}$$

4 Leading higher-twist correction at $s \gg Q^2 \gg Q_{\perp}^2 \gg m^2$

As we mentioned in the Introduction, our method is relevant for calculation of higher-twist corrections at any $s, Q^2 \gg Q_{\perp}^2, m^2$. However, the expressions become manageable in the physically interesting case $s \gg Q^2 \gg Q_{\perp}^2 \gg m^2$ which we consider in this section.⁹ We will demonstrate that the leading correction in this region comes from the following part of eq. (3.41)

$$g^2F^2(x) = \frac{8}{s}U_*^{ai}(x)V_{\bullet i}^a(x) + 2f^{mac}f^{mbd}\Delta^{ij,kl}U_i^a(x)U_j^b(x)V_k^c(x)V_l^d(x) + \dots \tag{4.1}$$

where

$$\Delta^{ij,kl} \equiv g^{ij}g^{kl} - g^{ik}g^{jl} - g^{il}g^{jk} \tag{4.2}$$

⁹We also assume that the scalar particle is emitted in the central region of rapidity so $\alpha_{qs} \sim \beta_{qs} \gg Q^2$.

The higher-twist correction coming from the second term in the r.h.s. will be $\sim \frac{Q_\perp^2}{Q^2}$ whereas other terms in the r.h.s. of eq. (3.41) yield contributions $\sim \frac{Q_\perp^2}{s}$, $\sim \frac{Q_\perp^2}{\alpha_q s}$, or $\sim \frac{Q_\perp^2}{\beta_q s}$ all of which are small (see the footnote 9). In this approximation we get

$$\begin{aligned}
 g^4 F^2(x) F^2(0) &= \frac{64}{s^2} U_*^{mi}(x) V_{\bullet i}^m(x) U_*^{nj}(0) V_{\bullet j}^n(0) \\
 &+ \frac{16}{s} f^{mac} f^{mbd} \Delta^{ij,kl} [U_i^a(x) U_j^b(x) V_k^c(x) V_l^d(x) U_*^{nr}(0) V_{\bullet r}^n(0) \\
 &+ U_*^{nr}(x) V_{\bullet r}^n(x) U_i^a(0) U_j^b(0) V_k^c(0) V_l^d(0)]
 \end{aligned} \tag{4.3}$$

where the first term is the leading order and the second is the higher-twist correction.

Substituting our approximation (4.1) to eq. (2.3) and promoting background fields to operators as discussed in section 2 we get (note that $\alpha_q \beta_q s = Q_{\parallel}^2 \simeq Q^2$):

$$\begin{aligned}
 W(p_A, p_B, q) &= \frac{64/s^2}{N_c^2 - 1} \int d^2 x_\perp \frac{2}{s} \int dx_\bullet dx_* \cos(\alpha_q x_\bullet + \beta_q x_* - (q, x)_\perp) \\
 &\times \left\{ \langle p_A | \hat{U}_*^{mi}(x_\bullet, x_\perp) \hat{U}_*^{mj}(0) | p_A \rangle \langle p_B | \hat{V}_{\bullet i}^n(x_*, x_\perp) \hat{V}_{\bullet j}^n(0) | p_B \rangle \right. \\
 &- \frac{4N_c^2}{N_c^2 - 4} \frac{\Delta^{ij,kl}}{Q^2} \frac{2}{s} \int_{-\infty}^{x_\bullet} dx'_\bullet d^{abc} \langle p_A | \hat{U}_{*i}^a(x_\bullet, x_\perp) \hat{U}_{*j}^b(x'_\bullet, x_\perp) \hat{U}_{*r}^c(0) | p_A \rangle \\
 &\times \left. \frac{2}{s} \int_{-\infty}^{x_*} dx'_* d^{mpq} \langle p_B | \hat{V}_{\bullet k}^m(x_*, x_\perp) \hat{V}_{\bullet l}^p(x'_*, x_\perp) \hat{V}_{\bullet r}^q(0) | p_B \rangle \right\}
 \end{aligned} \tag{4.4}$$

where we used formula [25, 26]

$$f^{acm} f^{bdm} d^{abn} d^{cdn} = \frac{1}{2} (N_c^2 - 1)(N_c^2 - 4) \tag{4.5}$$

Since an extra U_{*k} (or $V_{\bullet k}$) brings $s \frac{x_i}{x_\perp^2}$ ¹⁰ we see that the higher-twist correction in the r.h.s. of eq. (4.4) is $\sim \frac{q_\perp^2}{Q^2}$ so it gives the leading power correction in the region $s \gg Q^2 = m_\Phi^2 \gg q_\perp^2 \gg m^2$. The TMD factorization formula with the higher-twist correction (4.4) is the main result of the present paper.

We parametrize gluon TMD for unpolarized protons as (cf. ref. [27])

$$\begin{aligned}
 &\frac{4}{s^2 g^2} \int dx_* \int d^2 x_\perp e^{-i\beta_q x_* + i(k, x)_\perp} \langle p_B | V_{\bullet i}^a(x_*, x_\perp) V_{\bullet j}^a(0) | p_B \rangle \\
 &= -\pi \beta_q \left[g_{ij} D_g(\beta_q, k_\perp^2; \sigma_b) - \left(2 \frac{k_i k_j}{m^2} + g_{ij} \frac{k_\perp^2}{m^2} \right) H_g(\beta_q, k_\perp^2; \sigma_b) \right]
 \end{aligned} \tag{4.6}$$

where σ_b is the cutoff in α integration in the target matrix elements, see the discussion in ref. [18]. The normalization here is such that $D_g(\beta_q, k_\perp^2; \sigma_b)$ is an unintegrated gluon distribution:

$$\int \bar{d}^2 k_\perp D_g(\beta_q, k_\perp^2; \sigma_b) = D_g(\beta_q, \mu^2 = \sigma_b \beta_q s) \tag{4.7}$$

¹⁰To see this, we compared matrix elements of leading-twist operator $\langle p_A | U_*^{mi}(x_\bullet, x_\perp) U_*^{mj}(0) | p_A \rangle$ and higher-twist operator $\langle p_A | U_{*i}^a(x_\bullet, x_\perp) U_{*j}^b(x'_\bullet, x_\perp) U_{*r}^c(0) | p_A \rangle$ between quark states which gives an extra $s \frac{x_i}{x_\perp^2}$ modulo some logarithms.

where $D_g(\beta_q, \mu^2)$ is the usual gluon parton density (this formula is correct in the leading log approximation, see the discussion in ref. [18]).

Next, the three-gluon matrix element in eq. (4.4) for unpolarized hadrons can be parametrized as

$$\begin{aligned} & \frac{4}{s^2 g^3} \int dx_* \int d^2 x_\perp e^{-i\beta_q x_* + i(k, x)_\perp} \int_{-\infty}^{x_*} d\frac{x'_*}{s} d^{abc} \langle p_B | V_{\bullet i}^a(x_*, x_\perp) V_{\bullet j}^b(x'_*, x_\perp) V_{\bullet r}^c(0) | p_B \rangle + i \leftrightarrow j \\ & = -\pi \beta_q \left[(k_i g_{jr} + k_j g_{ir}) D_1^g(\beta_q, k_\perp^2; \sigma_b) + k_r g_{ij} D_2^g(\beta_q, k_\perp^2; \sigma_b) \right. \\ & \quad \left. - \left[k_i k_j k_r + \frac{k_\perp^2}{4} (k_r g_{ij} + k_i g_{jr} + k_j g_{ir}) \right] \frac{1}{m^2} H_1^g(\beta_q, k_\perp^2; \sigma_b) \right] \end{aligned} \quad (4.8)$$

At large k_\perp^2 gluon TMDs in the r.h.s. of eq. (4.6) behave as $D_g(\beta_q, k_\perp^2) \sim \frac{1}{k_\perp^2}$ and $H_g(\beta_q, k_\perp^2) \sim \frac{1}{k_\perp^4}$. Similarly, one should expect that $D_i^g(\beta_q, k_\perp^2) \sim \frac{1}{k_\perp^2}$ and $H_1^g(\beta_q, k_\perp^2) \sim \frac{1}{k_\perp^4}$.

It is well known that in our kinematic region $s \gg Q^2 \gg Q_\perp^2$ gluon TMDs (4.6) possess Sudakov logs of the type

$$\frac{4}{s^2 g^2} \int dx_* \int d^2 x_\perp e^{-i\beta_q x_* + i(k, x)_\perp} \langle p_B | V_{\bullet i}^n(x_*, x_\perp) V_{\bullet n}^i(0) | p_B \rangle \sim e^{-\frac{\alpha_s N_c}{2\pi} \ln^2 \frac{\sigma_b s}{k_\perp^2}} D_g \left(\beta_q, k_\perp^2; \ln \frac{k_\perp^2}{s} \right) \quad (4.9)$$

One should expect double-logs of this type in $D_i^g(\beta_q, k_\perp^2; \sigma_b)$ and $H_1^g(\beta_q, k_\perp^2; \sigma_b)$, too.

Let us now demonstrate that the terms in $(F^2(x))^{(0)}$ (see eq. (3.42)) which we neglected give small contributions. For example, consider the following contribution to $F^2(x)F^2(0)$:

$$-\frac{64i}{s^2} U_*^{ai}(x) V_{\bullet i}^a(x) V_{\bullet j}^{bj}(0) V_*^{bc}(0) U_j^c(0) \quad (4.10)$$

The corresponding contribution to hadronic tensor W has the form

$$\begin{aligned} & -\frac{64/s^2}{N_c^2 - 1} \int d^2 x_\perp e^{i(q, x)_\perp} \frac{2}{s} \int dx_\bullet dx_* e^{-i\alpha_q x_\bullet - i\beta_q x_*} \\ & \quad \times \frac{2}{\alpha_q s} \langle p_A | U_*^{ai}(x_\bullet, x_\perp) U_*^{aj}(0) | p_A \rangle \langle p_B | V_{\bullet i}^b(x_*, x_\perp) V_*^{bc}(0) V_{\bullet j}^c(0) | p_B \rangle \end{aligned} \quad (4.11)$$

Note that unlike eq. (4.4), the factor in the denominator is $\alpha_q s \gg Q^2$ so the contribution (4.11) is power suppressed in comparison to eq. (4.4) in our kinematic region.¹¹

As a less trivial example, consider the following term in $F^2(x)F^2(0)$

$$-\frac{64}{s^3} U_{*i}^a(x) V_{\bullet i}^{ai}(x) V_{\bullet j}^{bj}(0) \frac{1}{\beta} (V_{\bullet j} U^k)^{bc} \frac{1}{\beta} V_k^c(0) \quad (4.12)$$

The corresponding contribution to hadronic tensor W reads

$$\begin{aligned} & \frac{64/s^2}{N_c^2 - 1} \int d^2 x_\perp e^{i(q, x)_\perp} \frac{2}{s} \int dx_\bullet dx_* e^{-i\alpha_q x_\bullet - i\beta_q x_*} \left\{ \frac{i}{\alpha_q s} \langle p_A | U_*^{mi}(x_\bullet, x_\perp) U_*^{mj}(0) | p_A \rangle \right. \\ & \quad \left. \times \frac{4}{s^2} \int_{-\infty}^{0_*} dz_* \int_{-\infty}^{z_*} dz'_* (z - z'_*) \langle p_B | V_{\bullet i}^a(x_*, x_\perp) V_{\bullet k}^{bk}(0) (V_{\bullet k}(z_*, 0_\perp) T^a)^{bc} V_{\bullet j}^c(z'_*, 0_\perp) | p_B \rangle \right\} \end{aligned} \quad (4.13)$$

¹¹Of course, this power suppression may be moderated by difference in logarithmic evolution of operators in the r.h.s.'s of eqs. (4.4) and (4.11), but one should expect the evolution of these operators to be of the same order of magnitude.

where we used

$$\frac{1}{\beta + i\epsilon} V_k(x) = -i \int_{-\infty}^{x_*} dx'_* V_k(x'_*, x_\perp) = -\frac{2i}{s} \int_{-\infty}^{x_*} dx'_* (x - x'_*) V_{\bullet k}(x'_*, x_\perp)$$

In both examples (4.11) and (4.13) the factor $\frac{1}{\alpha_q}$ comes from an extra integration over x'_\bullet in U_i , see eq. (3.20):

$$\begin{aligned} \int dx_\bullet e^{-i\alpha_q x_\bullet} \langle U_i(x_\bullet, x_\perp) U_j(0) \rangle &= \frac{2}{s} \int dx_\bullet \int_{-\infty}^{x_\bullet} dx'_\bullet e^{-i\alpha_q x_\bullet} \langle U_{*i}(x'_\bullet, x_\perp) U_j(0) \rangle \\ &= -\frac{2i}{\alpha_q s} \int dx_\bullet e^{-i\alpha_q x_\bullet} \langle U_{*i}(x_\bullet, x_\perp) U_j(0) \rangle \end{aligned} \quad (4.14)$$

The way to figure out such integrations is very simple: take $\alpha_q \rightarrow 0$ and check if there is an infinite integration of the type $\int_{-\infty}^{x_\bullet} dx'_\bullet$. Evidently, it may happen if we have a single $U_i(x)$ (without any additional U -operators) at the point x , or a single $U_i(0)$.

Similarly, the factor $\frac{1}{\beta_q}$ comes from an extra integration over x'_* in V_i in eq. (3.20) so an indication of such contribution is the infinite integration $\int_{-\infty}^{x_*} dx'_*$ in the limit $\beta_q \rightarrow 0$ which translates to the condition of a single V_i at the point x or at the point 0.

Thus, to get the terms $\sim \frac{1}{Q^2}$ we need to find contributions which satisfy both of the above conditions which singles out the contribution (4.3).

5 Small-x limit and scattering of shock waves

Let us consider the hadronic tensor

$$\langle p_A, p_B | g^4 F^2(x) F^2(y) | p_A, p_B \rangle \quad (5.1)$$

in the small-x limit $s \rightarrow \infty$, Q^2 and q_\perp^2 - fixed. At first, let us not impose the condition $Q^2 \gg q_\perp^2$ which means that the relation between x_\parallel^2 and x_\perp^2 is arbitrary (later we will see that $Q^2 \gg q_\perp^2$ corresponds to $x_\parallel^2 \ll x_\perp^2$).

The small-x limit may be obtained by rescaling $s \rightarrow \lambda^2 s \Leftrightarrow p_1 \rightarrow \lambda p_1, p_2 \rightarrow \lambda p_2$. As discussed in refs. [7, 20, 28], the only components of field strength surviving in this rescaling are $U_{*i}(x_\bullet, x_\perp)$ and $V_{\bullet i}(x_*, x_\perp)$. Moreover, if we study classical fields at longitudinal distances which does not scale with λ , we can replace the projectile and target fields by infinitely thin “shock waves”

$$U_{*i}(x_\bullet, x_\perp) \rightarrow \frac{s}{2} \delta(x_\bullet) \mathcal{U}_i(x_\perp) \quad \text{and} \quad V_{\bullet i}(x_*, x_\perp) \rightarrow \frac{s}{2} \delta(x_*) \mathcal{V}_i(x_\perp) \quad (5.2)$$

However, since we need to compare the classical fields in the small- x limit to our expressions (3.40) at small longitudinal distances, we will keep x_* and x_\bullet dependence for a while.

As described above, to find the classical fields we can start with the trial configuration

$$\begin{aligned} g\mathcal{A}_i^{[0]}(x) &= U_i(x_\bullet, x_\perp) + V_i(x_*, x_\perp), & \mathcal{A}_*^{[0]} &= \mathcal{A}_\bullet^{[0]} = 0 \\ \Psi^{[0]}(x) &= \Sigma_a(x_\bullet, x_\perp) + \Sigma_b(x_*, x_\perp), & \not{p}_1 \Sigma_a &= \not{p}_2 \Sigma_b = \gamma_i \Sigma_a = \gamma_i \Sigma_b = 0 \end{aligned} \quad (5.3)$$

with the linear term

$$gL_\mu^a = \frac{2ip_{1\mu}}{s} V^{jab} U_{*j}^b + \frac{2ip_{2\mu}}{s} U^{jab} V_{\bullet j}^b - i\mathcal{D}_j^{ab}(U^{jbc} V_\mu^{\perp c} + V^{jbc} U_\mu^{\perp c}) \quad (5.4)$$

and improve it order by order in L_μ . In this way we'll get a set of Feynman diagrams in the background field (5.3). Unfortunately, in the general case of arbitrary relation between q_\parallel and q_\perp we no longer have a small parameter $\frac{p_\perp^2}{p_\parallel^2}$ so we need explicit expressions for propagators in the background fields, and, in addition, we need all orders in the expansion of linear term (5.4). Still, we can compare our calculations with the perturbative expansion of classical fields in powers of the ‘‘parameter’’ $[U_i, V_j]$ carried out in refs. [7, 8]. In the leading order in perturbation theory only the first line of eq. (3.28) survives and we get

$$\begin{aligned} gA_\bullet &= \frac{i}{p^2 + i\epsilon p_0} [U^j, V_{\bullet j}], & gA_* &= \frac{i}{p^2 + i\epsilon p_0} [V^j, U_{*j}] \\ gA_i &= U_i + V_i + \frac{p^j}{p^2 + i\epsilon p_0} ([U_i, V_j] - i \leftrightarrow j) \end{aligned} \quad (5.5)$$

The corresponding expressions for field strengths are

$$\begin{aligned} gF_{\bullet i} &= V_{\bullet i} - \frac{p^j}{p^2 + i\epsilon p_0} (g_{ij}[U^k, V_{\bullet k}] + [U_j, V_{\bullet i}] - [U_i, V_{\bullet j}]) \\ gF_{*i} &= U_{*i} - \frac{p^j}{p^2 + i\epsilon p_0} (g_{ij}[V^k, U_{*k}] + [V_j, U_{*i}] - [V_i, U_{*j}]) \\ gF_{*\bullet} &= \frac{2i}{p^2 + i\epsilon p_0} [U_*^j, V_{\bullet j}] \\ gF_{ij} &= -i[U_i, V_j] - \frac{ip_i p^k}{p^2 + i\epsilon p_0} ([U_j, V_k] - j \leftrightarrow k) - i \leftrightarrow j = \frac{4i/s}{p^2 + i\epsilon p_0} ([U_{*i}, V_{\bullet j}] - i \leftrightarrow j) \end{aligned} \quad (5.6)$$

In the last line we used the identity

$$p_i([U_j, V_k] - j \leftrightarrow k) - i \leftrightarrow j = -p_k([U_i, V_j] - i \leftrightarrow j) \quad (5.7)$$

and the fact that in the small- x limit $\partial_i U_j - \partial_j U_i - i[U_i, U_j] = \partial_i V_j - \partial_j V_i - i[V_i, V_j] = 0$.

Let us discuss now how our approximation $\frac{p_\perp^2}{p_\parallel^2} \ll 1$ looks in the coordinate space. The explicit expressions for fields (5.6) are

$$\begin{aligned} gF_{\bullet i}(x) &= V_{\bullet i}(x_*, x_\perp) + \frac{i}{4\pi} \int dz \frac{1}{(x-z)_*} \frac{\partial}{\partial x_j} \theta[(x-z)_\parallel^2 - (x-z)_\perp^2] \theta(x-z)_* gL_{ij}^-(z) \\ gF_{*i}(x) &= U_{*i}(x_\bullet, x_\perp) - \frac{i}{4\pi} \int dz \frac{1}{(x-z)_\bullet} \frac{\partial}{\partial x_j} \theta[(x-z)_\parallel^2 - (x-z)_\perp^2] \theta(x-z)_\bullet gL_{ij}^+(z) \\ gF_{*\bullet}(x) &= -\frac{i}{\pi} \int dz \delta[(x-z)_\parallel^2 - (x-z)_\perp^2] \theta(x-z)_* [U_*^j(z_\bullet, z_\perp), V_{\bullet j}(z_*, z_\perp)] \\ gF_{ij}(x) &= -\frac{2i}{\pi s} \int dz \delta[(x-z)_\parallel^2 - (x-z)_\perp^2] \theta(x-z)_* ([U_{*i}(z_\bullet, z_\perp), V_{\bullet j}(z_*, z_\perp)] - i \leftrightarrow j) \end{aligned} \quad (5.8)$$

where

$$gL_{ij}^\pm(z) \equiv g_{ij}[U_*^k, V_{\bullet k}] \pm [U_{*i}, V_{\bullet j}] \mp [U_{*j}, V_{\bullet i}] \quad (5.9)$$

At longitudinal distances $x_\bullet, x_* \sim 1$ these expressions agree with eq. (52) from ref. [7] after the replacement (5.2).

Now let us compare the fields (5.8) at small longitudinal distances to our approximate solution (3.40). Let us start with $F_{ij}(x)$ in the last line in eq. (5.8). If $(x-z)_\parallel^2$ is smaller than the characteristic transverse distances in the integral over z_\perp one can replace $[U_{*i}(z_\bullet, z_\perp), V_{\bullet j}(z_*, z_\perp)]$ by $[U_{*i}(z_\bullet, x_\perp), V_{\bullet j}(z_*, x_\perp)]$ and get

$$\begin{aligned}
 gF_{ij}(x) &= -\frac{2i}{s} \int d^2 z_\parallel \theta(x-z)_* \theta(x-z)_\bullet ([U_{*i}(z_\bullet, x_\perp), V_{\bullet j}(z_*, x_\perp)] - i \leftrightarrow j) \\
 &= -i[U_i(x_\bullet, x_\perp), V_j(x_*, x_\perp)] + i[U_j(x_\bullet, x_\perp), V_i(x_*, x_\perp)]
 \end{aligned}
 \tag{5.10}$$

which is exactly the last line in eq. (3.40). Similarly, the third line in eq. (5.8) reproduces $F_{*\bullet}$ in the fourth line in eq. (3.40).

Next, $gF_{\bullet i}^{(0)a}$ in second line in eq. (3.40) in the leading order in perturbation theory turns to

$$-\frac{\partial^j}{2\alpha} (g_{ij}[U^k, V_k] - [U_i, V_j] + [U_j, V_i]) = \frac{2i}{s^2} \int_{-\infty}^{x_*} dz_* \int_{-\infty}^{x_\bullet} dz_\bullet (x-z)_\bullet \partial^j L_{ij}^-(z_*, z_\bullet, x_\perp)
 \tag{5.11}$$

On the other hand, the first line in eq. (5.8) at small $(x-z)_\parallel$ gives

$$\begin{aligned}
 &\frac{i}{4\pi} \int dz \frac{\theta(x-z)_*}{(x-z)_*} \theta[(x-z)_\parallel^2 - (x-z)_\perp^2] \frac{\partial}{\partial z_j} L_{ij}^-(z) \\
 &\simeq \frac{i}{4\pi} \int dz \frac{\theta(x-z)_*}{(x-z)_*} \theta\left[\frac{4}{s}(x-z)_*(x-z)_\bullet - (x-z)_\perp^2\right] \partial^j L_{ij}^-(z_*, z_\bullet, x_\perp)
 \end{aligned}
 \tag{5.12}$$

which agrees with eq. (5.11) after integration over z_\perp . Similarly, one can check the consistency of two expressions for F_{*i} .

6 Conclusions and outlook

We have formulated the approach to TMD factorization based on the factorization in rapidity and found the leading higher-twist contribution to the production of a scalar particle (e.g. Higgs) by gluon-gluon fusion in the hadron-hadron scattering. Up to now our results are obtained in the tree-level approximation when the question of exact matching of cutoffs in rapidity does not arise. However, this question will become crucial starting from the first loop. In our previous papers we calculated the evolution of gluon TMD with respect to our rapidity cutoff so we need to match it to the coefficient functions in front of TMD operators. The work is in progress.

Also, we obtained power corrections for particle production only in the case of gluon-gluon fusion. It would be interesting (and we plan) to find power corrections to Drell-Yan process. There is a statement that for semi-inclusive deep inelastic scattering (SIDIS) the leading-order TMDs have different directions of Wilson lines: one to $+\infty$ and another to $-\infty$. We think that the same directions of Wilson lines will be in the case of power corrections and we plan to study this question in forthcoming publications.

The authors are grateful to J.C. Collins, S. Dawson, A. Kovner, D. Neill, A. Prokudin, T. Rogers, and R. Venugopalan for valuable discussions. This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under contracts DE-AC02-98CH10886 and DE-AC05-06OR23177.

A Diagrams with retarded propagators

In this section we prove that the field C_μ created by a source J_μ in the presence of external fields \bar{A}_μ and \bar{B}_μ ¹²

$$\begin{aligned} \langle C_\mu^a(x) \rangle_J \equiv & \int D\tilde{C}DC C_\mu^a(x) \exp \left\{ \int dz \left[\frac{i}{2} \tilde{C}^{m\xi} \square_{\xi\eta}^{mn} \tilde{C}^{m\eta} \right. \right. \\ & + ig f^{mnl} \bar{D}^\xi \tilde{C}^{m\eta} \tilde{C}_\xi^n \tilde{C}_\eta^l + \frac{ig^2}{4} f^{abm} f^{cdm} \tilde{C}^{a\xi} \tilde{C}^{b\eta} \tilde{C}_\xi^c \tilde{C}_\eta^d - iJ_\xi^m \tilde{C}^{m\xi} - \frac{i}{2} C^{m\xi} \square_{\xi\eta}^{mn} C^{m\eta} \\ & \left. \left. - ig f^{mnl} \bar{D}^\xi C^{m\eta} C_\xi^n C_\eta^l - \frac{ig^2}{4} f^{abm} f^{cdm} C^{a\xi} C^{b\eta} C_\xi^c C_\eta^d + iJ_\xi^m C^{m\xi} \right] \right\} \end{aligned} \quad (\text{A.1})$$

is given by a set of Feynman diagrams with retarded Green functions (note that eq. (A.1) implies that J_μ, \bar{A}_μ , and \bar{B}_μ are the same in the right and left part of the amplitude). Hereafter we use the notation $\square_{\mu\nu} \equiv \bar{P}^2 g_{\mu\nu} + 2i\bar{G}_{\mu\nu}$.

First, we consider gluon propagators for the double functional integral over C fields in the background fields $\bar{A} = \bar{\tilde{A}}, \bar{B} = \bar{\tilde{B}}$ and prove that

$$\begin{aligned} \langle C_\mu^a(x) C_\nu^b(y) \rangle - \langle C_\mu^a(x) \tilde{C}_\nu^b(y) \rangle &= (x | \frac{-i}{\square_{\mu\nu} + i\epsilon p_0} | y)^{ab} \\ \langle \tilde{C}_\mu^a(x) C_\nu^b(y) \rangle - \langle \tilde{C}_\mu^a(x) \tilde{C}_\nu^b(y) \rangle &= (x | \frac{-i}{\square_{\mu\nu} + i\epsilon p_0} | y)^{ab} \end{aligned} \quad (\text{A.2})$$

Note that we define $\langle \mathbb{O} \rangle$ in this section as

$$\langle \mathbb{O} \rangle \equiv \int D\tilde{C}DC \mathbb{O} e^{\int dz \left(\frac{i}{2} \tilde{C}^{a\mu} \square_{\mu\nu}^{ab} \tilde{C}^{b\nu} - \frac{i}{2} C^{a\mu} \square_{\mu\nu}^{ab} C^{b\nu} \right)} \quad (\text{A.3})$$

To prove eq. (A.2), we write down

$$\square_{\mu\nu} = p^2 g_{\mu\nu} + \mathcal{O}_{\mu\nu}, \quad \mathcal{O}_{\mu\nu} \equiv (\{p^\xi, \bar{A}_\xi + \bar{B}_\xi\} + (\bar{A} + \bar{B})^2) g_{\mu\nu} + 2i\bar{G}_{\mu\nu} \quad (\text{A.4})$$

and expand in powers of $\mathcal{O}_{\mu\nu}$.

In the trivial order eqs. (A.2) immediately follow from the bare propagators for the double functional integral (A.3)

$$\begin{aligned} \langle C_\mu^a(x) C_\nu^b(y) \rangle_{\text{bare}} &= (x | \frac{-ig_{\mu\nu} \delta^{ab}}{p^2 + i\epsilon} | y), & \langle \tilde{C}_\mu^a(x) \tilde{C}_\nu^b(y) \rangle_{\text{bare}} &= (x | \frac{ig_{\mu\nu} \delta^{ab}}{p^2 - i\epsilon} | y) \\ \langle C_\mu^a(x) \tilde{C}_\nu^b(y) \rangle_{\text{bare}} &= -g_{\mu\nu} \delta^{ab} (x | 2\pi \delta(p^2) \theta(-p_0) | y) \end{aligned} \quad (\text{A.5})$$

where

$$\langle \mathbb{O} \rangle_{\text{bare}} \equiv \int D\tilde{C}DC \mathbb{O} e^{\int dz \left(\frac{i}{2} C^{a\mu} \partial^2 C_\mu^a - \frac{i}{2} \tilde{C}^{a\mu} \partial^2 \tilde{C}_\mu^a \right)} \quad (\text{A.6})$$

¹²For simplicity, in this section we disregard quarks so in our case J_μ is eq. (3.26) without quark terms.

In the first order in $\mathcal{O}_{\mu\nu}$ we get

$$\begin{aligned}
 \langle C_\mu^a(x) C_\nu^b(y) \rangle^{(1)} &= i \int dz \left[- \langle C_\mu^a(x) C^{c\xi}(z) \rangle_{\text{bare}} \mathcal{O}_{\xi\eta}^{cd}(z) \langle C^{d\eta}(z) C_\nu^b(y) \rangle_{\text{bare}} \right. \\
 &\quad \left. + \langle C_\mu^a(x) \tilde{C}^{c\xi}(z) \rangle_{\text{bare}} \mathcal{O}_{\xi\eta}^{cd}(z) \langle \tilde{C}^{d\eta}(z) C_\nu^b(y) \rangle_{\text{bare}} \right] \\
 \langle C_\mu^a(x) \tilde{C}_\nu^b(y) \rangle^{(1)} &= i \int dz \left[- \langle C_\mu^a(x) C^{c\xi}(z) \rangle_{\text{bare}} \mathcal{O}_{\xi\eta}^{cd}(z) \langle C^{d\eta}(z) \tilde{C}_\nu^b(y) \rangle_{\text{bare}} \right. \\
 &\quad \left. + \langle C_\mu^a(x) \tilde{C}^{c\xi}(z) \rangle_{\text{bare}} \mathcal{O}_{\xi\eta}^{cd}(z) \langle \tilde{C}^{d\eta}(z) \tilde{C}_\nu^b(y) \rangle_{\text{bare}} \right] \quad (\text{A.7})
 \end{aligned}$$

so

$$\langle C_\mu^a(x) C_\nu^b(y) \rangle^{(1)} - \langle C_\mu^a(x) \tilde{C}_\nu^b(y) \rangle^{(1)} = i(x) \left| \frac{1}{p^2 + i\epsilon p_0} \mathcal{O}_{\mu\nu}^{ab} \frac{1}{p^2 + i\epsilon p_0} \right| y \quad (\text{A.8})$$

Similarly, it is easy to see that

$$\begin{aligned}
 \langle \tilde{C}_\mu^a(x) C_\nu^b(y) \rangle^{(1)} &= i \int dz \left[- \langle \tilde{C}_\mu^a(x) C^{c\xi}(z) \rangle_{\text{bare}} \mathcal{O}_{\xi\eta}^{cd}(z) \langle C^{d\eta}(z) C_\nu^b(y) \rangle_{\text{bare}} \right. \\
 &\quad \left. + \langle \tilde{C}_\mu^a(x) \tilde{C}^{c\xi}(z) \rangle_{\text{bare}} \mathcal{O}_{\xi\eta}^{cd}(z) \langle \tilde{C}^{d\eta}(z) C_\nu^b(y) \rangle_{\text{bare}} \right] \\
 \langle \tilde{C}_\mu^a(x) \tilde{C}_\nu^b(y) \rangle^{(1)} &= i \int dz \left[- \langle \tilde{C}_\mu^a(x) C^{c\xi}(z) \rangle_{\text{bare}} \mathcal{O}_{\xi\eta}^{cd}(z) \langle C^{d\eta}(z) \tilde{C}_\nu^b(y) \rangle_{\text{bare}} \right. \\
 &\quad \left. + \langle \tilde{C}_\mu^a(x) \tilde{C}^{c\xi}(z) \rangle_{\text{bare}} \mathcal{O}_{\xi\eta}^{cd}(z) \langle \tilde{C}^{d\eta}(z) \tilde{C}_\nu^b(y) \rangle_{\text{bare}} \right] \quad (\text{A.9})
 \end{aligned}$$

so

$$\langle \tilde{C}_\mu^a(x) C_\nu^b(y) \rangle^{(1)} - \langle \tilde{C}_\mu^a(x) \tilde{C}_\nu^b(y) \rangle^{(1)} = i(x) \left| \frac{1}{p^2 + i\epsilon p_0} \mathcal{O}_{\mu\nu}^{ab} \frac{1}{p^2 + i\epsilon p_0} \right| y \quad (\text{A.10})$$

In the second order in $\mathcal{O}_{\mu\nu}$

$$\begin{aligned}
 \langle C_\mu^a(x) C_\nu^b(y) \rangle^{(2)} &= i \int dz \left[- \langle C_\mu^a(x) C^{c\xi}(z) \rangle^{(1)} \mathcal{O}_{\xi\eta}^{cd}(z) \langle C^{d\eta}(z) C_\nu^b(y) \rangle_{\text{bare}} \right. \\
 &\quad \left. + \langle C_\mu^a(x) \tilde{C}^{c\xi}(z) \rangle^{(1)} \mathcal{O}_{\xi\eta}^{cd}(z) \langle \tilde{C}^{d\eta}(z) C_\nu^b(y) \rangle_{\text{bare}} \right] \\
 \langle C_\mu^a(x) \tilde{C}_\nu^b(y) \rangle^{(2)} &= i \int dz \left[- \langle C_\mu^a(x) C^{c\xi}(z) \rangle^{(1)} \mathcal{O}_{\xi\eta}^{cd}(z) \langle C^{d\eta}(z) \tilde{C}_\nu^b(y) \rangle_{\text{bare}} \right. \\
 &\quad \left. + \langle C_\mu^a(x) \tilde{C}^{c\xi}(z) \rangle^{(1)} \mathcal{O}_{\xi\eta}^{cd}(z) \langle \tilde{C}^{d\eta}(z) \tilde{C}_\nu^b(y) \rangle_{\text{bare}} \right] \quad (\text{A.11})
 \end{aligned}$$

so using the results (A.8) and (A.10) we get

$$\langle C_\mu^a(x) C_\nu^b(y) \rangle^{(2)} - \langle C_\mu^a(x) \tilde{C}_\nu^b(y) \rangle^{(2)} = -i(x) \left| \frac{1}{p^2 + i\epsilon p_0} \mathcal{O}_{\mu\xi} \frac{1}{p^2 + i\epsilon p_0} \mathcal{O}_\nu^\xi \frac{1}{p^2 + i\epsilon p_0} \right| y^{ab} \quad (\text{A.12})$$

Similarly, it is easy to demonstrate that

$$\langle \tilde{C}_\mu^a(x) C_\nu^b(y) \rangle^{(2)} - \langle \tilde{C}_\mu^a(x) \tilde{C}_\nu^b(y) \rangle^{(2)} = -i(x) \left| \frac{1}{p^2 + i\epsilon p_0} \mathcal{O}_{\mu\xi} \frac{1}{p^2 + i\epsilon p_0} \mathcal{O}_\nu^\xi \frac{1}{p^2 + i\epsilon p_0} \right| y^{ab} \quad (\text{A.13})$$

One can prove now eq. (A.2) by induction using formulas

$$\begin{aligned}
 \langle C_\mu^a(x) C_\nu^b(y) \rangle^{(n)} &= i \int dz \left[- \langle C_\mu^a(x) C^{c\xi}(z) \rangle^{(n-1)} \mathcal{O}_{\xi\eta}^{cd}(z) \langle C^{d\eta}(z) C_\nu^b(y) \rangle_{\text{bare}} \right. \\
 &\quad \left. + \langle C_\mu^a(x) \tilde{C}^{c\xi}(z) \rangle^{(n-1)} \mathcal{O}_{\xi\eta}^{cd}(z) \langle \tilde{C}^{d\eta}(z) C_\nu^b(y) \rangle_{\text{bare}} \right] \\
 \langle C_\mu^a(x) \tilde{C}_\nu^b(y) \rangle^{(n)} &= i \int dz \left[- \langle C_\mu^a(x) C^{c\xi}(z) \rangle^{(n-1)} \mathcal{O}_{\xi\eta}^{cd}(z) \langle C^{d\eta}(z) \tilde{C}_\nu^b(y) \rangle_{\text{bare}} \right. \\
 &\quad \left. + \langle C_\mu^a(x) \tilde{C}^{c\xi}(z) \rangle^{(n-1)} \mathcal{O}_{\xi\eta}^{cd}(z) \langle \tilde{C}^{d\eta}(z) \tilde{C}_\nu^b(y) \rangle_{\text{bare}} \right] \quad (\text{A.14})
 \end{aligned}$$

Now we are in a position to prove eq. (A.1). In the leading order in g it is trivial: using eqs. (A.2) one immediately sees that

$$\begin{aligned}
 \langle C_\mu^a(x) \rangle_J^{[0]} &= \int D\tilde{C}DC C_\mu^a(x) e^{\int dz \left(\frac{i}{2} \tilde{C}^{a\xi} \square_{\xi\eta}^{ab} \tilde{C}^{b\eta} - i J_\xi^a \tilde{C}^{a\xi} - \frac{i}{2} C^{a\xi} \square_{\xi\eta}^{ab} C^{b\eta} + i J_\xi^a C^{a\xi} \right)} \\
 &= \int D\tilde{C}DC \tilde{C}_\mu^a(x) e^{\int dz \left(-\frac{i}{2} \tilde{C}^{a\xi} \square_{\xi\eta}^{ab} \tilde{C}^{b\eta} - i J_\xi^a \tilde{C}^{a\xi} + \frac{i}{2} C^{a\xi} \square_{\xi\eta}^{ab} C^{b\eta} + i J_\xi^a C^{a\xi} \right)} \\
 &= \int dz (x | \frac{1}{\square_{\mu\nu} + i\epsilon p_0} | z)^{ab} J^{b\nu}(z) \tag{A.15}
 \end{aligned}$$

In the first order in g (with one three-gluon vertex) we obtain

$$\begin{aligned}
 \langle C_\mu^a(x) \rangle_J^{[1]} &= -ig f^{mnl} \int D\tilde{C}DC C_\mu^a(x) \int dz [\bar{D}^\xi C^{m\eta} C_\xi^m C_\eta^l(z) - \bar{D}^\xi \tilde{C}^{m\eta} \tilde{C}_\xi^m \tilde{C}_\eta^l(z)] \\
 &\quad \times \exp \left\{ \int dz' \left[\frac{i}{2} \tilde{C}^{a\xi} \square_{\xi\eta}^{ab} \tilde{C}^{b\eta} - i J_\xi^a \tilde{C}^{a\xi} - \frac{i}{2} C^{a\xi} \square_{\xi\eta}^{ab} C^{b\eta} + i J_\xi^a C^{a\xi} \right] (z') \right\} \\
 &= \frac{ig}{2} f^{mnl} \int dz dz' dz'' \langle C_\mu^a(x) \left[\bar{D}^\xi C^{m\eta} C_\xi^m C_\eta^l(z) - \bar{D}^\xi \tilde{C}^{m\eta} \tilde{C}_\xi^m \tilde{C}_\eta^l(z) \right] \\
 &\quad \times [J_\alpha^c C^{c\alpha}(z') - J_\alpha^c \tilde{C}^{c\alpha}(z')] [J_\beta^d C^{d\beta}(z'') - J_\beta^d \tilde{C}^{d\beta}(z'')] \rangle \\
 &= -ig f^{mnl} \int dz \left\{ (\langle C_\mu^a(x) \bar{D}^\xi C^{m\eta}(z) \rangle - \langle C_\mu^a(x) \bar{D}^\xi \tilde{C}^{m\eta}(z) \rangle) \langle C_\xi^m(z) \rangle_J^{[0]} \langle C_\eta^l(z) \rangle_J^{[0]} \right. \\
 &\quad \left. + (\langle C_\mu^a(x) C_\xi^m(z) \rangle - \langle C_\mu^a(x) \tilde{C}_\xi^m(z) \rangle) (\langle \bar{D}^\xi C^{m\eta}(z) \rangle_J^{[0]} - \xi \leftrightarrow \eta) \langle C_\eta^l(z) \rangle_J^{[0]} \right\} \\
 &= -ig f^{mnl} \int dz \left\{ (x | \frac{1}{\square_{\mu\eta} + i\epsilon p_0} \bar{P}^\xi | z)^{am} \langle C_\xi^m(z) \rangle_J^{[0]} \langle C_\eta^l(z) \rangle_J^{[0]} \right. \\
 &\quad \left. - i (x | \frac{1}{\square_{\mu\xi} + i\epsilon p_0} | z)^{an} (\langle \bar{D}^\xi C^{m\eta}(z) \rangle_J^{[0]} - \xi \leftrightarrow \eta) \langle C_\eta^l(z) \rangle_J^{[0]} \right\} \tag{A.16}
 \end{aligned}$$

which is the desired result.

Similarly, in the g^2 order one obtains after some algebra

$$\begin{aligned}
 \langle C_\mu^a(x) \rangle_J^{[2]} & \tag{A.17} \\
 &= -ig f^{mnl} \int dz \left\{ (x | \frac{1}{\square_{\mu\eta} + i\epsilon p_0} \bar{P}^\xi | z)^{am} \left[\langle C_\xi^m(z) \rangle_J^{[1]} \langle C_\eta^l(z) \rangle_J^{[0]} + \langle C_\xi^m(z) \rangle_J^{[0]} \langle C_\eta^l(z) \rangle_J^{[1]} \right] \right. \\
 &\quad \left. + i (x | \frac{1}{\square_{\mu\xi} + i\epsilon p_0} | z)^{am} \left[(\langle \bar{D}^\xi C^{m\eta}(z) \rangle_J^{[1]} - \xi \leftrightarrow \eta) \langle C_\eta^l(z) \rangle_J^{[0]} + (\langle \bar{D}^\xi C^{m\eta}(z) \rangle_J^{[0]} - \xi \leftrightarrow \eta) \right. \right. \\
 &\quad \left. \left. \times \langle C_\eta^l(z) \rangle_J^{[1]} \right] \right\} + g^2 \int d^4z (x | \frac{1}{\square_{\mu\xi} + i\epsilon p_0} | z)^{am} f^{mnb} f^{cdn} \langle C^{b\eta}(z) \rangle_J^{[0]} \langle C^{c\xi}(z) \rangle_J^{[0]} \langle C_\eta^d(z) \rangle_J^{[0]}
 \end{aligned}$$

At arbitrary order in g the structure similar to eq. (A.17) can be proved by induction.

Thus, we see that eq. (A.1) is given by a set of Feynman diagrams with retarded Green functions. In a similar way, one can demonstrate that

$$\begin{aligned}
 &\int D\tilde{C}DC \tilde{C}_\mu^a(x) \exp \left\{ \int dz \left[\frac{i}{2} \tilde{C}^{m\xi} \square_{\xi\eta}^{mn} \tilde{C}^{n\eta} \right. \right. \\
 &\quad \left. \left. + ig f^{mnl} \bar{D}^\xi \tilde{C}^{m\eta} \tilde{C}_\xi^m \tilde{C}_\eta^l + \frac{ig^2}{4} f^{abm} f^{cdm} \tilde{C}^{a\xi} \tilde{C}^{b\eta} \tilde{C}_\xi^c \tilde{C}_\eta^d - i J_\xi^m \tilde{C}^{m\xi} \right. \right. \\
 &\quad \left. \left. - \frac{i}{2} C^{m\xi} \square_{\xi\eta}^{mn} C^{n\eta} - ig f^{mnl} \bar{D}^\xi C^{m\eta} C_\xi^m C_\eta^l - \frac{ig^2}{4} f^{abm} f^{cdm} C^{a\xi} C^{b\eta} C_\xi^c C_\eta^d + i J_\xi^m C^{m\xi} \right] \right\} \\
 &= \text{r.h.s. of eq. (A.1)} = \langle C_\mu^a(x) \rangle_J \tag{A.18}
 \end{aligned}$$

B Solution of Yang-Mills equations in two dimensions

To find matrix $\Omega(x)$ satisfying eqs. (3.18) we will solve the following auxiliary problem: we fix x_\perp as a parameter and find the solution of Yang-Mills equations

$$\mathcal{D}^\nu \mathcal{F}_{\mu\nu}^a(x_*, x_\bullet) = 0 \quad (\text{B.1})$$

in 2-dimensional gluodynamics with initial conditions

$$\mathcal{A}_\mu(x_*, x_\bullet) \stackrel{x_* \rightarrow -\infty}{\equiv} \bar{A}_\mu(x_\bullet), \quad \mathcal{A}_\mu(x_*, x_\bullet) \stackrel{x_\bullet \rightarrow -\infty}{\equiv} \bar{B}_\mu(x_*) \quad (\text{B.2})$$

Since 2-dimensional gluodynamics is a trivial theory, the solution of the equation (B.1) will be a pure-gauge field $\mathcal{A}_\mu = \Omega i \partial_\mu \Omega^\dagger$ with $\Omega(x_*, x_\bullet)$ being the sought-for matrix satisfying eqs. (3.18).

Let us first demonstrate that the solution $\mathcal{A}_\mu(x_*, x_\bullet)$ of the YM equations (B.1) with boundary conditions (B.2) in two longitudinal dimensions is a pure gauge. To this end, we will construct $\mathcal{A}_\mu(x_*, x_\bullet)$ order by order in perturbation theory (see figure 3, but now in two dimensions) and prove that $F_{\mu\nu}^a(\mathcal{A}) = 0$.

We are looking for the solution of eq. (B.1) in the form

$$\mathcal{A}_\mu(x_*, x_\bullet) = \bar{A}_\mu(x_*, x_\bullet) + \bar{C}_\mu(x_*, x_\bullet), \quad \bar{A}_*(x_*, x_\bullet) = \bar{A}_*(x_\bullet), \quad \bar{A}_\bullet(x_*, x_\bullet) = \bar{B}_\bullet(x_*) \quad (\text{B.3})$$

Imposing the background-gauge condition

$$\bar{D}^\mu \bar{C}_\mu(x_*, x_\bullet) = 0 \quad (\text{B.4})$$

we get the equation

$$(\bar{P}^2 g_{\mu\nu} + 2ig \bar{F}_{\mu\nu})^{ab} \bar{C}^{b\nu} = \bar{D}^{ab\xi} \bar{F}_{\xi\mu}^b + g f^{abc} (2\bar{C}_\nu^b \bar{D}^\nu \bar{C}_\mu^c - \bar{C}_\nu^b \bar{D}_\mu \bar{C}^{c\nu}) - g^2 f^{abm} f^{cdm} \bar{C}^{b\nu} \bar{C}_\mu^c \bar{C}_\nu^d \quad (\text{B.5})$$

where $\bar{D}_\mu \equiv (\partial_\mu - ig[\bar{A}_\mu, \cdot])$ and $\bar{F}_{*\bullet} = -ig[\bar{A}_*, \bar{B}_\bullet]$. The boundary conditions (B.2) in terms of C fields read

$$C_\mu(x_*, x_\bullet) \stackrel{x_* \rightarrow -\infty}{\equiv} 0, \quad C_\mu(x_*, x_\bullet) \stackrel{x_\bullet \rightarrow -\infty}{\equiv} 0 \quad (\text{B.6})$$

It is convenient to rewrite the equation (B.5) in components as

$$\begin{aligned} & 2(\bar{P}_\bullet \bar{P}_*)^{ab} \bar{C}_\bullet^b \\ &= \bar{D}_\bullet^{ab} \bar{F}_{*\bullet}^b + ig \bar{F}_{*\bullet}^{ab} \bar{C}_\bullet^b + g \bar{D}_\bullet^{aa'} (f^{a'bc} \bar{C}_*^b \bar{C}_\bullet^c) + 2g f^{abc} \bar{C}_\bullet^b \bar{D}_* \bar{C}_\bullet^c - g^2 f^{abm} f^{cdm} \bar{C}_\bullet^b \bar{C}_\bullet^c \bar{C}_*^d \\ & 2(\bar{P}_* \bar{P}_\bullet)^{ab} \bar{C}_*^b \\ &= -\bar{D}_*^{ab} \bar{F}_{*\bullet}^b - ig \bar{F}_{*\bullet}^{ab} \bar{C}_*^b - g \bar{D}_*^{aa'} (f^{a'bc} \bar{C}_*^b \bar{C}_\bullet^c) + 2g f^{abc} \bar{C}_*^b \bar{D}_\bullet \bar{C}_*^c - g^2 f^{abm} f^{cdm} \bar{C}_*^b \bar{C}_*^c \bar{C}_\bullet^d \end{aligned} \quad (\text{B.7})$$

We will solve this equation by iterations in $\bar{F}_{*\bullet}$ and prove that $\mathcal{F}_{*\bullet} = 0$ in all orders.

In the first order we get the equation

$$2(\bar{P}_\bullet \bar{P}_*)^{ab} \bar{C}_\bullet^b = \bar{D}_\bullet^{ab} \bar{F}_{*\bullet}^b, \quad 2(\bar{P}_* \bar{P}_\bullet)^{ab} \bar{C}_*^b = -\bar{D}_*^{ab} \bar{F}_{*\bullet}^b \quad (\text{B.8})$$

The solution satisfying boundary conditions (B.6) has the form

$$\begin{aligned}\bar{C}_\bullet^{(1)} &= -\frac{i/2}{\bar{P}_* + i\epsilon} \bar{F}_{*\bullet} & \Leftrightarrow & \quad \bar{C}_\bullet^{(1)a}(x) = -\frac{i}{2} \int d^2 z_{\parallel} (x | \frac{1}{\bar{P}_* + i\epsilon} | z)^{ab} \bar{F}_{*\bullet}^b(z) \\ \bar{C}_*^{(1)} &= \frac{i/2}{\bar{P}_\bullet + i\epsilon} \bar{F}_{*\bullet} & \Leftrightarrow & \quad \bar{C}_*^{(1)a}(x) = \frac{i}{2} \int d^2 z_{\parallel} (x | \frac{1}{\bar{P}_\bullet + i\epsilon} | z)^{ab} \bar{F}_{*\bullet}^b(z)\end{aligned}\quad (\text{B.9})$$

Using the explicit form of the propagators in external \bar{A}_* and \bar{B}_\bullet fields

$$\begin{aligned}(x | \frac{1}{\bar{P}_\bullet + i\epsilon} | z) &= -i\delta(x_\bullet - z_\bullet)\theta(x_* - z_*)[x_*, z_*]^{\bar{B}_\bullet} \\ (x | \frac{1}{\bar{P}_* + i\epsilon} | z) &= -i\delta(x_* - z_*)\theta(x_\bullet - z_\bullet)[x_\bullet, z_\bullet]^{\bar{A}_*}\end{aligned}\quad (\text{B.10})$$

we get $\bar{C}^{(1)}$ in the form

$$\begin{aligned}\bar{C}_*^{(1)}(x) &= -\frac{i}{s} \int_{-\infty}^{x_*} dz_* [x_*, z_*]^{A_\bullet} [\bar{A}_*(x_\bullet), \bar{A}_*(z_*)][z_*, x_*]^{A_\bullet} \\ \bar{C}_\bullet^{(1)}(x) &= \frac{i}{s} \int_{-\infty}^{x_\bullet} dz_\bullet [x_\bullet, z_\bullet]^{A_*} [\bar{A}_*(z_\bullet), \bar{A}_*(x_*)][z_\bullet, x_\bullet]^{A_*}\end{aligned}\quad (\text{B.11})$$

From this equation it is clear that $C_\mu^{(1)}(x_*, x_\bullet)$ vanishes if $x_* \rightarrow -\infty$ and/or $x_\bullet \rightarrow -\infty$ (recall that we assume $\bar{A}_*(x_\bullet) \xrightarrow{x_\bullet \rightarrow \pm\infty} 0$ and $\bar{B}_\bullet(x_*) \xrightarrow{x_* \rightarrow \pm\infty} 0$).

Also, from eq. (B.9) we see that

$$\bar{D}_* \bar{C}_\bullet^{(1)} = -\frac{1}{2} \bar{F}_{*\bullet}, \quad \bar{D}_\bullet \bar{C}_*^{(1)} = \frac{1}{2} \bar{F}_{*\bullet}\quad (\text{B.12})$$

and therefore

$$\mathcal{F}_{*\bullet} = \bar{F}_{*\bullet} + \bar{D}_* \bar{C}_\bullet^{(1)} - \bar{D}_\bullet \bar{C}_*^{(1)} + O(\bar{F}^2) = O(\bar{F}^2)\quad (\text{B.13})$$

so in the first order in \bar{F} the field strength of the solution of classical equation (B.5) vanishes.

In the second order the equations for the field C_μ take the form

$$\begin{aligned}2(\bar{P}_\bullet \bar{P}_*)^{ab} \bar{C}_\bullet^{(2)b} &= g \bar{D}_*^{aa'} (f^{a'bc} \bar{C}_*^{(1)b} \bar{C}_\bullet^{(1)c}) & \Rightarrow & \quad \bar{C}_\bullet^{(2)a} = -\frac{ig}{2} \left(\frac{1}{\bar{P}_* + i\epsilon} \right)^{aa'} f^{a'bc} \bar{C}_*^{(1)b} \bar{C}_\bullet^{(1)c} \\ 2(\bar{P}_* \bar{P}_\bullet)^{ab} \bar{C}_*^{(2)b} &= -g \bar{D}_*^{aa'} (f^{a'bc} \bar{C}_*^{(1)b} \bar{C}_\bullet^{(1)c}) & \Rightarrow & \quad \bar{C}_*^{(2)a} = \frac{ig}{2} \left(\frac{1}{\bar{P}_\bullet + i\epsilon} \right)^{aa'} f^{a'bc} \bar{C}_*^{(1)b} \bar{C}_\bullet^{(1)c}\end{aligned}\quad (\text{B.14})$$

where we used eq. (B.12) to reduce the r.h.s. Again, from the explicit form of the propagators (B.10) we get

$$\begin{aligned}\bar{C}_*^{(2)}(x) &= -\frac{ig}{s} \int_{-\infty}^{x_*} dz_* [x_*, z_*]^{A_\bullet} [\bar{C}_*^{(1)}(z_*, x_\bullet), \bar{C}_\bullet^{(1)}(z_*, x_\bullet)][z_*, x_*]^{A_\bullet} \\ \bar{C}_\bullet^{(2)}(x) &= \frac{ig}{s} \int_{-\infty}^{x_\bullet} dz_\bullet [x_\bullet, z_\bullet]^{A_*} [\bar{C}_*^{(1)}(x_*, z_\bullet), \bar{C}_\bullet^{(1)}(x_*, z_\bullet)][z_\bullet, x_\bullet]^{A_*}\end{aligned}\quad (\text{B.15})$$

from which it is clear that $\bar{C}_\mu^{(2)}$ satisfy boundary conditions (B.6) (recall that we already proved that $\bar{C}_\mu^{(1)}$ satisfy eq. (B.6)). Next, we use

$$\bar{D}_* \bar{C}_\bullet^{(2)a} = -\frac{g}{2} f^{abc} \bar{C}_*^{(1)b} \bar{C}_\bullet^{(1)c}, \quad \bar{D}_\bullet \bar{C}_*^{(2)a} = \frac{g}{2} f^{abc} \bar{C}_*^{(1)b} \bar{C}_\bullet^{(1)c} \quad (\text{B.16})$$

to prove that $\mathcal{F}_{*\bullet}$ vanishes in the second order:

$$\begin{aligned} \mathcal{F}_{*\bullet}^a &= F_{*\bullet}^a (\bar{A} + C^{(1)} + C^{(2)}) + O(\bar{F}^3) \\ &= \bar{F}_{*\bullet}^a + (\bar{D}_* \bar{C}_\bullet^{(1)} - \bar{D}_\bullet \bar{C}_*^{(1)})^a + (\bar{D}_* \bar{C}_\bullet^{(2)} - \bar{D}_\bullet \bar{C}_*^{(2)})^a + g f^{abc} \bar{C}_*^{(1)b} \bar{C}_\bullet^{(1)c} + O(\bar{F}^3) \\ &= O(\bar{F}^3) \end{aligned} \quad (\text{B.17})$$

In the third order we get

$$\begin{aligned} 2(\bar{P}_* \bar{P}_*)^{ab} \bar{C}_\bullet^{(3)b} &= g \bar{D}_\bullet^{aa'} f^{a'bc} (\bar{C}_*^{(1)b} \bar{C}_\bullet^{(2)c} + \bar{C}_*^{(2)b} \bar{C}_\bullet^{(1)c}) \\ 2(\bar{P}_* \bar{P}_\bullet)^{ab} \bar{C}_*^{(3)b} &= -g \bar{D}_*^{aa'} f^{a'bc} (\bar{C}_*^{(1)b} \bar{C}_\bullet^{(2)c} + \bar{C}_*^{(2)b} \bar{C}_\bullet^{(1)c}) \end{aligned} \quad (\text{B.18})$$

where again we used eqs. (B.12) and (B.16) to reduce the r.h.s. The solution is

$$\begin{aligned} \bar{C}_\bullet^{(3)a} &= -\frac{ig}{2} \left(\frac{1}{\bar{P}_* + i\epsilon} \right)^{aa'} f^{a'bc} (\bar{C}_*^{(1)b} \bar{C}_\bullet^{(2)c} + \bar{C}_*^{(2)b} \bar{C}_\bullet^{(1)c}) \\ \bar{C}_*^{(3)a} &= \frac{ig}{2} \left(\frac{1}{\bar{P}_\bullet + i\epsilon} \right)^{aa'} f^{a'bc} (\bar{C}_*^{(1)b} \bar{C}_\bullet^{(2)c} + \bar{C}_*^{(2)b} \bar{C}_\bullet^{(1)c}) \end{aligned} \quad (\text{B.19})$$

Again, from the explicit form of propagators (B.10) it is clear that $\bar{C}_\mu^{(3)}$ satisfy boundary conditions (B.2) if $\bar{C}_\mu^{(1)}$ and $\bar{C}_\mu^{(2)}$ do (which we already proved). Next, from

$$\bar{D}_* \bar{C}_\bullet^{(3)a} = -\frac{g}{2} f^{abc} (\bar{C}_*^{(1)b} \bar{C}_\bullet^{(2)c} + \bar{C}_*^{(2)b} \bar{C}_\bullet^{(1)c}), \quad \bar{D}_\bullet \bar{C}_*^{(3)a} = \frac{g}{2} f^{abc} (\bar{C}_*^{(1)b} \bar{C}_\bullet^{(2)c} + \bar{C}_*^{(2)b} \bar{C}_\bullet^{(1)c}) \quad (\text{B.20})$$

we see that $\mathcal{F}_{*\bullet}$ vanishes in the third order:

$$\begin{aligned} \mathcal{F}_{*\bullet}^a &= F_{*\bullet}^a (\bar{A} + \bar{C}^{(1)} + \bar{C}^{(2)} + \bar{C}^{(3)}) + O(\bar{F}^4) \\ &= \bar{F}_{*\bullet}^a + (\bar{D}_* \bar{C}_\bullet^{(1)} - \bar{D}_\bullet \bar{C}_*^{(1)})^a + (\bar{D}_* \bar{C}_\bullet^{(2)} - \bar{D}_\bullet \bar{C}_*^{(2)})^a + (\bar{D}_* \bar{C}_\bullet^{(3)} - \bar{D}_\bullet \bar{C}_*^{(3)})^a \\ &\quad + g f^{abc} (\bar{C}_*^{(1)b} \bar{C}_\bullet^{(1)c} + \bar{C}_*^{(1)b} \bar{C}_\bullet^{(2)c} + \bar{C}_*^{(2)b} \bar{C}_\bullet^{(1)c}) + O(\bar{F}^4) = O(\bar{F}^4) \end{aligned} \quad (\text{B.21})$$

Note also that eqs. (B.12), (B.16) and (B.20) illustrate self-consistency check for the background-field condition (B.4).

One can continue and prove by induction that $\mathcal{F}_{*\bullet}$ vanishes in an arbitrary order in $\bar{F}_{*\bullet}^n$ and therefore the field \mathcal{A}_μ is a pure gauge

$$\begin{aligned} \mathcal{A}_*(x_*, x_\bullet) &= \bar{A}_*(x_\bullet) + \bar{C}_*(x_*, x_\bullet) = \Omega(x_*, x_\bullet) i \partial_* \Omega^\dagger(x_*, x_\bullet) \\ \mathcal{A}_\bullet(x_*, x_\bullet) &= \bar{B}_\bullet(x_*) + \bar{C}_\bullet(x_*, x_\bullet) = \Omega(x_*, x_\bullet) i \partial_\bullet \Omega^\dagger(x_*, x_\bullet) \end{aligned} \quad (\text{B.22})$$

Now we shall demonstrate that the matrix Ω satisfies our requirement (3.18). Since $C_*(x_* \rightarrow -\infty, x_\bullet) = 0$ due to eq. (B.2), we get

$$\Omega(-\infty, x_\bullet) i \partial_* \Omega^\dagger(-\infty, x_\bullet) = \bar{A}_*(x_\bullet) \quad \Rightarrow \quad \Omega(-\infty, x_\bullet) = [x_\bullet, -\infty_\bullet]^{\bar{A}_*} \quad (\text{B.23})$$

Similarly,

$$\Omega(x_*, -\infty) i\partial_\bullet \Omega^\dagger(x_*, -\infty) = \bar{B}_\bullet(x_*) \Rightarrow \Omega(x_*, -\infty) = [x_*, -\infty_*]^{\bar{B}_\bullet} \quad (\text{B.24})$$

One can also construct the expansion of matrix Ω in powers of \bar{A}_* and \bar{B}_\bullet . For example, up to the fifth power of the \bar{A}_μ fields

$$\begin{aligned} \Omega(x_*, x_\bullet) = & \frac{1}{2} \{ [x_*, -\infty_*]^{\bar{B}_\bullet}, [x_\bullet, -\infty_\bullet]^{\bar{A}_*} \} - \frac{1}{4} \left([[x_\bullet, -\infty_\bullet]^{\bar{A}_*}, [x_*, -\infty_*]^{\bar{B}_\bullet}] \right)^2 \\ & - \frac{4g^4}{s^4} \int_{-\infty}^{x_*} dx'_* \int_{-\infty}^{x'_*} dx''_* \int_{-\infty}^{x_\bullet} dx'_\bullet \int_{-\infty}^{x'_\bullet} dx''_\bullet [[\bar{A}_\bullet(x'_*), \bar{A}_*(x'_\bullet)], [\bar{A}_\bullet(x''_*), \bar{A}_*(x''_\bullet)]] \end{aligned} \quad (\text{B.25})$$

Now, for each x_\perp we solve auxiliary 2-dimensional classical problem (B.1) and find $\Omega(x_*, x_\bullet, x_\perp)$ satisfying the requirement (3.18).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License (CC-BY 4.0), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J. Collins, *Foundations of perturbative QCD*, Cambridge University Press, Cambridge U.K., (2013) [[INSPIRE](#)].
- [2] J.C. Collins and D.E. Soper, *Parton distribution and decay functions*, *Nucl. Phys. B* **194** (1982) 445 [[INSPIRE](#)].
- [3] J.C. Collins, D.E. Soper and G.F. Sterman, *Transverse momentum distribution in Drell-Yan pair and W and Z boson production*, *Nucl. Phys. B* **250** (1985) 199 [[INSPIRE](#)].
- [4] X.-D. Ji, J.-P. Ma and F. Yuan, *QCD factorization for semi-inclusive deep-inelastic scattering at low transverse momentum*, *Phys. Rev. D* **71** (2005) 034005 [[hep-ph/0404183](#)] [[INSPIRE](#)].
- [5] M.G. Echevarria, A. Idilbi and I. Scimemi, *Factorization theorem for Drell-Yan at low q_T and transverse momentum distributions on-the-light-cone*, *JHEP* **07** (2012) 002 [[arXiv:1111.4996](#)] [[INSPIRE](#)].
- [6] J. Collins and T. Rogers, *Understanding the large-distance behavior of transverse-momentum-dependent parton densities and the Collins-Soper evolution kernel*, *Phys. Rev. D* **91** (2015) 074020 [[arXiv:1412.3820](#)] [[INSPIRE](#)].
- [7] I. Balitsky, *Factorization and high-energy effective action*, *Phys. Rev. D* **60** (1999) 014020 [[hep-ph/9812311](#)] [[INSPIRE](#)].
- [8] I. Balitsky, *High-energy effective action from scattering of QCD shock waves*, *Phys. Rev. D* **72** (2005) 074027 [[hep-ph/0507237](#)] [[INSPIRE](#)].
- [9] Y. Hatta, E. Iancu, L. McLerran, A. Stasto and D.N. Triantafyllopoulos, *Effective Hamiltonian for QCD evolution at high energy*, *Nucl. Phys. A* **764** (2006) 423 [[hep-ph/0504182](#)] [[INSPIRE](#)].
- [10] Y. Hatta, E. Iancu, L. McLerran and A. Stasto, *Color dipoles from Bremsstrahlung in QCD evolution at high energy*, *Nucl. Phys. A* **762** (2005) 272 [[hep-ph/0505235](#)] [[INSPIRE](#)].

- [11] T. Altinoluk, A. Kovner, M. Lublinsky and J. Peressutti, *QCD reggeon field theory for every day: pomeron loops included*, *JHEP* **03** (2009) 109 [[arXiv:0901.2559](#)] [[INSPIRE](#)].
- [12] B.A. Kniehl and M. Spira, *Low-energy theorems in Higgs physics*, *Z. Phys. C* **69** (1995) 77 [[hep-ph/9505225](#)] [[INSPIRE](#)].
- [13] M.A. Shifman, A.I. Vainshtein, M.B. Voloshin and V.I. Zakharov, *Low-energy theorems for Higgs boson couplings to photons*, *Sov. J. Nucl. Phys.* **30** (1979) 711 [*Yad. Fiz.* **30** (1979) 1368] [[INSPIRE](#)].
- [14] J.F. Gunion, H.E. Haber, G.L. Kane and S. Dawson, *The Higgs hunter's guide*, *Front. Phys.* **80** (2000) 1 [[INSPIRE](#)].
- [15] I.I. Balitsky and V.M. Braun, *Nonlocal operator expansion for structure functions of e^+e^- annihilation*, *Phys. Lett. B* **222** (1989) 123 [[INSPIRE](#)].
- [16] I.I. Balitsky and V.M. Braun, *The nonlocal operator expansion for inclusive particle production in e^+e^- annihilation*, *Nucl. Phys. B* **361** (1991) 93 [[INSPIRE](#)].
- [17] I.I. Balitsky and V.M. Braun, *Valleys in Minkowski space and instanton induced cross-sections*, *Nucl. Phys. B* **380** (1992) 51 [[INSPIRE](#)].
- [18] I. Balitsky and A. Tarasov, *Rapidity evolution of gluon TMD from low to moderate x* , *JHEP* **10** (2015) 017 [[arXiv:1505.02151](#)] [[INSPIRE](#)].
- [19] I. Balitsky and A. Tarasov, *Gluon TMD in particle production from low to moderate x* , *JHEP* **06** (2016) 164 [[arXiv:1603.06548](#)] [[INSPIRE](#)].
- [20] I. Balitsky, *Scattering of shock waves in QCD*, *Phys. Rev. D* **70** (2004) 114030 [[hep-ph/0409314](#)] [[INSPIRE](#)].
- [21] F. Gelis and R. Venugopalan, *Large mass $q\bar{q}$ production from the color glass condensate*, *Phys. Rev. D* **69** (2004) 014019 [[hep-ph/0310090](#)] [[INSPIRE](#)].
- [22] F. Gelis and R. Venugopalan, *Particle production in field theories coupled to strong external sources*, *Nucl. Phys. A* **776** (2006) 135 [[hep-ph/0601209](#)] [[INSPIRE](#)].
- [23] F. Gelis, T. Lappi and R. Venugopalan, *High energy scattering in quantum chromodynamics*, *Int. J. Mod. Phys. E* **16** (2007) 2595 [[arXiv:0708.0047](#)] [[INSPIRE](#)].
- [24] I.Z. Rothstein and I.W. Stewart, *An effective field theory for forward scattering and factorization violation*, *JHEP* **08** (2016) 025 [[arXiv:1601.04695](#)] [[INSPIRE](#)].
- [25] L.M. Kaplan and M. Resnikoff, *Matrix products and the explicit 3, 6, 9, and 12- j coefficients of the regular representation of $SU(n)$* , *J. Math. Phys.* **8** (1967) 2194.
- [26] A.J. MacFarlane, A. Sudbery and P.H. Weisz, *On Gell-Mann's λ -matrices, d - and f -tensors, octets and parametrizations of $SU(3)$* , *Commun. Math. Phys.* **11** (1968) 77 [[INSPIRE](#)].
- [27] P.J. Mulders and J. Rodrigues, *Transverse momentum dependence in gluon distribution and fragmentation functions*, *Phys. Rev. D* **63** (2001) 094021 [[hep-ph/0009343](#)] [[INSPIRE](#)].
- [28] I. Balitsky, *Operator expansion for high-energy scattering*, *Nucl. Phys. B* **463** (1996) 99 [[hep-ph/9509348](#)] [[INSPIRE](#)].