

Old Dominion University
ODU Digital Commons

Biological Sciences Faculty Publications

Biological Sciences

2014

Ehrlichia and Spotted Fever Group Rickettsiae Surveillance in *Amblyomma americanum* in Virginia Through Use of a Novel Six-Plex Real-Time PCR Assay

David N. Gaines

Darwin J. Operario


Suzanne Stroup

Ellen Stromdahl

Chelsea Wright
Old Dominion University

See next page for additional authors

Follow this and additional works at: https://digitalcommons.odu.edu/biology_fac_pubs

 Part of the [Environmental Public Health Commons](#), and the [Immunology and Infectious Disease Commons](#)

Repository Citation

Gaines, David N.; Operario, Darwin J.; Stroup, Suzanne; Stromdahl, Ellen; Wright, Chelsea; Gaff, Holly; Broyles, James; Smith, Joshua; Norris, Douglas E.; Henning, Tyler; Lucas, Agape; and Houpt, Eric, "Ehrlichia and Spotted Fever Group Rickettsiae Surveillance in *Amblyomma americanum* in Virginia Through Use of a Novel Six-Plex Real-Time PCR Assay" (2014). *Biological Sciences Faculty Publications*. 299.
https://digitalcommons.odu.edu/biology_fac_pubs/299

Original Publication Citation

Gaines, D. N., Operario, D. J., Stroup, S., Stromdahl, E., Wright, C., Gaff, H., . . . Houpt, E. (2014). *Ehrlichia* and spotted fever group rickettsiae surveillance in *Amblyomma americanum* in Virginia through use of a novel six-plex real-time PCR assay. *Vector-Borne and Zoonotic Diseases*, 14(5), 307-316. doi:10.1089/vbz.2013.1509

Authors

David N. Gaines, Darwin J. Operario, Suzanne Stroup, Ellen Stromdahl, Chelsea Wright, Holly Gaff, James Broyhill, Joshua Smith, Douglas E. Norris, Tyler Henning, Agape Lucas, and Eric Houpt

Ehrlichia and Spotted Fever Group Rickettsiae Surveillance in *Amblyomma americanum* in Virginia Through Use of a Novel Six-Plex Real-Time PCR Assay

David N. Gaines,^{1,*} Darwin J. Operario,^{2,*} Suzanne Stroup,² Ellen Stromdahl,³ Chelsea Wright,⁴ Holly Gaff,^{4,5} James Broyhill,¹ Joshua Smith,⁶ Douglas E. Norris,⁷ Tyler Henning,⁷ Agape Lucas,⁸ and Eric Houpt²

Abstract

The population of the lone star tick *Amblyomma americanum* has expanded in North America over the last several decades. It is known to be an aggressive and nondiscriminatory biter and is by far the most common human-biting tick encountered in Virginia. Few studies of human pathogen prevalence in ticks have been conducted in our state since the mid-twentieth century. We developed a six-plex real-time PCR assay to detect three *Ehrlichia* species (*E. chaffeensis*, *E. ewingii*, and Panola Mountain *Ehrlichia*) and three spotted fever group Rickettsiae (SFGR; *R. amblyommii*, *R. parkeri*, and *R. rickettsii*) and used it to test *A. americanum* from around the state. Our studies revealed a presence of all three *Ehrlichia* species (0–24.5%) and a high prevalence (50–80%) of *R. amblyommii*, a presumptively nonpathogenic SFGR, in all regions surveyed. *R. parkeri*, previously only detected in Virginia's *Amblyomma maculatum* ticks, was found in *A. americanum* in several surveyed areas within two regions having established *A. maculatum* populations. *R. rickettsii* was not found in any sample tested. Our study provides the first state-wide screening of *A. americanum* ticks in recent history and indicates that human exposure to *R. amblyommii* and to Ehrlichiae may be common. The high prevalence of *R. amblyommii*, serological cross-reactivity of all SFGR members, and the apparent rarity of *R. rickettsii* in human biting ticks across the eastern United States suggest that clinical cases of tick-borne disease, including ehrlichiosis, may be commonly misdiagnosed as Rocky Mountain spotted fever, and that suspicion of other SFGR as well as *Ehrlichia* should be increased. These data may be of relevance to other regions where *A. americanum* is prevalent.

Key Words: *Rickettsia*—*Ehrlichia*—Ticks—Real-time RT-PCR—Vector borne.

Introduction

VIRGINIA IS A STATE with diverse and changing tick-borne disease epidemiology that includes human ehrlichiosis, Rocky Mountain spotted fever (RMSF), Tidewater spotted fever (an illness caused by *Rickettsia parkeri*; Wright et al. 2011), Lyme disease, and anaplasmosis. The lone star tick *Amblyomma americanum*, the most common tick to bite

people in the southeastern United States, is increasingly a cause of disease transmission (Paddock and Yabsley 2007, Apperson et al. 2008, Smith et al. 2010, Stromdahl and Hickling 2012), and is likely among the most important arthropod vectors of disease to humans in Virginia.

A. americanum potentially transmits ehrlichiosis and rickettsiosis. The case rates of these diseases vary across Virginia (Fig. 1). Human ehrlichiosis attributed to *Ehrlichia chaffeensis*

¹Virginia Department of Health, Richmond, Virginia.

²Division of Infectious Diseases and International Health, University of Virginia Health System, Charlottesville, Virginia.

³US Army Public Health Command, Aberdeen Proving Ground, Maryland.

⁴Department of Biological Sciences, Old Dominion University, Norfolk, Virginia.

⁵School of Mathematics, Statistics and Computer Science, University of KwaZulu-Natal, Durban, South Africa.

⁶Fairfax County Health Department, Fairfax, Virginia.

⁷Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland.

⁸North Carolina Agricultural and Technical State University, Greensboro, North Carolina.

*These authors contributed equally to this work.




FIG. 1. Mean number of human cases of ehrlichiosis or spotted fever rickettsiosis in Virginia counties and cities averaged over a 3-year period from 2009–2011.

may cause a mild to fatal illnesses, with a hospitalization rate of 49% and a fatality rate of 1.9% (Dahlgren et al. 2011). Ehrlichiosis caused by *E. ewingii* is less commonly reported and tends to be associated with previously immune-suppressed patients (Paddock and Yabsley 2007). In 2006, a new *Ehrlichia* species was discovered in *A. americanum* in Georgia (Loftis et al. 2006). This new species, known as Panola Mountain *Ehrlichia* (PME), was subsequently associated with human illness (Reeves et al. 2008) and has been detected in *A. americanum* collected from 10 different states, including Virginia (Loftis et al. 2008, Yabsley et al. 2008).

During the past decade, a number of state health departments in the southeastern United States, including the Virginia Department of Health (VDH), have seen large increases in diagnosed and reported cases of RMSF despite a decline in the mortality rate (*i.e.*, a rise in national incidence from 1.7 to 7 cases per million but decrease in mortality from 2.2% to 0.3%) (Chapman et al. 2006, Openshaw et al. 2010, Virginia Department of Health 2012). The agent of RMSF is *Rickettsia rickettsii*, and *Dermacentor variabilis*, the American dog tick, is thought to be the primary vector. However, *R. rickettsii* has rarely been found in the thousands of *D. variabilis* tested over the past two decades (Ammerman et al. 2004, Dergousoff et al. 2009, Moncayo et al. 2010, Stromdahl et al. 2011). This leaves open the possibility that *A. americanum* may play a role as a vector (Stromdahl et al. 2011).

A. americanum is also recognized as an important host of *Rickettsia amblyommii*, a spotted fever group *Rickettsia* (SFGR) with no apparent pathogenicity for humans. Mixson et al. (2006) found an *R. amblyommii* infection rate in adult *A. americanum* of 41.2% from nine states across the eastern United States. Several recent tick surveys in Virginia also found *R. amblyommii* to be common in *A. americanum*, with rates of 26–70% (J.S. and H.G., unpublished data) (Jiang et al. 2010).

Another SFGR potentially transmitted by *A. americanum* is *R. parkeri*. Although identified over 50 years ago (Lackman et al. 1949, Lackman et al. 1965), *R. parkeri* was only recognized as a human pathogen in Virginia in 2002 (Paddock et al. 2004). Since that time, human illness caused by *R. parkeri* has been identified across the southeastern United States (Paddock et al. 2008). The primary vector of *R. parkeri* is *Amblyomma maculatum*, the Gulf Coast tick, with high

infection rates (41–43%) in ticks collected in Virginia (Fornadel et al. 2011, Wright et al. 2011). However, *R. parkeri* has also been detected in adult *A. americanum* from Tennessee and Georgia (Cohen et al. 2009). *R. parkeri* can infect *A. americanum*, be transmitted transstadially and transovarially, and subsequently be transmitted to guinea pigs by feeding *A. americanum* (Goddard 2003).

Few studies of tick species distribution and human pathogen prevalence have been conducted in Virginia since the mid-twentieth century, when research focused on *D. variabilis* and *R. rickettsii* (Sonenshine et al. 1966) and more recent surveys have focused on ticks other than *A. americanum* (Fornadel et al. 2011, Nadolny et al. 2011, Wright et al. 2011). In the intervening decades, deer populations have increased and have led to the subsequent increase and range expansion of the ticks they host, including *A. americanum* (Paddock and Yabsley 2007, Virginia Department of Game and Inland Fisheries 2007). To investigate the epidemiological role of *A. americanum*, we first developed a multiplex real-time PCR assay for Ehrlichiae (*E. chaffeensis*, *E. ewingii*, and PME) and three species of Rickettsiae (*R. amblyommii*, *R. parkeri*, and *R. rickettsii*) that could be applied to tick samples. We then collected tick samples from across the state, tested a total of 2545 nymph and adult *A. americanum*, and compared these findings with state human disease statistics.

Materials and Methods

Study areas and sample collection

The ticks collected for this study came from the VDH, the Fairfax County Department of Health (FCDH), the Old Dominion University (ODU) Department of Biology, and the US Army Public Health Command (USAPHC). Ticks collected by the VDH were obtained between June 1 and August 15, 2012, by cloth drags, primarily from sample sites at selected municipal, county, or state park lands within each of five survey regions of Virginia, including the Richmond City Area, an area in Southside Virginia, the Charlottesville Area, the Lynchburg Area, and the Fredericksburg Area (Fig. 2). Tick samples were prepared as single adult samples ($n = 112$) or as pooled nymphs ($n = 1206$ nymphs in 388 pools; Table 1). Small pools (two ticks/pool) were created to permit the estimation of infection rates for agents that were likely to be




FIG. 2. *Amblyomma americanum* collection sites across Virginia. Richmond City area: 1, Henrico County; 2, Goochland Co.; 3, Chesterfield Co. Charlottesville area: Sites 4–6, Albemarle Co. Southside Virginia area: 7, Brunswick Co.; 8, Mecklenberg Co., and Site 28, Fort Pickett, Nottoway County. Lynchburg area: Sites 9–14, Lynchburg City. Fredericksburg area: 15, Fredericksburg; 16, Stafford Co.; 17, Spotsylvania Co.; 18, King George Co.; Site 27, Fort A.P. Hill, Caroline Co.; Fairfax County Sites 19–23 and Site 28, Fort Belvoir (Fairfax Co. and adjacent counties/cities). Hampton Roads Area: 24, Portsmouth City; 25, Chesapeake City.

common (*i.e.*, >50% infection rates). Large pools contained up to eight ticks and served to increase the number of ticks tested. Ticks collected by VDH were processed and tested at the University of Virginia (UVA).

Ticks collected by the FCDH were obtained from five sites within Fairfax County (Fig. 2) by means of cloth drags, flags, or CO₂-baited traps. Ticks were initially submitted to the Johns Hopkins Bloomberg School of Public Health (JHSPH) for DNA extraction, and, subsequently, VDH personnel made a random sampling of tick samples collected between June 4 and August 15, 2012, to be sent to UVA for testing. Nymph extracts were pooled (two to nine extracts per pool), while adults were tested as single samples. A total of 49 single adult ticks and 163 nymphs (in 50 pools) were tested (Table 1).

Ticks from ODU were collected by use of drag cloths from March 1 to July 18, 2012, and came from one site in the City

of Chesapeake and one in the City of Portsmouth (Fig. 2). Tick specimens were initially extracted at ODU, and included 45 single adult tick extracts and 12 single nymph extracts from 2012 collections (Table 1). ODU also provided tick extracts (24 adults and 18 nymphs) collected from these two sites in 2010 and 2011.

Ticks from the USAPHC were removed from humans at three army installations: Fort Pickett in Southside Virginia, Fort A.P. Hill in the Fredericksburg Area, and Fort Belvoir in Fairfax County (Fig. 2), and submitted to the USAPHC Tick-borne Disease Laboratory, Aberdeen Proving Ground, Maryland for *Ehrlichia* testing. Ticks from Fort Pickett and Fort A.P. Hill were removed from soldiers training on those installations, whereas ticks from Fort Belvoir were collected from military personnel, civilian employees, and dependents that worked at that installation but lived in the surrounding areas. USAPHC had “superpooled” extracts from the Virginia ticks with tick extracts from installations in other states. USAPHC also maintained extracts of each individual tick so they could be individually retested when a superpool tested positive. Among these pools, a total of 370 adult ticks (351 pools) and 588 nymphs (471 pools) from Virginia, submitted to USAPHC from March 21, 2012 to September 13, 2012, were tested. Because these superpools were not originally tested for SFGR, it was not possible to back trace the original geographic origin of contributing samples whose superpool later tested positive for a SFGR.

DNA extraction

The Qiagen DNeasy Blood and Tissue Kit (Qiagen Inc., Valencia, CA) was used for ticks processed at UVA or at ODU, with some modifications to the manufacturer’s protocol.

TABLE 1. TICK DEMOGRAPHICS

	Total samples tested	Nymphs		
		Adults	Pools	Total
Richmond Area	103	40	63	192
Charlottesville Area	103	19	84	261
Southside Virginia	102	41	61	194
Lynchburg Area	100	9	91	284
Fredricksburg Area	92	3	89	275
Hampton Roads Area	57	45	12	12
Fairfax Area	99	49	50	163
US Army Public Health Command	822	370 (351 pools)	471	588
Overall	1276	576	871	1969

Briefly, adult ticks (extracted individually) were bisected longitudinally with a razor or surgical blade. Nymphs were extracted individually or in pools. Ticks were placed into 180 μ L of Buffer ALT and homogenized with glass beads. Samples were centrifuged at 18,000 relative centrifugal force (rcf) for 30 s at room temperature and placed in a dry block. For some samples, an additional 180 μ L of Buffer ALT was added followed by the addition of 40 μ L of Proteinase K Solution and 6 μ L of carrier RNA (1 μ g/ μ L; Qiagen). Samples were placed on the side in a shaker incubator at 56°C for a 1- to 3-h digestion. DNA was isolated using the spin-column protocol for animal tissues, eluted, and stored at -20°C prior to testing by PCR.

At JHSPH, DNA was extracted from individual ticks using a MasterPure DNA Purification Kit (Epicentre Biotechnologies, Madison, WI), with a modified procedure. Briefly, a tick was placed in Tissue and Cell Lysis Solution, disrupted with a 5-mm stainless steel bead using a TissueLyser II (Qiagen) for 3 min, then centrifuged for 1 min at 16,100 rcf at room temperature in a microcentrifuge. To each sample, 250 μ L of Tissue and Cell Lysis Solution containing 1 μ L of Proteinase K (50 g/ μ L) was added. Beads were removed magnetically, and each sample was incubated at 65°C for 60 min followed by a 5-min incubation on ice. DNA was isolated from debris by addition of 150 μ L of MPC Protein Precipitation Reagent and centrifugation at 4°C for 10 min at 17,000 rcf. The DNA was then pelleted using isopropanol, air dried, and then re-suspended in 30 μ L of molecular-grade water.

At USAPHC, genomic tick DNA was extracted by using a Zymo Genomic DNA II Kit™ (Zymo Research Corporation, Orange, California) according to manufacturers' instructions (Stromdahl et al. 2011).

Detection of bacterial DNA by real-time PCR

All multiplex testing for this study was performed at UVA using the Applied Biosystems ViiA 7 Real-Time PCR system (Applied Biosystems, Foster City, CA). Reactions were performed with Bio-Rad iQ Multiplex Powermix (Bio-Rad, Hercules, CA) in a 25- μ L reaction volume with 5 μ L of DNA extract. Primers and TaqMan probes (Eurofins MWG Operon, Huntsville, AL) were used at concentrations listed in Table 2. Probes for *R. rickettsii* and *R. amblyomii* were adapted from molecular beacon designs. All reactions were run using the following program: 95°C for 3 min, followed by 40 cycles of 95°C for 15 s, and 61°C for 1 min. Each run included a "No template control" (nuclease-free water) and positive controls for each of the six bacterial targets. Post-reaction baselines and thresholds were adjusted manually as needed.

Sequencing of *R. parkeri* samples

Samples positive for *R. parkeri* in the multiplex assay were retested by singleplex PCR containing the same *R. parkeri* primers and probe used in the multiplex. Amplicons from the singleplex were cloned into the PCR 2.1 TOPO vector using the TOPO TA cloning kit (Invitrogen, Carlsbad, CA), and resulting plasmids were sequenced using GeneWiz (South Plainfield, NJ) or Eurofins MWG Operon DNA sequencing services. Sequence data were compared to the corresponding *ompB* gene sequence of *R. parkeri* strain At24 (Genbank EF102239).

TABLE 2. PRIMERS AND PROBES

Bacterial target	Gene target	Oligo name, type	Sequence, 5' → 3'	Final conc., μ M	Amplicon size	Reference
<i>R. rickettsii</i>	<i>OmpB</i>	RR1370F, F primer	ATAACCCAAAGACTCAAACCTTTGGTA	0.4	124 bp	Smith et al. 2010
		RR1494R, R primer	GCAGTGTACCGGGATITGCT	0.4		
		RR1425B, Probe	FAM-TTAAAGTTCCTAATGCTATAACCCCTTACC-BHQ1	0.2		
<i>R. amblyomii</i>	<i>OmpB</i>	Ra477F, F primer	GGTGTGCGGGTCTACATTAG	0.2	141 bp	Smith et al. 2010
		Ra618R, R primer	CTGAAACTTGAATAAATCCATTAGTAACAT	0.2		
		Ra532P, Probe	HEX-TCCTCTTACACTGGACAGAAATGCT-BHQ2	0.2		
<i>R. parkeri</i>	<i>OmpB</i>	Rpa129F, F primer	CAAAATGTTGCAGTTCCTCTAAA	0.2	96 bp	Jiang et al. 2010
		Rpa224R, R primer	AAAAACAAACCGTTAAAACACTACCG	0.2		
		RpaTM-P-TAMRA, Probe	TAMRA-AATTAATACCTTATGARCASCAGCAG-BHQ2	0.2		
<i>E. chaffeensis</i> / <i>E. ewingii</i>	16S rRNA	ECH16S-17, F primer	GCGGCAAGCCTAACACATG	0.4	81 bp	Loftis et al. 2003
		ECH16S-97, R primer	CCCGTGTGCCACTAACAAATTATT	0.4		
		ECH16S-38, Probe	Texas Red-AGTCGAACGGACAATTGCTTATAACCTTTTGGT-BHQ2	0.2		
<i>E. ewingii</i>	16S rRNA	EEW16S-97, R primer	CCCGTGTGCCACTAACAACTATC	0.2	83 bp	This study
		EEW16S-P, Probe	Cy5.5-AGTCGAACGAAACAATTCCTAAATAGTCTCTGAC-BHQ2	0.4		
		Ehr3CS-214F, F primer	TGTCATTTCCACAGCAATTCATC	0.4		
Panola Mountain <i>Ehrlichia</i>	<i>gltA</i>	PME344R, R primer	ATTAGCGCAATCATACTTGCAA	0.4	145 bp	Loftis et al. 2008
		PME266P, Probe	Cy5-TGCCCTTAGCTGCACATTATTGTGAT-BHQ2	0.2		

Statistics and calculations

Minimum infection rate (MIR) for pooled adults or nymphs is the number of positive pools per total number of pooled nymphs and assumes that each positive pool of nymphs contains only one infected tick. Maximum likelihood estimation (MLE) was calculated using a the Poole-InfRate, ver. 4.0 add-on to Microsoft Excel (Biggerstaff 2009). MLE calculations are based on the number of pools, pool sizes (number of ticks per pool), and number of positive pools. This calculation provides a 95% confidence interval (CI) on the estimate (Biggerstaff 2008). Adult detection rates by region were compared using the Fisher exact test. All *p* values were two-tailed, and *p* < 0.05 was considered significant.

Results

Test results from ticks collected from the environment by use of dragging or trapping are presented in Tables 3A–C. Test results for ticks collected from personnel on Army Installations are presented in Tables 4A and B.

Detection of Ehrlichia species

We observed variable *E. chaffeensis* infection rates in ticks across regions and sites with ranges of 0–24.5% (average 7.3%) in adult ticks and 0–13.4% (average 3.4%) in nymphs. The Charlottesville Area (Table 3A and B) was unique in that *E. chaffeensis* infection was absent in both nymph and adult ticks at each of the three sites sampled. *E. chaffeensis* was also absent in the two sites sampled in Southside Virginia, but was detected in a relatively high proportion of ticks from Fort Pickett within that same region. Adult ticks from Fairfax County had a statistically higher rate of *E. chaffeensis* infection than was seen from any other region (*p* < 0.05, Fisher exact test) (Table 3A and B), but a surprisingly low *E. chaffeensis* infection rate (0.4% in adult ticks and 0.5% for nymphs) was observed in the ticks collected from Fort Belvoir personnel who resided in Fairfax County and surrounding areas. The geographic distribution of *E. ewingii* was more uniform, occurring in adult ticks in all regions except for the Lynchburg and Fredericksburg areas, where the numbers of adult ticks tested was relatively low. The *E. ewingii* detection rate by

TABLE 3A. VIRGINIA *AMBLIOMMA AMERICANUM* ADULTS BY SURVEY REGIONS, COLLECTION SITES AND DATES, AND TEST RESULTS

Tick collection regions, sites and site numbers	Collection dates	Ticks tested	Ehrlichia chaffeensis % Positive	Ehrlichia ewingii % Positive	Panola Mountain Ehrlichia % Positive	Rickettsia amblyommii % Positive	Rickettsia parkeri % Positive
Richmond City		40	5.0	7.5	2.5	72.5	0
Henrico Co. (1)	7/4–6/2012	33	3.0	9.1	3.0	78.8	0
Goochland Co. (2)	7/11–23/2012	6	16.7	0	0	50.0	0
Chesterfield Co. (3)	7/23/12	1	0	0	0	0	0
Charlottesville		19	0	5.3	0	68.4	0
Albemarle Co. (4)	6/10 to 7/20/2012	6	0	0	0	33.3	0
Albemarle Co. (5)	7/19/2012	3	0	0	0	100	0
Albemarle Co. (6)	7/19/2012	10	0	10.0	0	80.0	0
Southside Virginia		41	0	9.8	0	75.6	0
Brunswick Co. (7)	7/25/2012	38	0	10.5	0	79.0	0
Mecklenberg Co. (8)	7/25/2012	3	0	0	0	33.3	0
Lynchburg		9	0	0	0	55.6	0
Lynchburg (9)	7/31/2012	0	—	—	—	—	—
Lynchburg (10)	7/31/2012	3	0	0	0	100	0
Lynchburg (11)	7/31/2012	1	0	0	0	0	0
Lynchburg (12)	7/31/2012	0	—	—	—	—	—
Lynchburg (13)	8/3/2012	3	0	0	0	66.7	0
Lynchburg (14)	8/3/2012	2	0	0	0	0	0
Fredericksburg		3	0	0	0	100	0
Fredericksburg (15)	8/7/2012	1	0	0	0	100	0
Stafford Co. (16)	8/7/2012	2	0	0	0	100	0
Spotsylvania Co. (17)	8/15/2012	0	—	—	—	—	—
King George Co. (18)	8/15/2012	0	—	—	—	—	—
Fairfax County		49	24.5*	14.3	6.1	81.6	2.04
Fairfax Co. (19)	6/15–28/2012	3	33.3	0	0	100	0
Fairfax Co. (20)	6/4 to 7/25/2012	2	100	0	50.0	100	0
Fairfax Co. (21)	6/7 to 7/11/2012	19	26.3	26.3	0	57.9	0
Fairfax Co. (22)	6/4 to 7/25/2012	11	0	18.2	9.1	100	9.1
Fairfax Co. (23)	6/4 to 7/30/2012	14	28.6	0	7.1	92.9	0
Hampton Roads		45	2.2	2.2	0	64.4	2.2
Portsmouth (24)	6/1 to 7/18/2012	28	3.6	3.60	0	71.4	0
Chesapeake (25)	3/23 to 7/18/2012	17	0	0	0	52.9	5.9
Total Across All Sites		206	7.3	7.8	1.9	72.8	1.0

**p* < 0.05, for detection rate of *E. chaffeensis* in Fairfax County area vs. Richmond, Charlottesville, Southside, and Hampton Roads areas.

TABLE 3B. VIRGINIA AMBLYOMMA AMERICANUM NYMPHS BY SURVEY REGIONS, COLLECTION SITES, COLLECTION DATES, AND TEST RESULTS FOR EHRLICHIA SPECIES

Tick collection regions, sites, and site numbers	Collection dates	Nymphs tested	Ehrlichia chaffeensis			Ehrlichia ewingii			Panola Mountain Ehrlichia		
			Pooled nymphs			Pooled nymphs			Pooled nymphs		
			MIR ^a % pos.	MLE ^b (95% CI)	% pos.	MIR ^a % pos.	MLE ^b (95% CI)	% pos.	MIR ^a % pos.	MLE ^b (95% CI)	% pos.
Richmond City		192	3.7	3.8	0.5	0.5	2.1	2.2	2.1	2.2	(0.7-5.1)
Henrico Co. (1)	7/4-6/2012	82	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	(0.1-6.0)
Goochland Co. (2)	7/11-23/2012	54	1.9	1.8	0	0	1.9	1.8	1.9	1.8	(0.1-8.4)
Chesterfield Co. (3)	7/23/12	56	8.9	10.5	0	0	3.6	3.7	3.6	3.7	(0.7-11.9)
Charlottesville		261	0	0	0.4	0.4	0.8	0.8	0.8	0.8	(0.1-2.5)
Albemarle Co. (4)	6/10 to 7/20/2012	14	0	0	0	0	0	0	0	0	—
Albemarle Co. (5)	7/19/2012	101	0	0	0	0	2.0	2.0	2.0	2.0	(0.4-6.2)
Albemarle Co. (6)	7/19/2012	146	0	0	0.7	0.7	0	0	0	0	—
Southside Virginia		194	0	0	0.5	0.5	1.0	1.0	1.0	1.0	(0.2-3.4)
Brunswick Co. (7)	7/25/2012	116	0	0	0.9	0.9	0.9	0.9	0.9	0.9	(0.1-4.2)
Mecklenberg Co. (8)	7/25/2012	78	0	0	0	0	1.3	1.3	1.3	1.3	(0.1-5.9)
Lynchburg		284	4.9	5.3	4.6	5.1	2.8	3.0	2.8	3.0	(1.4-5.6)
Lynchburg (9)	7/31/2012	14	7.1	8.3	0	0	0	0	0	0	—
Lynchburg (10)	7/31/2012	82	0	0	0	0	2.4	2.6	2.4	2.6	(0.5-8.6)
Lynchburg (11)	7/31/2012	12	0	0	0	0	25.0	27.8	25.0	27.8	(8.1-60.6)
Lynchburg (12)	7/31/2012	80	1.0	12.0	11.3	15.7	2.5	2.6	2.5	2.6	(0.5-8.2)
Lynchburg (13)	8/3/2012	48	8.3	8.3	2.1	2.2	2.1	2.2	2.1	2.2	(0.1-10.4)
Lynchburg (14)	8/3/2012	48	2.1	2.0	6.3	6.7	0	0	0	0	—
Fredericksburg		275	0.7	0.7	0.7	0.7	1.5	1.5	1.5	1.5	(0.5-3.5)
Fredericksburg (15)	8/7/2012	14	0	0	0	0	0	0	0	0	—
Stafford Co. (16)	8/7/2012	20	0	0	0	0	0	0	0	0	—
Spotsylvania Co. (17)	8/15/2012	96	2.1	2.1	0	0	0	0	0	0	—
King George Co. (18)	8/15/2012	145	0	0	1.4	1.4	2.7	2.9	2.7	2.9	(0.9-6.8)
Fairfax County		163	11.6	13.4	4.3	4.5	1.2	1.3	1.2	1.3	(0.2-4.0)
Fairfax Co. (19)	6/15-28/2012	22	4.5	5.1	9.1	8.8	0	0	0	0	—
Fairfax Co. (20)	6/4 to 7/25/2012	28	10.7	11.4	0	0	3.6	3.8	3.6	3.8	(0.2-18.4)
Fairfax Co. (21)	6/7 to 7/11/2012	41	7.3	7.2	2.4	2.5	2.4	2.3	2.4	2.3	(0.1-10.7)
Fairfax Co. (22)	6/4 to 7/25/2012	38	13.2	14.0	5.3	5.2	0	0	0	0	—
Fairfax Co. (23)	6/4 to 7/30/2012	34	20.6	28.6	5.9	6.4	0	0	0	0	—
Hampton Roads		12	8.3	—^c	0	0	0	0	0	0	—
Portsmouth (24)	6/11/2012	4	25.0	— ^c	0	0	0	0	0	0	—
Chesapeake (25)	6/4 to 10/15/2012	8	0	— ^c	0	0	0	0	0	0	—
Total across all sites		1381	3.1	3.4^e	1.8	1.9	1.6	1.6	1.6	1.6	(1.1-2.4)

^aMIR estimates for Fairfax County area higher than other regions.

^bMIR is minimum infection rate for pooled nymphs; number of positive pools per total number of pooled nymphs; assumes that each positive pool of nymphs contains only one infected nymph.

^cMLE is maximum likelihood estimation from Pooled Infection Rate calculator, Excel add-in; estimates the maximum likely infection rate (%) based on the number of pools, number of ticks per pool, and number of positive pools; provides a 95% confidence interval (CI) on the estimate (Biggerstaff 2009).

^dCalculation of MLE was not necessary because each tested sample consisted of a single tick.

^eAll pools in this group tested positive, making it impossible to estimate a MLE for the testing outcome.

^fThe calculation for MLE-% positive across sites includes the samples from all sites.

TABLE 3C. VIRGINIA *AMBLYOMMA AMERICANUM* NYMPHS BY SURVEY REGIONS, COLLECTION SITES, COLLECTION DATES, AND TEST RESULTS FOR *RICKETTSIA* SPECIES

Tick collection regions, sites, and site numbers	Collection dates	Nymphs tested	Rickettsia amblyommii			Rickettsia parkeri		
			Pooled nymphs			Pooled nymphs		
			MIR ^a % Pos.	MLE ^b		MIR ^a % Pos.	MLE ^b	
	% Pos.	(95% CI)		% Pos.	(95% CI)			
Richmond City		192	30.2	68.3	(55.4–80.9)	0	0	—
Henrico Co. (1)	7/4–6/2012	82	29.3	67.4	(47.6–88.6)	0	0	—
Goochland Co. (2)	7/11–23/2012	54	29.6	61.5	(39.6–84.3)	0	0	—
Chesterfield Co. (3)	7/23/12	56	32.1	72.1	(48.7–94.0)	0	0	—
Charlottesville		261	26.8	54.4	(44.4–65.1)	0	0	—
Albemarle Co. (4)	6/10 to 7/20/2012	14	28.6	44.8	(16.4–87.8)	0	0	—
Albemarle Co. (5)	7/19/2012	101	27.7	56.1	(40.3–73.1)	0	0	—
Albemarle Co. (6)	7/19/2012	146	26.0	53.0	(39.8–67.4)	0	0	—
Southside Virginia		194	24.7	48.4	(37.3–60.5)	0	0	—
Brunswick Co. (7)	7/25/2012	116	25.0	47.9	(34.2–63.6)	0	0	—
Mecklenberg Co. (8)	7/25/2012	78	24.4	47.9	(31.2–67.6)	0	0	—
Lynchburg		284	24.7	46.8	(37.9–56.6)	0	0	—
Lynchburg (9)	7/31/2012	14	21.4	35.8	(10.4–85.6)	0	0	—
Lynchburg (10)	7/31/2012	82	23.2	41.9	(27.2–60.1)	0	0	—
Lynchburg (11)	7/31/2012	12	33.3	39.9	(14.8–74.3)	0	0	—
Lynchburg (12)	7/31/2012	80	25.0	49.1	(32.5–68.5)	0	0	—
Lynchburg (13)	8/3/2012	48	25.0	48.4	(28.0–73.4)	0	0	—
Lynchburg (14)	8/3/2012	48	25.0	48.4	(28.0–73.4)	0	0	—
Fredericksburg		275	27.6	57.5	(47.4–67.9)	0	0	—
Fredericksburg (15)	8/7/2012	14	28.8	44.8	(16.4–87.7)	0	0	—
Stafford Co. (16)	8/7/2012	20	20.0	28.2	(9.9–60.8)	0	0	—
Spotsylvania Co. (17)	8/15/2012	96	26.0	50.4	(35.1–67.7)	0	0	—
King George Co. (18)	8/15/2012	145	29.7	70.9	(55.5–85.6)	0	0	—
Fairfax County		163	28.0	64.4	(49.8–79.8)	3.7	3.8	(1.6–7.7)
Fairfax Co. (19)	6/15–28/2012	22	31.8	— ^d	—	0	0	—
Fairfax Co. (20)	6/4 to 7/25/2012	28	32.1	— ^d	—	3.6	3.4	(0.2–15.5)
Fairfax Co. (21)	6/7 to 7/11/2012	41	24.4	49.9	(26.7–78.8)	4.9	5.0	(1.0–15.6)
Fairfax Co. (22)	6/4 to 7/25/2012	38	26.3	49.0	(26.8–78.6)	7.9	8.1	(2.3–20.3)
Fairfax Co. (23)	6/4 to 7/30/2012	34	29.4	— ^d	—	0	0	—
Hampton Roads		12	50.0	— ^c	—	0	0	—
Portsmouth (24)	6/11/2012	4	75.0	— ^c	—	0	0	—
Chesapeake (25)	6/4 to 10/15/2012	8	37.5	— ^c	—	0	0	—
Total across all sites		1381	27.3	55.9^e	(51.4–60.5)	0.4	0.4	(0.2–0.9)

^aMIR is minimum infection rate for pooled nymphs; number of positive pools per total number of pooled nymphs; assumes that each positive pool of nymphs contains only one infected nymph.

^bMLE is maximum likelihood estimation from Pooled Infection Rate calculator, Excel add-in; Estimates the maximum likely infection rate (%) based on the number of pools, number of ticks per pool, and number of positive pools; provides a 95% confidence interval (CI) on the estimate (Biggerstaff 2009).

^cCalculation of MLE was not necessary because each tested sample consisted of a single tick.

^dAll pools in this group tested positive, making it impossible to estimate a MLE for the testing outcome.

^eThe calculation for MLE-% positive across sites includes the samples from all sites.

region ranged from 2.2% to 14.3% for adult ticks (average 7.8%) and 0.4% to 5.1% (average 1.9%) for nymphs (Tables 3A and B). The statewide adult tick infection rate for *E. ewingii* averaged 7.8%, similar to the 7.3% statewide *E. chaffeensis* rate. PME was found in adult ticks at sites in only two regions (Fairfax County and the Richmond City Area), but was found in nymphs at all regions except for the Hampton Roads region. Unexpectedly, although rates of PME infection were relatively high in adult and nymph-stage ticks from most sites in Fairfax County, it was not seen in any of the adult or nymph ticks taken from Fort Belvoir personnel who reside in that same region. Compared to *E. chaffeensis* and *E. ewingii*, the percentage of PME-positive samples by region

was lower, between 2.5% and 6.1% (average 1.9) for adult ticks and 0.8% and 2.8% (average 1.6) for nymphs.

Detection of Rickettsia species

In addition to the three *Ehrlichia* species, we detected *R. amblyommii* and *R. parkeri* (Table 3A and C), but not *R. rickettsii*. Adult ticks from all regions and all but three sites tested positive for *R. amblyommii*, and it was found in nymphs from every region and site sampled. *R. amblyommii* was found in much greater abundance than any bacterial species tested for, with regional detection rates of between 55.6 and 81.6% (average 72.8%) in adult ticks and 46.8 and 68.3% (average

TABLE 4A. *AMBLYOMMA AMERICANUM* ADULTS SUBMITTED TO THE US ARMY PUBLIC HEALTH COMMAND^a

Submitting installation	Dates received	Adults tested	Ehrlichia chaffeensis	Ehrlichia ewingii	Panola Mountain Ehrlichia
			Adult ticks % Positive ^b	Adult ticks % Positive ^b	Adult ticks % Positive ^b
Ft. A. P. Hill	4/3–7/24, 2012	84	1.2	7.1	0
Ft. Belvoir	3/26–8/21, 2012	226	0.4	0.9	0
Ft. Pickett	5/18–8/21, 2012	60	6.7	3.3	0
Total across sites		370	2.8	3.8	0

^aData from USAPHC samples is presented separately due to differences in collection methods.

^bAlthough some of the adult ticks from each installation pooled, none of the pooled adult samples tested positive, permitting the calculation of infection rates as a percentage of the total number of adult ticks tested.

TABLE 4B. *AMBLYOMMA AMERICANUM* NYMPHS SUBMITTED TO US ARMY PUBLIC HEALTH COMMAND^a

Submitting installation	Dates received	Nymphs tested	Ehrlichia chaffeensis			Ehrlichia ewingii			Panola Mountain Ehrlichia		
			MIR % Pos.	Pooled nymphs		MIR % Pos.	Pooled nymphs		MIR % Pos.	Pooled nymphs	
				MLE			MLE			MLE	
				% Pos.	(95% CI)		% Pos.	(95% CI)		% Pos.	(95% CI)
Ft. A. P. Hill	3/21–7/24, 2012	127	0.8	0.8	(0.1–3.8)	0			0.8	0.8	(0.1–3.8)
Ft. Belvoir	3/27–9/13, 2012	194	0.5 ^b			0			0		
Ft. Pickett	5/18–8/21, 2012	267	1.5	1.5	(0.5–3.6)	1.5	1.5	(0.5–3.5)	1.1	1.1	(0.3–3.0)
Total across sites		588	0.9	1.0^c	(0.4–2.1)	0.5	0.7^c	(0.2–1.6)	0.6	0.7^c	(0.2–1.6)

^aData from the US Army Public Health Command (USAPHC) samples is presented separately due to differences in collection methods.

^bAlthough some of the nymphs from Fort Belvoir pooled, none of the pooled samples tested positive, so the MIR value represents a percentage of the total number of nymphs tested.

^cThe calculation for MLE was % positive across sites includes the samples from all sites.

MIR, minimum infection rate; MLE, maximum likelihood estimation; CI, confidence interval.

55.9%) in nymphs. Unexpectedly, *R. parkeri* was found in *A. americanum* in two regions of Virginia—in the Hampton Roads region and Fairfax County. Additional tick samples from 2010 and 2011 from the Hampton Roads region were tested to determine if the 2012 *R. parkeri*-positive ticks were a unique finding. *R. parkeri* was detected in samples from both the previous 2 years, and samples from all 3 years were confirmed for *R. parkeri* by singleplex PCR and sequence analysis.

Discussion

This bacterial survey in *A. americanum* ticks represents the first statewide screening of this species in the state of Virginia since the mid-twentieth century (Sonenshine et al. 1966). We developed a multiplex real-time PCR assay for six bacteria of public health relevance and uncovered several findings.

First, the distribution of *E. chaffeensis* was focal, but where present rates could be relatively high—up to 24.5% in adults and up to 13.4% in nymphs. Across our study, tick infection rates averaged 7.3% in adults and 3.1% in nymphs. *E. ewingii* and PME were generally found at lower rates in nymphs but were present at multiple sites in most regions. Therefore, given the preponderance of *A. americanum*, and its aggressive predilection for biting humans, clinicians in this region should maintain ehrlichiosis high in the differential diagnosis for tick-borne disease. Human ehrlichiosis

cases caused by *E. chaffeensis* have steadily increased in Virginia each year, from 23 cases in 2007 to 130 cases in 2012. This study provides a baseline, whereby future tick surveillance activities could examine whether increases in human illness are paralleled by increases in the tick population and/or its infection rate.

Second, *R. amblyommii* was found consistently at sites across the state, with regional rates ranging from 56% to 82% of adults and 47% to 68% of nymphs (MLE estimate, Tables 3A and C). In recent years, Virginia has seen substantial increases in RMSF case detection rates. Specifically, in 2010 a total of 145 cases of RMSF were reported, increasing to 231 cases in 2011 and to 461 cases in 2012; these cases are typically reported to health departments by physicians and/or commercial testing laboratories as RMSF, but have been listed under “Spotted Fever Rickettsiosis” in the VDH annual reports since 2011 (Virginia Department of Health 2012). We think it likely that much of this increase in reported RMSF cases is caused by other illnesses, perhaps *Ehrlichia*, with “false” cross-reacting seropositivity due to the prevalence of *R. amblyommii* (Apperson et al. 2008, Smith et al. 2010). The rarity of *R. rickettsii* detection in *D. variabilis* (Ammerman et al. 2004, Dergousoff et al. 2009, Moncayo et al. 2010, Stromdahl et al. 2011) and its absence in the lone star tick populations we tested further supports this notion and begs the question of what is the RMSF vector? Again, we would advocate for more aggressive consideration

of ehrlichiosis in individuals with clinically suggestive tick-borne illness. Of course this distinction does not change treatment (e.g., doxycycline for either), but could greatly affect public health case reporting and more clearly focus the disease prevention message on *A. americanum*.

We suspect that deer populations are a significant driver of the *A. americanum* prevalence in Virginia. Virginia deer populations have risen in the last half-century due to several factors, expanding from an estimated 150,000 in the early 1950s, to approximately 422,000 in 1980, to an estimated 945,000 in 2004 (Virginia Department of Game and Inland Fisheries 2007). This expanding population has resulted in some deer becoming adapted to suburban environments, and control efforts in these settings have been largely unsuccessful. The increase in deer has come with an increase in associated ticks, including *A. americanum*.

A few other findings are worth noting. The discovery of *R. parkeri* in *A. americanum* in Virginia was surprising, but strengthens evidence that *A. americanum* could be a vector. To our knowledge, this is the first study to identify this pathogen in multiple *A. americanum* from each of several regions of within a state. The two regions where these *R. parkeri*-positive *A. americanum* ticks were found are the only regions known in Virginia to have established *A. maculatum* populations, suggesting that *A. americanum* and *A. maculatum* share one or more reservoir hosts in these two areas of Virginia. Additional studies will be required to determine the identity of these hosts, assess the extent of host sharing, and determine the shared reservoir host range.

We expected to detect PME within Virginia, but we were surprised to detect it in so many regions of the state. A study by Yabsley et al. demonstrated that PME had entered the Virginia white-tailed deer population as early as 2002 and showed these deer to be a natural reservoir host for PME (Yabsley et al. 2008). PME is not only a disease concern in humans, but in dogs as well (Qurollo et al. 2013). Further surveillance for illness caused by PME is needed to understand its significance as a pathogen.

In sum, we developed a six-plex real-time PCR assay to interrogate bacterial populations within the lone star tick, a vector of substantial increase and public health significance in the state of Virginia. We used the assay in tick populations from around the state, resulting in a substantial detection of *Ehrlichia*, (including rarely considered species), massive detection of *R. amblyomii*, and occasional detection of *R. parkeri*. We advocate use of such methods in the future to track changes in pathogen prevalence in this important tick.

Acknowledgements

The project described was supported by grant number K25AI067791 to H.G. from the National Institute Of Allergy and Infectious Diseases (NIAID). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIAID or the National Institutes of Health (NIH). This work was also supported by the Henry M. Jackson Foundation to C.W. Support to D.E.N. for this project comes from Fairfax County, Virginia, and support to T.H. comes from NIH grant number T32 AI 007471, the Johns Hopkins Malaria Research Institute, and the A. Ralph and Silvia E. Barr Fellowship.

Author Disclosure Statement

All authors declare no competing financial interests exist that may create a conflict of interest in connection with this article.

References

- Ammerman NC, Swanson KI, Anderson JM, Schwartz TR, et al. Spotted-fever group *Rickettsia* in *Dermacentor variabilis*, Maryland. *Emerg Infect Dis* 2004; 10:1478–1481.
- Apperson CS, Engber B, Nicholson WL, Mead DG, et al. Tick-borne diseases in North Carolina: Is “*Rickettsia amblyomii*” a possible cause of rickettsiosis reported as Rocky Mountain spotted fever? *Vector Borne Zoonotic Dis* 2008; 8:597–606.
- Biggerstaff BJ. PooledInfRate, Version 4.0: a Microsoft® Office Excel® Add-in to compute prevalence estimates from pooled samples. Centers for Disease Control and Prevention, Fort Collins, CO, USA; 2009.
- Chapman AS, Murphy SM, Demma LJ, Holman RC, et al. Rocky Mountain spotted fever in the United States, 1997–2002. *Vector Borne Zoonotic Dis* 2006; 6:170–178.
- Cohen SB, Yabsley MJ, Garrison LE, Freye JD, et al. *Rickettsia parkeri* in *Amblyomma americanum* ticks, Tennessee and Georgia, USA. *Emerg Infect Dis* 2009; 15:1471–1473.
- Dahlgren FS, Mandel EJ, Krebs JW, Massung RF, et al. Increasing incidence of *Ehrlichia chaffeensis* and *Anaplasma phagocytophilum* in the United States, 2000–2007. *Am J Trop Med Hyg* 2011; 85:124–131.
- Dergousoff SJ, Gajadhar AJ, Chilton NB. Prevalence of *Rickettsia* species in Canadian populations of *Dermacentor andersoni* and *D. variabilis*. *Appl Environ Microbiol* 2009; 75:1786–1789.
- Fornadel CM, Zhang X, Smith JD, Paddock CD, et al. High rates of *Rickettsia parkeri* infection in Gulf Coast ticks (*Amblyomma maculatum*) and identification of “*Candidatus Rickettsia andeanae*” from Fairfax County, Virginia. *Vector Borne Zoonotic Dis* 2011; 11:1535–1539.
- Goddard J. Experimental infection of lone star ticks, *Amblyomma americanum* (L.), with *Rickettsia parkeri* and exposure of guinea pigs to the agent. *J Med Entomol* 2003; 40:686–689.
- Jiang J, Yarina T, Miller MK, Stromdahl EY, et al. Molecular detection of *Rickettsia amblyomii* in *Amblyomma americanum* parasitizing humans. *Vector Borne Zoonotic Dis* 2010; 10:329–340.
- Lackman DB, Parker RR, Gerloff RK. Serological characteristics of a pathogenic rickettsia occurring in *Amblyomma maculatum*. *Public Health Rep* 1949; 64:1342–1349.
- Lackman DB, Bell EJ, Stoenner HG, Pickens EG. The Rocky Mountain spotted fever group of rickettsias. *Health Lab Sci* 1965; 2:135–141.
- Loftis AD, Massung RF, Levin ML. Quantitative real-time PCR assay for detection of *Ehrlichia chaffeensis*. *Journal of clinical microbiology*, 2003; 41:3870–3872.
- Loftis AD, Reeves WK, Spurlock JP, Mahan SM, et al. Infection of a goat with a tick-transmitted *Ehrlichia* from Georgia, U.S.A., that is closely related to *Ehrlichia ruminantium*. *J Vector Ecol* 2006; 31:213–223.
- Loftis AD, Mixson TR, Stromdahl EY, Yabsley MJ, et al. Geographic distribution and genetic diversity of the *Ehrlichia* sp. from Panola Mountain in *Amblyomma americanum*. *BMC Infect Dis* 2008; 8:54.
- Mixson TR, Campbell SR, Gill JS, Ginsberg HS, et al. Prevalence of *Ehrlichia*, *Borrelia*, and *Rickettsial* agents in

- Amblyomma americanum* (Acari: Ixodidae) collected from nine states. *J Med Entomol* 2006; 43:1261–1268.
- Moncayo AC, Cohen SB, Fritzen CM, Huang E, et al. Absence of *Rickettsia rickettsii* and occurrence of other spotted fever group rickettsiae in ticks from Tennessee. *Am J Trop Med Hyg* 2010; 83:653–657.
- Nadolny RM, Wright CL, Hynes WL, Sonenshine DE, et al. *Ixodes affinis* (Acari: Ixodidae) in southeastern Virginia and implications for the spread of *Borrelia burgdorferi*, the agent of Lyme disease. *J Vector Ecol* 2011; 36:464–467.
- Openshaw JJ, Swerdlow DL, Krebs JW, Holman RC, et al. Rocky mountain spotted fever in the United States, 2000–2007: Interpreting contemporary increases in incidence. *Am J Trop Med Hyg* 2010; 83:174–182.
- Paddock CD, Yabsley MJ. Ecological havoc, the rise of white-tailed deer, and the emergence of *Amblyomma americanum*-associated zoonoses in the United States. *Curr Top Microbiol Immunol* 2007; 315:289–324.
- Paddock CD, Sumner JW, Comer JA, Zaki SR, et al. *Rickettsia parkeri*: A newly recognized cause of spotted fever rickettsiosis in the United States. *Clin Infect Dis* 2004; 38:805–811.
- Paddock CD, Finley RW, Wright CS, Robinson HN, et al. *Rickettsia parkeri* rickettsiosis and its clinical distinction from Rocky Mountain spotted fever. *Clin Infect Dis* 2008; 47:1188–1196.
- Quorollo BA, Davenport AC, Sherbert BM, Grindem CB, et al. Infection with Panola Mountain *Ehrlichia* sp. in a dog with atypical lymphocytes and clonal T-cell expansion. *J Vet Intern Med* 2013; 27:1251–1255.
- Reeves WK, Loftis AD, Nicholson WL, Czarkowski AG. The first report of human illness associated with the Panola Mountain *Ehrlichia* species: A case report. *J Med Case Rep* 2008; 2:139.
- Smith MP, Ponnusamy L, Jiang J, Ayyash LA, et al. Bacterial pathogens in ixodid ticks from a Piedmont County in North Carolina: Prevalence of rickettsial organisms. *Vector Borne Zoonotic Dis* 2010; 10:939–952.
- Sonenshine DE, Atwood EL, Lamb JT, Jr. The ecology of ticks transmitting Rocky Mountain spotted fever in a study area in Virginia. *Ann Entomol Soc Am* 1966; 59:1234–1262.
- Stromdahl EY, Hickling GJ. Beyond Lyme: Aetiology of tick-borne human diseases with emphasis on the South-Eastern United States. *Zoonoses and Public Health* 2012; 59:48–64.
- Stromdahl EY, Jiang J, Vince M, Richards AL. Infrequency of *Rickettsia rickettsii* in *Dermacentor variabilis* removed from humans, with comments on the role of other human-biting ticks associated with spotted fever group Rickettsiae in the United States. *Vector Borne Zoonotic Dis* 2011; 11:969–977.
- Virginia Department of Game and Inland Fisheries. Virginia Deer Management Plan 2006–2015. Virginia Department of Game and Inland Fisheries, 2007.
- Virginia Department of Health (VDH). Reportable Disease Surveillance in Virginia, 2012.
- Wright CL, Nadolny RM, Jiang J, Richards AL, et al. *Rickettsia parkeri* in gulf coast ticks, southeastern Virginia, USA. *Emerg Infect Dis* 2011; 17:896–898.
- Yabsley MJ, Loftis AD, Little SE. Natural and experimental infection of white-tailed deer (*Odocoileus virginianus*) from the United States with an *Ehrlichia* sp. closely related to *Ehrlichia ruminantium*. *J Wildl Dis* 2008; 44:381–387.

Address correspondence to:

David Gaines

Division of Environmental Epidemiology

Virginia Department of Health

109 Governor Street

PO Box 2448

Richmond, VA 23219

E-mail: David.Gaines@vdh.virginia.gov