

Virginia Journal of Science
Volume 63, Issue 3 & 4
Fall & Winter 2012

A Centennial Record of Paleosalinity Change in the Tidal Reaches of the Potomac and Rappahannock Rivers, Tributaries to Chesapeake Bay

Neil E. Tibert², Lindsay J. Walker¹, William P. Patterson², J. Bradford Hubeny³, Emma Jones¹, Olivia R. Cooper¹

¹Department of Earth and Environmental Sciences, University of Mary Washington, Fredericksburg, Virginia 22401.

²Department of Geological Sciences, University of Saskatchewan, Saskatoon Saskatchewan, Canada.

³Department of Geological Sciences, Salem State University, Salem, Massachusetts 01970

ABSTRACT

Gravity and push cores from the Potomac and Rappahannock Rivers (Virginia Tidewater) were collected from central and proximal estuarine zones with known seasonal salinity stratification. The lowermost microfossil associations in the cores comprise alternating ostracode populations of *Cyprideis salebrosa* and *Cytheromorpha*. This microfossil association gives way to an oligohaline association dominated by the freshwater ostracode *Darwinula stevensoni*. Stable oxygen isotope values ($\delta^{18}\text{O}$) of Rappahannock *Cyprideis salebrosa* are highly variable ranging between -6.6 to -3.2‰ VPDB. $\delta^{18}\text{O}$ values for Potomac *Cytheromorpha fuscata* range from -8.2 to -3.2‰ VPDB. Positive excursions in $\delta^{18}\text{O}$ values are synchronous with population peaks for both *Cyprideis* and *Cytheromorpha* indicative of increased marine influence and/or higher salinities. Microfossil paleoecology coupled with oxygen isotope values record a marked shift towards gradual freshening and deterioration of the salinity structure in the tidal tributaries during the mid-to late 19th century. We attribute these trends to both decadal climate trends and aggressive land use practices in the Chesapeake Bay watershed during the late 19th to middle 20th centuries.

INTRODUCTION

Estuaries are physically, chemically, and biologically complex environments at the convergence of continental and marine processes. In the Chesapeake Bay, the recent combination of anthropogenic watershed modification and sea-level rise are forcing mechanisms that have potentially influenced mixing of fresh and marine waters in the tidal reaches of the major tributaries (Colman and Bratton 2003; Boon 2012). To test this hypothesis, sediment cores were collected from the proximal and central reaches

Corresponding author: Neil E. Tibert, ntibert@umw.edu




FIGURE 1. Sediment cores were collected from the Potomac and Rappahannock estuaries, both of which are tributaries to Chesapeake Bay, the largest estuary in the eastern United States (Colman and Mixon, 1988).

of the Potomac and Rappahannock estuaries for microfossil and stable isotopic analyses. Paleosalinity indicators were established on the basis of ostracode paleoecology (population abundances and pore morphometrics) and oxygen isotope values ($\delta^{18}\text{O}$). The paleosalinity trends were considered in the context of sedimentation history based on ^{137}Cs dating, organic matter concentrations, and magnetic susceptibility of collected cores. Cumulative results of these analyses indicate that salinity gradients in both estuaries have changed markedly since the beginning of the 19th century which is suggestive of anthropogenic influence on estuarine processes in the Chesapeake Bay.

BACKGROUND

Geographic Location

The Chesapeake Bay is the largest estuarine system in the United States that is located between Virginia and Maryland on the Atlantic Coastal Plain (Colman and Mixon 1988). This study focuses on the upper tidal reaches of the Potomac and Rappahannock Rivers where they transition from estuarine-to fluvial conditions (Fig. 1). Sediment cores were collected near the boundary between proximal (oligohaline) and central estuary (mesohaline) near Aquia Creek in the Potomac estuary and Blandfield Point in the Rappahannock estuary (Ellison and Nichols 1970, 1976; Ellison 1972; USEPA 1998). Since the 1980s, these estuaries have produced the highest annual




FIGURE 2. Historic sea level data for Washington, D.C. illustrates a constant rise in sea level over the last century for the Potomac estuary (NOAA 2008).

sediment yields for all major Chesapeake Bay tributaries (Langland and Cronin 2003). In general, the relatively high sedimentation rates in Chesapeake Bay are related to ongoing sea-level rise and anthropogenic watershed modification (Fig. 2) (Colman and Mixon 1988; Brush 1989; Colman and Bratton 2003; Boon and others 2010; Boon 2012).

Estuarine Circulation

The convergence of fluvial and marine waters creates a dynamic circulation pattern that impacts sedimentary processes in estuarine environments. Where freshwater outflow meets incoming saltwater in a partially mixed estuary like the Chesapeake Bay, the seasonal halocline (salinity gradient) forms a relatively impermeable surface to sediment transport that frequently coincides with the estuarine turbidity maximum (ETM) (Langland and Cronin 2003). In the Potomac estuary, the maximum extent of saltwater intrusion occurs near Aquia Creek (Elliott 1976). In the Rappahannock estuary, the maximum extent of saltwater intrusion occurs near Blandfield Point (Elliott and Nichols 1970). Our preliminary assessment of modern salinity structure revealed oligohaline conditions and/or weak haloclines near Aquia Creek and Blandfield Point during June 2009 (Fig. 3).

Microfossils

Ostracodes are aquatic crustaceans that are sensitive to changes in salinity and temperature (Frenzel and Boomer 2005) and their ecological associations have been used to make inferences about past environmental conditions in the Chesapeake Bay (Elliott and others 1966; Cronin and Grinbaum 1999; Cronin and others 2005, 2010). When used in conjunction with $\delta^{18}\text{O}$ values of their calcite carapaces, ostracodes can be used to develop paleosalinity proxies in the context of both climate induced evaporation and mixing of marine and freshwaters (Anderson and Arthur 1983; Anadón




FIGURE 3. Late spring salinity (mg/L) and temperature (°C) profiles measured from the (A) Potomac and (B) Rappahannock estuaries. This analysis was part of an unpublished assessment of water quality during June 2009.

and others 2002; Holmes and Chivas 2002; Ito and others 2003). Studies by Medley and others (2008) demonstrated that the shape of the sieve pores on the external surface of the carapace (Type C of Puri 1974) varies significantly with salinity which further improves the potential for past salinity determinations using ostracoda.

METHODS

Sediment cores were collected from the Rappahannock and Potomac estuaries with Ogeechee and Gravity (Wildco Inc.) corers (Fig. 1). The longer Ogeechee cores (>100

cm) were analyzed with a Bartington MS2C Core Logging Sensor, and subsequently partitioned into 1 cm segments for loss on ignition and microfossil processing. The shorter gravity cores (40 cm) were divided into 2 cm segments and samples sent to Core Scientific International (Winnipeg, MB, Canada) for ^{137}Cs analysis *via* gamma spectrometry.

Microfossil census counts were completed at 2 cm intervals using the methods outlined in Medley and others (2008). Sediment samples were rinsed on a 125 μm mesh sieve and wet samples were examined with a stereoscopic zoom microscope (Nikon SMZ1500). Select ostracodes were photographed using a variable pressure scanning electron microscope (Hitachi S-3400N). Morphometric shape analysis was completed according to the methods described by Medley and others (2008). Sieve pores on external valves of *Cyprideis* were traced to determine the areas and shapes using *ImageJ 1.3.1v* (National Institutes of Health by Wayne Rasband). Values for circularity were calculated using the following formula: $\text{Circularity} = 4\pi (\text{area}/\text{perimeter}^2)$. The best-fit trendline of circularity versus area crossplots was used to determine the pore slope values.

Ostracode carapaces were bathed in deionized water and sent to the Saskatchewan Isotope Laboratory (Saskatoon, SK, Canada) for oxygen and carbon isotope analysis *via* stable isotope ratio mass spectrometry (SIRMS). SIRMS preparation entails the heating of samples *in vacuo* to dissipate contaminants (e.g. organic matter and water that may influence isotope values) prior to analysis with a Finnigan Kiel-IV carbonate preparation device directly coupled to a Finnigan MAT 253 isotope ratio mass spectrometer. Data is expressed relative to the VPDB scale and calibrated to the NBS-19 standard ($\delta^{13}\text{C}=1.95\text{‰}$ VPDB; $\delta^{18}\text{O}=-2.2\text{‰}$ VPDB).

Sediment accumulation rates were determined following the ^{137}Cs analysis method of Robbins and Edgington (1975). Cesium-137 is a radioactive isotope (half life= \sim 30 years) that was released into the atmosphere during nuclear testing and simple gamma spectrometry can measure its concentration in sediment (USEPA 2010). Given that atmospheric levels of ^{137}Cs peaked in 1963, a sampling site's average sedimentation rate can be calculated using the simple stratigraphic thickness above the peak divided by the time in years since 1963. To ensure accurate gamma spectroscopy results, at least 2 g of sediment were removed from the center of each core segment. Forty samples from two Potomac estuary cores were analyzed by Core Scientific International (Winnipeg, Manitoba). Select samples were pretreated for radiocarbon dating at the University of Pittsburgh following the methods outlined by Abbott and Stafford (1996). AMS ^{14}C analyses were performed at the University of Arizona's Accelerator Mass Spectrometry Laboratory.

RESULTS

Physical Stratigraphy

Core PT-09-C3 from Aquia Creek (Potomac River) comprises 134 cm of dark grey clay with a layer of fine sand in the basal 10 cm (Figs. 1, 4). Magnetic susceptibility values average 38.7 SI Units with a minimum value of 18.7 (124 cm) and a maximum value of 166.7 (134 cm). Total organic matter (TOM) averages 6.59% with a minimum value of 1.42% (132 cm) and a maximum value of 9.68% (11 cm). The maximum concentration of ^{137}Cs occurs between the 20-22 cm cored interval (Fig. 5).




FIGURE 4. (A) Potomac Estuary and (B) Rappahannock Estuary: physical stratigraphy, ostracode populations, magnetic susceptibility, total organic matter, ostracode oxygen isotope values, ostracode carbon isotope values, and *Cypridella* pore morphometric data. These results are interpreted in the context of three distinct paleosalinity intervals (see Figure 7).




FIGURE 5. Results of $^{137}\text{Cesium}$ analysis of sediment collected near (A) the Harry Nice Bridge and (B) Aquia Creek. ^{137}Cs disintegrations per minute (DPM) were greatest between 20 and 22 cm in both cores.

Core RA-09-C3 at Blandfield Point (Rappahannock River) comprises 141 cm of brown-to-grey clay (Figs. 1, 4). Magnetic susceptibility in this core averages 86.3 SI Units with a minimum value of 8.9 (141 cm) and a maximum value of 206.3 (37 cm). Total organic matter averages 8.47% with a minimum value of 6.02% (81 cm) and a maximum value of 16.2% (37 cm). The maximum concentration of ^{137}Cs occurs between the 20-22 cm cored interval (Fig. 5).

Ostracode Paleoecology (Census and Pore Morphometrics)

Ostracodes observed at Aquia Creek include *Cyprideis salebrosa*, *Cytheromorpha fuscata*, and *Darwinula stevensoni* (Figs. 4A, 6). *Cyprideis salebrosa* dominates the 128-20 cm interval with a relative abundance maximum at 84 cm (n=146).




FIGURE 6. Oligohaline (*Darwinula stevensoni*), euryhaline (*Cyprideis salebrosa*), and mesohaline (*Cytheromorpha fuscata* and *Cytheromorpha curta*) ostracodes observed in sediment cores from the Rappahannock and Potomac Rivers. Salinity classification adopted from Belt and others (2005).

Cytheromorpha fuscata is relatively common within the 118-20 cm interval with a relative abundance maximum at 70 cm (n=30). *Darwinula stevensoni* is sparsely distributed with single valves observed above the 116 cm position (Fig. 4A).

Ostracodes at Blandfield Point include *Cyprideis salebrosa*, *Cytheromorpha curta* and *Darwinula stevensoni*. *Cyprideis salebrosa* is relatively common within the 139-101 cm interval with a relative abundance maximum at 107 cm (n=52) (Fig. 4B). *Cytheromorpha curta* is sparsely distributed with relative abundance peaks at 123 cm and 121 cm. *Darwinula stevensoni* is sparsely distributed with occurrences above the 57 cm position (Fig. 4B).

Valves of *Cyprideis salebrosa* from the Potomac River were analyzed for pore morphometric shape at nine stratigraphic positions within the 118-46 cm interval (Fig. 4A). An average of 39 pores were measured per ostracode valve whereas the calculated pore slope values that range from -0.0062 at 60 cm to -0.0011 at 76 cm. Valves of *C. salebrosa* from Blandfield Point (RA-09-C3) were selected for pore analyses at nine stratigraphic positions between 139-101 cm (Fig. 4A). An average of 43 pores were measured per ostracode whereas pore slope values range from -0.0054 at 139 cm to -0.0009 at 105 cm and exhibit a trend toward more positive slope values in the uppermost core.

Oxygen and Carbon Stable Isotope Values

Cytheromorpha fuscata

Oxygen isotope values measured in valves of *Cytheromorpha fuscata* at Aquia Creek average -6.1‰ VPDB. The minimum $\delta^{18}\text{O}$ value is -8.2‰ VPDB (60 cm) and the maximum value is -3.2‰ VPDB (78 cm) (Table 1)(Fig. 4A). $\delta^{18}\text{O}$ values are variable with the highest values recorded within the 122-78 cm and 57-38 cm intervals. Values decrease significantly above the 36 cm position.

Carbon isotope values for *C. fuscata* average -9.1‰ VPDB. The minimum $\delta^{13}\text{C}$ value is -12.0‰ VPDB (60 cm) and the maximum value is -4.2‰ VPDB (106 cm)(Table 1)(Fig. 4A). $\delta^{13}\text{C}$ values are most positive at the core's base, and gradually decreasing from -7.0 to -12.0‰ VPDB above the 78 cm position. Prominent positive excursion peaks mark the 78 and 106 cm positions (Fig. 4A).

Cyprideis salebrosa

Oxygen isotope values of *Cyprideis salebrosa* at Aquia creek average -5.6‰ VPDB. The minimum $\delta^{18}\text{O}$ value is -7.6‰ VPDB (100 cm) and the maximum value is -4.2‰ VPDB (48 cm) (Table 2)(Fig. 4A). $\delta^{18}\text{O}$ values of *Cyprideis salebrosa* gradually become higher in the top of the core. $\delta^{18}\text{O}$ values of *Cyprideis salebrosa* valves from Blandfield Point average -5.1‰ VPDB. The minimum value is -6.6‰ VPDB (115 cm) and the maximum value is -3.2‰ VPDB (134 cm) (Table 3)(Fig. 4B).

Carbon isotope values of *Cyprideis salebrosa* at Aquia Creek are more positive at the base of the core and shift to more negative values above the 78 cm position (Fig. 4A). $\delta^{13}\text{C}$ values average -9.8‰ VPDB. The minimum value is -13.6‰ VPDB (100 cm) and the maximum value is -6.6‰ VPDB (92 cm) (Table 2) (Fig. 4A). $\delta^{13}\text{C}$ values at Blandfield Point average -6.4‰ VPDB. The minimum value is -9.7‰ VPDB (115 cm) and the maximum value is -3.6‰ VPDB (134 cm) (Figure 4B). The maxima of both $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ are synchronous with both trending to lower values within the 134-115 cm interval (Table 3) (Fig. 4B).

Core Geochronology and Sedimentation Rates

Gravity cores (PT-10-C1 and PT-10-C3) collected from the Potomac estuary yield maximum ^{137}Cs decay rates between 20 and 22 cm that indicates an average sedimentation rate of 0.45 cm/yr since 1963 (Fig. 5). An earlier ^{137}Cs study at Blandfield Point (Rappahannock River) (Baliwag and others 2010) yielded an average sedimentation rate of the 0.54 cm/yr (see Tibert and others 2013). A single AMS ^{14}C radiometric date from a core in the Rappahannock River (upstream of Blandfield Point) yielded an uncorrected date of 245 ± 15 that suggests a longer term sedimentation rate of 0.36 cm/yr. The collective summary of the sedimentation rates and the AMS ^{14}C in the Rappahannock and Potomac Rivers indicates basal core ages that approximate middle-to late 18th century colonial times (1767 ± 15).

DISCUSSION/INTERPRETATIONS

Paleosalinity Indicators

Ostracodes reported from the tidal reaches of the Potomac and Rappahannock Rivers have well known ecological tolerances in estuarine environments (Frenzel and Boomer 2005). Species of *Cytheromorpha*, including *C. fuscata* and *C. curta*, are often

TABLE 1. Stable isotope values of *Cytheromorpha* spp. from the Potomac River.

Stratigraphic Position (cm)	δ^{18}_o (‰ V-PDB)	δ^{13}_c (‰ V-PDB)
20	-6.46	-10.75
26	-7.80	-8.78
36	-7.75	-11.51
38	-3.98	-9.06
42	-5.34	-8.36
44	-8.05	-11.50
46	-6.08	-11.35
48	-4.70	-11.41
50	-7.35	-10.80
52	-5.42	-10.06
54	-6.02	-9.30
56	-3.90	-9.68
60	-8.19	-12.00
62	-7.16	-9.59
68	-7.09	-11.55
74	-6.69	-9.25
76	-3.94	-7.03
76	-6.73	-10.40
78	-5.74	-9.68
78	-3.22	-4.37
80	-5.86	-8.52
64	-6.32	-9.17
70	-7.38	-11.42
72	-6.74	-7.96
82	-6.57	-9.24

TABLE 1. Continued

Stratigraphic Position (cm)	$\delta^{18}\text{O}$ (‰ V-PDB)	$\delta^{13}\text{C}$ (‰ V-PDB)
84	-7.38	-10.14
86	-7.86	-10.12
88	-4.32	-8.63
90	-3.83	-6.40
92	-6.97	-8.49
94	-5.61	-8.75
96	-5.69	-7.28
100	-6.99	-8.71
104	-5.90	-7.31
106	-4.47	-4.21
108	-7.53	-9.34
110	-6.62	-6.97
112	-6.93	-8.74
116	-5.04	-8.92
120	-6.35	-10.61
122	-5.79	-7.05

associated with mesohaline conditions (mesohaline = ~7-15 ‰), while *Darwinula stevensoni* is restricted to oligohaline waters (oligohaline = 0-5 ‰)(Elliot and others 1966; Cronin and Vann 2003). *Cyprideis salebrosa*, on the other hand, is common in both hyposaline and hypersaline environments, and can therefore have a wide salinity tolerance (euryhaline = 0-60‰)(Sandberg and Plusquellec 1974; Carbonnel 1983).

Cyprideis salebrosa has been used to reconstruct Holocene paleosalinity histories in lagoonal and lacustrine deposits in the Dominican Republic (Medley and others 2008). This studies show that synchronous positive excursions of the $\delta^{18}\text{O}$ values coupled with increased sieve pore irregularity are viable indicators for times of increased aridity and salinity. Although the $\delta^{18}\text{O}$ values of Potomac River water are more lower than those reported from Lago Enriqueillo (a hydrologically closed lake), they do closely match $\delta^{18}\text{O}$ values of foraminiferal calcite reported in the Chesapeake Bay (Cronin and others 2005). The isotope values considered in the context of sieve shape variability serve as confirmation for significant changes in salinity spanning the past several centuries.

TABLE 2. Stable isotope values of *Cyprideis salebrosa* from the Potomac River.

Stratigraphic Position (cm)	$\delta^{18}\text{O}$ (‰ V-PDB)	$\delta^{13}\text{C}$ (‰ V-PDB)
46	-4.58	-11.05
48	-4.22	-11.21
52	-5.02	-9.49
74	-5.91	-8.56
76	-4.51	-7.99
78	-4.83	-10.01
80	-5.68	-10.43
92	-5.25	-6.60
100	-7.61	-13.57
108	-6.04	-7.26
116	-7.47	-10.99
118	-5.77	-10.51

A Centennial Record of Paleosalinity for the Virginia Tidewater

Basal core intervals in the Potomac River are characterized by euryhaline ostracodes with inconsistent oxygen isotope values that support our interpretation of a highly variable annual salinity range during late colonial times (Fig. 7). Trends in $\delta^{18}\text{O}$ values reflect the ostracodes' preference to live close to the landward extent of the marine salt wedge in the partial mixed estuary. Given that this interval pre-dates the industrial period, the seasonal stratification in this partially mixed Chesapeake Bay estuary maintained a salinity gradient that was not significantly impacted by natural storm events.

The middle intervals in the cores exhibit $\delta^{18}\text{O}$ values that trend toward their maximum values. This pronounced variability in the isotopic variability is coincident with a paleoecological shift to a *Cytheromorpha fuscata*-dominated ostracode association, thus recording the transition from near-oligohaline (~5-10 ‰) to increased mesohaline (~7-15 ‰) salinities. Within this interval, $\delta^{18}\text{O}$ values vary over 5‰ highlighting the magnitude of the salinity gradient at this time (Fig. 7). The relative abundance of *Cyprideis salebrosa* with irregular pore shapes supports our interpretation for elevated salinities likely due to increasing marine influence due to sea-level rise during the early-late 19th century.

The uppermost core intervals show a sharp decline in *Cytheromorpha fuscata* and a marked decrease in $\delta^{18}\text{O}$ values. Because fresh water has lower $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values than marine water (Patterson and Walter 1994, Patterson 1998), the lower $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values of ostracode carapaces were likely forced by a transition to less saline water.

TABLE 3. Stable isotope values of *Cyprideis salebrosa* from the Rappahannock River.

Stratigraphic Position (cm)	$\delta^{18}\text{O}$ (‰ V-PDB)	$\delta^{13}\text{C}$ (‰ V-PDB)
101	-5.25	-6.33
103	-5.17	-6.39
109	-5.48	-5.93
111	-5.98	-7.51
113	-5.34	-7.60
115	-6.55	-9.67
123	-6.12	-7.13
127	-4.47	-6.28
129	-3.98	-5.44
131	-4.32	-5.20
133	-4.66	-5.73
134	-3.18	-3.62
137	-4.30	-6.31
139	-6.02	-6.84

Sedimentation rates during this interval indicate an average rate of 0.45 cm/yr that also corresponds to a paleoecological shift to the freshwater ostracode *Darwinula stevensoni*. Population shifts, pore shape analyses, and isotope trends indicate decreased salinity and weakening salinity gradient near Aquia Creek during the early 20th century (Figs. 4, 7).

Climate Forcing and Sediment Loading

Cronin and others (2005) evaluated oxygen and carbon isotopes from calcareous foraminifera, and magnesium-calcium ratios from ostracode carapaces that generally oscillate in-phase for the last ~150 years in the Chesapeake Bay. Similar in-phase variations characterize the isotope record from Rappahannock and Potomac ostracodes (e.g., $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ maxima at 78 cm) (Fig. 4). The variability in our paleosalinity proxy indicators suggests that global climatic events, including the last vestiges of the Little Ice Age (Cronin and others 2003), may have influenced late 18th and early 19th century salinity variability in both the Rappahannock and Potomac estuaries. Cronin and Vann (2003) and Cronin and others (2005, 2010) analyzed ostracode and foraminifera populations collected from the mouth of the Patuxent estuary. Their results indicate that decadal climate variability in the Chesapeake Bay is potentially influenced by the




FIGURE 7. Model of paleosalinity change in based on microfossil populations, geochemistry, and stratigraphy of the Potomac estuary, and supported by supplementary results from the Rappahannock estuary. Three stages of halocline (A) development/strengthening, (B) establishment, and (C) deterioration were reconstructed for the last two centuries in the tidal reaches of these estuaries.

North Atlantic Oscillation and possibly El Niño Southern Oscillation (ENSO). The apparent decadal variability in our paleosalinity data substantiates their hypothesis.

Between 1983 and 2000, the Rappahannock and Potomac Rivers produced the greatest average annual sediment yields for all major Chesapeake Bay tributaries (Langland and Cronin 2003). Sedimentation rates and patterns are variable across the bay due to its tributaries' variable physiographic identities and land use histories (Brush 1989; Langland and Cronin 2003). Brush (1989) linked recently altered sedimentation rates and patterns in Chesapeake Bay tributaries to aggressive post-colonial land development. However, the amount of this "legacy sediment" stored in Chesapeake Bay watersheds, as well as time required for it to reach the bay, currently remains unquantified (Langland and Cronin 2003).

CONCLUSIONS

The paleosalinity history for the Potomac and Rappahannock Rivers is likely related to both natural climate change and anthropogenic forcing mechanisms. The initial shift from euryhaline-to mesohaline dominated ostracode associations was attributed to increasing salinity associated with ongoing Holocene sea-level rise. Despite the increasing marine influence due sea-level rise, sediment deposited in the tidal reaches during the 20th century was characterized by increased abundances of oligohaline ostracodes with decreasing $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values. While the exact cause of decreasing salinity during a period of 20th century is unknown, our results corroborates the hypothesis that anthropogenically-induced salinity change, post-Industrial times, has had a significant impact on seasonal salinity gradient development in the tidal reaches of the Potomac and Rappahannock Rivers.

ACKNOWLEDGEMENTS

The research presented herein is the result of a senior theses completed by student author LJW completed at the University of Mary Washington. We'd like to thank

Olivia Cooper, Nate Winston, Fila Baliwag, and Henry Talley for their contributions to this data set. We acknowledge financial support provided by the University of Mary Washington with Faculty Development Grants awarded to NET and Undergraduate Research Grants awarded to LIW, ORC, and HT. The UMW Summer Science Institute provided funding to LJW and EJ to support their research during the summer of 2009. We also would like to thank Dr. Ben Kisila (UMW) for his help with ^{137}Cs methodologies and the many discussions regarding the sedimentation history for our sites. Thanks to Dinka Besic at the Saskatchewan Isotope Laboratory for loading and analyzing samples for stable isotope values. A special thanks to Joe Cutry and his crew at the UMW Auto Shop who helped us maintain safe and operational conditions on the UMW research boat. Thanks to the anonymous reviewers for their thoughtful comments that helped to improve the final version of this manuscript.

LITERATURE CITED

- Abbott, M.B., and Stafford, T. W. 1996. Radiocarbon Geochemistry of Modern and Ancient Arctic Lake Systems, Baffin Island, Canada: *Quaternary Research* 45:300-311.
- Anadón P., Gliozzi E., and Mazzini I. 2002. Paleoenvironmental reconstruction of marginal marine environments from combined paleoecological and geochemical analyses on ostracods. In: J Holmes and A Chivas (Eds.), *The Ostracoda: Applications in Quaternary research geophysical monograph*, 131 (pp. 227-247). Washington, DC: American Geophysical Union.
- Anderson T.F., and Arthur M.A., 1983. Stable isotopes of oxygen and carbon and their applications to sedimentologic and paleoenvironmental problems. In: MA Arthur, TF Anderson, IR Kaplan, and LS Land (Eds.), *Stable Isotopes in Sedimentary Geology SEPM Short Course No. 10* (pp. 1-151). Tulsa, OK: SEPM.
- Baliwag F., Wilson L., and Odhiambo B.K. 2010. Analysis of long term effects of mining on fluvial systems: A case study on the Rappahannock River system and the fate of old tailings of the Kelly Gold Mine, Virginia. *GSA Abstracts with Programs* 42(1):182.
- Belt E.S., Tibert N.E., Curran H.A., Diemer J.A., Hartman J.H., Kroeger T.J., and Harwood D.M. 2005. Evidence for marine influence on a low-gradient coastal plain: Ichnology and invertebrate paleontology of the lower Tongue River Member (Fort Union Formation, middle Paleocene), western Williston Basin, U.S.A. *Rocky Mountain Geology* 40(1):1-24.
- Boon, J.D. 2012. Evidence of Sea Level Acceleration at U.S. and Canadian Tide Stations, Atlantic Coast, North America: *Journal of Coastal Research* 28(6):1437-1445.
- Boon J.D., Brubaker J.M., and Forrest D.R. 2010. Chesapeake Bay land subsidence and sea level change: An evaluation of past and present trends and future outlook. Special Report No. 424 in *Applied Marine Science and Ocean Engineering*, Virginia Institute of Marine Science, Gloucester Point, VA.
- Bratton J.F., Colman S.M., and Seal R.S. II. 2003. Eutrophication and carbon sources in Chesapeake Bay over the last 2700 years: Human impacts in context. *Geochimica et Cosmochimica Acta*, 67(18):3385-3402.

- Brush G.S. 1989. Rates and patterns of estuarine sediment accumulation: *Limnology and Oceanography*, 34(7):1235-1246.
- Carbonnel G. 1983. Morphométrie et hypersalinité chez *Cyprideis torosa* (Jones) (Ostracoda, actuel) dans les salines de Santa-Pola (Alicante, Espagne): *Scientifique Géologie Bulletin* 36(4):211-219. Strasbourg.
- Colman, S.M. and Bratton, J.F. 2003. Anthropogenically induced changes in sediment and biogenic silica fluxes in Chesapeake Bay: *Geology* 31(1):71-74.
- Colman S.M. and Mixon R.B. 1988. The record of major Quaternary sea-level changes in a large coastal plain estuary, Chesapeake Bay, eastern United States: *Palaeogeography, Palaeoclimatology, and Palaeoecology* 68:99-116.
- Cronin T.M., Hayo K., Thunell R.C., Dwyer G.S., Saenger C., Willard D.A. 2010. The Medieval Climate Anomaly and Little Ice Age in Chesapeake Bay and the North Atlantic Ocean: *Palaeogeography, Palaeoclimatology, Palaeoecology* 297:299–310.
- Cronin T.M., Dwyer G.S., Kamiya T., Schwede S., and Willard D.A. 2003. Medieval Warm Period, Little Ice Age and 20th century temperature variability from Chesapeake Bay: *Global and Planetary Change* 36:17-29.
- Cronin T.M., Thunell R., Dwyer G.S., Saenger C., Mann M.E., Vann C., and Seal II R.R. 2005. Multiproxy evidence of Holocene climate variability from estuarine sediments, eastern North America: *Paleoceanography* 20: PA4006, doi:10.1029/2005PA001145.
- Cronin T.M. and Vann C.D. 2003. The sedimentary record of climatic and anthropogenic influence on the Patuxent estuary and Chesapeake Bay ecosystems: *Estuaries* 26:196-209.
- Dalrymple R.W., Zaitlin B.A., and Boyd R. 1992. Estuarine facies models; Conceptual basis and stratigraphic implications: *Journal of Sedimentary Research* 62(6):1130-1146.
- Elliott A.J. 1976. The circulation and salinity distribution of the upper Potomac estuary: *Chesapeake Science* 17(3):141-147.
- Elliott H.A., Ellison R., and Nichols M.M. 1966. Distribution of recent Ostracoda in the Rappahannock Estuary, Virginia: *Chesapeake Science*, 7(4):203-207.
- Ellison R.L. 1972. *Ammobaculites*, foraminiferal proprietor of the Chesapeake Bay estuaries. In: BW Nelson (Ed.), *Environmental Framework of Coastal Plain Estuaries*: Geological Society of America Memoir 133:247-262.
- Ellison R.L. and Nichols M.M. 1970. Estuarine foraminifera from the Rappahannock River, Virginia: *Contributions from the Cushman Foundation for Foraminiferal Research* 21:1-17.
- Ellison, R.L. and Nichols, M. M. 1976. Modern and Holocene Foraminifera in the Chesapeake Bay Region: *Maritime Sediments* 1:131-151.
- Frenzel P. and Boomer I. 2005. The use of ostracods from marginal marine, brackish waters as bioindicators of modern and Quaternary environmental change: *Palaeogeography, Paleoclimatology, Palaeoecology* 225: 68-92.
- Holmes JA, and Chivas AR, (2002). Ostracod shell chemistry—Overview. In: J Holmes and A Chivas (Eds.), *The Ostracoda: Applications in Quaternary research geophysical monograph*, 131 (pp. 185-204). Washington, DC: American Geophysical Union.

- Ito E., De Deckker R, and Eggins M. 2003. Ostracodes and their shell chemistry: Implications for paleohydrologic and paleoclimatologic applications. In: LE Park and LE Smith (Eds.), *Bridging the Gap: Trends in the Ostracode Biological and Geological Sciences* (pp. 116-152). Paleontological Society, New Haven, CT.
- Kemp, A.C., Horton, B.P., Culver, S.J., Corbett, D.R., van de Plassche, O., Gehrels, W.R., Douglas, B.C. and Parnell, A.C. 2009. Timing and magnitude of recent accelerated sea-level rise (North Carolina, United States): *Geology* 37(11):1035-1038.
- Langland M, and Cronin T (Eds.), (2003). A summary report of sediment processes in Chesapeake Bay and watershed. U.S. Geological Survey Water-Resources Investigations Report 03-4123. New Cumberland, PA.
- Medley P., Tibert N.E., Patterson W.P., Curran H.A., Greer L., and Colin J.P. 2008. Paleosalinity history of a middle Holocene lagoonal and lacustrine deposits in the Enriquillo Valley, Dominican Republic based on pore morphometrics and isotope geochemistry of Ostracoda: *Micropaleontology* 53(5):409-419.
- Patterson W.P. 1999. Oldest isotopically characterized fish otoliths provide insight to Jurassic continental climate of Europe: *Geology* 27(3):199-202.
- Patterson, W.P. and Walter L.M. 1994. Depletion of $d^{13}C$ in seawater ΣCO_2 on modern carbonate platforms: significance for the carbon isotopic record of carbonates: *Geology* 22:885-888.
- Puri H.S. 1974. Normal pores and the phylogeny of Ostracoda: *Geoscience and Man* 6:137-151.
- National Oceanic and Atmospheric Administration (NOAA). 2008. Mean sea level trends for stations in Virginia: Tides and Currents <http://tidesandcurrents.noaa.gov/index.shtml>.
- Robbins J.A. and Edgington D.N. 1975. Determination of recent sedimentation rates in Lake Michigan using Pb-210 and Cs-137: *Geochimica et Cosmochimica Acta* 39:285-304.
- Rosenfeld A. and Vesper, B. 1977. The variability of the sieve-pores in Recent and fossil species of *Cyprideis torosa* (Jones 1850) as an indicator for salinity and palaeosalinity. In: Loeffler, H., and Danielopol, D. L., Eds., *Aspects of Ecology and Zoogeography of Recent and Fossil Ostracoda*. The Hague: Dr W. Junk b.v., p. 55-67.
- Sandberg P.A. and Plusquellec P.L. 1974. Notes on the anatomy and passive dispersal of Cyprideis (Cytheracea, Ostracoda): *Geoscience and Man* 6:1-26.
- Scott D.B., Medioli F.S., and Schafer C.T. 2001. *Monitoring in coastal environments using foraminifera and thecameobian indicators*: Cambridge University Press, Cambridge, New York.
- Tibert N.E., Hubeny J.B., Abbott M., Kiker J.M., Walker, L.J., and McKenzie S. 2013. Latest Holocene sedimentation rates and sea level history for the tidal marshes of the Potomac and Rappahannock rivers, tributaries to Chesapeake Bay: *Virginia Journal of Science* 63(3):91-109.
- United States Environmental Protection Agency (USEPA), 2010. Cesium. Retrieved from <http://www.epa.gov/rpdweb00/radionuclides/cesium.html>

United States Environmental Protection Agency (USEPA), 1998. Condition of the Mid-Atlantic estuaries (Publication No. EPA 600-R-98-147): U.S. Government Printing Office, Washington, DC.

Wolman MG, (1967). A cycle of sedimentation and erosion in urban river channels. Geografiska Annaler Series A, Physical Geography 49(2): 385-395.