

Old Dominion University ODU Digital Commons

Computer Science Faculty Publications

Computer Science

1992

On Sources in Comparability Graphs, With Applications

Stephan Olariu
Old Dominion University

Follow this and additional works at: https://digitalcommons.odu.edu/computerscience_fac_pubs

 Part of the [Applied Mathematics Commons](#), and the [Computer Sciences Commons](#)

Repository Citation

Olariu, Stephan, "On Sources in Comparability Graphs, With Applications" (1992). *Computer Science Faculty Publications*. 128.
https://digitalcommons.odu.edu/computerscience_fac_pubs/128

Original Publication Citation

Olariu, S. (1992). On sources in comparability graphs, with applications. *Discrete Mathematics*, 110(1-3), 289-292. doi:10.1016/0012-365x(92)90721-q

This Article is brought to you for free and open access by the Computer Science at ODU Digital Commons. It has been accepted for inclusion in Computer Science Faculty Publications by an authorized administrator of ODU Digital Commons. For more information, please contact digitalcommons@odu.edu.

Note

On sources in comparability graphs, with applications

S. Olariu

Department of Computer Science, Old Dominion University, Norfolk, VA 23529-0162, USA

Received 31 October 1989

Abstract

Olariu, S., On sources in comparability graphs, with applications, *Discrete Mathematics* 110 (1992) 289–292.

We characterize sources in comparability graphs and show that our result provides a unifying look at two recent results about interval graphs.

An orientation Θ of a graph G is obtained by assigning unique directions to its edges. To simplify notation, we write $xy \in \Theta$, whenever the edge xy receives the direction from x to y . A vertex w is called a *source* whenever $vw \in \Theta$ for no vertex v in G . An orientation Θ is termed *transitive* if for every vertices x, y, z , $xy \in \Theta$ and $yz \in \Theta$ implies $xz \in \Theta$. It is well known that a graph G that admits a transitive orientation is a *comparability* graph.

In this context, it makes sense to ask the following natural question:

for what vertices of a comparability graph can we find a
transitive orientation that makes them into a source? (1)

The purpose of this note is to provide an answer to (1). As it turns out, our result provides a unifying look at two recent results concerning interval graphs [4, 6].

All the graphs in this work are simple with no self-loops nor multiple edges.

Familiarity with standard graph theoretical terminology compatible with Golumbic [5] is assumed. To specify our results, however, we need to define some new terms. For an arbitrary vertex w of G , the graph G^w is obtained from G by adding a new vertex w' and by making w' adjacent to w only. We claim that

w is a source in some transitive orientation of G if,
and only if, the graph G^w is a comparability graph. (2)

To justify (2), we note first that if G^w is a comparability graph then, by reversing the orientation on all the edges if necessary, we guarantee that w is a source in some transitive orientation Θ of G^w . In particular, w is a source in the restriction of Θ to G .

Conversely, given a transitive orientation Θ of G that makes w into a source, the orientation $\Theta^w = \Theta \cup \{ww'\}$ of G^w is transitive, and so G^w is a comparability graph, as claimed.

A vertex w of a graph G is called *special* if w coincides with one of the highlighted vertices in some graph F_i , ($1 \leq i \leq 4$) or in the complement \bar{F}_5 of the graph F_5 featured in Fig. 1. A vertex that is not special is referred to as *regular*. As it turns out, regular vertices play an important role in the answer to (1). More precisely, we state the following result.

Theorem 1. *A vertex w of a comparability graph G is a source in some transitive orientation Θ of G if, and only if, w is regular.*

Proof. First, let w be a regular vertex of G . If w fails to be a source in any transitive orientation of G then, by (2), the graph G^w is not a comparability graph. Hence G^w must contain an induced subgraph H isomorphic to one of Gallai's forbidden graphs (for a list see Gallai [2] or Duchet [1]). Since, by




Fig. 1.

assumption, G is a comparability graph, w' must belong to H . But now, it is a straightforward observation that w must be a special vertex, a contradiction.

Conversely, let w be a source in some transitive orientation Θ of G . Again, it is routine to check that if w is special, then G^w contains one of the forbidden graphs in Gallai's catalog, contradicting (2). This completes the proof of Theorem 1. \square

A graph $G = (V, E)$ is termed an *interval graph* if there exists a family $\{I_v\}_{v \in V}$ of intervals such that for distinct vertices u, v in G

$$uv \in E \text{ if, and only if, } I_u \cap I_v \neq \emptyset.$$

Such a family $\{I_v\}_{v \in V}$ of intervals is commonly referred to as an *interval representation* of G . An interval $I_x = [a_x, b_x]$, is called an *end interval* if $a_x \leq a_y$ for every $y \in V$; the vertex x itself is termed an *end vertex*.

An early characterization of interval graphs was proposed by Gilmore and Hoffman [3]: they showed that a graph G is an interval graph if, and only if, G itself is triangulated and its complement \bar{G} is a comparability graph.

Let Θ be a transitive orientation of a graph G . A linear order $<$ on the vertex-set of G is said to be *consistent* with Θ if

$$u < v \text{ whenever } uv \in \Theta.$$

(Note that such a linear order is readily available: we only need apply a topological sort to G . Furthermore, every source in Θ can be placed first in $<$.)

We are now in a position to show that Theorem 1 implies the following result.

Corollary 1.1 (Gimbel [4]). *A vertex w in an interval graph G is an end vertex if, and only if, G contains an induced subgraphs none of the graphs $\bar{F}_1, \bar{F}_2,$ or F_5 featured in Fig. 1, with w one of the highlighted vertices.*

Proof. The ‘only if’ implication is easy: we only need observe that if G contains an induced subgraph isomorphic to one of the graphs, $\bar{F}_1, \bar{F}_2,$ or F_5 featured in Fig. 1, then the highlighted vertices cannot correspond to an end interval in any interval representation of G .

To prove the ‘if’ implication, assume that G contains no induced subgraph isomorphic to one of the graphs $\bar{F}_1, \bar{F}_2,$ or F_5 featured in Fig. 1. Since G must be a triangulated graph, G cannot contain an induced subgraph isomorphic to the complement of the graph F_3 or F_4 of Fig. 1. Consequently, w is a regular vertex in \bar{G} . By Theorem 1, w is a source in some transitive orientation Θ of \bar{G} . Now a result of Gilmore and Hoffman [3] guarantees that

for every transitive orientation Θ of the edges of \bar{G} , there exists a linear order $<$ on the set of the maximal cliques of G such that $<$ is consistent with Θ and such that for every vertex x of G the maximal cliques containing x occur consecutively in $<$.

(For a proof the interested reader is referred to Golombic [5, pp. 172–173].)

Note, furthermore, that by virtue of (3), the set I_x of all the maximal clique containing x is an *interval*. Since w is a source in Θ , I_w becomes an end interval in $\{I_v\}_{v \in V}$, as claimed. \square

Recently, Skrien and Gimbel [6] proposed to call an interval graph G *homogeneously representable* if for every vertex v of G , there exists an interval representation of G in which I_v is an end interval.

Theorem 1 implies the following characterization of the homogeneously representable interval graphs.

Corollary 1.2 (Skrien and Gimbel [6]). *An interval graph G is homogeneously representable if, and only if, G contains no induced subgraph isomorphic to one of the graphs \bar{F}_1 and F_5 with $n = 3$.*

Proof. The ‘only if’ implication is immediate; to settle the ‘if’ implication, we only need show that every vertex of \bar{G} can be a source in some transitive orientation of \bar{G} , for then the conclusion follows by (3). Since G is triangulated, \bar{G} cannot have an induced subgraph isomorphic to one of the graphs F_3 and F_4 ; further, F_1 is an induced subgraph of F_2 . Consequently, every vertex of \bar{G} must be regular, and Theorem 1 implies that every vertex of \bar{G} is a source in some transitive orientation of \bar{G} . \square

References

- [1] P. Duchet, Classical Perfect Graphs, Ann. Discrete Math. 21 (North-Holland, Amsterdam, 1984).
- [2] T. Gallai, Transitiv orientierbare Graphen, Acta Math. Acad. Sci. Hungar. 18 (1967) 25–66.
- [3] P.C. Gilmore and A.J. Hoffman, A characterization of comparability graphs and interval graphs, Canad. J. of Math. 16 (1964) 539–548.
- [4] J. Gimbel, End vertices in interval graphs, Discrete Appl. Math. 21 (1988) 257–259.
- [5] M.C. Golumbic, Algorithmic Graph Theory and Perfect Graphs (Academic Press, New York, 1980).
- [6] D. Skrien and J. Gimbel, Homogeneously representable interval graphs, Discrete Math. 55 (1985) 213–216.