

Ursidae: The Undergraduate Research Journal at the University of Northern Colorado

Volume 2 | Number 1

Article 11

January 2012

Divergence of Sulfur-Flower Buckwheat using DNA Analyses

Taylor Brevik

Mitchell McGlaughlin

Follow this and additional works at: <http://digscholarship.unco.edu/urj>

 Part of the [Biology Commons](#)

Recommended Citation

Brevik, Taylor and McGlaughlin, Mitchell (2012) "Divergence of Sulfur-Flower Buckwheat using DNA Analyses," *Ursidae: The Undergraduate Research Journal at the University of Northern Colorado*: Vol. 2 : No. 1 , Article 11.

Available at: <http://digscholarship.unco.edu/urj/vol2/iss1/11>

This Poster Abstract is brought to you for free and open access by Scholarship & Creative Works @ Digital UNC. It has been accepted for inclusion in *Ursidae: The Undergraduate Research Journal at the University of Northern Colorado* by an authorized editor of Scholarship & Creative Works @ Digital UNC. For more information, please contact Jane.Monson@unco.edu.

Divergence of Sulfur-Flower Buckwheat using DNA Analyses

Biological Sciences

Presenter(s): Brevik, Taylor, McGlaughlin, Mitchell

Faculty Sponsor(s): McGlaughlin, Mitchell

Eriogonum umbellatum Torr. (Polygonaceae), the sulfur-flower buckwheat, is among the most diverse plant species in North America with 40 different recognized varieties. Several of these recognized varieties are rare, 24 of which are termed vulnerable, imperiled, or critically imperiled by NatureServe (natureserve.org). *Eriogonum umbellatum* species can be found inhabiting 10 states and 2 Canadian provinces in the mountainous west. Varieties display broad morphological, ecological, and geographic differences which make each taxonomically distinct. For this study, I am analyzing and comparing nuclear (ITS) DNA, inherited from both parents, and chloroplast DNA, inherited maternally, among five different varieties of *Eriogonum umbellatum* from several distinct populations residing in both Colorado and Oregon. Tissue samples were collected from several individuals from each population. The five varieties of interest include *E. umbellatum* var. *hausknechtii*, *E. umbellatum* var. *modocensis*, *E. umbellatum* var. *majus*, *E. umbellatum* var. *aureum*, and *E. umbellatum* var. *umbellatum*. Genetic analyses of these varieties using molecular markers should help to evaluate gene flow between populations and among varieties, mechanisms of differentiation, and evolutionary history. Understanding these will allow us to determine whether these varieties are monophyletic, share a common ancestor, or if they should be considered distinct evolutionary entities. Implications of this research could affect land management and conservation efforts, as well as our understanding of the mechanisms of evolutionary differentiation within diverse plant taxa.