

Touro Scholar

NYMC Faculty Posters

Faculty

Spring 3-23-2017

ROCK2 Gene Single Nucleotide Polymorphisms and Association with Bronchopulmonary Dysplasia in Extremely Low Birth Weight Infants

Sharina Rajbhandari
New York Medical College

Asma Amin
New York Medical College

Shaili Amatya
New York Medical College

Leewen Hsu
New York Medical College

Anna Zylak
New York Medical College

See next page for additional authors

Follow this and additional works at: https://touro scholar.touro.edu/nymc_fac_posters



Part of the [Enzymes and Coenzymes Commons](#), and the [Respiratory Tract Diseases Commons](#)

Recommended Citation

Rajbhandari, S., Amin, A., Amatya, S., Hsu, L., Zylak, A., Paudel, U., & Parton, L. A. (2017). ROCK2 Gene Single Nucleotide Polymorphisms and Association with Bronchopulmonary Dysplasia in Extremely Low Birth Weight Infants. Retrieved from https://touro scholar.touro.edu/nymc_fac_posters/26

This Poster is brought to you for free and open access by the Faculty at Touro Scholar. It has been accepted for inclusion in NYMC Faculty Posters by an authorized administrator of Touro Scholar. For more information, please contact touro.scholar@touro.edu.

Authors

Sharina Rajbhandari, Asma Amin, Shaili Amatya, Leewen Hsu, Anna Zylak, Umesh Paudel, and Lance A. Parton

Sharina Rajbhandari, Asma Amin, Shaili Amatya, Leewen Hsu, Anna Zylak, Umesh Paudel, Lance A. Parton
Neonatology, Maria Fareri Children's Hospital at Westchester Medical Center and New York Medical College,
Valhalla, NY, United States

Background

Bronchopulmonary dysplasia (BPD) leads to significant morbidity in ELBW infants.




Fig 1. Pathogenesis of BPD


Rho associated coiled-coil containing protein kinase 2 (ROCK-2) gene

- On chromosome 2p24
- ROCK: Serine/threonine protein kinase



RBD = Rho binding domain
PH = Plekstrin homology
CRD = Cysteine-rich zinc finger like motif Domain

Fig 2. ROCK-2 functional domains



*ROCK inhibitor decreases pulmonary edema, microvascular permeability and lung injury

Fig 3. ROCK pathways (eNOS, endothelial Nitric Oxide Synthase; ERM, ezrin/radixin/moesin; MLCPh, Myosin Light Chain Phosphatase)

ROCK-2 SNP rs2290156

- In intron; MAF (Minor Allele Frequency), C = 0.30
- G allele and GC genotype more in RDS

ROCK-2 SNP rs726843

- In intron; MAF, A = 0.35
- TC genotype more in RDS compared to controls

ROCK-2 SNP rs978906

- In 3' UTR; MAF, C = 0.39
- Affects ROCK2 expression by interfering with microRNA-1183 binding
- Association with stiffer arteries and with high altitude essential hypertension

Hypothesis

We hypothesize that ROCK2 gene SNP variants rs2290156, rs726843 and rs978906 are associated with development of BPD in ELBW infants.

Methods

Inclusion criteria

- ELBW infants (birth weight < 1kg)
- Informed parental consent

SNP Analysis

DNA was isolated from buccal swabs of 137 ELBW infants and analyzed via real-time PCR using Taqman probes for ROCK2 gene SNP variants rs2290156, rs726843 and rs978906.

BPD was defined by the need for Oxygen supplementation at 36 weeks postmenstrual age.

Statistics

Chi-square test, Fisher's exact test, Mann-Whitney Rank Sum test and t-test were performed for statistical analysis; p < 0.05 was considered significant.

Results

Demographic Characteristics

	No BPD (n = 57)	BPD (n = 80)	p value
Gestational age, wks, median (IQR)	26 (24, 27)	25 (24, 26)	0.22
Birth weight, g, mean (SD)	792.9 (123.8)	752.0 (147.3)	0.09
Female Gender, n (%)	32 (56)	48 (60)	0.78
Race, n (%)	Non Hispanic White	18 (33)	0.89
	Non Hispanic Black	16 (29)	
	Hispanic	16 (29)	
	Other	5 (9)	
Antenatal steroids, n (%)	43 (81)	68 (91)	0.18

Genotype Distributions

	Genotype	No BPD n (%)	BPD n (%)	P value
rs2290156	Wild allele	25 (64)	32 (71)	0.13
	Heterozygous	7 (18)	11 (24)	
	Minor allele	7 (18)	2 (4)	
	GG	25 (64)	32 (71)	0.64
	Any c	14 (36)	13 (28)	
rs726843	Wild allele	21 (55)	24 (42)	0.42
	Heterozygous	12 (32)	25 (44)	
	Minor allele	5 (13)	8 (14)	
	GG	21 (55)	24 (42)	0.22
	Any a	17 (45)	33 (58)	
rs978906	Wild allele	22 (52)	20 (33)	0.09
	Heterozygous	11 (26)	28 (47)	
	Minor allele	9 (21)	12 (20)	
	TT	22 (52)	20 (33)	0.04*
	Any c	20 (47)	40 (67)	

Conclusions

- ROCK2 gene SNP rs978906 shows association with BPD
- We speculate that this variant may play a role in the development of BPD by influencing smooth muscle tone in the pulmonary vasculature

References

- Liao YC, et al. Two functional polymorphisms of ROCK2 enhance arterial stiffening through inhibiting its activity and expression. Journal of Molecular and Cellular Cardiology. 2015; 79: 180-186
- Noma K, Kihara Y, Higashi Y. Striking crosstalk of ROCK signaling with endothelial function. Journal of Cardiology. 2012; 60:1-6
- Kaya G, et al. Association of Rho-kinase Gene Polymorphisms with Respiratory Distress Syndrome in Preterm Neonates. Pediatrics and Neonatology. 2017; 58: 36-42
- Pandey P, et al. Polymorphisms and haplotype of ROCK2 associated with high altitude essential hypertension in native high altitude Ladakhi Indian population: a preliminary study. Clin Exp Hypertens. 2016;38(2):238-44

