

Touro Scholar

Lander College for Women - The Anna Ruth and Mark Hasten School Publications and Research

Lander College for Women - The Anna Ruth and Mark Hasten School

2018

Interactive Map Portal of Some Important Fossil Localities Along the Southern Tethyan Margin of the Sephardic (Triassic) and Ethiopian (Jurassic) Faunal Provinces

Marcelo Rosensaft

Howard R. Feldman Touro College, howard.feldman@touro.edu

Follow this and additional works at: https://touroscholar.touro.edu/lcw_pubs



Part of the Paleontology Commons

Recommended Citation

Rosensaft, M., & Feldman, H. R. (2018). Interactive map portal of some important fossil localities along the southern Tethyan margin of the Sephardic (Triassic) and Ethiopian (Jurassic) faunal provinces. Poster presented at the 2018 Geological Society of America Northeastern Section Meeting, Portland, ME.

This Abstract is brought to you for free and open access by the Lander College for Women - The Anna Ruth and Mark Hasten School at Touro Scholar. It has been accepted for inclusion in Lander College for Women - The Anna Ruth and Mark Hasten School Publications and Research by an authorized administrator of Touro Scholar. For more information, please contact Timothy J Valente timothy.valente@touro.edu.

Northeastern Section - 53rd Annual Meeting - 2018

Paper No. 10-17

Presentation Time: 8:00 AM-12:00 PM

INTERACTIVE MAP PORTAL OF SOME IMPORTANT FOSSIL LOCALITIES ALONG THE SOUTHERN TETHYAN MARGIN OF THE SEPHARDIC (TRIASSIC) AND ETHIOPIAN (JURASSIC) FAUNAL PROVINCES

ROSENSAFT, Marcelo, Division of Geological Mapping, Geological Survey of Israel, 30 Malkhei Israel St., Jerusalem, 95501, Israel and FELDMAN, Howard R., Division of Paleontology (Invertebrates), American Museum of Natural History, 79th Street at Central Park West, New York, NY 10024-5192

The use of a "map portal" allows researchers to publish their maps and the data incorporated therein in a simple way that will reach a broad audience. Colleagues, as well as other workers worldwide, can connect to the portal through the internet and display the maps on desktop computers. The process is dynamic; any change made in the source map is immediately displayed on the user's desktop. In this report we illustrate a map of Triassic and Jurassic brachiopod collecting localities on the southern Tethyan margin in the Middle East. The locations are shown as red points that overlie a base map. By clicking on a point, a small box is opened that contains the textual data of the specimens collected at that locality: species names, the name of the author or collector, GIS data and one or more related field photos (if available). Brachiopod-bearing strata include some classic sites such as Douvillé's (1916) and Cossmann's (1925) localities in northern Sinai, Egypt, Muir-Wood's (1925) report on the fauna from Wadi Zarqa, Jordan, and Cooper's (1989) study of brachiopods from the Tuwaiq Mountains, Saudi Arabia. More modern studies are also noted such as Feldman's brachiopod studies from Gebel El-Maghara and Gebel El-Minshera, northern Sinai and from Hamakhtesh Hagadol and Makhtesh Ramon in southern Israel.

Session No. 10--Booth# 42

D3. Paleontology (Posters)

Sunday, 18 March 2018: 8:00 AM-12:00 PM

Lake Champlain Exhibition Hall (DoubleTree by Hilton)

Geological Society of America Abstracts with Programs. Vol. 50, No. 2

doi: 10.1130/abs/2018NE-309654

© Copyright 2018 The Geological Society of America (GSA), all rights reserved. Permission is hereby granted to the author(s) of this abstract to reproduce and distribute it freely, for noncommercial purposes. Permission is hereby granted to any individual scientist to download a single copy of this electronic file and reproduce up to 20 paper copies for noncommercial purposes advancing science and education, including classroom use, providing all reproductions include the complete content shown here, including the author information. All other forms of reproduction and/or transmittal are prohibited without written permission from GSA Copyright Permissions.