

Air pollution and heart failure: Relationship with the ejection fraction

Alberto Dominguez-Rodriguez, Javier Abreu-Afonso, Sergio Rodríguez, Ruben A Juarez-Prera, Eduardo Arroyo-Ucar, Yenny Gonzalez, Pedro Abreu-Gonzalez, Pablo Avanzas

Alberto Dominguez-Rodriguez, Ruben A Juarez-Prera, Eduardo Arroyo-Ucar, Department of Cardiology, Hospital Universitario de Canarias, E-38320 Santa Cruz de Tenerife, Spain

Alberto Dominguez-Rodriguez, Universidad Europea de Canarias, Facultad de Ciencias de la Salud, E-38300 Santa Cruz de Tenerife, Spain

Alberto Dominguez-Rodriguez, Instituto Universitario de Tecnologías Biomédicas, E-38320 Santa Cruz de Tenerife, Spain

Javier Abreu-Afonso, Department of Applied Physics, Universidad de Valencia, E-46900 Valencia, Spain

Sergio Rodríguez, Yenny Gonzalez, Center for Atmospheric Research Izaña, AEMET-CSIC, E-38071 Santa Cruz de Tenerife, Spain

Pedro Abreu-Gonzalez, Department of Physiology, Universidad de la Laguna, E-38320 Santa Cruz de Tenerife, Spain

Pablo Avanzas, Department of Cardiology, Hospital Universitario Central de Asturias, E-33006 Oviedo, Spain

Author contributions: Dominguez-Rodriguez A and Rodríguez S had contributed to the conception, design, analysis and interpretation of data, drafting and final approval of the manuscript submitted; Abreu-Afonso J, Juarez-Prera RA, Arroyo-Ucar E and Gonzalez Y had contributed to the analysis and interpretation of data, drafting and final approval of the manuscript submitted; Abreu-Gonzalez P and Avanzas P had contributed to the drafting of the manuscript, revising it critically for important intellectual content and final approval of the manuscript submitted.

Supported by The framework of one research projects funded by the Spanish Society of Cardiology (Daiichi-Sankyo project 2011)

Correspondence to: Alberto Dominguez-Rodriguez, MD, PhD, FESC, Department of Cardiology, Hospital Universitario de Canarias, Ofra s/n La Cuesta, E-38320 La Laguna, Santa Cruz de Tenerife, Spain. adrvdg@hotmail.com

Telephone: +34-922-679040 Fax: +34-922-678460

Received: December 4, 2012 Revised: December 7, 2012

Accepted: January 17, 2013

Published online: March 26, 2013

admission due to heart failure in patients with heart failure with preserved ejection fraction and reduced ejection fraction.

METHODS: We studied 353 consecutive patients admitted into a tertiary care hospital with a diagnosis of heart failure. Patients with ejection fraction of $\geq 45\%$ were classified as having heart failure with preserved ejection fraction and those with an ejection fraction of $< 45\%$ were classified as having heart failure with reduced ejection fraction. We determined the average concentrations of different sizes of particulate matter (< 10 , < 2.5 , and $< 1 \mu\text{m}$) and the concentrations of gaseous pollutants (carbon monoxide, sulphur dioxide, nitrogen dioxide and ozone) from 1 d up to 7 d prior to admission.

RESULTS: The heart failure with preserved ejection fraction population was exposed to higher nitrogen dioxide concentrations compared to the heart failure with reduced ejection fraction population ($12.95 \pm 8.22 \mu\text{g}/\text{m}^3$ vs $4.50 \pm 2.34 \mu\text{g}/\text{m}^3$, $P < 0.0001$). Multivariate analysis showed that nitrogen dioxide was a significant predictor of heart failure with preserved ejection fraction (odds ratio ranging from (1.403, 95%CI: 1.003-2.007, $P = 0.04$) to (1.669, 95%CI: 1.043-2.671, $P = 0.03$).

CONCLUSION: This study demonstrates that short-term nitrogen dioxide exposure is independently associated with admission in the heart failure with preserved ejection fraction population.

© 2013 Baishideng. All rights reserved.

Key words: Air pollution; Heart failure; Preserved ejection fraction; Reduced ejection fraction; Nitrogen dioxide

Dominguez-Rodriguez A, Abreu-Afonso J, Rodríguez S, Juarez-Prera RA, Arroyo-Ucar E, Gonzalez Y, Abreu-Gonzalez P, Avanzas

Abstract

AIM: To study whether the concentrations of particulate matter in ambient air are associated with hospital

zas P. Air pollution and heart failure: Relationship with the ejection fraction. *World J Cardiol* 2013; 5(3): 49-53 Available from: URL: <http://www.wjgnet.com/1949-8462/full/v5/i3/49.htm> DOI: <http://dx.doi.org/10.4330/wjc.v5.i3.49>

INTRODUCTION

Ambient air pollution is a recognized risk factor for cardiovascular morbidity and mortality^[1-3]. Nitrogen dioxide (NO₂) is a strong respiratory irritant gas originating from high-temperature combustion. Main outdoor sources of NO₂ include vehicle exhausts (particularly those equipped with diesel engines) and fossil-fuel power plants, whereas the most important indoor sources are gas heaters, stoves, and environmental tobacco smoke^[4].

Large meta-analyses of studies on the short-term health effects of NO₂ have been carried out in Europe^[5,6], the United States^[7,8], and Canada^[9]. The results indicate a positive association between daily increases of NO₂, cardiovascular and respiratory mortality. Several studies using administrative databases have shown a positive association between short-term increases in respirable or fine particles and the risk of hospitalization for congestive heart failure (HF)^[10-12].

The aim of this investigation was to study whether the concentrations of particulate matter in ambient air are associated with hospital admission due to HF in patients with HF with preserved ejection fraction (HF-PEF) and reduced ejection fraction (HF-REF).

MATERIALS AND METHODS

Study population

We prospectively enrolled 458 consecutive patients admitted into a tertiary care hospital with a diagnosis of HF. The diagnosis of HF had to be established according to the clinical Framingham criteria^[13]. We did not include patients with severe primary valve heart disease ($n = 13$), chronic obstructive pulmonary disease ($n = 30$), airway hyperresponsiveness ($n = 25$), asthma ($n = 16$) and presence of respiratory infection 15 d before admission ($n = 21$). Hence, 353 patients were included in the study. Patients with ejection fraction of $\geq 45\%$ were classified as having HF-PEF and those with an ejection fraction of $< 45\%$ were classified as having HF-REF^[14].

The study was planned according to the Declaration of Helsinki and approved by the local ethics committee, and all patients provided signed informed consent. Clinical data, including age, sex, arterial hypertension ($> 140/90$ mmHg), hypercholesterolemia (> 5.17 mmol/L), smokers, diabetes and left ventricular ejection fraction, were analyzed as baseline variables on admission. The left ventricular ejection fraction was measured using the modified Simpson's rule^[15].

Air pollution measurements

The atmospheric pollutants were measured in an urban

background monitoring station using reference methods (Directive 2008/50/EC). Concentrations of particulate matter (PM) smaller than 10, 2.5 and 1 μm (PM₁₀, PM_{2.5} and PM₁ respectively) were measured with automatic analyzer and the gravimetric method^[16].

The concentrations of gaseous pollutants were measured using different methods: (1) sulphur dioxide was measured using ultraviolet fluorescence (Thermo Electron CorporationTM, model 43C); (2) NO₂ was measured using chemiluminescence (Thermo Electron CorporationTM, model 42C); (3) ozone was measured using ultraviolet absorption (Thermo Electron CorporationTM, model 49C); and (4) carbon monoxide was measured using the technique NDIR-Gas Correlation Filter Analyser (Thermo Electron CorporationTM, model 48C). The analyzers were calibrated every 3 mo and they always had a high linearity ($r^2 = 0.99$)^[17]. Meteorological variables (temperature, relative humidity and wind speed) were measured using standard techniques. These variables were measured with 1 min resolution. Then, 24 h averages from the previous day up to 7 d prior to admission were calculated.

Statistical analysis

Results for normally distributed continuous variables are expressed as mean \pm SD. Continuous variables with non-normal distribution are presented as median values and interquartile intervals; categorical data are expressed as percentages. Analysis of normality of the continuous variables was performed with the Kolmogorov-Smirnov test. Differences between groups were assessed by unpaired 2-tailed *t* test and the Mann-Whitney *U* test for continuous variables, as appropriate. Categorical data and proportions were analyzed by use of χ^2 or Fisher's exact test when required. In our study, all of the pollutants were expressed as the 24 h average concentrations from the previous day up to 7 d prior to admission.

A multivariate analysis was carried out using a binary logistic regression model to estimate the risk of admission for HF-PEF compared to admission for HF-REF, according to sizes of particulate matter and concentrations of gaseous pollutants during 7 d prior to admission. All of the variables with a value of $P < 0.05$ in the univariate analysis were included in the model. Differences were considered statistically significant if the null hypothesis could be rejected with $> 95\%$ confidence. All probability values are 2 tailed. The SPSS 15 statistical software package (SPSS Inc, Chicago, IL, United States) was used for all calculations.

RESULTS

According to the pre-established criteria, 124 patients were classified as HF-PEF. The baseline characteristics of the patients with HF-PEF and HF-REF are listed in Table 1. The HF-PEF population was significantly older and included a larger proportion of women. There were no significant differences between groups regarding presence of conventional coronary risk factors for coronary

Table 1 Clinical variables of 353 consecutive patients with heart failure: Comparison between patients with heart failure and preserved ejection fraction and patients with heart failure and reduced ejection fraction *n* (%)

Variables	HF-PEF (<i>n</i> = 124)	HF-REF (<i>n</i> = 229)	<i>P</i> value
Age (yr)	69 ± 8	66 ± 12	0.01
Male gender	56 (45.2)	154 (67.2)	< 0.001
Hypertension	75 (60.5)	84 (36.7)	< 0.001
Hypercholesterolemia	35 (28.2)	52 (22.7)	0.25
Smokers	14 (11.3)	40 (17.5)	0.12
Diabetes	45 (36.3)	104 (45.4)	0.09
LVEF (%)	55 ± 9	33 ± 6	< 0.001

Data are expressed as mean ± SD, and *n* (%) for categorical variables. HF-PEF: Heart failure with preserved ejection fraction; HF-REF: Heart failure with reduced ejection fraction; LVEF: Left ventricular ejection fraction.

Table 2 Data on atmospheric pollution in ambient air and meteorological variables between the previous day and the 7 d prior to admission for both of the study group

	HF-PEF (<i>n</i> = 124)	HF-REF (<i>n</i> = 229)	<i>P</i> value
Meteorological variables			
Wind speed (m/s)	2.72 ± 0.68	2.62 ± 0.78	0.21
Temperature (°C)	19.76 ± 2.52	20.08 ± 2.82	0.32
Relative humidity (%)	67.68 ± 6.10	66.85 ± 7.02	0.29
Gaseous pollutants (mg/m ³)			
CO	172.44 ± 23.89	177 ± 27.10	0.11
SO ₂	8.1 ± 4.40	7.33 ± 3.46	0.06
NO ₂	12.95 ± 8.22	4.50 ± 2.34	< 0.0001
O ₃	59.80 ± 12.52	61.25 ± 11	0.26
Atmospheric particles (mg/m ³)			
PM10	21 (13-30)	25 (17.5-32)	0.02
PM2.5	13.5 (9-21)	16.5 (11-21)	0.12
PM1	8 (6-16)	9.5 (7-13)	0.42

All of the pollutants are expressed as the average concentration of the pollutant. Data are expressed as mean ± SD, and median values (interquartile intervals). HF-PEF: Heart failure with preserved ejection fraction; PM: Particulate material with an aerodynamic diameter; PM10: PM < 10 μm; PM2.5: PM < 2.5 μm; PM1: PM < 1 μm.

artery disease, with the exception of hypertension, which were higher in the patients with HF-PEF. Left ventricular ejection fraction was significantly reduced in the patients with HF-REF.

No statistically significant differences were found in the meteorological variables between both groups. Regarding gaseous pollutants, we found no statistically significant differences, except that there were higher concentrations of NO₂ exposure in patients with HF-PEF. When comparing, exposure to concentrations of sizes of particulate matter, between patients with HF-PEF and HF-REF, the first group tended to have lower values of PM10 (Table 2). We carried out partial multivariable binary logistic regression analyses, using a stepwise selection model. This analysis showed that exposure to NO₂ was a significant predictor of HF-PEF [odds ratio ranging from (1.403, 95%CI: 1.003-2.007, *P* = 0.04) to (1.669, 95%CI: 1.043-2.671, *P* = 0.03); Table 3].

Table 3 Multivariate binary logistic regression analysis including nitrogen dioxide as the main independent variable

	OR	95%CI	<i>P</i> value
Model 1 (unadjusted)			
NO ₂	1.428	1.001-2.055	0.04
Model 2			
NO ₂	1.669	1.043-2.671	0.03
Age	1.278	0.912-1.618	0.33
Model 3			
NO ₂	1.429	0.992-2.058	0.05
Gender	0.948	0.111-8.067	0.96
Model 4			
NO ₂	1.403	1.003-2.007	0.04
Hypertension	0.36	0.037-3.482	0.37
Model 5			
NO ₂	1.516	1.005-2.397	0.01
LVEF	1.024	0.791-1.324	0.85
Model 6			
NO ₂	1.489	1.009-2.453	0.01
PM10	1.124	0.997-1.974	0.94

LVEF: Left ventricular ejection fraction; NO₂: Nitrogen dioxide; PM: Particulate material with an aerodynamic diameter; PM10: PM < 10 μm.

DISCUSSION

Short-term exposure to air pollution is associated with acute cardiovascular events^[18-20]. Our results show that HF-PEF is common and accounts for a significant proportion of admissions in patients with HF, 35% of our patients. This rate of patients is similar to that reported in previous studies^[14,21]. This group of patients had different characteristics from those of patients with HF-REF, including older population, higher proportion of women, and more frequent history of hypertension. In the present study, we demonstrated that short-term exposure to raised NO₂ levels are an independent risk factor for admission to hospital for HF-PEF population, even superior to classical described predictors, such as age, sex, hypertension and left ventricular ejection fraction^[21,22].

Despite the large body of evidence linking NO₂ with daily mortality, few studies have addressed the issue of susceptibility to NO₂ by performing analyses by age, sex, and chronic morbidity^[6]. Recent epidemiology studies have focused on cardiopulmonary dysregulation, including the role of air pollutant exposure in provoking decompensated congestive HF^[3,10,23]. A number of mechanisms have been proposed to explain the cardiovascular effects of air pollutant. At the cellular level, these various mechanisms involve free radical production, oxidative stress, cytokine release, inflammation, endotoxin-mediated damage, stimulation of capsaicin receptors, autonomic nervous system activity and covalent modification of key cellular molecules^[24-27].

Ongoing investigation suggests that, although diastolic abnormalities may be present in many patients with HF-PEF, other aspects of pathophysiology likely also contribute to symptoms. Previous studies have concluded that inflammation contributes to diastolic abnormalities in HF-PEF^[28]. In our study, we discovered that NO₂ may

be a precipitating factor for admission for HF-PEF rather than the cause of this condition. The short-term elevations of NO₂ could play an important pathophysiologic role in HF-PEF population, perhaps through of the activation of molecular inflammatory pathways that could be transduced systemically even in the absence of obvious alveolitis or interstitial pneumonitis^[29]. In this way, Briet et al demonstrated that exposure to urban gaseous pollutant (including NO₂) affect artery endothelial function in patients as well as healthy control subjects^[30].

Our study has some limitations. We did not use time series analysis in our study to examine the short-term relationship between the variations in atmospheric pollution and HF. This was because daily variations in the pollutants during the 7 d prior to admission were small enough to allow us to exclude the time series analysis^[29]. Moreover, the sample size could be small, but the association between NO₂ and HF-PEF was highly significant.

This is the first study that demonstrates that short-term exposure to NO₂ is independently associated with the HF-PEF population, when compared to the HF-REF population.

COMMENTS

Background

Several studies using administrative databases have shown a positive association between short-term increases in respirable or fine particles and the risk of hospitalization for congestive heart failure.

Research frontiers

Heart failure is of growing incidence and prevalence and is now the main cause for hospital admission among the elderly and increasing expenditure in medicine. Ambient air pollution is a recognized risk factor for cardiovascular morbidity and mortality. Despite the large body of evidence linking nitrogen dioxide with daily mortality, few studies have addressed the issue of susceptibility to nitrogen dioxide by performing analyses by age, sex, and risk of admission for heart failure with preserved ejection fraction and reduced ejection fraction.

Innovations and breakthroughs

Nitrogen dioxide is a strong respiratory irritant gas originating from high-temperature combustion. Main outdoor sources of nitrogen dioxide include motor vehicles (particularly those equipped with diesel engines) and fossil-fuel power plants, whereas the most important indoor sources are gas heaters, stoves, and environmental tobacco smoke. In this study, authors found statistically significant association between nitrogen dioxide and admission in the heart failure with preserved ejection fraction population.

Applications

Several precautionary recommendations can be made for healthcare providers who interact with individuals who are at risk for cardiovascular diseases. Although they have not been clinically tested or proven to reduce mortality, they are practical and feasible measures that may help to reduce exposures to air pollution and therefore potentially lower the associated cardiovascular risk. These recommendations can be: (1) all patients with cardiovascular disease should be educated about the cardiovascular risks posed by air pollution; (2) part of patient education should include the provision of information regarding the available sources (local and national newspapers) that provide a daily air quality index; and (3) on the basis of the forecast air quality index, prudent recommendations for reducing exposure and limiting activity should be provided based on the patient's level of risk.

Terminology

Heart failure is a condition that is usually caused by a reduction of the contractile function of the ventricular chambers or an impairment of the relaxation properties of the cardiac chambers. Air pollution is the introduction into the atmosphere of chemicals, particulate matter, or biological materials that cause discomfort, disease, or death to humans.

Peer review

This is a good descriptive study in which the authors analyze effect of short-term exposure of nitrogen dioxide in patients with the clinical syndrome of heart failure with preserved and depressed left ventricular ejection fraction. The results are interesting and suggest that other aspects, as the exposure of nitrogen dioxide can contribute to pathophysiology of the heart failure with preserved ejection fraction. These data are of public health importance.

REFERENCES

- 1 **Pope CA**, Burnett RT, Thurston GD, Thun MJ, Calle EE, Krewski D, Godleski JJ. Cardiovascular mortality and long-term exposure to particulate air pollution: epidemiological evidence of general pathophysiological pathways of disease. *Circulation* 2004; **109**: 71-77 [PMID: 14676145 DOI: 10.1161/01.CIR.0000108927.80044.7F]
- 2 **Pope CA**, Renlund DG, Kfoury AG, May HT, Horne BD. Relation of heart failure hospitalization to exposure to fine particulate air pollution. *Am J Cardiol* 2008; **102**: 1230-1234 [PMID: 18940298 DOI: 10.1016/j.amjcard.2008.06.044]
- 3 **Wellenius GA**, Schwartz J, Mittleman MA. Particulate air pollution and hospital admissions for congestive heart failure in seven United States cities. *Am J Cardiol* 2006; **97**: 404-408 [PMID: 16442405 DOI: 10.1016/j.amjcard.2005.08.061]
- 4 **Kraft M**, Eikmann T, Kappos A, Künzli N, Rapp R, Schneider K, Seitz H, Voss JU, Wichmann HE. The German view: effects of nitrogen dioxide on human health—derivation of health-related short-term and long-term values. *Int J Hyg Environ Health* 2005; **208**: 305-318 [PMID: 16078645 DOI: 10.1016/j.ijheh.2005.04.002]
- 5 **Dingwall C**. Transport across the nuclear envelope: enigmas and explanations. *Bioessays* 1991; **13**: 213-218 [PMID: 1654049 DOI: 10.1183/09031936.06.00143905]
- 6 **Chiusolo M**, Cadum E, Stafoggia M, Galassi C, Berti G, Faustini A, Bisanti L, Vigotti MA, Dessi MP, Cernigliaro A, Mallone S, Pacelli B, Minerba S, Simonato L, Forastiere F. Short Term Effects of Nitrogen Dioxide on Mortality and Susceptibility Factors in Ten Italian Cities: the EpiAir Study. *Environ Health Perspect* 2011; Epub ahead of print [PMID: 21586369 DOI: 10.1289/ehp.1002904]
- 7 **Stieb DM**, Judek S, Burnett RT. Meta-analysis of time-series studies of air pollution and mortality: effects of gases and particles and the influence of cause of death, age, and season. *J Air Waste Manag Assoc* 2002; **52**: 470-484 [PMID: 12002192 DOI: 10.1080/10473289.2002.10470794]
- 8 **Stieb DM**, Judek S, Burnett RT. Meta-analysis of time-series studies of air pollution and mortality: update in relation to the use of generalized additive models. *J Air Waste Manag Assoc* 2003; **53**: 258-261 [PMID: 12661685 DOI: 10.1080/10473289.2003.10466149]
- 9 **Shin HH**, Stieb DM, Jessiman B, Goldberg MS, Brion O, Brook J, Ramsay T, Burnett RT. A temporal, multicity model to estimate the effects of short-term exposure to ambient air pollution on health. *Environ Health Perspect* 2008; **116**: 1147-1153 [PMID: 18795155 DOI: 10.1289/ehp.11194]
- 10 **Wellenius GA**, Bateson TF, Mittleman MA, Schwartz J. Particulate air pollution and the rate of hospitalization for congestive heart failure among medicare beneficiaries in Pittsburgh, Pennsylvania. *Am J Epidemiol* 2005; **161**: 1030-1036 [PMID: 15901623 DOI: 10.1093/aje/kwi135]
- 11 **Morris RD**, Naumova EN. Carbon monoxide and hospital admissions for congestive heart failure: evidence of an increased effect at low temperatures. *Environ Health Perspect* 1998; **106**: 649-653 [PMID: 9755140 DOI: 10.1289/ehp.98106649]
- 12 **Burnett RT**, Smith-Doiron M, Stieb D, Cakmak S, Brook JR. Effects of particulate and gaseous air pollution on cardiorespiratory hospitalizations. *Arch Environ Health* 1999; **54**: 130-139 [PMID: 10094292 DOI: 10.1080/0003989990602248]

- 13 **Anguita M**, Comin J, Almenar L, Crespo M, Delgado J, Gonzalez-Costello J, Hernandez-Madrid A, Manito N, Perez de la Sota E, Segovia J, Segura C, Alonso-Gomez AM, Anguita M, Cequier A, Comin J, Diaz-Buschmann I, Fernandez-Lozano I, Fernandez-Ortiz A, Gomez de Diego JJ, Pan M, Worner F, Alonso-Pulpon L, Bover R, Castro A, Diaz-Molina B, Gomez-Bueno M, Gonzalez-Juanatey JR, Lage E, Lopez-Granados A, Lupon J, Martinez-Dolz L, Munoz R, Pascual D, Ridocci F, Roig E, Varela A, Vazquez de Prada JA. Comments on the ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012. A report of the Task Force of the Clinical Practice Guidelines Committee of the Spanish Society of Cardiology. *Rev Esp Cardiol (Engl Ed)* 2012; **65**: 874-878 [PMID: 22999110 DOI: 10.1016/j.recesp.2012.07.013]
- 14 **Magaña-Serrano JA**, Almahmeed W, Gomez E, Al-Shamiri M, Adgar D, Sosner P, Herpin D. Prevalence of heart failure with preserved ejection fraction in Latin American, Middle Eastern, and North African Regions in the I PREFER study (Identification of Patients With Heart Failure and PREServed Systolic Function: an epidemiological regional study). *Am J Cardiol* 2011; **108**: 1289-1296 [PMID: 22000627 DOI: 10.1016/j.amjcard.2011.06.044]
- 15 **Kan G**, Visser CA, Lie KI, Durrer D. Early two-dimensional echocardiographic measurement of left ventricular ejection fraction in acute myocardial infarction. *Eur Heart J* 1984; **5**: 210-217 [PMID: 6723690]
- 16 **Rodriguez S**, Cuevas E, Gonzalez Y, Ramos R, Romero PM, Perez N, Querol X, Alastuey A. Influence of sea breeze circulation and road traffic emissions on the relationship between particle number, black carbon, PM₁, PM_{2.5} and PM_{2.5-10} concentrations in a coastal city. *Atmos Environ* 2008; **42**: 6523-6534 [DOI: 10.1016/j.atmosenv.2008.04.022]
- 17 **Gonzalez Y**, Rodriguez S, Guerra JC, Trujillo JL, Garcia R. Ultrafine particles pollution in urban coastal air due to ship emissions. *Atmos Environ* 2011; **45**: 4907-4914 [DOI: 10.1016/j.atmosenv.2011.06.002]
- 18 **Peters A**, Dockery DW, Muller JE, Mittleman MA. Increased particulate air pollution and the triggering of myocardial infarction. *Circulation* 2001; **103**: 2810-2815 [PMID: 11401937 DOI: 10.1161/01.CIR.103.23.2810]
- 19 **Peters A**, von Klot S, Heier M, Trentinaglia I, Hörmann A, Wichmann HE, Löwel H. Exposure to traffic and the onset of myocardial infarction. *N Engl J Med* 2004; **351**: 1721-1730 [PMID: 15496621 DOI: 10.1056/NEJMoa040203]
- 20 **Dominguez-Rodriguez A**, Abreu-Afonso J, Rodríguez S, Juárez-Prera RA, Arroyo-Ucar E, Jiménez-Sosa A, González Y, Abreu-González P, Avanzas P. Comparative study of ambient air particles in patients hospitalized for heart failure and acute coronary syndrome. *Rev Esp Cardiol* 2011; **64**: 661-666 [PMID: 21652134 DOI: 10.1016/j.recesp.2010.12.017]
- 21 **Yang XS**, Sun JP. Advances in diastolic heart failure. *World J Cardiol* 2010; **2**: 58-63 [PMID: 21160757 DOI: 10.4330/wjc.v2.i3.58]
- 22 **González-Juanatey JR**, Alegría Ezquerro E, Bertoméu Martínez V, Conthe Gutiérrez P, de Santiago Nocito A, Zsolt Fradera I. [Heart failure in outpatients: comorbidities and management by different specialists. The EPISERVE Study]. *Rev Esp Cardiol* 2008; **61**: 611-619 [PMID: 18570782 DOI: 10.1157/13123067]
- 23 **Peel JL**, Metzger KB, Klein M, Flanders WD, Mulholland JA, Tolbert PE. Ambient air pollution and cardiovascular emergency department visits in potentially sensitive groups. *Am J Epidemiol* 2007; **165**: 625-633 [PMID: 17194748 DOI: 10.1093/aje/kwk051]
- 24 **Shofer S**, Chen TM, Gokhale J, Kuschner WG. Outdoor air pollution: counseling and exposure risk reduction. *Am J Med Sci* 2007; **333**: 257-260 [PMID: 17435421 DOI: 10.1097/MAJ.0b013e31803b913c]
- 25 **Mills NL**, Donaldson K, Hadoke PW, Boon NA, MacNee W, Cassee FR, Sandström T, Blomberg A, Newby DE. Adverse cardiovascular effects of air pollution. *Nat Clin Pract Cardiovasc Med* 2009; **6**: 36-44 [PMID: 19029991 DOI: 10.1038/ncp-cardio1399]
- 26 **Chen TM**, Shofer S, Gokhale J, Kuschner WG. Outdoor air pollution: overview and historical perspective. *Am J Med Sci* 2007; **333**: 230-234 [PMID: 17435417 DOI: 10.1097/MAJ.0b013e31803b8c91]
- 27 **Araujo JA**. Are ultrafine particles a risk factor for cardiovascular diseases? *Rev Esp Cardiol* 2011; **64**: 642-645 [PMID: 21723025 DOI: 10.1016/j.recesp.2011.05.002]
- 28 **Alves AJ**, Eynon N, Oliveira J, Goldhammer E. RAAS and adrenergic genes in heart failure: Function, predisposition and survival implications. *World J Cardiol* 2010; **2**: 187-197 [PMID: 21160750 DOI: 10.4330/wjc.v2.i7.187]
- 29 **Brook RD**, Rajagopalan S, Pope CA, Brook JR, Bhatnagar A, Diez-Roux AV, Holguin F, Hong Y, Luepker RV, Mittleman MA, Peters A, Siscovick D, Smith SC, Whitsel L, Kaufman JD. Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association. *Circulation* 2010; **121**: 2331-2378 [PMID: 20458016 DOI: 10.1161/CIR.0b013e3181d8bec1]
- 30 **Briet M**, Collin C, Laurent S, Tan A, Azizi M, Agharazii M, Jeunemaitre X, Alhenc-Gelas F, Boutouyrie P. Endothelial function and chronic exposure to air pollution in normal male subjects. *Hypertension* 2007; **50**: 970-976 [PMID: 17875820]

P- Reviewer Aronow W S- Editor Wen LL L- Editor A
E- Editor Zhang DN

