

Temporal variability of summer-time ozone and aerosols in the free troposphere over the eastern North Atlantic

J. M. Prospero¹, R. Schmitt², E. Cuevas³, D. L. Savoie¹, W. C. Graustein⁴,
K. K. Turekian⁴, A. Volz-Thomas⁵, A. Díaz⁶, S. J. Oltmans⁷, H. Levy II⁸

Abstract. In the free troposphere over Tenerife in the summer, O₃ concentrations are anti-correlated with major pollutant aerosols (nss-SO₄⁻ and NO₃⁻) and with ²¹⁰Pb, a tracer for boundary layer sources. In contrast, O₃ is highly correlated with ⁷Be, a product of cosmic ray interactions in the upper troposphere and stratosphere. This suggests that natural O₃ sources (i.e. the stratosphere) might be playing an important role. Nonetheless our results do not preclude the possibility that substantial amounts of pollution-related O₃ could be transported in the free troposphere. However, to be consistent with our results, the transport mechanisms would have to incorporate efficient processes for the removal of pollutant aerosol species and ²¹⁰Pb.

Introduction

Tropospheric O₃ is both a greenhouse gas and a strong oxidant and is a precursor to the formation of highly reactive radicals such as OH which play a critical role in controlling the lifetimes and fates of many chemical species [Levy, 1971; Thompson, 1992]. Tropospheric O₃ is principally derived from two sources: episodic injection of air from the stratosphere where most of the Earth's O₃ column inventory resides; and *in situ* photochemical production in the polluted continental boundary layer where most of the nitrogen oxides (NO_x) and reactive organic carbon compounds (VOC's) are emitted. Conversely, there is the potential for net chemical destruction of O₃ in the free troposphere where NO_x levels are low. Results from numerical simulations suggest that increasing anthropogenic emissions of both NO_x and VOC's are leading to higher concentrations of tropospheric O₃ throughout the northern hemisphere [e.g., Thompson, 1992]. Direct measurements indicate that O₃ has been increasing at an annual rate of 1-2% per year for the past several decades at some relatively remote European surface sites and may have more than doubled over the last 100 years [Janach, 1989; Volz, 1993]. In contrast, a recent analysis of two decades of ozonesonde data from a wide range of sites [Logan, 1994] finds that, while O₃ appears to have increased in the free

troposphere over a number of stations in Europe, it has remained relatively constant in most other regions [Oltmans, 1993; Logan, 1994], e.g. at Mauna Loa Observatory (MLO), Hawaii (19.53°N, 155.57°W) [Oltmans and Levy, 1994].

The export of O₃ from continental regions to the marine atmosphere involves complex meteorological [Moody *et al.*, 1995] as well as chemical [Jacob *et al.*, 1993] processes. This report focuses on data from Izaña, Tenerife, Canary Islands (28°18'N, 16°29'W) where, as a part of the Tropospheric Ozone Research (TOR) program, O₃ has been measured at a Spanish meteorological observatory located on a mountain ridge (elevation 2370 m). This region is affected by a variety of air masses [Sancho *et al.*, 1992], including relatively clean air from the central North Atlantic and more polluted air from Europe or North America. The data from Izaña are interesting because of the strong and coherent temporal changes in species concentration and their clear association with specific meteorological conditions.

Izaña experiences a pronounced orographic flow regime [Schmitt *et al.*, 1988]. At night downslope winds prevail and samples should be representative of free tropospheric air. Night-time O₃ at Izaña exhibits a marked seasonal cycle [Schmitt *et al.*, 1988; Schmitt and Carretero, 1995]. From August through February, the composited monthly mean concentrations (1987-1994) are relatively constant, with values of 38-42 ppbv. Mean concentrations increase rapidly in the spring to a maximum of about 57 ppbv in June and then decrease through the summer. The monthly means are similar to those at MLO [Oltmans and Levy, 1992] except from June through September when the Izaña means exceed those at MLO by 8-15 ppbv.

The high summer means at Izaña have previously been attributed to the impact of anthropogenic species [Schmitt *et al.*, 1988; Schmitt and Carretero, 1995]. Relatively high concentrations of O₃ are generally associated with back-trajectories from north of Tenerife that pass over or close to Europe and North America. In some cases, by backtracking along trajectories, the O₃ concentration measured at Izaña could be matched with good agreement to ozonesonde measurements over continental stations [Schmitt and Carretero, 1995]. Indeed, the Izaña monthly mean O₃ concentrations are similar to those at non-urban sites in Europe throughout the year [Janach, 1989].

In this paper we use the concentrations of non-sea-salt (nss) SO₄⁻, NO₃⁻, ²¹⁰Pb, and ⁷Be, to more thoroughly investigate the potential sources of the high summertime O₃ concentrations at Izaña. Over the North Atlantic, NO₃⁻ and nss-SO₄⁻ are largely derived from anthropogenic sources [Savoie *et al.*, 1992]. Pb-210 (τ_{1/2} = 22 years) is a decay product of ²²²Rn (τ_{1/2} = 3.8 days) in soils; ⁷Be (τ_{1/2} = 53 days) is produced in the stratosphere and upper troposphere by cosmic ray interactions. Consequently these nuclides can serve as excellent tracers for the transport history of air parcels [Brost

¹Div. of Marine and Atmospheric Chemistry, Univ. of Miami

²Meteorologie Consult GmbH, Königstein, Deutschland

³Inst. Nacional de Meteorología, Tenerife, Islas Canarias, España

⁴Dept. of Geology and Geophysics, Yale University

⁵Inst. Chemie Dynamik der Geosphäre, Jülich, Deutschland

⁶Universidad de La Laguna, Tenerife, Islas Canarias, España

⁷CMDL, NOAA, Boulder, Colorado

⁸GFDL, NOAA, Princeton, New Jersey

Copyright 1995 by the American Geophysical Union.

Paper number 95GL02791

0094-8534/95/95GL-02791\$03.00

et al., 1991; Balkanski et al., 1993]. Although PAN (peroxyacetyl nitrate), also measured at the site, can help in assessing the influence of anthropogenic sources, its concentrations during summer at Izaña are too low for this purpose, probably due to its dissociation at the typical summertime temperatures [Schmitt et al., 1988; Schmitt and Carretero, 1995; Schmitt, 1995].

Results and Discussion

Ozone is continuously monitored with a Dasibi UV absorption instrument whose calibration is tied to the U.S. National Institute of Standards and Technology reference O_3 photometer maintained at Gaithersburg, Maryland [Oltmans and Levy, 1992]. Daily high-volume ($1 \text{ m}^3 \text{ min}^{-1}$) aerosol filter samples are collected at night (i.e., during downslope wind conditions) as a part of the Atmosphere/Ocean Chemistry Experiment (AEROCE). Samples for chemical analysis are collected on Whatman-41 filters and analyzed in Miami for nss-SO_4^- (total aerosol SO_4^- minus Na^+ times 0.2517, the $\text{SO}_4^-/\text{Na}^+$ ratio in bulk seawater) and NO_3^- using ion chromatography [Savoie et al., 1992]. The radio-tracer samples are collected on quartz filters and analyzed at Yale using a high-purity germanium well gamma counter [Turekian et al., 1989].

Our filters quantitatively collect aerosol NO_3^- and also gas-phase HNO_3 , a major end-product of PAN decomposition. Thus our NO_3^- measurements should yield a good estimate of the total of aerosol NO_3^- , HNO_3 , and PAN that might have been present. At Izaña the NO_3^-/O_3 volume mixing ratio during high O_3 events is typically in the range of 0.004 to 0.008 (i.e., NO_3^- concentrations of $0.5\text{--}1 \mu\text{g m}^{-3}$ and O_3 of 50–60 ppbv). These ratios are in the range of values for the NO_3^-/O_3 ratios measured in the upper troposphere over northern Europe [Murphy et al., 1993].

A daily time series of O_3 and aerosol (night-time) data from July through September (henceforth, "summer") of 1991 shows that the day-to-day changes in O_3 concentrations are strongly correlated with those of ^7Be (Fig. 1a). In every case, the maxima and minima in the O_3 concentration data are associated with corresponding maxima and minima in ^7Be . Note the series of sharp one-day changes around mid-July and just after mid-August. In contrast, the O_3 time series is strongly anticorrelated with that of ^{210}Pb (Fig. 1b). Often ^{210}Pb changes by a factor of two or more in a one-day period while O_3 undergoes an equally sharp opposite change. There are also extended periods of low O_3 that are accompanied by high concentrations of ^{210}Pb (e.g., 6 days in mid-July and 12 days in early September).

Lower nss-SO_4^- and NO_3^- concentrations (Fig. 1c and 1d, respectively) are almost invariably associated with peaks in O_3 ; conversely aerosol maxima coincide with O_3 minima. The temporal changes in nss-SO_4^- are particularly remarkable. Many of the nss-SO_4^- values that are associated with the O_3 peaks are in the range of $0.1\text{--}0.4 \mu\text{g m}^{-3}$ (23–93 pptv), comparable to those at remote South Pacific sites [Savoie et al., 1989]. In contrast, the peak concentrations of nss-SO_4^- (usually associated with low O_3 values) are often a factor of ten or more higher. The NO_3^- data (Fig. 1d) show a pattern that is similar to that for nss-SO_4^- . Samples having high concentrations of nss-SO_4^- , NO_3^- , and ^{210}Pb usually contained high concentrations of mineral dust as reflected in the aluminum concentrations (e.g. Fig. 2c; Arimoto et al., 1995); these filters also had a very pronounced red-brown coloration that is characteristic of Saharan dust [Prospero and Nees, 1986]. Dust from large-scale dust storms, which occur often during the summer months in North Africa, is transported across the west coast of Africa, typically reaching altitudes of 6–8 km and covering large areas of the tropical North Atlantic [Carlson and Prospero, 1972; Prospero and Nees, 1986].

A series of scatterplots shows that the relationships in the summer of 1991 discussed above are generally valid for the entire data set from Izaña. In these plots, the data are partitioned into two groups, those for which O_3 concentrations are greater than 50 ppbv ("high O_3 ") and those for concentrations less than 35 ppbv ("low O_3 "). The scatterplot of ^{210}Pb versus ^7Be (Fig. 2a) shows that high O_3 is associated with high ^7Be and low ^{210}Pb and, conversely, that low O_3 is associated with high ^{210}Pb and low ^7Be . A scatterplot of nss-SO_4^- versus ^{210}Pb (Fig. 2b) shows that nss-SO_4^- is well correlated with ^{210}Pb . Within this distribution, the low O_3 values are tightly clustered at the high end of the $\text{nss-SO}_4^-/^{210}\text{Pb}$ distribution whereas the high O_3 values are clustered at the low end.

As previously stated, high concentrations of nss-SO_4^- and NO_3^- are usually associated with high concentrations of aluminum from mineral dust. Figure 2c shows that nss-SO_4^- concentrations above about $1 \mu\text{g m}^{-3}$ are usually associated




Figure 1. Daily mean night-time concentrations during July-September 1991 of O_3 with: (a) ^7Be ; (b) ^{210}Pb ; (c) nss-SO_4^- ; and (d) NO_3^- . Data joined by lines indicate contiguous samples.




Figure 2. Scatterplot of July–September night-time aerosol concentrations at Izaña. Data are selected and partitioned for O_3 concentrations greater than 50 ppbv (Δ) and less than 35 ppbv (O). (a) ^{210}Pb vs. ^7Be , 1 July 1989–31 August 1993; (b) nss-SO_4^- vs. ^{210}Pb , 1 July 1989–31 August 1993; (c) aluminum vs. nss-SO_4^- , 2 July 1989–30 September 1992; and (d) NO_3^- vs. nss-SO_4^- , 25 July 1987–30 September 1994.

with Al concentrations of several $\mu\text{g m}^{-3}$ or greater (i.e., dust concentrations of 10's of $\mu\text{g m}^{-3}$ or more). These samples are generally characterized by low O_3 concentrations. Conversely, the high O_3 group is primarily associated with extremely low Al concentrations and with nss-SO_4^- concentrations less than $1 \mu\text{g m}^{-3}$.

Although the time sequence of high-low aerosol NO_3^- concentrations (Fig. 1d) is similar to that for nss-SO_4^- (Fig. 1c), it differs in one major respect. The low NO_3^- values (associated with O_3 peaks) are typically between 0.5 to $1 \mu\text{g m}^{-3}$ (180–360 pptv). These values are 5 to 10 times higher than South Pacific values [Savoie *et al.*, 1989] where anthropogenic impacts are minimal. At Izaña the mass ratio of NO_3^- to nss-SO_4^- changes markedly with respect to the O_3 concentration. Figure 2d shows the entire July–September Izaña data set for 1987–1994 partitioned according to O_3 concentration. When aerosol concentrations are high (e.g., above about $1 \mu\text{g m}^{-3}$), the $\text{NO}_3^-/\text{nss-SO}_4^-$ mass ratio falls between 0.5 to 1, values that are typical for polluted air masses; these conditions are usually associated with low O_3 . When aerosol concentrations are low (less than about $1 \mu\text{g m}^{-3}$), the $\text{NO}_3^-/\text{nss-SO}_4^-$ mass ratio is usually much greater than 1, values that are high compared to typical pollution aerosols; these occasions usually coincide with relatively high O_3 values (in Fig. 2d, over 50 ppbv).

Because of the similarities between the temporal variability of ^{210}Pb and that of nss-SO_4^- and NO_3^- and because of the high concentrations of nss-SO_4^- and NO_3^- during periods of high ^{210}Pb , we conclude that the nss-SO_4^- and NO_3^- are transported from continental pollution sources. This interpretation is supported by isentropic trajectories [J. Merrill, personal

communication; see Moody *et al.*, 1995] which, on days when the concentration of nss-SO_4^- and NO_3^- are high, usually track back across North Africa, often hooking north to Europe. These trajectories are consistent with the fact that these samples usually contain very high concentrations of mineral dust as previously discussed. In contrast, the high O_3 (and low aerosol) events were generally associated with trajectories that came from the north in the middle and upper troposphere but which generally do not pass over continental Europe. The tight clustering of high- O_3 events at the low end of the Al versus nss-SO_4^- scatterplot (Fig. 2c) is compatible with this interpretation. Our trajectory results are consistent with the Izaña trajectory climatology [Sancho *et al.*, 1992] which shows that, during July–September, 700 mb trajectories (the pressure altitude applicable to Izaña) fall largely into two classes: one group (26–37% of the cases by month) comes from over the North Atlantic (and possibly North America) north of 28°N (the latitude of Izaña); a second group (23–35%) passes over North Africa (and possibly Europe). At 700 mb, only a very small percentage of the trajectories (2–3%) come directly from Europe or England.

Conclusions

During the summer, high O_3 events at Izaña are associated with air masses that contain very low concentrations of aerosol species that are usually characteristic of "polluted" air masses. Conversely, samples that had high concentrations of pollution aerosols were usually associated with relatively low concentrations of O_3 and with trajectories that had passed over North Africa (and, usually, Europe). If one assumes that

these polluted air masses initially contained high concentrations of O₃, then much of the O₃ must have been subsequently destroyed in transit. The net balance of O₃ production and destruction depends critically on the concentration of NO_x and the concentration of water vapor. Our data suggest that for the cases with high aerosol concentrations, losses of O₃ appear to dominate over production on the time scales that it takes to transport air from pollution sources in Africa and Europe (i.e., a few days). The African-European trajectories generally lie at higher altitudes (e.g., 700 mb [Sancho *et al.*, 1992]) and the air is extremely dry [Carlson and Prospero, 1972] so that the water-vapor-related destruction processes for O₃ should be relatively inefficient. The large surface area available on dust might be a factor in the O₃ loss.

While our data can be interpreted to imply a substantial impact from upper tropospheric (and, perhaps, stratospheric) O₃ sources, they do not preclude the possibility that significant amounts of O₃ might be photochemically produced in the free troposphere or transported there from surface sources via convective clouds [Pickering *et al.*, 1992]. If boundary layer sources are playing an important role in free-troposphere O₃, then the transport mechanism from the boundary layer must efficiently remove pollutant nss-SO₄⁻ and NO₃⁻ aerosols and their gaseous precursors, especially SO₂ along with the tracer ²¹⁰Pb. In contrast, O₃, NO_x, PAN, organic nitrates, and VOC's, being relatively insoluble, would be vented to the free troposphere where subsequent chemistry could slowly produce O₃ [e.g., Liu *et al.*, 1987] and oxidize the remaining NO_x to NO₃⁻. As the destruction rates of O₃ are also slow in the free troposphere (because of the low water vapor content) the net effect of the production and destruction processes could be to maintain relatively high levels of surface-derived O₃ along with high NO₃⁻ values. The removal of ²¹⁰Pb, would obscure the signature for the surface origin of the air parcel [Balkanski *et al.*, 1993].

Acknowledgments. We thank the Spanish Meteorological Service (INM) for their cooperation in this program, especially C. Rus. We are particularly grateful to the Izaña Observatory staff who carried out many of the routine operations. This work is supported by National Science Foundation grants ATM-8703411, ATM-9013125, ATM-9414808, ATM-8701292, and ATM-9012950 and the National Atmospheric and Oceanic Administration, Dept. of Commerce. TOR activities are supported by the German Ministry of Research and Technology (BMFT) under grant 07EU7649/A.

The editor thanks the referees for their assistance in evaluating this paper.

References

- Arimoto, R., R. A. Duce, B. J. Ray, W. G. Ellis, Jr., J. D. Cullen, and J. T. Merrill, Trace elements in the atmosphere over the North Atlantic, *J. Geophys. Res.*, **100**, 1199-1213, 1995.
- Balkanski, Y. J., D. J. Jacob, G. M. Gardner, W. C. Graustein, and K. K. Turekian, Transport and residence times of tropospheric aerosols inferred from a global three-dimensional simulation of ²¹⁰Pb, *J. Geophys. Res.*, **98**, 20,573-20,586, 1993.
- Brost, R. A., J. Feichter, and M. Heimann, Three-dimensional simulation of ⁷Be in a global climate model, *J. Geophys. Res.*, **96**, 22,423-22,445, 1991.
- Carlson, T. N., and J. M. Prospero, The large-scale movement of Saharan air outbreaks over the northern equatorial Atlantic, *J. Appl. Meteorol.*, **11**, 283-297, 1972.
- Jacob, D.J., J.A. Logan, G.M. Gardner, R.M. Yevich, C.M. Spivakovsky, S.C. Wofsy, S. Sillman, and M.J. Prather, Factors regulating ozone over the United States and its export to the global atmosphere, *J. Geophys. Res.*, **98**, 14,817-14,826, 1993.
- Janach, W. E., Surface ozone: trend details, seasonal variations, and interpretation, *J. Geophys. Res.*, **94**, 18,289-18,295, 1989.
- Levy, H., II, Normal atmosphere: large radical and formaldehyde concentrations predicted, *Science*, **173**, 141-143, 1971.
- Liu, S. C., M. Trainer, F. C. Fehsenfeld, D. D. Parrish, E. J. Williams, D. W. Fahey, G. Hübler, and P. C. Murphy, Ozone production in the rural troposphere and the implications for regional and global ozone distributions, *J. Geophys. Res.*, **92**, 4191-4207, 1987.
- Logan, J. A., Trends in the vertical distribution of ozone: An analysis of ozonesonde data, *J. Geophys. Res.*, **99**, 25,553-25,586, 1994.
- Moody, J. L., S. J. Oltmans, H. Levy II, and J. T. Merrill, Transport climatology of tropospheric ozone: Bermuda, 1988-1991, *J. Geophys. Res.*, **100**, 7179-7194, 1995.
- Murphy, D. M., D. W. Fahey, M. H. Proffitt, S. C. Liu, K. R. Chan, C. S. Eubank, S. R. Kawa, and K. K. Kelly, Reactive nitrogen and its correlation with ozone in the lower stratosphere and upper troposphere, *J. Geophys. Res.*, **98**, 8751-8773, 1993.
- Oltmans, S. J., Climatology of Arctic and Antarctic ozone, in *The Tropospheric Chemistry of Ozone in the Polar Regions*, edited by H. Niiki, pp. 25-40, Springer, Berlin, 1993.
- Oltmans, S. J., and H. Levy, II, Seasonal cycle of surface ozone over the western North Atlantic, *Nature*, **358**, 392-394, 1992.
- Oltmans, S. J., and H. Levy, II, Surface ozone measurements from a global network, *Atmos. Environ.*, **28**, 9-24, 1994.
- Pickering, K. E., A. M. Thompson, J. R. Scala, W.-K. Tao, R. R. Dickerson, and J. Simpson, Free tropospheric ozone production following entrainment of urban plumes into deep convection, *J. Geophys. Res.*, **97**, 17,985-18,000, 1992.
- Prospero, J. M., and R. T. Nees, Impact of the North African drought and El Niño on mineral dust in the Barbados trade wind, *Nature*, **320**, 735-738, 1986.
- Sancho, P., J. de la Cruz, A. Díaz, F. Martín, E. Hernández, F. Valero, and B. Albarrán, A five-year climatology of back-trajectories from the Izaña baseline station, Tenerife, Canary Islands, *Atmos. Environ.*, **26A**, 1081-1096, 1992.
- Savoie, D. L., J. M. Prospero, J. T. Merrill, and M. Uematsu, Nitrate in the atmospheric boundary layer of the tropical South Pacific: Implications regarding sources and transport, *J. Atmos. Chem.*, **8**, 391-415, 1989.
- Savoie, D. L., J. M. Prospero, S. J. Oltmans, W. C. Graustein, K. K. Turekian, J. T. Merrill, and H. Levy II, Sources of nitrate and ozone in the marine boundary layer of the tropical North Atlantic, *J. Geophys. Res.*, **97**, 11,575-11,589, 1992.
- Schmitt, R., B. Schreiber, and I. Levin, Effects of long-range transport on atmospheric trace constituents at the Baseline Station Tenerife (Canary Islands), *J. Atmos. Chem.*, **7**, 335-351, 1988.
- Schmitt, R., and M. Carretero, Ozone in the free troposphere over the North Atlantic: Production and long range transport, in *Proceedings of EUROTRAC Symposium, 1995* (in press).
- Thompson, A. M., The oxidizing capacity of the earth's atmosphere: probable past and future changes, *Science*, **256**, 1157-1165, 1992.
- Turekian, K. K., W. C. Graustein, and J. K. Cochran, Lead-210 in the SEAREX program: An aerosol tracer across the Pacific, in *Chemical Oceanography*, vol. 10, edited by J. P. Riley, R. Chester, and R. A. Duce, pp. 51-81, Academic, San Diego, Calif., 1989.
- Volz, A., Trends in photo-oxidant concentrations, in *Proceedings of EUROTRAC Symposium '92*, edited by P.M. Burrell *et al.*, pp. 59-64, Academic Pub., The Hague, 1993.
- E. Cuevas, Instituto Nacional de Meteorología, Tenerife, Islas Canarias, España (e-mail: ecuevas@ull.es)
- A. Díaz, Universidad de La Laguna, Tenerife, Islas Canarias, España (e-mail: adiaz@ull.es)
- W. C. Graustein and K. K. Turekian, Department of Geology and Geophysics, Yale University, New Haven, CT (e-mail: billg@lamb.geology.yale.edu; kkt@milne.geology.yale.edu)
- H. Levy II, Geophysical Fluid Dynamics Laboratory, National Oceanic and Atmospheric Administration, Princeton, NJ (e-mail: hl@gfdl.gov)
- S. J. Oltmans, Climate Monitoring and Diagnostics Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO (e-mail: soltmans@cmdl.noaa.gov)
- J. M. Prospero and D. L. Savoie, Division of Marine and Atmospheric Chemistry, University of Miami, Miami, FL 33149 (e-mail: jprospero@rsmas.miami.edu; dsavoie@rsmas.miami.edu)
- R. Schmitt, Meteorologie Consult GmbH, Königstein, Deutschland (e-mail: 100060.2416@CompuServe.com)
- A. Volz-Thomas, Inst. Chemie Dynamik der Geosphäre, Jülich, Deutschland (e-mail: a.volz-thomas@kfa-juelich.de)

(Received March 28, 1995; revised July 18, 1995; accepted September 5, 1995)